University of Groningen

Biofilm formation and dispersal in Gram-positive bacteria

Abee, Tjakko; Kovacs, Akos T.; Kuipers, Oscar P.; van der Veen, Stijn

Published in:
Current Opinion in Biotechnology

DOI:
[10.1016/j.copbio.2010.10.016](https://doi.org/10.1016/j.copbio.2010.10.016)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Abee, T., Kovacs, A. T., Kuipers, O. P., & van der Veen, S. (2011). Biofilm formation and dispersal in Gram-positive bacteria. *Current Opinion in Biotechnology*, 22(2), 172-179.
<https://doi.org/10.1016/j.copbio.2010.10.016>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Biofilm formation and dispersal in Gram-positive bacteria

Tjakko Abee^{1,2}, Ákos T Kovács³, Oscar P Kuipers^{1,3,4} and Stijn van der Veen^{1,2}

Biofilms are structured communities of bacteria, which are adhered to a surface and embedded in a self-produced matrix of extracellular polymeric substances. Since biofilms are very resistant to antimicrobial agents, they are at the basis of a range of problems, including quality and safety issues in food industry. Recently, major advances have been made in elucidating the different structural components of the biofilm matrix, the regulatory pathways involved in biofilm formation, and signaling molecules involved in biofilm formation and dispersal, which provide opportunities for prevention and control of these biofilms in the food industry.

Addresses

¹ Top Institute Food and Nutrition (TIFN), Nieuwe Kanaal 9A, 6709 PA Wageningen, The Netherlands

² Laboratory of Food Microbiology, Wageningen University and Research Centre, Bomenweg 2, 6703 HD Wageningen, The Netherlands

³ Molecular Genetics Group, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Kerklaan 30, 9751NN Haren, The Netherlands

⁴ Kluysver Centre for Genomics of Industrial Fermentation, Kerklaan 30, 9751NN Haren, The Netherlands

Corresponding author: Abee, Tjakko (Tjakko.Abee@wur.nl)

Current Opinion in Biotechnology 2011, **22**:172–179

This review comes from a themed issue on
 Food biotechnology
 Edited by Oscar Kuipers and Tjakko Abee

Available online 23rd November 2010

0958-1669/\$ – see front matter

© 2010 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.copbio.2010.10.016](https://doi.org/10.1016/j.copbio.2010.10.016)

Introduction

Biofilms are complex microbial communities established on a wide range of surfaces that are generally encapsulated by an extracellular protective matrix composed of various types of biopolymers. Since biofilms are very difficult to eradicate, the ability of bacteria to form biofilms poses a major problem in various industrial and medical settings, being a persistent source of (re)contamination and/or infection, respectively. Mechanisms that have been proposed to explain the observed increased resistance of biofilms to antimicrobial agents are the impenetrable character of the biofilm, the slow growth rate of organisms, and the induction of resistance mechanisms [1]. The molecular mechanisms and factors involved in biofilm formation and subsequent dispersal of bacteria from the biofilm are starting to be unraveled

(Figure 1). This review provides an overview of the recent insights in the factors and mechanisms involved in biofilm formation and dispersal focusing on a selection of Gram-positive bacteria including *Bacillus* spp, *Listeria monocytogenes*, *Staphylococcus* spp, and lactic acid bacteria (LAB).

Diversity in biofilms

Depending on the specific systems investigated and the nature of the microorganism, biofilms can display a wide range of phenotypes. Biofilms of *L. monocytogenes*, grown under static conditions, generally consist of a homogeneous layer of cells and/or microcolonies, with the biofilm cells displaying a morphology similar to that of planktonic cells. By contrast, *L. monocytogenes* biofilms grown under continuous flow conditions consist of spherically shaped microcolonies that are surrounded by a network of knitted chains composed of elongated cells [2]. Interestingly, the formation of this type of *L. monocytogenes* biofilms is dependent on the activation of the SOS response factor YneA, which is responsible for cell elongation [3]. *Staphylococcus aureus* biofilms grown under both static and continuous flow conditions consist of a dense layer of cells with an elaborate matrix harboring various types of polymers. Biofilm formation and maturation in *S. aureus* is dependent on the interplay between various regulators including those encoded by *sarA*, *agr*, *ica*, and *sigB* [4–6]. Also, static and continuous flow biofilms of LAB such as *Lactobacillus plantarum* and *Lactococcus lactis* consist of a dense layer of cells that are frequently embedded in polymeric substances [7,8]. Biofilms of *Bacilli* are generally studied at the air–liquid interface where they form structured pellicles floating on culture media or as architecturally complex colonies on solid agar media [9–11]. The latter type of biofilms develops aerial structures called ‘fruiting bodies’ that serve as sites for sporulation [9]. The formation of these fruiting bodies is dependent on the temporal and spatial regulation of several distinctive pathways. In addition, *Bacilli* can form submersed biofilms both under static and continuous flow conditions [10,11].

In most conditions bacteria will generally grow on surfaces in competition with other microorganisms in a mixed species biofilm. Mixed species biofilm formation of several *S. aureus* strains with *L. monocytogenes* has been demonstrated and generally the number of cells in the mixed species biofilm is similar to the number of cells in single species biofilms [12]. A study on mixed biofilm formation of *L. monocytogenes* in combination with various secondary species showed that mixed species biofilms were established and depending on the specific combination, they

Figure 1

Schematic presentation showing mechanisms and components involved in biofilm formation and dispersal. Biofilms can contain various extracellular biopolymers like extracellular DNA (eDNA), extracellular polysaccharides, amyloid fibers, and biofilm-associated proteins (BAP). These matrix components might be good targets for (combinations of) putative enzymes such as DNases, proteases, and extracellular polysaccharide degrading enzymes to prevent formation of biofilms or to stimulate dispersal of already formed biofilms. Communication between cells during biofilm formation and dispersal of biofilms is dependent on quorum sensing systems and molecules like autoinducer 2 (AI-2), D-amino acids, and *cis*-2-decenoic acid. Furthermore, motility is an important factor in the establishment of new biofilms and the dispersal of cells from mature biofilms. Also, aerial structures of the biofilm serve as specific sites for the generation of spores (see text for details and corresponding references).

showed increased, reduced, or no effect on the number of *L. monocytogenes* cells in the biofilm [13]. Furthermore, an elaborate study on the formation of mixed species biofilms of *L. monocytogenes* and *L. plantarum* showed that mixed species biofilms have the capacity to be more resistant against disinfectant treatments than single species biofilm or planktonic cells [14]. Another interesting study on the adherence of *L. monocytogenes* to preformed *L. lactis* biofilms with different architectures, matrices, and cell surface properties showed that *L. monocytogenes* biofilm formation can be influenced by resident biofilms [7[•]]. The impact of secondary species on *L. monocytogenes* settlement, biofilm formation and persistence in food processing environments remains to be characterized.

The biofilm matrix

When forming a biofilm, bacteria produce various biopolymers mediating cell-to-cell and cell-to-surface attachments. However, diverse polymers are used by different species or strains of the same species. These extracellular polymeric substances (EPS) are mainly polysaccharides, proteins, nucleic acids, and lipids, that provide on the one

hand mechanical stability of biofilms and adhesion to surfaces, and on the other hand form a scaffold for the three-dimensional architecture that interconnects and immobilizes biofilm cells [15]. For example, *Bacillus subtilis* NCIB3610 produces an exopolysaccharide [16], while *B. subtilis* RO-FF-1 synthesizes poly- γ -D,L-glutamate [17] as matrix component. *S. aureus* biofilms contain the polymer poly-N-acetyl glucosamine (PNAG), which is also referred to as polysaccharide intercellular adhesion (PIA), and is synthesized by enzymes encoded on the *ica*-operon [18]. A functional biofilm matrix often also requires single protein or several protein components. The *B. subtilis* matrix contains amyloid fibers of the protein TasA [19[•]], while for *L. monocytogenes* the presence of a biofilm-associated protein, BapL, in the matrix has been described [20]. BAP proteins are also involved in biofilm formation in various *S. aureus* strains [21–23], but in this species several other extracellular biofilm promoting proteins have also been described [24,25]. Recently, an extracellular protein, MabA, was described for *Lactobacillus rhamnosus*, which appeared to be very important for biofilm formation

[26]. Furthermore, a function for extracellular DNA (eDNA) in cell adhesion and biofilm formation was recently shown for *Bacillus cereus* [27], *S. aureus* [28,29^{*}], and *L. monocytogenes* [30]. The release of genomic DNA from a subpopulation of cells of *S. aureus* during biofilm development is functionally analogous to the role of apoptosis in eukaryotic development [31]. Although the Cid/Lrg system was revealed to be involved in the activation of programmed cell death (PCD) of *S. aureus*, and although it is not known how PCD is induced only in a subpopulation of the cells, it provides an intriguing example of differentiation in bacterial biofilms. The identification of these novel matrix components may provide clues to the identification and application of matrix-degrading enzymes that prevent formation and/or activate dispersal of biofilms [32,33] (Figure 1).

Involvement of flagella and motility in biofilm formation

Biofilms generally consist of cells that are not motile. However, for several bacteria motility has been indicated to be an important factor during biofilm formation or attachment of cells to a surface (Figure 1). For *B. subtilis*, flagella and swarming motility was important, but not essential for normal progression of pellicle type biofilm formation [34]. For *L. monocytogenes* static biofilm formation, flagellar based motility appeared to be essential to propel cells towards the surface before attachment [35], while under continuous flowing conditions loss of flagellar motility resulted in lower initial cell attachment but increased biofilm formation [36]. Similar observations were made for the role of flagella-based motility in *B. cereus* biofilm formation [10]. Motility appeared to be important for static biofilm formation in this organism but not for biofilm formation under continuous flow conditions. These results show that it is important for bacteria to control the transition between the motile planktonic mode of growth and the biofilm mode of growth. For *L. monocytogenes*, it has been shown that this transition was under control of the DegU response regulator, that appeared not to be associated with a cognate histidine protein kinase, and its phosphorylation status seems to be directly controlled by Acetyl-phosphate levels. This may connect biofilm formation to the metabolic state of cells, which may be highly relevant for biofilm formation control [37–39]. DegU also plays an important role in biofilm formation in *B. subtilis*. Gradual phosphorylation of DegU determines whether cells activate swarming (low DegU~P level), biofilm development (medium DegU~P level), or exoprotease production (high DegU~P level) [40,41]. Furthermore, a very sophisticated mechanism to control motility of *B. subtilis* cells in biofilms was recently identified. EpsE, which is encoded within an operon required for biofilm matrix synthesis, is capable of arresting flagellar rotation in biofilm cells by disengaging the motor force generating elements [42].

Spore formation in biofilms: intertwinement of regulatory pathways

Control of sporeformers and resultant spores is of eminent importance for food quality and safety, owing to the resistance of spores against processing regimes such as heating. Notably, biofilms provide an optimal environment for sporulation of *Bacilli*. Developmental processes, like motility, biofilm formation, and sporulation are strictly connected and share global regulators. Furthermore, these processes are spatio-temporally coordinated, while genes facilitating motility are first expressed, matrix production is activated thereafter, and finally spores appear on the upper aerial structures of the biofilm (Figure 1) [43^{*}]. This complex coordination of developmental processes is achieved by an interplay of several regulators. Also the activity states of various regulators are important for proper modulation of phenotypic traits. The coordination of the formation of different cell types is achieved through cell–cell communication. In case of high cell density, quorum sensing molecules accumulate and induce the production of surfactin in a subpopulation of cells [44]. The produced surfactin is in turn sensed by distinct cells, but not by the surfactin producing cells themselves, and triggers potassium leakage and subsequent activation of the KinC sensor kinase localized in membrane microdomains analogous to lipid rafts in eukaryotic cells [45^{**},46]. In spite of differences in the regulatory networks that control biofilm formation in *B. subtilis* and *S. aureus*, compounds that inhibit the formation of lipid rafts such as zaragozic acid, also inhibited the development of *S. aureus* biofilm [46]. The generation of a signal by some cells to which only certain target cells respond presents a novel paracrine signaling pathway that requires a proper distribution of signal producing and sensing cell types. Activated KinC modulates activity of the master regulator Spo0A through a phosphorelay pathway. Intermediate levels of Spo0A~P activate the production of SinI, a small protein that antagonizes SinR, the repressor of several biofilm operons [47]. Upon alleviating the inhibitory effect of SinR, the operons involved in matrix production are expressed. As Spo0A phosphorylation and expression of matrix operons are activated only in a subpopulation of cells, biofilm components are produced by part of the population as a division of labor [48]. Recently, the new regulator SlrR was identified that plays a role in biofilm formation by repressing *sinR* and by forming a SinR–SlrR complex. It titrates SinR and prevents it from repressing *slrR* itself [49]. Furthermore, this complex represses autolysin and motility genes. Thus, this epigenetic switch controls cell separation and helps the formation of long chains of cells that is a prerequisite for biofilm development. Further increase of the Spo0A~P level drives a subpopulation of cells in the maturing biofilm to the formation of spores. However, in matrix-deficient mutants the KinD membrane sensor protein acts rather as a phosphatase than a kinase and inhibits the activity of Spo0A that results in a delay of

sporulation [50]. Notably, most of these regulators that assist in *B. subtilis* biofilm formation are conserved in *Bacilli*, but their role in biofilm formation and regulatory cascades involved, remain to be elucidated.

In *B. cereus* the production of virulence factors, including enterotoxins, is regulated by PlcR, a pleiotropic transcriptional regulator that downregulates biofilm formation presumably via altering production of biosurfactants [51]. Interestingly, master regulator Spo0A was shown to downregulate *plcR* expression [52], suggesting that enterotoxin production is hampered in biofilms. By contrast, the putative cell wall peptidase, CwpFM, was shown to contribute to virulence and biofilm formation [53]. The cells and spores accumulated in biofilms may disperse and recontaminate foods, where they are delivered to the gastrointestinal (GI) tract after consumption, followed by germination, in case of spores, growth and production of enterotoxins [54].

Communication to build and break biofilms

Single and mixed species biofilm development, maturation, survival of different physicochemical conditions encountered in the range of micro niches in biofilms, and the release of cells from mature biofilms, require complex sequential mechanisms for which cross-species cell to cell communication and/or interspecies quorum sensing (QS) might be important factors (Figure 1). Although the specific QS molecules fill an important role in affecting the developmental regulatory networks like sporulation and competence, there are examples of broadly conserved QS systems that control biofilm formation. One of these systems involves the LuxS enzyme that synthesizes autoinducer-2 (AI-2). For several Gram-positive bacteria it has recently been shown that a functional *luxS* gene and AI-2 signal were present and that this system is involved in biofilm formation [55]. For *L. monocytogenes*, it was shown that inactivation of *luxS* resulted in increased biofilm formation [56,57]. However, addition of *in vitro* synthesized AI-2 did not restore wild-type biofilm levels. Instead, biofilm formation appeared to be increased after addition of S-ribosyl homocysteine (SRH), which is one of the precursors in the AI-2 pathway [56]. In *B. cereus*, the addition of *in vitro* synthesized AI-2 inhibited biofilm formation and stimulated the release of cells from a mature biofilm [58]. Increased biofilm formation for a *luxS* mutant was also shown in *Lactobacillus reuteri* [59] and *Staphylococcus epidermidis* [60]. However, in *S. epidermidis* the *luxS* mutant showed increased expression of the *ica*-operon and increased production of biofilm polysaccharide PIA, which could explain the observed biofilm phenotype. By contrast, biofilm formation of the *luxS* mutant was reduced in *L. rhamnosus* [61] and *B. subtilis* [62]. In *B. subtilis*, AI-2 production was furthermore regulated by the sporulation and biofilm regulators Spo0A and SinR and spatial activity of LuxS appeared to be important for the development of specific biofilm

structures [62]. The other major quorum sensing system is the peptide-mediated accessory gene regulator (Agr) system, which consists of an operon containing four genes (*agrBDCA*). In *L. monocytogenes* a specific role for the *agr*-system in the positive regulation of adherence and biofilm formation was identified by mutant analyses and expression studies [63,64]. Similar observations were made for *L. plantarum*, although interestingly, this organism contains two homologous Agr systems. However, only the regulatory two-component part of the four-gene module is duplicated, while only a single copy of the peptide generating gene-set is present in this species. The corresponding peptide is proposed to communicate via both two-component regulatory Agr-like systems mediating control of cell adherence and biofilm formation [65]. By contrast, *S. aureus* biofilm formation was shown to be negatively regulated by the Agr-system, since its activation stimulated biofilm dispersal [66]. Other communication molecules that are involved in dispersal of biofilm cells have also been identified (Figure 1). The fatty acid *cis*-2-decenoic acid, which is produced by *Pseudomonas aeruginosa*, induced dispersal in various bacteria, including *B. subtilis* and *S. aureus* [67]. Another communication or dispersal molecule was identified in *B. subtilis* biofilms. As *B. subtilis* biofilms age, D-amino acids are produced just before its disassembly. Addition of individual or combinations of D-amino acids (e.g. D-leucine, D-methionine, D-tyrosine, and D-tryptophan) before biofilm formation inhibits the functional assembly of the matrix [68**]. The D-amino acids modify the connection between the TasA protein component and the cell wall, acting through the YqxM protein. Particular strains harboring mutations in the protein sequence of YqxM that is required for the association of TasA are resistant to the presence of D-amino acids and delayed in disassembly. It is hypothesized that a domain near the C terminus of YqxM could trigger the release of TasA in response to the presence of D-amino acids in the cell wall. Interestingly, D-amino acids are effective against biofilms of *S. aureus* and *P. aeruginosa*, in spite of the lack of TasA and YqxM homologs in these species, suggesting a general strategy for biofilm disassembly. The production of D-amino acids by bacteria can be a common signal for biofilm disassembly and may provide clues for the search and development of effective inhibitors of biofilm formation in food processing environments.

Tools in biofilm analysis

Advancements in imaging techniques for studying the formation of single and mixed species biofilms have contributed significantly to the current status of the biofilm research community. These new techniques, which include laser scanning microscopy, scanning transmission X-ray microscopy, and magnetic resonance imaging, made it possible to study the formation and dynamics of biofilms *in situ* (reviewed in [69]). In particular, imaging techniques that use fluorescent reporters have proven to be an excellent tool to study activation of specific genes in biofilms, to

Figure 2

Application of fluorescent reporters to study the expression of specific genes in biofilms (a,b) and to investigate the formation of single and mixed species biofilms (c,d). Phase contrast (a) and fluorescent microscopy images (b) of a *L. monocytogenes* continuous flow biofilm expressing EGFP from the *yneA* promoter. Fluorescent microscopy image of *L. monocytogenes* constitutively expressing EGFP in a static biofilm on stainless steel (c), and confocal microscopy image of *L. monocytogenes* constitutively expressing DsRed and *L. plantarum* constitutively expressing EGFP in a mixed dual species biofilm (d).

investigate biofilm formation on surfaces that are not penetrable by light, or to study the distribution of organisms in mixed species biofilms (Figure 2). Fluorescent imaging furthermore proved very useful for investigating the dynamics in biofilm matrix structures. This is highlighted with the example of programmed cell death in a subpopulation of the *S. aureus* biofilm cells, which resulted in the release of genomic DNA that subsequently becomes an important component for biofilm maturation [28,29]. Future tool developments in biofilm research will most probably focus on capturing the role and fate of single cells in biofilm formation and maturation.

Concluding remarks

The majority of bacteria are able to form biofilms displaying a large variety in architecture, phenotypes, and matrix components. Novel insights include factors contributing to phenotypic heterogeneity within biofilms, the identification and characterization of a range of matrix building

blocks such as extracellular polysaccharides, eDNA, and amyloid fibers, and the identification of components that activate dispersal such as D-amino acids. Notably, most information has been obtained from studies with single species biofilms, and there is an urgent need to extend our knowledge on mixed species biofilms, since these presumably also develop in food processing environments displaying features different from that of the respective single species biofilms. The advances in our understanding of the different factors and mechanisms involved in biofilm formation and dispersal of various Gram-positive bacteria might provide clues and stimulate developments in the search for (natural) compounds and combinations thereof, for prevention and control of spoilage and pathogenic bacteria in industrial settings.

Acknowledgements

A.T.K. was financially supported by grant 818.02.004 from ALW-NWO Open programma. O.P.K. is supported by the research program of the

Kluyver Centre for Genomics of Industrial Fermentation that is part of the Netherlands Genomics Initiative/Netherlands Organization for Scientific Research.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Donlan RM, Costerton JW: **Biofilms: survival mechanisms of clinically relevant microorganisms.** *Clin Microbiol Rev* 2002, **15**:167-193.
 2. Rieu A, Briandet R, Habimana O, Garmyn D, Guzzo J, Piveteau P: ***Listeria monocytogenes* EGD-e biofilms: no mushrooms but a network of knitted chains.** *Appl Environ Microbiol* 2008, **74**:4491-4497.
 3. van der Veen S, Abee T: **Dependence of continuous-flow biofilm formation by *Listeria monocytogenes* EGD-e on SOS response factor YneA.** *Appl Environ Microbiol* 2010, **76**:1992-1995.
- Using fluorescent reporters and targeted *L. monocytogenes* mutants, the authors show that the formation of knitted-chain biofilms in a continuous flow system is dependent on cell elongation mediated by the SOS regulon member YneA, providing new insights in the role of the SOS response in pathogen biofilm formation.
4. Cue D, Lei MG, Luong TT, Kuechenmeister L, Dunman PM, O'Donnell S, Rowe S, O'Gara JP, Lee CY: **Rbf promotes biofilm formation by *Staphylococcus aureus* via repression of icaR, a negative regulator of icaADBC.** *J Bacteriol* 2009, **191**:6363-6373.
 5. Lauderdale KJ, Boles BR, Cheung AL, Horswill AR: **Interconnections between Sigma B, agr, and proteolytic activity in *Staphylococcus aureus* biofilm maturation.** *Infect Immun* 2009, **77**:1623-1635.
 6. Tamber S, Cheung AL: **SarZ promotes the expression of virulence factors and represses biofilm formation by modulating SarA and agr in *Staphylococcus aureus*.** *Infect Immun* 2009, **77**:419-428.
 7. Habimana O, Meyrand M, Meylheuc T, Kulakauskas S, Briandet R: **Genetic features of resident biofilms determine attachment of *Listeria monocytogenes*.** *Appl Environ Microbiol* 2009, **75**:7814-7821.
- Using *L. lactis* cell wall mutants that affect bacterial chain formation, exopolysaccharide (EPS) synthesis, and surface hydrophobicity, significant differences were seen in *L. monocytogenes* settlement as a function of the genetic background of resident lactococcal biofilm cells. In particular, biofilms of the *L. lactis* chain-forming mutant resulted in a marked increase in *L. monocytogenes* settlement, while biofilms of the EPS-secreting mutant efficiently prevented pathogen fixation. These results offer new insights into the role of resident biofilms in governing the settlement of pathogens on surfaces.
8. Sturme MH, Nakayama J, Molenaar D, Murakami Y, Kunugi R, Fujii T, Vaughan EE, Kleerebezem M, de Vos WM: **An agr-like two-component regulatory system in *Lactobacillus plantarum* is involved in production of a novel cyclic peptide and regulation of adherence.** *J Bacteriol* 2005, **187**:5224-5235.
 9. Branda SS, Gonzalez-Pastor JE, Ben-Yehuda S, Losick R, Kolter R: **Fruiting body formation by *Bacillus subtilis*.** *Proc Natl Acad Sci USA* 2001, **98**:11621-11626.
 10. Houry A, Briandet R, Aymerich S, Gohar M: **Involvement of motility and flagella in *Bacillus cereus* biofilm formation.** *Microbiology* 2010, **156**:1009-1018.
 11. Wijman JG, de Leeuw PP, Moezelaar R, Zwietering MH, Abee T: **Air-liquid interface biofilms of *Bacillus cereus*: formation, sporulation, and dispersal.** *Appl Environ Microbiol* 2007, **73**:1481-1488.
 12. Rieu A, Lemaître JP, Guzzo J, Piveteau P: **Interactions in dual species biofilms between *Listeria monocytogenes* EGD-e and several strains of *Staphylococcus aureus*.** *Int J Food Microbiol* 2008, **126**:76-82.
 13. Carpentier B, Chassaing D: **Interactions in biofilms between *Listeria monocytogenes* and resident microorganisms from food industry premises.** *Int J Food Microbiol* 2004, **97**:111-122.
 14. van der Veen S, Abee T: **Mixed species biofilms of *Listeria monocytogenes* and *Lactobacillus plantarum* show enhanced resistance to benzalkonium chloride and peracetic acid.** *Int J Food Microbiol* 2010, doi:10.1016/j.ijfoodmicro.2010.10.029.
 15. Flemming HC, Wingender J: **The biofilm matrix.** *Nat Rev Microbiol* 2010, **8**:623-633.
 16. Branda SS, Chu F, Kearns DB, Losick R, Kolter R: **A major protein component of the *Bacillus subtilis* biofilm matrix.** *Mol Microbiol* 2006, **59**:1229-1238.
 17. Stanley NR, Lazazzera BA: **Defining the genetic differences between wild and domestic strains of *Bacillus subtilis* that affect poly-gamma-dl-glutamic acid production and biofilm formation.** *Mol Microbiol* 2005, **57**:1143-1158.
 18. O'Gara JP: **ica and beyond: biofilm mechanisms and regulation in *Staphylococcus epidermidis* and *Staphylococcus aureus*.** *FEMS Microbiol Lett* 2007, **270**:179-188.
 19. Romero D, Aguilar C, Losick R, Kolter R: **Amyloid fibers provide structural integrity to *Bacillus subtilis* biofilms.** *Proc Natl Acad Sci USA* 2010, **107**:2230-2234.
- The authors describe the physicochemical properties of purified TasA and show that it forms fibers of variable length and 10-15 nm in width. Biochemical analyses, in combination with the use of specific dyes and microscopic analyses, indicate that TasA forms amyloid fibers that bind cells together in the biofilm.
20. Jordan SJ, Perni S, Glenn S, Fernandes I, Barbosa M, Sol M, Tenreiro RP, Chambel L, Barata B, Zilhao I et al.: ***Listeria monocytogenes* biofilm-associated protein (BapL) may contribute to surface attachment of *L. monocytogenes* but is absent from many field isolates.** *Appl Environ Microbiol* 2008, **74**:5451-5456.
 21. Lasa I, Penades JR: **Bap: a family of surface proteins involved in biofilm formation.** *Res Microbiol* 2006, **157**:99-107.
 22. Latasa C, Solano C, Penades JR, Lasa I: **Biofilm-associated proteins.** *C R Biol* 2006, **329**:849-857.
 23. Cucarella C, Solano C, Valle J, Amorena B, Lasa I, Penades JR: **Bap, a *Staphylococcus aureus* surface protein involved in biofilm formation.** *J Bacteriol* 2001, **183**:2888-2896.
 24. Corrigan RM, Rigby D, Handley P, Foster TJ: **The role of *Staphylococcus aureus* surface protein SasG in adherence and biofilm formation.** *Microbiology* 2007, **153**:2435-2446.
 25. Schroeder K, Jularic M, Horsburgh SM, Hirschhausen N, Neumann C, Bertling A, Schulte A, Foster S, Kehrel BE, Peters G et al.: **Molecular characterization of a novel *Staphylococcus aureus* surface protein (SasC) involved in cell aggregation and biofilm accumulation.** *PLoS ONE* 2009, **4**:e7567.
 26. Velez MP, Petrova MI, Lebeer S, Verhoeven TL, Claes I, Lambrechts I, Tynkynen S, Vanderleyden J, De Keersmaecker SC: **Characterization of MAbA, a modulator of *Lactobacillus rhamnosus* GG adhesion and biofilm formation.** *FEMS Immunol Med Microbiol* 2010, **59**:386-398.
 27. Vilain S, Pretorius JM, Theron J, Brozel VS: **DNA as an adhesin: *Bacillus cereus* requires extracellular DNA to form biofilms.** *Appl Environ Microbiol* 2009, **75**:2861-2868.
 28. Mann EE, Rice KC, Boles BR, Endres JL, Ranjit D, Chandramohan L, Tsang LH, Smeltzer MS, Horswill AR, Bayles KW: **Modulation of eDNA release and degradation affects *Staphylococcus aureus* biofilm maturation.** *PLoS ONE* 2009, **4**:e5822.
 29. Rice KC, Mann EE, Endres JL, Weiss EC, Cassat JE, Smeltzer MS, Bayles KW: **The cidA murein hydrolase regulator contributes to DNA release and biofilm development in *Staphylococcus aureus*.** *Proc Natl Acad Sci USA* 2007, **104**:8113-8118.
- To examine the biological role of cell death and lysis, the authors studied the impact of the cidA mutation on *S. aureus* biofilm development. It was demonstrated that cidA-controlled cell lysis plays a significant role during biofilm development and that released genomic DNA, that is, extracellular DNA, is an important structural component of *S. aureus* biofilm.

30. Harmsen M, Lappann M, Knochel S, Molin S: **Role of extracellular DNA during biofilm formation by *Listeria monocytogenes***. *Appl Environ Microbiol* 2010, **76**:2271-2279.
31. Bayles KW: **The biological role of death and lysis in biofilm development**. *Nat Rev Microbiol* 2007, **5**:721-726.
32. Karatan E, Watnick P: **Signals, regulatory networks, and materials that build and break bacterial biofilms**. *Microbiol Mol Biol Rev* 2009, **73**:310-347.
33. Landini P, Antoniani D, Burgess JG, Nijland R: **Molecular mechanisms of compounds affecting bacterial biofilm formation and dispersal**. *Appl Microbiol Biotechnol* 2010, **86**:813-823.
34. Kobayashi K: ***Bacillus subtilis* pellicle formation proceeds through genetically defined morphological changes**. *J Bacteriol* 2007, **189**:4920-4931.
35. Lemon KP, Higgins DE, Kolter R: **Flagellar motility is critical for *Listeria monocytogenes* biofilm formation**. *J Bacteriol* 2007, **189**:4418-4424.
36. Todhanakasem T, Young GM: **Loss of flagellum-based motility by *Listeria monocytogenes* results in formation of hyperbiofilms**. *J Bacteriol* 2008, **190**:6030-6034.
37. Gueriri I, Bay S, Dubrac S, Cyncynatus C, Msadek T: **The Pta-AckA pathway controlling acetyl phosphate levels and the phosphorylation state of the DegU orphan response regulator both play a role in regulating *Listeria monocytogenes* motility and chemotaxis**. *Mol Microbiol* 2008, **70**:1342-1357.
38. Gueriri I, Cyncynatus C, Dubrac S, Arana AT, Dussurget O, Msadek T: **The DegU orphan response regulator of *Listeria monocytogenes* autorepresses its own synthesis and is required for bacterial motility, virulence and biofilm formation**. *Microbiology* 2008, **154**:2251-2264.
39. Murray EJ, Kiley TB, Stanley-Wall NR: **A pivotal role for the response regulator DegU in controlling multicellular behaviour**. *Microbiology* 2009, **155**:1-8.
40. Kobayashi K: **Gradual activation of the response regulator DegU controls serial expression of genes for flagellum formation and biofilm formation in *Bacillus subtilis***. *Mol Microbiol* 2007, **66**:395-409.
41. Verhamme DT, Kiley TB, Stanley-Wall NR: **DegU co-ordinates multicellular behaviour exhibited by *Bacillus subtilis***. *Mol Microbiol* 2007, **65**:554-568.
42. Blair KM, Turner L, Winkelman JT, Berg HC, Kearns DB: **A molecular clutch disables flagella in the *Bacillus subtilis* biofilm**. *Science* 2008, **320**:1636-1638.
43. Vlamakis H, Aguilar C, Losick R, Kolter R: **Control of cell fate by the formation of an architecturally complex bacterial community**. *Genes Dev* 2008, **22**:945-953.
This paper shows that motile, matrix-producing, and sporulating *B. subtilis* cells localize to distinct regions within the biofilm, and that the localization and percentage of each cell type is dynamic throughout development of the community. Importantly, mutants that do not produce extracellular matrix form unstructured biofilms that are deficient in sporulation. It is proposed that sporulation is a culminating feature of biofilm formation, and that spore formation is coupled to the formation of an architecturally complex community of cells.
44. Lopez D, Vlamakis H, Losick R, Kolter R: **Paracrine signaling in a bacterium**. *Genes Dev* 2009, **23**:1631-1638.
45. Lopez D, Fischbach MA, Chu F, Losick R, Kolter R: **Structurally diverse natural products that cause potassium leakage trigger multicellularity in *Bacillus subtilis***. *Proc Natl Acad Sci USA* 2009, **106**:280-285.
This is a highly interesting paper that describes a new quorum-sensing mechanism for triggering multicellularity in *B. subtilis*. Evidence is provided that biofilm formation is stimulated by a variety of small molecules, including *B. subtilis* surfactin, that share the ability to induce potassium leakage subsequently activating KinC that governs expression of genes involved in biofilm formation. This phenomenon may also bear high impact on the development of mixed species biofilms, a feature that remains to be investigated.
46. Lopez D, Kolter R: **Functional microdomains in bacterial membranes**. *Genes Dev* 2010.
47. Chai Y, Chu F, Kolter R, Losick R: **Bistability and biofilm formation in *Bacillus subtilis***. *Mol Microbiol* 2008, **67**:254-263.
48. Kearns DB: **Division of labour during *Bacillus subtilis* biofilm formation**. *Mol Microbiol* 2008, **67**:229-231.
49. Chai Y, Norman T, Kolter R, Losick R: **An epigenetic switch governing daughter cell separation in *Bacillus subtilis***. *Genes Dev* 2010, **24**:754-765.
50. Aguilar C, Vlamakis H, Guzman A, Losick R, Kolter R: **KinD is a checkpoint protein linking spore formation to extracellular-matrix production in *Bacillus subtilis* biofilms**. *mBio* 2010, **1**: e00035-00010.
51. Hsueh YH, Somers EB, Lereclus D, Wong AC: **Biofilm formation by *Bacillus cereus* is influenced by PlcR, a pleiotropic regulator**. *Appl Environ Microbiol* 2006, **72**:5089-5092.
52. Lereclus D, Agaisse H, Grandvalet C, Salamitou S, Goiminet M: **Regulation of toxin and virulence gene transcription in *Bacillus thuringiensis***. *Int J Med Microbiol* 2000, **290**:295-299.
53. Tran SL, Guillemet E, Gohar M, Lereclus D, Ramarao N: **CwpFM (EntFM) is a *Bacillus cereus* potential cell wall peptidase implicated in adhesion, biofilm formation, and virulence**. *J Bacteriol* 2010, **192**:2638-2642.
54. Stenfor Arnesen LP, Fagerlund A, Granum PE: **From soil to gut: *Bacillus cereus* and its food poisoning toxins**. *FEMS Microbiol Rev* 2008, **32**:579-606.
55. Hardie KR, Heurlier K: **Establishing bacterial communities by 'word of mouth': LuxS and autoinducer 2 in biofilm development**. *Nat Rev Microbiol* 2008, **6**:635-643.
56. Challan Belval S, Gal L, Margiewes S, Garmyn D, Piveteau P, Guzzo J: **Assessment of the roles of LuxS, S-ribosyl homocysteine, and autoinducer 2 in cell attachment during biofilm formation by *Listeria monocytogenes* EGD-e**. *Appl Environ Microbiol* 2006, **72**:2644-2650.
57. Sela S, Frank S, Belausov E, Pinto R: **A Mutation in the luxS gene influences *Listeria monocytogenes* biofilm formation**. *Appl Environ Microbiol* 2006, **72**:5653-5658.
58. Auger S, Krin E, Aymerich S, Gohar M: **Autoinducer 2 affects biofilm formation by *Bacillus cereus***. *Appl Environ Microbiol* 2006, **72**:937-941.
59. Tannock GW, Ghazally S, Walter J, Loach D, Brooks H, Cook G, Surette M, Simmers C, Bremer P, Dal Bello F et al.: **Ecological behavior of *Lactobacillus reuteri* 100-23 is affected by mutation of the luxS gene**. *Appl Environ Microbiol* 2005, **71**:8419-8425.
60. Xu L, Li H, Vuong C, Vadyvaloo V, Wang J, Yao Y, Otto M, Gao Q: **Role of the luxS quorum-sensing system in biofilm formation and virulence of *Staphylococcus epidermidis***. *Infect Immun* 2006, **74**:488-496.
61. Lebeer S, De Keersmaecker SC, Verhoeven TL, Fadda AA, Marchal K, Vanderleyden J: **Functional analysis of luxS in the probiotic strain *Lactobacillus rhamnosus* GG reveals a central metabolic role important for growth and biofilm formation**. *J Bacteriol* 2007, **189**:860-871.
62. Lombardia E, Rovetto AJ, Arabolaza AL, Grau RR: **A LuxS-dependent cell-to-cell language regulates social behavior and development in *Bacillus subtilis***. *J Bacteriol* 2006, **188**:4442-4452.
63. Riedel CU, Monk IR, Casey PG, Waidmann MS, Gahan CG, Hill C: **AgrD-dependent quorum sensing affects biofilm formation, invasion, virulence and global gene expression profiles in *Listeria monocytogenes***. *Mol Microbiol* 2009, **71**:1177-1189.
64. Rieu A, Weidmann S, Garmyn D, Piveteau P, Guzzo J: **Agr system of *Listeria monocytogenes* EGD-e: role in adherence and differential expression pattern**. *Appl Environ Microbiol* 2007, **73**:6125-6133.

65. Fujii T, Ingham C, Nakayama J, Beerthuyzen M, Kunuki R, Molenaar D, Sturme M, Vaughan E, Kleerebezem M, de Vos W: **Two homologous Agr-like quorum-sensing systems cooperatively control adherence, cell morphology, and cell viability properties in *Lactobacillus plantarum* WCFS1.** *J Bacteriol* 2008, **190**:7655-7665.
66. Boles BR, Horswill AR: **Agr-mediated dispersal of *Staphylococcus aureus* biofilms.** *PLoS Pathog* 2008, **4**:e1000052.
67. Davies DG, Marques CN: **A fatty acid messenger is responsible for inducing dispersion in microbial biofilms.** *J Bacteriol* 2009, **191**:1393-1403.
68. Kolodkin-Gal I, Romero D, Cao S, Clardy J, Kolter R, Losick R: **•• D-amino acids trigger biofilm disassembly.** *Science* 2010, **328**:627-629.
An important paper that shows that a mixture of D-amino acids produced by *B. subtilis* initiates disassembly of the biofilm at nanomolar concentrations. Analysis of mutants revealed alterations in the protein YqxM, required for the formation and anchoring of amyloid fibers to the cell wall. The authors suggest that D-amino acids may act as widespread signal for biofilm disassembly.
69. Neu TR, Manz B, Volke F, Dynes JJ, Hitchcock AP, Lawrence JR: **Advanced imaging techniques for assessment of structure, composition and function in biofilm systems.** *FEMS Microbiol Ecol* 2010, **72**:1-21.