University of Groningen

Infrared Emitting and Photoconducting Colloidal Silver Chalcogenide Nanocrystal Quantum Dots from a Silylamide-Promoted Synthesis

Yarema, Maksym; Pichler, Stefan; Sytnyk, Mykhailo; Seyrkammer, Robert; Lechner, Rainer T.; Fritz-Popovski, Gerhard; Jarzab, Dorota; Szendrei, Krisztina; Resel, Roland; Korovyanko, Oleksandra

Published in:
Acs Nano

DOI:
[10.1021/nn2001118](https://doi.org/10.1021/nn2001118)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Yarema, M., Pichler, S., Sytnyk, M., Seyrkammer, R., Lechner, R. T., Fritz-Popovski, G., Jarzab, D., Szendrei, K., Resel, R., Korovyanko, O., Loi, M. A., Paris, O., Hesser, G., Heiss, W., & Hesser, G. (2011). Infrared Emitting and Photoconducting Colloidal Silver Chalcogenide Nanocrystal Quantum Dots from a Silylamide-Promoted Synthesis. *Acs Nano*, 5(5), 3758-3765. <https://doi.org/10.1021/nn2001118>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Supporting Information for

***Infrared Emitting and Photoconducting Colloidal
Silver Chalcogenide Nanocrystal Quantum Dots
from a Silylamide Promoted Synthesis***

Maksym Yarema*, Stefan Pichler, Mykhailo Sytnyk,
Robert Seyrkammer, Rainer T. Lechner, Gerhard Fritz-Popovski,
Dorota Jarzab, Krisztina Szendrei, Roland Resel, Oleksandra
Korovyanko, Maria Antonietta Loi, Oskar Paris, Günter Hesser,
and Wolfgang Heiss*

Institute for Semiconductor and Solid State Physics, University Linz, Altenbergerstr. 69, Linz 4040 (Austria)

Department of Inorganic Chemistry, University Chernivtsi, L. Ukrainky Str. 25, Chernivtsi, 58012 (Ukraine)

Center of Surface and Nanoanalytics, University Linz, Altenbergerstr. 69, Linz, 4040 (Austria)

Institute of Physics, Montanuniversitaet Leoben, Franz Josef Str. 18, Leoben, 8700 (Austria)

Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, Groningen, 9747 AG (The Netherlands)

Institute of Solid State Physics, Graz University of Technology, Petersgasse 16, Graz, 8010 (Austria)

* Email: maksym.yarema@jku.at, wolfgang.heiss@jku.at

Figure S1. TEM images of Ag_2Se nanocrystals synthesized by a direct reaction between AgTFA and TOPSe in (A) and with addition of tin (II) bis[bis(trimethylsilyl)amide] (B).

Figure S2. Absorbance and PL spectrum for Ag_2Se nanocrystals, synthesized with addition of tin (II) bis[bis(trimethylsilyl)amide]

Figure S3. Absorbance (A), and EDX spectrum (B) of Ag nanocrystals. The vertical lines in B give the theoretical positions of the chalcogene lines which are clearly not present in our colloidal solution. Inset (A): photo of the Ag NC solution.

Figure S4. Absorption spectra of differently sized Ag_2Se nanocrystals. Inset: linear dependence of the absorbance peak position in respect to the Ag/Se molar ratio, evidencing a size control by the Ag/Se molar ratio.

Figure S5. PL spectra of samples with different growth times (synthesis at 70 °C), evidencing a size focusing with time.

Figure S6. PL spectra of Ag_2Se (black lines) and $\text{Ag}_2\text{Se}/\text{ZnSe}$ core/shell nanocrystals (gray lines), measured after synthesis (A) and after 6 months of storing (B).

Figure S7. Selected area electron diffraction (SAED) of Ag_2Se nanocrystals, suggesting the presence of the low-temperature α - Ag_2Se crystal structure (solid arcs), and an absence of high-temperature α - Ag_2Se (dashed arcs).^{1,2}

Figure S8. Theoretical atom arrangement in β -type (low temperature modification) Ag_2Se (as simulated for Figure 2C).

Figure S9. EDX spectra of (A) Ag₂Se nanocrystals, and (B) Ag₂Se/ZnSe core shell nanocrystals with quantifications of elemental contents.

Figure S10. Wide angle X-ray scattering (WAXS) spectrum of the 2.0 nm Ag₂Se nanocrystals compared to theoretical spectrum for bulk Ag₂Se (low-temperature phase).¹

Figure S11. SAXS spectra of the 2.0 nm Ag_2Se nanocrystals (black dots), and 2.2 nm $\text{Ag}_2\text{Se}/\text{ZnSe}$ core/shell nanocrystals (gray dots), together with fits (red lines).

Figure S12. Absorption spectrum of PCBM solution in chlorobenzene (A), and photoresponse of PCBM film deposited on interdigitate gold electrodes (B).

Figure S13. EDX spectra of (A) Ag_2S nanocrystals, and (B) Ag_2Te nanocrystals with quantifications of elemental contents (silicon was used as substrate).

Figure S14. TEM images of (A) Ag_2S nanocrystals, and (B) Ag_2Te nanocrystals synthesized with addition of lithium bis(trimethylsilyl)amide.

Figure S15. Size histograms for Ag₂S (A-B), and Ag₂Te nanocrystals (C) taken from TEM images, and fitted by Gauss distributions.

References:

- 1 Billetter, H.; Ruschewitz, U. Structural Phase Transitions in Ag₂Se (Naumannite). *Z. Anorg. Allg. Chem.* **2008**, 634, 241-246.
- 2 Oliveria, M.; McMullan, R. K.; Wuensch, B. J. Single crystal neutron diffraction analysis of the cation distribution in the high-temperature phases α -Cu_{2-x}S, α -Cu_{2-x}Se, and α -Ag₂Se. *Solid State Ionics* **1988**, 28-30, 1332-1337.