University of Groningen

Interactions between large herbivores and litter removal by termites across a rainfall gradient in a South African savanna

Buitenwerf, Robert; Stevens, Nicola; Gosling, Cleo M.; Anderson, T. Michael; Olf, Han

Published in:
Journal of Tropical Ecology

DOI:
[10.1017/S0266467411000125](https://doi.org/10.1017/S0266467411000125)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Buitenwerf, R., Stevens, N., Gosling, C. M., Anderson, T. M., & Olf, H. (2011). Interactions between large herbivores and litter removal by termites across a rainfall gradient in a South African savanna. *Journal of Tropical Ecology*, 27(4), 375-382. <https://doi.org/10.1017/S0266467411000125>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Interactions between large herbivores and litter removal by termites across a rainfall gradient in a South African savanna

Robert Buitenwerf^{*,1}, Nicola Stevens[†], Cleo M. Gosling^{*}, T. Michael Anderson[‡] and Han Olff^{*}

^{*} Community and Conservation Ecology Group, Centre for Ecological and Evolutionary Studies (CEES), University of Groningen, P.O. Box 14, 9750 AA Haren, the Netherlands

[†] Natural Resources and the Environment, Council for Scientific and Industrial Research (CSIR), P.O. Box 91230, Auckland Park, 2006, South Africa

[‡] Wake Forest University, Department of Biology, 206 Winston Hall, Winston-Salem NC 27109, USA

(Accepted 14 February 2011)

Abstract: Litter-feeding termites influence key aspects of the structure and functioning of semi-arid ecosystems around the world by altering nutrient and material fluxes, affecting primary production, foodweb dynamics and modifying vegetation composition. Understanding these complex effects depends on quantifying spatial heterogeneity in termite foraging activities, yet such information is scarce for semi-arid savannas. Here, the amount of litter that was removed from 800 litterbags in eight plots (100 litterbags per plot) was measured in Hluhluwe–iMfolozi Park (HiP) South Africa. These data were used to quantify variation in litter removal at two spatial scales: the local scale (within 450-m² plots) and the landscape scale (among sites separated by 8–25 km). Subsequently, we attempted to understand the possible determinants of termites' foraging patterns by testing various ecological correlates, such as plant biomass and bare ground at small scales and rainfall and fences that excluded large mammalian herbivores at larger scales. No strong predictors for heterogeneity in termite foraging intensity were found at the local scale. At the landscape scale termite consumption depended on an interaction between rainfall and the presence of large mammalian herbivores: litter removal by termites was greater in the presence of large herbivores at the drier sites but lower in the presence of large herbivores at the wetter sites. The effect of herbivores on termite foraging intensity may indicate a switch between termites and large herbivore facilitation and competition across a productivity gradient. In general, litter removal decreased with increasing mean annual rainfall, which is in contrast to current understanding of termite consumption across rainfall and productivity gradients. These results generate novel insights into termite ecology and interactions among consumers of vastly different body sizes across spatial scales.

Key Words: Africa, decomposition, heterogeneity, Isoptera, nitrogen, nutrient cycling, nutrient hotspot, patchiness, spatial scale, termite mound

INTRODUCTION

Savannas are complex ecosystems in which various groups of organisms interact to create heterogeneity in vegetation structure and ecological processes (Anderson *et al.* 2008, Pickett *et al.* 2003, Turner 1989). The resulting heterogeneity, together with that created by variation in the physical landscape, is an important determinant of savanna functioning (Pickett *et al.* 2003, Scholes 1990). To gain a better understanding of savannas it is important to identify the biotic agents

of heterogeneity, understand their distribution and interactions, and quantify the scales at which they affect ecological processes. Important and often overlooked players in the ecology of savannas are litter-feeding termites, which move large quantities of litter, soil and nutrients through the landscape (Goudie 1988).

Termites are among the main macro-faunal organisms involved in litter decomposition in African savannas (Bignell & Eggleton 2000, Scholes & Walker 1993). Their total biomass can exceed large mammalian biomass in African savannas (Deshmukh 1989) and litter removal by termites can amount to 60% of total annual litter production (Wood & Sands 1978). By collecting live and dead plant material and herbivore dung (Freymann *et al.* 2008), and concentrating it in below-ground nest

¹ Corresponding author. Current address: Department of Botany, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa. Email: buitenwerfrobert@hotmail.com

structures their mounds become nutrient hotspots in the landscape (Grant & Scholes 2006, Holt & Lepage 2000, Zaady *et al.* 2003). Nutrients are also prevented from getting lost to the system through run-off or fire (Abenspergtraun & Milewski 1995). These actions, along with N₂ fixation and the potential to change soil structure, can alter plant community composition on the nutrient-enriched mounds (Fox-Dobbs *et al.* 2010, Glover *et al.* 1964, Jouquet *et al.* 2004, Loveridge & Moe 2004, Moe *et al.* 2009, Mordelet & Menaut 1995), increase plant N and P concentrations (Grant & Scholes 2006, Jouquet *et al.* 2005), and ultimately increase herbivory on the mounds (Dangerfield *et al.* 1998, Grant & Scholes 2006, Loveridge & Moe 2004).

By creating heterogeneity termites have the potential to significantly affect ecosystem functioning and food-web dynamics (Pringle *et al.* 2010), however without identifying the determinants of termite distribution and consumption, the extent of their impact cannot be fully understood. Rainfall has been identified as a major determinant for termite abundance and consumption across continental and regional scales, where abundance and consumption increase with rainfall (Buxton 1981, Deshmukh 1989, Picker *et al.* 2007, Pomeroy 1978). This pattern is suggested to result from an increase in food availability with rainfall, reflecting the well-known positive relation between primary production and annual rainfall in semi-arid ecosystems. However, primary production and peak herbaceous biomass in African savanna ecosystems are also significantly affected by other variables, such as fire and herbivores (Scholes & Walker 1993). While a handful of studies have quantified effects of fire and herbivores on termite species assemblages and abundance (Abenspergtraun 1992, Abenspergtraun & Milewski 1995, Tracy *et al.* 1998), it is unclear how termite consumption changes across a rainfall gradient in a system dominated by herbivores.

The goal of our research was to study the relationship between rainfall and termite consumption in the presence and absence of large herbivores, while controlling for the effects of fire. We expect termite activity to be higher in areas with high rainfall and in the absence of herbivores, as litter production should be highest under these conditions. Importantly, we quantify foraging intensity of the entire grass- and litter-feeding functional group without separating effects among species, as opposed to previous studies that selected species with large, above-ground mounds (Buxton 1981, Deshmukh 1989, Ferrar 1982, Meyer 2001, Meyer *et al.* 1999, Picker *et al.* 2007, Pomeroy 1978) and therefore excluded the majority of termite species (Uys 2002). However, because the factors that determine variation in termite consumption across a rainfall or productivity gradient are expected to be different than those that determine termite foraging activity on a local level, we conducted research at two

spatial scales: within 450-m² plots and across sites separated by 8–25 km. Within sites, where climate and soils are similar, we expect termite activity to be highest in patches with high vegetation cover as food availability and physical protection from predators and harmful solar radiation is highest there.

STUDY SITE

The study was conducted at Hluhluwe–iMfolozi Park (HiP) (28°00′–28°26′S, 31°43′–32°00′E) an 897-km² reserve in KwaZulu-Natal, South Africa. Within HiP annual rainfall ranges between 630 mm in the low-altitude areas and 1000 mm on the highest peaks, resulting in a strong rainfall gradient over a relatively short distance (Balfour & Howison 2001). The Hluhluwe (northern) part of the reserve is characterized by mixed patches of forest, grassland, thicket and savanna. Vegetation in the iMfolozi (southern) part mainly consists of open savanna woodland (Whateley & Porter 1983).

Termite consumption was studied at four sites within HiP: Mona, Gqoyeni, Ledube and Nombali. Nombali and Ledube are situated in a high-rainfall area (628 and 707 mm y⁻¹, respectively) on nutrient-poor substrate (sandstone and shale) while Gqoyeni and Mona are situated in a low-rainfall area (561 and 551 mm y⁻¹, respectively) on more mineral-nutrient-rich substrate (dolerite) (Table 1). Each site contains a long-term 40 × 40-m fenced herbivore enclosure (Figure 1) which excludes all herbivores larger than a scrub hare (*Lepus saxatilis* F. Cuvier, 1823). Common large herbivores visiting the sites were white rhino (*Ceratotherium simum* Burchell, 1817), buffalo (*Syncerus caffer* Sparrman, 1779), zebra (*Equus quagga burchellii* Gray, 1824), impala (*Aepyceros melampus* Lichtenstein, 1812) and warthog (*Phacochoerus africanus* Gmelin, 1718). Each enclosure was paired with an open area of similar size to control for the effects of grazing. Controlled burns were applied every second year both inside the enclosure and to the adjacent control areas.

METHODS

Quantification of termite activity

Termite consumption rates were measured by quantifying litter removal from mesh bags placed at the sites (Bodine & Ueckert 1975). Bags were filled with 5.0 g of dried (48 h at 60 °C) *Themeda triandra* Forssk. grass harvested from a single location to control for variation in forage quality. Since the grass placed within the bags was mostly moribund, it was assumed that nutrients had been largely resorbed by the plants (Ratnam *et al.* 2008) and thus

Table 1. Mean annual rainfall, interpolated from 11 rainfall stations in HiP, between 2001 and 2007 and parent geological material (King 1970) are given for the four study sites. Mean biomass from 200 disc-pasture meter measurements per plot ($\pm$ SE) and the mean proportion of bare ground from 200 visual estimates per plot ($\pm$ SD) are given for each treatment plot within site.

	Rain (mm)	Parent material	Treatment plot	Biomass (g m^{-2})	Bare ground (%)
Mona	551	Dolerite, basalt	Grazed	13 ± 0	95 ± 3
			Ungrazed	115 ± 4	63 ± 11
Gqoyeni	561	Dolerite, basalt, alluvium	Grazed	13 ± 0	93 ± 11
			Ungrazed	129 ± 4	72 ± 9
Ledube	705	Sandstone, shale	Grazed	360 ± 12	6 ± 4
			Ungrazed	307 ± 12	4 ± 3
Nombali	628	Sandstone, shale	Grazed	163 ± 6	35 ± 20
			Ungrazed	305 ± 18	5 ± 2

Figure 1. Hluhluwe–iMfolozi Park, South Africa, with the location of the four study sites where litter removal by termites was measured. The inset shows the schematic layout of two plots (40×40 m) within a site: an ungrazed plot from which mammalian herbivores were excluded and an equally sized control plot open to herbivores. The dots within the plots indicate the location of 100 litter bags used to quantify litter removal by termites.

the harvested material was functionally equivalent to senesced grass litter. Grass was cut into segments of approximately 5 cm, mixing leaves and stems. Bags were constructed from aluminium mesh with a pore size of 2×2 mm, although the loosely woven structure of the material allowed slightly larger-bodied termites (i.e. *Hodotermes mossambicus* Hagen and *Macrotermes natalensis* Haviland) into the litterbags. The bags were approximately 10×10 cm and were secured to the substrate with a nail.

Litter removal rates were measured within 450-m^2 plots (15×30 m) that were established in each enclosure and adjacent control area. Plots were divided in a regular grid of 200 cells measuring 1.5×1.5 m. A litterbag was placed in every second cell, starting in the first cell of each odd numbered row and in the second cell of each even numbered row to obtain an optimal coverage. The 100 litterbags in each plot were collected after 1 mo, dried at 60°C for 48 h and then emptied. Grass was separated from soil that had accumulated in the bags as a result of soil sheathing by termites, and both were weighed separately.

Within each grid cell grass biomass was measured using a disc-pasture meter (DPM) (Bransby & Tainton 1977) from which biomass under the disc was calculated by: $\text{grass biomass } (\text{g m}^{-2}) = 12.6 + 26.1 \text{ DPM}$ ($R^2 = 0.73$, $N = 1745$) (Waldram *et al.* 2008). Proportion of bare ground was estimated visually within grid cells.

Statistical analyses

Local-scale patterns of variation in termite activity within plots were assessed by calculating Moran's I statistic for spatial autocorrelation. Our a priori expectation was that termite foraging would be patchy and that patches of high foraging activity would coincide with patches of high resource availability, such as herbaceous biomass (food) and the proportion of bare ground. Patchiness within plots of herbaceous biomass and bare ground was assessed with Moran's I. Within-plot correlations between litter removal and herbaceous biomass and bare ground were used to assess the spatial association of litter removal with resource availability. Inverse distance-weighted interpolation surfaces (power = 2, extent = 12 closest points) of litter removal were created using Spatial Analyst in ArcMap 9.2 (ESRI 2006, Redlands, CA, USA).

To test the effects of rainfall and herbivory on litter removal at the landscape scale, a linear mixed-effects model was constructed using the LME function in the NLME library version 3.1–89 (R Foundation for Statistical Computing, Vienna, Austria) for R (R Development Core Team, R Foundation for Statistical Computing, Vienna, Austria). The LME function employs a restricted maximum likelihood (REML) estimator. The main advantage of this method is that it is not sensitive to unbalanced designs or missing observations (Pinheiro & Bates 2000). While our design was balanced, some observations (litter bags) were missing (Appendix 1). Close proximity of litterbags within a site could result in correlated measurements between them (spatial autocorrelation), however the LME function allows such correlated within-group errors to be estimated explicitly

Table 2. Moran's I statistics for within-plot spatial autocorrelation of litter removal rates, herbaceous biomass and proportion of bare ground. *denotes significance at $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Site	Treatment plot	Litter removal	Biomass	Bare ground
Mona	Grazed	0.39	0.00	0.00
Mona	Ungrazed	1.83*	0.02***	0.07***
Gqoyeni	Grazed	1.71*	0.00	0.00
Gqoyeni	Ungrazed	0.71	0.01*	0.06***
Ledube	Grazed	-0.21	0.10***	0.16***
Ledube	Ungrazed	-0.19	0.02*	0.05***
Nombali	Grazed	0.33	0.05***	0.11***
Nombali	Ungrazed	-0.37	0.00	0.01

in the model. LME also allows unequal variances between groups to be modelled explicitly by adding a variance structure to the model. The model included rainfall and herbivory as fixed effects, while site was included as a random variable. The best model was selected using Akaike information criterion (AIC) where smaller values indicate a better model.

RESULTS

Within plots

In general, we found no clear spatial association between termite mounds and spatial patterns of litter removal within sites. Within-plot values of litter removal revealed statistically significant spatial autocorrelation (patchiness) in only two of the eight plots (Figure 2, Table 2). Both plots were located in dry sites, however one (Mona) was inaccessible to herbivores, while the other (Gqoyeni), was exposed to herbivores. In those two plots the local litter removal rate was negatively correlated with proportion of bare ground: significantly in Mona ungrazed ($r = -0.26$, $P < 0.01$, $n = 100$) and with a small but not significant P-value in Gqoyeni grazed ($r = -0.20$, $P = 0.06$, $n = 85$). There were no significant correlations between proportion of bare ground and litter removal in the other plots, nor were there significant correlations between herbaceous biomass and litter removal in any of the plots. The small variance of biomass and proportion of bare ground within most plots (Table 1) indicates a fairly homogeneous vegetation structure at the measured scale, even though Moran's I for spatial clustering was significant for biomass and proportion of bare ground in five of the eight plots (Table 2).

Between plots

We found a significant interactive effect of rainfall and grazing ($F_{1,2} = 71.7$, $P < 0.05$) on litter removal, where grazing increased litter removal in the dry area but

Figure 2. Inverse distance-weighted interpolation surface (15×30 m) of litter removal within two out of eight study plots. The plots shown here were the only plots with significant spatial autocorrelation in litter removal by termites. The location of termite mounds within plots is indicated to show the poor relationship between patterns of litter removal and location of termite mounds.

Figure 3. Mean percentage of litter removal by termites in wet and dry areas, separated between herbivore exclusion plots and plots open to herbivores. Error bars indicate 1 SE.

decreased litter removal in the wet area (Figure 3). The main effect of grazing on litter removal was also highly significant ($F_{1,2} = 144$; $P < 0.01$) while the effect of rainfall had a small but not significant P-value ($F_{1,2} = 16.2$; $P = 0.057$) (more litter removal in the drier sites).

Of the initial variation in litter removal, approximately 30% was associated with differences between sites, while the remaining 70% was associated with differences within sites. Of the 30% between-site variance, 90% was explained by adding rainfall as a predictor variable.

Adding grazing and its interaction with rainfall to the model explained 10% of the initial 70% within-site variance.

A model including a random site effect had a significantly smaller AIC (−843) than the equivalent model without a random site effect (AIC = −818), indicating that common (unstudied) site characteristics had a significant effect on litter removal.

As can be seen from Figure 3, the variance in litter removal in the dry area was greater than in the wet area, and therefore modelling the variance for dry and wet areas separately significantly improved the model. The variance in the wet area was modelled as 46% of the variance in the dry area, which improved the AIC from −843 to −1059. Adding a correlation structure did not significantly improve the model (AIC was not reduced), indicating that there was no significant spatial auto-correlation within sites.

DISCUSSION

As litter-feeding termites contribute substantially to spatial heterogeneity and ecological processes in savannas (Pringle *et al.* 2010) it is important to identify determinants of their distribution and foraging intensity. In this study we show that termite foraging intensity varies with rainfall and herbivory across relatively large scales (among sites separated by 8–25 km), while termite foraging showed little spatial pattern and did not correlate well with herbaceous vegetation structure at local scales (within 450-m² plots). The significant interaction effect between rainfall and herbivory on termite foraging intensity may indicate a switch between facilitation and competition between large herbivores and termites across the rainfall gradient in HiP. Additionally, an overall negative relation between rainfall and litter removal transpired from the rainfall main effect that explained the largest portion of variation in litter removal. Such a negative relationship between rainfall and termite consumption is inconsistent with previous studies (Buxton 1981, Deshmukh 1989). While we acknowledge that our small sample size and restricted rainfall range limits our ability to extrapolate the results to other ecosystems, we feel that our study highlights ecological interactions which may exist in other savannas.

Local scale

Our initial expectation was that epigeal mound placement within sites would serve as an indicator of termite foraging as they are useful indicators of termite distribution for species that construct them, e.g. *Macrotermes* spp. (Glover *et al.* 1964, Meyer *et al.* 1999). One possible reason for the

lack of association between mounds and termite activity on the local scale is that we made no attempt to test mound occupancy. Mound occupancy can however not be used to explain differences between grazed and ungrazed plots within a site, as these plots are in close enough proximity to overlap with the foraging range of a single termite colony (Coaton & Sheasby 1972, Heidecker & Leuthold 1984). Another possibility is that our plots were too small to capture termite foraging patterns at local scales. A final explanation is that some termite species do not construct (obvious) mounds and therefore mounds may be poor predictors of the foraging intensity of an entire functional feeding group (Abenspergtraun 1992).

Based on termite biology, foraging intensity on the local scale was expected to be higher under vegetation cover, e.g. from higher soil moisture that facilitates the construction of protective sheeting (Belsky *et al.* 1989, Dangerfield & Schuurman 2000, Whitford *et al.* 1982) or through interception of harmful solar radiation (Holt & Lepage 2000). The proportion of bare ground correlated negatively with termite activity but only in plots with significant spatial clustering of termite activity. In these plots, as expected, termites therefore concentrate foraging in covered patches, however this pattern was not present in plots without spatial clustering of termite activity at the scale of measurement. It is possible that termite activity in these plots is clustered at smaller or larger spatial scales than the measured scale and the negative relationship with bare ground might be present at the scale of clustering. Small-scale heterogeneity in termite activity found in other ecosystems was correlated to vegetation structure and attributed to litter availability (Tracy *et al.* 1998). Therefore, the weak correlation we found between litter removal and vegetation structure in two of our eight plots, may indicate that stronger relationships might be obtained by adjusting the scale of observation. This could be achieved by either increasing the spatial extent (plot size) to include more heterogeneity in vegetation structure, or by decreasing the spatial grain (cell size) to measure heterogeneity at a smaller scale.

Landscape scale

In contrast to the local scale, we identified strong predictors of termite foraging intensity on the landscape scale. Foraging intensity was much higher in dry sites compared with wetter sites, which contradicts previous studies that find an increase of termite consumption with increasing rainfall (Buxton 1981, Deshmukh 1989). Deshmukh (1989) compiled consumption rates from studies across the African continent and suggests that the increase in termite consumption with increased rainfall is driven by increased food availability, due to the increase of herbaceous primary production with rainfall (Rosenzweig

1968, Rutherford 1981). Buxton (1981) also reports increased termite consumption with rainfall in Tsavo National Park, Kenya, even though the two driest sites (total sites = 9), which also had the highest termite consumption, were left out of the regression because they did not fit the trend.

We offer some possible explanations for the apparent mismatch between termite consumption and food availability in our study. Firstly, litter quality may be higher in the drier area. While litter N concentrations are likely to be higher in the dry area as a result of the nutrient-rich geological substrate, termites are highly adapted to food with extremely high C:N ratios and are therefore not likely to be attracted to more nutrient-rich litter (Rouland *et al.* 2003). However, grasses in the wetter part of HiP have significantly higher concentrations of lignin and secondary metabolites such as phenolics (Masumelele 2007), which may make them less palatable and hence decrease litter quality to termites and the *Termitomyces* sp. R. Heim fungus that is cultivated by Macrotermitinae species.

Secondly, food accessibility for termites may be facilitated in the dry part of HiP by the high abundance of mammalian grazers (Cromsigt *et al.* 2009), potentially resulting in a higher carrying capacity for termites and an overall higher termite foraging intensity in this area. Grazers increase litter-fall by dropping plant fragments whilst grazing and trample the vegetation (Cumming & Cumming 2003, Deshmukh 1989), making it more accessible to termites. In addition, herbivore dung contains a large proportion of undigested plant material and is readily exploited by litter-feeding termites (Freyman *et al.* 2008). Within the dry sites, plots with large herbivores had higher termite foraging intensity, supporting the proposed positive effects of large herbivores on termites. It remains unclear why the opposite pattern is observed in the wet area, where termite consumption is higher in ungrazed plots. Possibly this is due to a less favourable microclimate within the tall-grass vegetation of the exclosures. Contrasting effects of large herbivores on termites are also reported in other studies. In a Chihuahuan desert ecosystem, termite activity was higher in ungrazed sites compared to grazed sites, which was attributed to changed litter availability in the grazed area (Tracy *et al.* 1998). In an Australian *Eucalyptus* woodland and a Burkina Faso savanna, no effects of grazing on termite diversity and abundance were found, although foraging activity was not measured directly (Abenspergtraun 1992, Traore & Lepage 2008). While herbivores clearly have effects on termites, the mechanisms by which they do so remain unclear and may be interactive, e.g. with rainfall, and location-specific.

The positive association between herbivores and termite consumption that we found may lead to food competition during droughts, as reported for African

rangelands where harvester termites (*Hodotermes* spp.) consumed up to 60% of standing grass biomass and all the litter, resulting in stock mortality (Coaton & Sheasby 1972, Mitchell 2002). No such dramatic events have been reported for systems with wild herbivores, however a detailed understanding of interactions between herbivores and termites will improve understanding of ecosystem functioning and is likely to benefit the management of protected areas and large herbivores.

CONCLUSIONS

To fully understand ecosystem structure and functioning it is essential to identify determinants of termite distribution and foraging intensity and quantify relations with other ecosystem components such as herbivores. This study provides novel insights into the relationship of termites, the main litter decomposers and primary agents of nutrient and vegetation heterogeneity in savannas, with rainfall and mammalian herbivores. The exact mechanisms that produce the observed patterns and correlations need to be identified in order to improve understanding and management of savanna ecosystems.

ACKNOWLEDGEMENTS

The authors wish to thank Khanyi Mpandza and Aafke van Erk for assistance with data collection. We thank Ezemvelo KwaZulu-Natal Wildlife for allowing us to do research in HiP. The Marco Polo Fund and the Groninger Universiteits Fonds partially funded this research.

LITERATURE CITED

- ABENSPERGTRAUN, M. 1992. The effects of sheep-grazing on the subterranean termite fauna (Isoptera) of the Western-Australian wheat-belt. *Australian Journal of Ecology* 17:425–432.
- ABENSPERGTRAUN, M. & MILEWSKI, A. V. 1995. Abundance and diversity of termites (Isoptera) in unburnt versus burnt vegetation at the Barrens in mediterranean Western-Australia. *Australian Journal of Ecology* 20:413–417.
- ANDERSON, T. M., DEMPEWOLF, J., METZGER, K. L., REED, D. N. & SERNEELS, S. 2008. Generation and maintenance of heterogeneity in the Serengeti ecosystem. Pp. 135–182 in Sinclair, A. R. E., Packer, C., Mduma, S. A. R. & Fryxell, J. M. (eds.). *Serengeti III: human impacts on ecosystem dynamics*. The University of Chicago Press, Chicago.
- BALFOUR, D. A. & HOWISON, O. E. 2001. Spatial and temporal variation in a mesic savanna fire regime: responses to variation in annual rainfall. *African Journal of Range and Forage Science* 19:43–51.
- BELSKY, A. J., AMUNDSON, R. G., DUXBURY, J. M., RIHA, S. J., ALL, A. R. & MWONGA, S. M. 1989. The effects of trees on their physical, chemical, and biological environments in a semi-arid savanna in Kenya. *Journal of Applied Ecology* 26:1005–1024.

- BIGNELL, D. E. & EGGLETON, P. 2000. Termites in ecosystems. Pp. 363–387 in Abe, T., Bignell, D. & Higashi, M. (eds.). *Termites: evolution, sociality, symbioses, ecology*. Kluwer Academic Publishers, Dordrecht.
- BODINE, M. C. & UECKERT, D. N. 1975. Effect of desert termites on herbage and litter in a shortgrass ecosystem in west Texas. *Journal of Range Management* 28:353–358.
- BRANSBY, D. I. & TAINTON, N. M. 1977. The disk pasture meter: possible applications in grazing management. *Proceedings of the Grassland Society of Southern Africa* 5:115–118.
- BUXTON, R. D. 1981. Changes in the composition and activities of termite communities in relation to changing rainfall. *Oecologia* 51:371–378.
- COATON, W. G. H. & SHEASBY, J. L. 1972. *Preliminary report on a survey of the termites (Isoptera) of South West Africa*. Plant Protection Research Institute, Pretoria. 129 pp.
- CROMSIGT, J. P. G. M., PRINS, H. H. T. & OLFF, H. 2009. Habitat heterogeneity as a driver of ungulate diversity and distribution patterns: interaction of body mass and digestive strategy. *Diversity and Distributions* 15:513–522.
- CUMMING, D. H. M. & CUMMING, G. S. 2003. Ungulate community structure and ecological processes: body size, hoof area and trampling in African savannas. *Oecologia* 134:560–568.
- DANGERFIELD, J. M. & SCHUURMAN, G. 2000. Foraging by fungus-growing termites (Isoptera: Termitidae, Macrotermitinae) in the Okavango Delta, Botswana. *Journal of Tropical Ecology* 16:717–731.
- DANGERFIELD, J. M., MCCARTHY, T. S. & ELLERY, W. N. 1998. The mound-building termite *Macrotermes michaelseni* as an ecosystem engineer. *Journal of Tropical Ecology* 14:507–520.
- DESHMUKH, I. 1989. How important are termites in the production ecology of African savannas? *Sociobiology* 15:155–168.
- FERRAR, P. 1982. Termites of a South African savanna 1. List of species and subhabitat preferences. *Oecologia* 52:125–132.
- FOX-DOBBS, K., DOAK, D. F., BRODY, A. K. & PALMER, T. M. 2010. Termites create spatial structure and govern ecosystem function by affecting N₂ fixation in an East African savanna. *Ecology* 91:1296–1307.
- FREYMAN, B. P., BUITENWERF, R., DESOUZA, O. & OLFF, H. 2008. The importance of termites (Isoptera) for the recycling of herbivore dung in tropical ecosystems: a review. *European Journal of Entomology* 105:165–173.
- GLOVER, P. E., TRUMP, E. C. & WATERIDGE, L. E. D. 1964. Termitaria and vegetation patterns on the Loita plains of Kenya. *Journal of Ecology* 52:367–377.
- GOUDIE, A. S. 1988. The geomorphological role of termites and earthworms in the tropics. Pp. 166–191 in Viles, H. A. (ed.). *Biogeomorphology*. Blackwell, Oxford.
- GRANT, C. C. & SCHOLES, M. C. 2006. The importance of nutrient hotspots in the conservation and management of large wild mammalian herbivores in semi-arid savannas. *Biological Conservation* 130:426–437.
- HEIDECKER, J. L. & LEUTHOLD, R. H. 1984. The organisation of collective foraging in the harvester termite *Hodotermes mossambicus* (Isoptera). *Behavioral Ecology and Sociobiology* 14:195–202.
- HOLT, J. A. & LEPAGE, M. 2000. Termites and soil properties. Pp. 389–407 in Abe, T., Bignell, D. E. & Higashi, M. (eds.). *Termites: evolution, sociality, symbioses, ecology*. Kluwer Academic Publishers, Dordrecht.
- JOUQUET, P., TESSIER, D. & LEPAGE, M. 2004. The soil structural stability of termite nests: role of clays in *Macrotermes bellicosus* (Isoptera, Macrotermitinae) mound soils. *European Journal of Soil Biology* 40:23–29.
- JOUQUET, P., TAVERNIER, V., ABBADIE, L. & LEPAGE, M. 2005. Nests of subterranean fungus-growing termites (Isoptera, Macrotermitinae) as nutrient patches for grasses in savannah ecosystems. *African Journal of Ecology* 43:191–196.
- KING, L. 1970. The geology of the Hluhluwe Game Reserve. *Petros* 2:16–19.
- LOVERIDGE, J. P. & MOE, S. R. 2004. Termitaria as browsing hotspots for African megaherbivores in miombo woodland. *Journal of Tropical Ecology* 20:337–343.
- MASUMELELE, M. L. 2007. *Decomposition of grasses in a South African savanna*. M.Sc. thesis, University of Cape Town, Cape Town. 124 pp.
- MEYER, V. W. 2001. *Intracolony demography, biomass and food consumption of Macrotermes natalensis (Haviland) (Isoptera: Termitidae) colonies in the northern Kruger National Park, South Africa*. Ph.D. thesis, University of Pretoria, Pretoria. 84 pp.
- MEYER, V. W., BRAACK, L. E. O., BIGGS, H. C. & EBERSOHN, C. 1999. Distribution and density of termite mounds in the northern Kruger National Park, with specific reference to those constructed by *Macrotermes Holmgren* (Isoptera: Termitidae). *African Entomology* 7:123–130.
- MITCHELL, J. D. 2002. Termites as pests of crops, forestry, rangeland and structures in southern Africa and their control. *Sociobiology* 40:47–69.
- MOE, S. R., MOBAEK, R. & NARMO, A. K. 2009. Mound building termites contribute to savanna vegetation heterogeneity. *Plant Ecology* 202:31–40.
- MORDELET, P. & MENAUT, J. C. 1995. Influence of trees on aboveground production dynamics of grasses in a humid savanna. *Journal of Vegetation Science* 6:223–228.
- PICKER, M. D., HOFFMAN, M. T. & LEVERTON, B. 2007. Density of *Microhodotermes viator* (Hodotermitidae) mounds in southern Africa in relation to rainfall and vegetative productivity gradients. *Journal of Zoology* 271:37–44.
- PICKETT, S. T. A., CADENASSO, M. L. & BENNING, T. L. 2003. Biotic and abiotic variability as key determinants of savanna heterogeneity at multiple spatiotemporal scales. Pp. 22–40 in du Toit, J. T., Rogers, K. H. & Biggs, H. C. (eds.). *The Kruger experience: ecology and management of savanna heterogeneity*. Island Press, Washington.
- PINHEIRO, J. C. & BATES, D. M. 2000. *Mixed-effects models in S and S-PLUS*. Springer Verlag, New York. 528 pp.
- POMEROY, D. E. 1978. The abundance of large termite mounds in Uganda in relation to their environment. *Journal of Applied Ecology* 15:51–63.
- PRINGLE, R. M., DOAK, D. F., BRODY, A. K., JOCQUE, R. & PALMER, T. M. 2010. Spatial pattern enhances ecosystem functioning in an African savanna. *PLoS Biology* 8:e1000377.
- RATNAM, J., SANKARAN, M., HANAN, N. P., GRANT, R. C. & ZAMBATIS, N. 2008. Nutrient resorption patterns of plant functional

- groups in a tropical savanna: variation and functional significance. *Oecologia* 157:141–151.
- ROSENZWEIG, M. L. 1968. Net primary productivity of terrestrial communities: prediction from climatological data. *American Naturalist* 102:67–74.
- ROULAND, C., LEPAGE, M., CHOTTE, J. L., DIOUF, M., NDIAYE, D., NDIAYE, S., SEUGE, C. & BRAUMAN, A. 2003. Experimental manipulation of termites (Isoptera, Macrotermitinae) foraging patterns in a Sahelo-Sudanese savanna: effect of litter quality. *Insectes Sociaux* 50:309–316.
- RUTHERFORD, M. C. 1981. Annual plant-production precipitation relations in arid and semi-arid regions. *South African Journal of Science* 76:53–56.
- SCHOLES, R. J. 1990. The influence of soil fertility on the ecology of southern African dry savannas. *Journal of Biogeography* 17:415–419.
- SCHOLES, R. J. & WALKER, B. H. 1993. *An African savanna, synthesis of the Nylsvley study*. Cambridge University Press, Cambridge. 306 pp.
- TRACY, K. N., GOLDEN, D. M. & CRIST, T. O. 1998. The spatial distribution of termite activity in grazed and ungrazed Chihuahuan desert grassland. *Journal of Arid Environments* 40:77–89.
- TRAORE, S. & LEPAGE, M. 2008. Effects of controlled livestock grazing and annual prescribed fire on epigeal termite mounds in a savannah woodland in Burkina Faso. *Insectes Sociaux* 55:183–189.
- TURNER, M. G. 1989. Landscape ecology: the effect of pattern on process. *Annual Review of Ecology and Systematics* 20:171–197.
- UYS, V. 2002. *A guide to the termite genera of southern Africa*. ARC-Plant protection Research Institute, Pretoria. 116 pp.
- WALDRAM, M. S., BOND, W. J. & STOCK, W. D. 2008. Ecological engineering by a mega-grazer: white rhino impacts on a South African savanna. *Ecosystems* 11:101–112.
- WHATELEY, A. & PORTER, R. N. 1983. The woody vegetation communities of the Hluhluwe-Corridor-Umfolozi Game Reserve complex. *Bothalia* 14:745–758.
- WHITFORD, W. G., STEINBERGER, Y. & ETTERS HANK, G. 1982. Contributions of subterranean termites to the economy of Chihuahuan desert ecosystems. *Oecologia* 55:298–302.
- WOOD, T. G. & SANDS, W. A. 1978. The role of termites in ecosystems. Pp. 245–292 in Brian, M. V. (ed.). *Production ecology of ants and termites*. Cambridge University Press, Cambridge.
- ZAADY, E., GROFFMAN, P. M., SHACHAK, M. & WILBY, A. 2003. Consumption and release of nitrogen by the harvester termite *Anacanthotermes ubachi* navas in the northern Negev desert, Israel. *Soil Biology and Biochemistry* 35:1299–1303.

Appendix 1. While in each plot 100 litterbags were laid out initially, some bags were not retrieved as a result of disturbance by animals. Here the actual number of litterbags per plot that were analysed is shown.

Site	Treatment plot	Number of litterbags
Mona	Grazed	92
Mona	Ungrazed	100
Gqoyeni	Grazed	85
Gqoyeni	Ungrazed	99
Ledube	Grazed	99
Ledube	Ungrazed	100
Nombali	Grazed	99
Nombali	Ungrazed	100