University of Groningen

Oral isotretinoin and pregnancy prevention programmes

Crijns, H. J. M. J.; Straus, S. M.; de Jong-van den Berg, L. T. W.

Published in:
BRITISH JOURNAL OF DERMATOLOGY

DOI:
[10.1111/j.1365-2133.2011.10687.x](https://doi.org/10.1111/j.1365-2133.2011.10687.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Crijns, H. J. M. J., Straus, S. M., & de Jong-van den Berg, L. T. W. (2012). Oral isotretinoin and pregnancy prevention programmes: reply from authors. *BRITISH JOURNAL OF DERMATOLOGY*, 166(2), 467-468.
<https://doi.org/10.1111/j.1365-2133.2011.10687.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Oral isotretinoin has had a major positive impact on the lives of countless patients since its introduction in the early 1980s. Of course, we feel it is imperative to provide oral isotretinoin to our patients in the safest manner possible. But we also believe that instituting increasingly restrictive programmes around the use of the drug may not be the optimal approach in view of the limitations of such programmes. In particular, restrictive programmes such as iPLEDGE may have reached the limit of influencing human behaviour. More restrictive programmes, apart from mandated use of injectable long-acting contraceptives, cannot prevent unintended pregnancies resulting from failure to use contraceptives appropriately. We think that the situation is similar for Europe. Until there are improved options for the patients who are now isotretinoin candidates, it is important to take a balanced approach that allows for appropriate use of this important drug. We are saddened that the article by Shin et al. suggests that women selectively may be receiving fewer prescriptions for isotretinoin, despite the fact that prescription levels for males have remained relatively constant. We agree with a recent editorial by Maloney and Stone¹⁵ suggesting that a team or committee evaluation of prescribing patterns and fetal exposure would be a valuable addition to the current system with the goal of improving patient access and outcomes.

Department of Dermatology,
Pennsylvania State University College of Medicine,
Hershey, PA 17033, U.S.A.

*Department of Dermatology and Venereology,
Medical Faculty, Otto-Von-Guericke-Universität,
Magdeburg, Germany

†Department of Dermatology, Arcispedale S. Anna,
University of Ferrara, Ferrara, Italy

‡Department of Dermatology,
Hotel Dieu, Nantes, France

§Department of Dermatology,
Johns Hopkins Medicine,
Baltimore, MD, U.S.A.

¶Department of Dermatology,
University of Pennsylvania
School of Medicine,
Philadelphia, PA, U.S.A.

**Department of Dermatology,
State University of New York
Downstate Medical Center, Brooklyn, NY, U.S.A.

††Department of Dermatology,
Juarez Hospital, Mexico City, Mexico
E-mail: dthiboutot1@gmail.com

D. THIBOUTOT

H. GOLLNICK*

V. BETTOLI†

B. DRÉNO‡

S. KANG§

J.J. LEYDEN¶

A. SHALITA**

V. TORRES††

FOR THE GLOBAL
ALLIANCE TO IMPROVE
OUTCOMES IN ACNE

- 2 Cunliffe WJ, Gould DJ. Prevalence of facial acne vulgaris in late adolescence and in adults. *Br Med J* 1979; **1**:1109–10.
- 3 Goulden V, McGeown CH, Cunliffe WJ. The familial risk of adult acne: a comparison between first-degree relatives of affected and unaffected individuals. *Br J Dermatol* 1999; **141**:297–300.
- 4 Collier CN, Harper JC, Cafardi JA et al. The prevalence of acne in adults 20 years and older. *J Am Acad Dermatol* 2008; **58**:56–9.
- 5 Koo J. The psychosocial impact of acne: patients' perceptions. *J Am Acad Dermatol* 1995; **32**:S26–30.
- 6 Kellett SC, Gawkrödger DJ. The psychological and emotional impact of acne and the effect of treatment with isotretinoin. *Br J Dermatol* 1999; **140**:273–82.
- 7 Pearl A, Arroll B, Lello J et al. The impact of acne: a study of adolescents' attitudes, perception and knowledge. *N Z Med J* 1998; **111**:269–71.
- 8 Tallab TM. Beliefs, perceptions and psychological impact of acne vulgaris among patients in the Assir region of Saudi Arabia. *West Afr J Med* 2004; **23**:85–7.
- 9 Tan JK, Balagurusamy M, Fung K et al. Effect of quality of life impact and clinical severity on adherence to topical acne treatment. *J Cutan Med Surg* 2009; **13**:204–8.
- 10 Tan JK, Li Y, Fung K et al. Divergence of demographic factors associated with clinical severity compared with quality of life impact in acne. *J Cutan Med Surg* 2008; **12**:235–42.
- 11 Hassan J, Grogan S, Clark-Carter D et al. The individual health burden of acne: appearance-related distress in male and female adolescents and adults with back, chest and facial acne. *J Health Psychol* 2009; **14**:1105–18.
- 12 Schaefer C. Drug safety in pregnancy: utopia or achievable prospect? Risk information, risk research and advocacy in Teratology Information Services. *Congenit Anom (Kyoto)* 2011; **51**:6–11.
- 13 Mendelsohn AB, Governale L, Trontell A et al. Changes in isotretinoin prescribing before and after implementation of the System to Manage Accutane Related Teratogenicity (SMART) risk management program. *Pharmacoevidenciol Drug Saf* 2005; **14**:615–18.
- 14 Shin J, Cheetham TC, Wong L et al. The impact of the iPLEDGE program on isotretinoin fetal exposure in an integrated health care system. *J Am Acad Dermatol* 2011; **65**:1117–25.
- 15 Maloney ME, Stone SP. Isotretinoin and iPLEDGE: a view of results. *J Am Acad Dermatol* 2011; **65**:418–19.

Funding sources: Galderma supported the writing of this manuscript by funding the services of a medical writer.

Conflicts of interest: Galderma International supports The Global Alliance to Improve Outcomes in Acne with an unrestricted educational grant and funded the services of a professional medical writer.

References

- 1 Crijns HJM, Straus SM, Gispen-de Wied C, de Jong-van den Berg LTW. Compliance with pregnancy prevention programmes of isotretinoin in Europe: a systematic review. *Br J Dermatol* 2011; **164**:238–44.

Oral isotretinoin and pregnancy prevention programmes: reply from authors

DOI: 10.1111/j.1365-2133.2011.10687.x

MADAM, We thank Professors Thiboutot, Gollnick, Bettoli, Dréno, Kang, Leyden, Shalita and Torres¹ for the Global Alliance to Improve Outcome in Acne, for their comments on our article.² Their letter points out an important aspect of having acne and the effectiveness of isotretinoin in treating this disorder.

Thiboutot et al. draw attention to the effectiveness of isotretinoin and the psychological burden of acne. Most patients with acne are in the age group of 12–20 years and approximately 50% are females of childbearing potential.

The effectiveness of isotretinoin was not considered the subject of discussion in our paper. We agree that the effectiveness of isotretinoin is well acknowledged³ and the benefit–risk balance is considered to be positive, provided that the pregnancy prevention programme (PPP) is adhered to.

The major concern for isotretinoin is its teratogenicity.

The risk of congenital anomalies with isotretinoin should be minimized in a way such that it can still be prescribed for the treatment of acne with the least restraints. This can only be achieved when both healthcare professionals and patients are aware of – and are compliant with – the PPP. By creating awareness of noncompliance with the PPP, it is hoped that prescribers, pharmacists and patients will adjust their routine and take responsibility for adhering to the PPP.

The iPLEDGE programme in the U.S.A. contains a PPP with stricter rules and requirements compared with the previous programme ‘System to Manage Accutane-Related Teratogenicity’ (SMART). Research showed that iPLEDGE did not decrease the pregnancy rate⁴ compared with the rate while SMART was in place. This information might suggest that other actions might be necessary to improve the prevention of pregnancies and the adherence to the programme.

We agree with Thiboutot et al. that the emotional aspect of acne should not be trivialized; however, the retinoid embryopathy might also have a major impact, as well as the emotional burden of induced or spontaneous abortion in patients getting pregnant while using isotretinoin. Based on the current experience with the programmes (PPPs), we also would like to support the suggestion of Maloney and Stone⁵ to develop a multidisciplinary team that will review all available data to detect where the system fails and where it over-regulates. Our joint aim is to keep an effective treatment for acne on the market while causing the least restraints but with a high com-

pliance for measures to minimize the risk of congenital anomalies due to isotretinoin.

*Medicines Evaluation Board,
The Hague, the Netherlands

H.J.M.J. CRIJNS*†

S.M. STRAUS*‡

†Department of

L.T.W. DE JONG-VAN DEN BERG†

PharmacoEpidemiology

& PharmacoEconomics,

University of Groningen,

Groningen, the Netherlands

‡Department of Medicinal Informatics,

Erasmus Medical Center,

Rotterdam, the Netherlands

E-mail: ineke.crijns@worldonline.nl

References

- 1 Thiboutot D, Gollnick H, Bettoli V et al. Oral isotretinoin and pregnancy prevention programmes. *Br J Dermatol* 2012; **166**:466–7.
- 2 Crijns HJM, Straus SM, Gispen-de Wied C, De Jong-van den Berg LTW. Compliance with pregnancy prevention programmes of isotretinoin in Europe: a systematic review. *Br J Dermatol* 2011; **164**:238–44.
- 3 European Agency for the Evaluation of Medicinal Products. Committee for Proprietary Medicinal Products (CPMP) Summary Information on a Referral Opinion Following an Arbitration Pursuant to Article 29 of Directive 2001/83/EC, for Isotretinoin/Lurantal/Trivane/Rexidal/Scheritonin. Available at: http://www.ema.europa.eu/docs/en_GB/document_library/Referrals_document/Isotretinoin_29/WC500010882.pdf (last accessed 31 October 2011).
- 4 Shin J, Cheetham TC, Wong L et al. The impact of the iPLEDGE program on isotretinoin fetal exposure in an integrated health care system. *J Am Acad Dermatol* 2011; **65**:1117–25.
- 5 Maloney ME, Stone SP. Isotretinoin and iPLEDGE: a view of results. *J Am Acad Dermatol* 2011; **65**:418–19.

Funding sources: none.

Conflicts of interest: none declared.

News and Notices

DOI: 10.1111/j.1365-2133.2012.10804.x

Annual Evidence Based Update Meeting 2012: Acne and Rosacea

Thursday 10th May 2012: 10am–4.15pm

Location: Holywell Park, Loughborough

Organiser: Professor Hywel Williams

Cost: £90/£100

For further information please contact: Carron Layfield on carron.layfield@nottingham.ac.uk

National Skin Centre Dermatology Update 2012 May 11th–13th 2012

Meritus Mandarin Orchard, Singapore.

For further information contact:

Conference Secretariat Mrs Alice Chew

National Skin Center (S) Pte Ltd

1 Mandalay Road, Singapore 308205

Republic of Singapore

Tel: (65) 63508405

Fax: (65) 62533225

Email: Training@nsc.gov.sg

Applied Photodermatology course 21st–22nd May 2012, BAD House, London