University of Groningen

Independence among physiological traits suggests flexibility in the face of ecological demands on phenotypes

Buehler, D. M.; Vezina, F.; Goymann, W.; Schwabl, I.; Versteegh, M.; Tieleman, B. I.; Piersma, T.

Published in:
Journal of Evolutionary Biology

DOI:
[10.1111/j.1420-9101.2012.02543.x](https://doi.org/10.1111/j.1420-9101.2012.02543.x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Buehler, D. M., Vezina, F., Goymann, W., Schwabl, I., Versteegh, M., Tieleman, B. I., & Piersma, T. (2012). Independence among physiological traits suggests flexibility in the face of ecological demands on phenotypes. *Journal of Evolutionary Biology*, 25(8), 1600-1613. <https://doi.org/10.1111/j.1420-9101.2012.02543.x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Physiological data

Data on physiological traits taken from individual birds (Red Knots, *Calidris canutus*) over an annual cycle. Birds were part of a year long experiment. This file contains experimental and temporal categories as well as physiological measures.

Dataset Key - Buehler et al 2012 J Evol Biol

Bird: Unique identifier for individual birds
Treatment: Experimental treatment group
Cage: Identifier for housing room/cage
DNAsex: Sex obtained via DNA analysis (1 = male, 2 = female)
ExpMonth: Categorical value for month during the experiment
BMRWATT: Basal metabolic rate (in watts)
MCEc: Microbicidal activity against *E.coli* (as proportion of colonies killed after a 10 min incubation)
MCCa: Microbicidal activity against *C. albicans* (as proportion of colonies killed after a 60 min incubation)
MCSa: Microbicidal activity against *S.aureus* (as proportion of colonies killed after a 120 min incubation)
Het: Heterophils per ul blood
Lym: Lymphocytes per ul blood
Mon: Monocytes per ul blood
Agg: log2 agglutination titer
Lys: log2 lysis titer
MASSBMR: Body mass during BMR measurement
Cort: Corticosterone in ng/ml
Hct: Hematocrit (proportion of RBCs in total volume)