University of Groningen

Clinical and Genetic Characterization of Patients with Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Caused by a Plakophilin-2 Splice Mutation

van der Smagt, Jasper J.; van der Zwaag, Paul A.; van Tintelen, J. Peter; Cox, Moniek G. P. J.; Wilde, Arthur A. M.; van Langen, Irene M.; Ummels, Amber; Hennekam, F. A. M.; Dooijes, Dennis; Gerbens, Frans

Published in:
Cardiology

DOI:
[10.1159/000342717](https://doi.org/10.1159/000342717)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Smagt, J. J., van der Zwaag, P. A., van Tintelen, J. P., Cox, M. G. P. J., Wilde, A. A. M., van Langen, I. M., Ummels, A., Hennekam, F. A. M., Dooijes, D., Gerbens, F., Bikker, H., Hauer, R. N. W., & Doevendans, P. A. (2012). Clinical and Genetic Characterization of Patients with Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Caused by a Plakophilin-2 Splice Mutation. *Cardiology*, 123(3), 181-189. <https://doi.org/10.1159/000342717>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Clinical and Genetic Characterization of Patients with Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Caused by a Plakophilin-2 Splice Mutation

Jasper J. van der Smagt^a Paul A. van der Zwaag^d J. Peter van Tintelen^{c, d}
Moniek G.P.J. Cox^b Arthur A.M. Wilde^{c, e} Irene M. van Langen^d
Amber Ummels^a F.A.M. Hennekam^a Dennis Dooijes^a Frans Gerbens^d
Hennie Bikker^f Richard N.W. Hauer^{b, c} Pieter A. Doevendans^{b, c}

Departments of ^aMedical Genetics and ^bCardiology, University Medical Center Utrecht, University of Utrecht, and ^cInteruniversity Cardiology Institute of The Netherlands, Utrecht, ^dDepartment of Genetics, University of Groningen, University Medical Center Groningen, Groningen, and ^eHeart Failure Research Center and ^fDepartment of Clinical Genetics, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

Key Words

Arrhythmia · Cardiomyopathy · Genetics · Desmosome · Arrhythmogenic right ventricular dysplasia/cardiomyopathy

Abstract

Objectives: Arrhythmogenic right ventricular dysplasia/cardiomyopathy (ARVD/C) is characterized by fibrofatty replacement of cardiomyocytes. In around 50% of index patients, a genetic predisposition is demonstrated. The purpose of this study was to examine a plakophilin-2 (*PKP2*) splice site mutation, c.2489+4A>C, identified in 4 separately ascertained Dutch ARVD/C families. **Methods:** Genealogical studies and comprehensive screening of 5 desmosomal genes were undertaken. Reverse transcriptase PCR (RT-PCR) and subsequent sequencing was performed. **Results:** An A-to-C change (c.2489+4A>C) near the splice donor site of intervening sequence 12 of *PKP2* was found in all 4 families.

Based on pedigree data and haplotype sharing, a common ancestor should be situated more than 7 generations ago. RT-PCR demonstrated the presence of aberrant messenger RNA. Clinical manifestations ranged from severe disease to nonpenetrance in elderly mutation carriers. **Conclusions:** This founder mutation in *PKP2* is predicted to lead to the presence of a dysfunctional *PKP2* protein, whereas most truncating mutations are expected to lead to loss of protein. Mutation carriers displayed a wide range of disease severity, suggesting that *PKP2* mutations alone are not sufficient to cause disease, which results in the variable expression and incomplete penetrance characteristic of ARVD/C mutations.

Copyright © 2012 S. Karger AG, Basel

J.J.v.d.S. and P.A.v.d.Z. contributed equally to this article.
F.G. passed away on May 15, 2008.

Introduction

Arrhythmogenic right ventricular dysplasia/cardiomyopathy (ARVD/C) is a disorder characterized by progressive replacement of cardiomyocytes by fibrofatty tissue, primarily in the right ventricle. Life-threatening ventricular arrhythmias originating from the right ventricle can be an early manifestation of the disease. Clinical diagnosis depends on the fulfillment of Task Force Criteria (TFC) [1], for which modifications have recently been proposed [2].

An important genetic contribution is evident from the fact that in 30–50% of cases one or more first-degree relatives also display signs of the disease [3, 4]. A large majority of mutations in ARVD/C patients has been found in genes encoding different components of the cardiac desmosome, i.e. plakophilin-2 (*PKP2*), desmocollin-2 (*DSC2*), desmoglein-2 (*DSG2*), desmoplakin (*DSP*), and plakoglobin (*JUP*) [5–12], suggesting that ARVD/C is primarily a disease of disturbed desmosomal function [13, 14]. Mutations in other genes have also been reported in ARVD/C, i.e. among others, transmembrane protein 43 (*TMEM43*), desmin (*DES*), and titin (*TTN*) [15–18], indicating genetic heterogeneity. In addition, several ARVD/C cases were found to be caused by multiple mutations in the same gene (compound heterozygosity) or mutations in different genes (digenic inheritance), which could result in an earlier onset and increased disease severity [19–21]. The precise mechanisms by which mutations in these genes lead to the distinct ARVD/C phenotype remain, to a large extent, to be elucidated. Decreased desmosome numbers, structurally abnormal desmosomes, secondary effects on other intercalated disk structures, or impaired Wnt/beta-catenin signaling, resulting in increased expression of adipogenic and fibrogenic genes and fat accumulation, may all contribute to the ARVD/C pathophysiology [22–24]. However, regardless of the gene that is mutated, translocation of plakoglobin from the desmosome to the cytosol may be an important common step in disease causation [22, 24, 25].

It has been shown that mutations in *PKP2* are frequent in ARVD/C patients in the northwestern part of Europe and the USA [5, 7, 26, 27]. In a cohort of Dutch ARVD/C patients up to 90% of *PKP2* mutations were found in cases of proven familial disease [21]. Eight different recurrent *PKP2* mutations have been found in the Dutch population, accounting for 84% (41/49) of separately ascertained *PKP2*-positive ARVD/C families. For some of them a shared *PKP2* haplotype was demonstrated, while other families could be traced back to a common ancestor

by genealogy, sometimes as far as 10 generations ago [7]. Therefore, it is likely that several different *PKP2* founder mutations segregate in The Netherlands, which may contribute to the high prevalence of *PKP2* mutations in Dutch ARVD/C patients.

Looking at the spectrum of different mutations reported in *PKP2*, it is evident that truncating mutations (nonsense and frameshift) are most frequent [28]. The classification of missense variants as pathogenic, in *PKP2* as well as in other genes, remains challenging without solid functional evidence [29]. Truncating mutations are predicted to lead to loss of *PKP2* protein by nonsense-mediated messenger RNA decay (NMD), although mRNA studies by Gerull et al. [5] showed that the identified splice mutations had detectable mRNA with frameshifts, resulting in premature termination codons, that may evade NMD. As a general rule, introduction of a stop codon more than 50–55 nucleotides upstream of the last exon-intron boundary of any gene will lead to the breakdown of messenger-RNA in vivo, resulting in only very little translation of the abnormal truncated protein [30]. This is an argument for haploinsufficiency (insufficient amounts of normal *PKP2*) as the predominant mechanism by which *PKP2* mutations predispose to disease, as opposed to a dominant negative effect due to the presence of abnormal *PKP2* protein. Indeed, a Western blot from a heart biopsy from a patient with *PKP2* mutation c.2076_2077delAA (p.Ser693CysfsX49) did not show detectable protein [5].

This report discusses both the clinical characteristics of patients from 4 different families that were found to have an identical splice mutation (c.2489+4A>C) at the C-terminal end of the *PKP2* gene, and the possible mechanism by which it exerts its effect.

Methods

Clinical Evaluation

All index patients and their relatives were evaluated at the Cardiology Departments of the University Medical Centers in Utrecht and Amsterdam. Family members were evaluated following a genetic counseling procedure and after informed consent had been obtained. The evaluation consisted of at least a physical examination, 12-lead ECG, and echocardiography. Diagnosis was based on the recently modified ARVD/C TFC [2].

Genealogy

Since 1997, genealogical studies have been performed in newly ascertained ARVD/C cases, in order to select patients that could contribute to the finding of new ARVD/C loci, using identity by descent strategies [31]. Investigations were carried out using community registries, starting from information on the grandparents as supplied by the index patients.

Fig. 1. Pedigrees of families 1–4. Squares denote males, circles females. Black squares/circles denote individuals that satisfy the 2010 TFC for a clinical diagnosis of ARVD/C (≥ 4 points). Upper right quadrant-blackened squares/circles denote individuals that did not fulfill the TFC but did have abnormalities compatible with

ARVD/C at their most recent cardiac examination. +/– indicates the presence or absence of the *PKP2* c.2489+4A>C mutation. Individuals denoted with ‘n’ had a normal cardiac evaluation. Numbers in squares/circles denote the number of individuals. * Genotype used for the haplotype sharing test, using SNP array data.

Molecular Genetic Analyses

DNA was isolated from peripheral lymphocytes according to standard protocols. In index patients the coding regions of the *PKP2*, *DSC2*, *DSG2*, *DSP*, *JUP*, and *TMEM43* genes were analyzed using denaturing high-performance liquid chromatography (*PKP2*, *DSC2*, and *DSG2*) as described previously [32], and/or direct sequencing (*DSP*, *JUP*, and *TMEM43*). Direct sequencing was performed with a BigDye Terminator DNA sequencing kit (version 2.0) on a 3730 automated sequencer (Applied Biosystems, Foster City, Calif., USA). All sequences were analyzed using SeqScape software (version 2.1.1, Applied Biosystems). In addition, multiplex ligation-dependent probe amplification analysis was performed to exclude large deletions in *PKP2* (SALSA multiplex ligation-dependent probe amplification kit P168 ARVC-*PKP2*; MRC Holland, Amsterdam, The Netherlands). Relatives of index patients were only analyzed for the presence of a previously identified mutation.

Total RNA was isolated from fresh blood samples from individual VII:6 and a control individual using a PAXgene kit (QIAGEN Benelux, Venlo, The Netherlands) and subjected to random hexamer primed reverse transcriptase PCR (RT-PCR).

The obtained cDNA products were amplified by PCR with primers specific for the *PKP2* coding sequence. *PKP2* cDNA fragments were separated according to size using 2% agarose gel electrophoresis along with a 100-bp DNA ladder (O’GeneRuler; Fermentas, Burlington, Ont., Canada). Both normal and aberrant fragments were gel purified and used for direct sequencing. Primers and PCR details are available upon request.

Results

Genealogy

In families 2 and 3 (fig. 1), common ancestors were found who were born around 1780 AD. Family 1 could be linked to family 2, but a link with the common ancestors in families 2 and 3 could not be established. Family 4 could not be traced back to the same ancestors, yet all ancestors lived in the same region.

Fig. 2. RT-PCR for the *PKP2* c.2489+4A>C mutation. **a** Two percent agarose gel electrophoresis showing cDNA products derived from mRNA of a control sample and index patient VII:6. The first lane shows a 100-bp ladder, followed by the control, the patient sample, and a blank with primers in exons 11 and 13. The expected normal cDNA product with primers in exons 11 and 13 is 335 bp in length. The shorter transcript in the patient sample (lane 3)

lacks exon 12 (190 bp) as shown in **b**. The bands in the patient sample show an ~1:1 ratio, indicating that preferred PCR of the shorter transcript, lacking exon 12, did not occur. This could indicate partial degradation of the mutated mRNA. **b** Sequencing of the cDNA derived from both the normally spliced transcript and the abnormally spliced transcript demonstrating the skipping of exon 12.

Table 1. Results of splice prediction programs

Splice prediction program	Wild type	c.2489+4A>C mutation	Notes
NetGene2	0.93	0.91	both ‘highly confident donor sites’ donor score cutoff: 0.40
NNSPLICE 0.9	0.98	0.57	
<i>Using Alamut mutation interpretation software</i>			
SpliceSiteFinder-Like	73.60	62.81	
MaxEntScan	9.73	6.22	
GeneSplicer	2.54	1.02	

Results of the different prediction programs for the normal exon 12 donor splice site are shown. Splice predictions are available from the following websites: NetGene2: www.cbs.dtu.dk/services/NetGene2, NNSPLICE 0.9: www.fruitfly.org/seq_tools/splice.html, MaxEntScan: genes.mit.edu/burgelab/maxent/Xmaxentscan_scoreseq.html, GeneSplicer: www.cbc.umd.edu/software/GeneSplicer, and Alamut: www.interactive-biosoftware.com.

Molecular Genetic Analyses

Analysis of the entire coding regions of *PKP2*, *DSC2*, *DSG2*, *DSP*, *JUP*, and *TMEM43* including exon-intron boundaries in the 4 index patients revealed a c.2489+4A>C mutation in the *PKP2* gene. Besides this intervening sequence mutation, no other mutations, including large deletions in *PKP2*, were found. The *PKP2* c.2489+4A>C mutation is within the splice donor consensus sequence. Splice prediction programs were inconclusive with respect to a predicted effect on splicing (table 1). The mutation was absent in over 150 ethnically matched controls (300 alleles) and not annotated in both dbSNP and the 1000 Genomes Project databases (ac-

cessed October 10, 2011). In addition to the 4 index patients, 10 relatives were also found to carry the *PKP2* c.2489+4A>C mutation.

RT-PCR demonstrated the presence of an abnormal *PKP2* messenger RNA (fig. 2a), indicating that the effect of the c.2489+4A>C mutation is indeed through aberrant splicing. The abnormal *PKP2* messenger RNA lacks the entire exon 12 (190 bp) (fig. 2b). In the predicted protein a frameshift occurs that leads to stop codon 99 residues downstream, well within the 3' untranslated region of the *PKP2* gene. Therefore, all residues normally encoded by exons 12, 13, and 14 are missing from the abnormal transcript, but the predicted abnormal protein is only

Fig. 3. Aberrant splicing as a result of the *PKP2* c.2489+4A>C mutation. A and B represent the C-terminus of the normal plakophilin-2 protein and the normally spliced mRNA, respectively. C depicts part of the genomic DNA of the *PKP2* gene [exons in white, introns (IVS) in grey, 3' untranslated region (UTR) in

black] with the position of the wild-type splice site above the bar and the mutated splice site below the bar. D and E show the aberrant splice product and the predicted abnormal protein, respectively. The abnormal protein is only 15 amino acid residues shorter than the wild-type protein.

15 residues shorter than the wild-type protein (fig. 3). Since the stop codon occurs in the 3' untranslated region, the abnormal protein cannot be subjected to NMD.

Clinical Evaluation

All index patients with the *PKP2* c.2489+4A>C mutation fulfilled the modified TFC for ARVD/C [2]. Figure 1 shows the pedigree of all families, and clinical data on the *PKP2* c.2489+4A>C mutation carriers are summarized in table 2. Index patient 1 (VI:1) was admitted to the hospital at the age of 33 years with sustained monomorphic ventricular tachycardia (VT) (240/min) with left bundle branch block (LBBB) morphology during exercise. He experienced at least 4 other episodes of VT, but no syncope. Supraventricular tachycardias also occurred, caused by AV nodal re-entry and provoked by premature ventricular complexes. Radiofrequency (RF) ablation was performed for the ventricular arrhythmias from the right ventricular outflow tract and for the AV nodal re-entry tachycardia (AVNRT). He has been doing well on disopyramide medication for 12 years. Index patient 2 (VI:5) has been known to have frequent VT episodes with LBBB morphology since the age of 20, which were treated with RF ablation. At the age of 36 she had an out-of-hospital

cardiac arrest (OHCA) due to ventricular fibrillation (VF) while on sotalol. Subsequently an implantable cardioverter defibrillator (ICD) was implanted, which has delivered appropriate anti-tachycardia pacing and shocks. Index patient 3 (VII:6) experienced syncope at the age of 26 during exercise. She was admitted with a wide QRS complex tachycardia (250/min). Treatment with disopyramide was started. An AVNRT was diagnosed at electrophysiological study, which was treated with RF ablation, after a second arrhythmia episode. She has been free of symptoms for 14 years. Index patient 4 (VI:10) had an OHCA with VF at the age of 29. He received an ICD and was treated with metoprolol for VTs. At the age of 33 he experienced two episodes with electrical storms possibly triggered by a viral infection. Metoprolol was changed to sotalol and amiodarone was added. The patient has been doing well for 9 years.

In addition to the 4 index patients, 2 of the mutation-positive family members developed clinical complaints likely related to ARVD/C, resulting in 6 symptomatic mutation carriers. In family 4, a 34-year-old male (VI:9), who had had no complaints at the initial evaluation, received an ICD after he suffered a near syncope while already taking sotalol. Nonsustained polymorphic VTs of

Table 2. Clinical data on *PKP2* c.2489+4A>C mutation carriers

Individual (fig. 1)	Sex	Onset age years	Age at first evalua- tion years	Follow- up years	Events	Modified task force criteria for the diagnosis of ARVD/C								Treatment	Other
						func- tion	tissue ^a	repolar- ization	depolar- ization	arrhyth- mias	family history	task force criteria ^b	clinical status		
Family 1															
VI:1 ^c	M	33	33	12	VT	+				++	++	5	Aff	RF ablation, disopyramide	atypical AVNRT, no progression
V:2	F		61	0	none						++	2	Pos	none	no follow-up
V:4	F		69	7	none			++		++	++	6	Aff	anticoagulant therapy	paroxysmal AF, no progression
VI:3	F		35	7	none				+		++	3	Bor	none	no progression
Family 2															
VI:5	F	20	20	32	VT/VF/ OHCA	++		++	++	++	++	10	Aff	RF ablation, ICD, sotalol	appropriate ICD~, good LV function
VII:3	M		17	11	none	++			+		++	5	Aff	none	mild progression
VII:4	F		15	11	none						++	2	Pos	none	no progression
V:7	F														not evaluated
Family 3															
VII:6 ^d	F	26	26	14	syncope	+		++	+	++	++	8	Aff	RF ablation	concurrent AVNRT, no progression
VI:7	F		62	3	none						++	2	Pos	none	no progression
Family 4															
VI:10	M	29	29	9	VF/ OHCA	++		++		++	++	8	Aff	ICD, sotalol, amiodarone	appropriate ICD~, mild progression
V:11	F		59	9	none						++	2	Pos	none	mild progression
VI:8	F	40	33	9	near syncope	++					++	4	Aff	ICD	evident progression
VI:9	M	34	32	9	near syncope					++	++	4	Aff	ICD, sotalol	mild progression

AF = Atrial fibrillation; Aff = affected; Bor = borderline; F = female; ICD~ = ICD discharge; LV = left ventricle; M = male; NA = not available; nsVT = nonsustained ventricular tachycardia; Pos = possibly; RV = right ventricle; sVT = sustained ventricular tachycardia.

^a Not available. ^b Minor criteria yield 1 point and major criteria 2 points; for a definite diagnosis of ARVD/C a minimum of 4 points is required, 3 points indicate a borderline diagnosis, and 2 points indicate a possible diagnosis. ^c For this patient a biopsy was available; it showed fibrosis but no fat and was considered inconclusive. ^d Syncope probably resulting from AVNRT; thus far no symptoms have recurred after RF ablation of AVNRT.

insufficient duration to warrant ICD therapy were subsequently recorded. His sister (VI:8) received an ICD shortly after she experienced a near syncope during a stressful event. Although without any signs of disease at initial evaluation at the age of 33, she developed aneurysmatic changes of the right ventricle in the course of 9 years.

Remarkably, 7 of the 13 mutation carriers (age range at first evaluation 15–69 years) who were evaluated were asymptomatic. In 4 of them only a major criterion for a positive family history was available, leading to a possible clinical status according to the modified TFC. In 3 of the asymptomatic carriers additional criteria were present, resulting in a borderline diagnosis in 1 and a definite diagnosis in 2 asymptomatic carriers [2].

Left ventricular (LV) involvement was not systematically assessed in these families. Cardiac MRI was performed in 5 mutation carriers; all of them showed normal LV volumes and LV ejection fraction. None of them

showed LV wall motion abnormalities or late gadolinium enhancement. Index patient VI:5 experienced VT with RBBB morphology, but she had a normal LV function. Finally, none of the mutation carriers had isolated inverted T-waves in the left precordial leads V4–V6.

Discussion

PKP2 mutations are a frequent cause of ARVD/C. The *PKP2* gene was first implicated in ARVD/C through a candidate gene approach [5]. The fact that the *PKP2* locus had not been previously found in linkage studies can be explained by low penetrance resulting in phenotype-negative family members who were classified as unaffected, although they were in fact mutation carriers. Previously, the haplotype sharing test identified a shared haplotype on chromosome 12p12.1–12q13.13, encompassing *PKP2*,

as the largest shared haplotype in these families [33]. The haplotype sharing test compares the genotypes from SNP arrays, using an affected-only strategy, to search for large shared haplotypes within pedigrees most likely to contain a causative mutation. In these families, the haplotype sharing test indicated a common founder, who must have lived well before the 19th century. Based on pedigree data, it was estimated that the most recent common ancestor should be situated over 7 generations ago. However, these are crude estimations since only 4 families were available. The actual age of the mutation could be considerably older. Little is known about the age of *PKP2* mutations. Our data show that mutations may have occurred many generations earlier, in the absence of a family history of cardiac disease. The fact that in some families we have been able to demonstrate common ancestors as far as 10 generations back by genealogy suggests that reproductive fitness in ARVD/C is not significantly reduced.

The *PKP2* c.2489+4A>C mutation seems to be relatively frequent in The Netherlands, having been detected in 4 independently ascertained ARVD/C families so far. Another 5 mutations (c.235C>T, c.397C>T, c.1211-1212dupT, c.2386T>C, and c.2489+1G>A) in *PKP2* occurred at least as frequently in the Dutch ARVD/C cohort. Haplotype analysis and genealogy suggest that there may also be common founders contributing to these mutations, explaining the high prevalence of *PKP2* mutations in Dutch ARVD/C patients [32, 34].

The frequent occurrence of nonsense and frameshift mutations in *PKP2* is indicative that the causative mechanism at the protein level is that of loss of normal *PKP2* function, as most of these mutations are predicted to lead to NMD [28]. Gerull et al. [5], while performing Western blot analysis on cardiac tissue in a patient with a *PKP2* c.2076_2077delAA mutation, demonstrated reduced wild-type *PKP2* but were unable to demonstrate the predicted abnormal protein. This is in keeping with the concept of NMD in the heart. As the *PKP2* c.2489+4A>C mutation cannot be subjected to NMD, it is predicted to lead to a dysfunctional *PKP2* protein. However, we cannot exclude degradation of the abnormal protein through other mechanisms, which could still lead to haploinsufficiency. If a dysfunctional protein is present, the last two conserved armadillo repeat regions would be disrupted, completely altering the C-terminal tail. The exact role of the C-terminal tail is unknown, while the N-terminal part of the protein is most important for binding other desmosomal proteins and targeting *PKP2* to the plasma membrane [35]. The pathogenic effect of splice mutations

may be ameliorated when the induced aberrant splicing is not absolute, as has been shown for the only recessive *PKP2* mutation thus far [36]. In the absence of heterozygous polymorphisms in exons 12–14 of the *PKP2* gene, the presence of a normal transcript from the diseased allele could not be assessed.

In total 14 patients with the *PKP2* c.2489+4A>C mutation (4 males, 10 females), including index patients, were identified in the 4 families. Despite the overrepresentation of female mutation carriers in this study, 2 out of 4 index patients were males. Nine out of 10 family members with a mutation had a cardiac workup. Only 2 of 9 evaluated mutation-positive family members developed clinical complaints (table 2). Five family members fulfilled the modified TFC for ARVD/C. Incomplete penetrance and clinical variability are characteristic for ARVD/C [4] and were also seen in the families with the *PKP2* c.2489+4A>C mutation. Compound heterozygosity or digenic inheritance for mutations in the desmosomal genes or in *TMEM43* had been excluded as an explanation for this variability, but other genetic, epigenetic, or nongenetic factors, such as lifestyle, could have contributed to the clinical differences observed in these families.

Which preventive treatment modalities are justified in asymptomatic *PKP2* mutation carriers is currently undetermined. We suggest that they be followed-up regularly in order to assess disease progression with annual echocardiography, Holter monitoring, and exercise testing. Besides, additional cardiac MRI at intervals no longer than 5 years (or before implantation of an ICD) is deemed appropriate in The Netherlands.

Although we find molecular diagnosis in ARVD/C to be a useful tool in identifying persons at risk for developing ARVD/C, its value is still limited by the inability to accurately predict phenotypes from genotypes, especially when a missense variant is identified [29]. The fact that many identified *PKP2* mutation carriers will never experience life-threatening arrhythmias emphasizes the need for identifying the truly pathogenic mutations as well as other genetic or environmental factors that play a role in disease causation. We believe that the described *PKP2* c.2489+4A>C mutation is pathogenic, based on its cosegregation with the ARVD/C phenotype, the absence of the mutation in controls as well as in the dbSNP, and the 1000 Genomes Project databases, the fact that the mutation is located on the largest shared haplotype, indicating a founder effect [33], and the RT-PCR which demonstrated the presence of an abnormal *PKP2* messenger RNA (fig. 2a). As is the case for many other desmosomal muta-

tions implicated in ARVD/C, the c.2489+4A>C mutation is characterized by incomplete penetrance, indicating that not all mutation carriers will develop the disease. The mutation is predicted to lead to the presence of a dysfunctional PKP2 protein, whereas most truncating mutations are expected to lead to loss of protein. Mutation carriers displayed a wide range of disease severity, suggesting that *PKP2* mutations alone are not sufficient to cause disease.

Acknowledgements

We sincerely thank the families for their cooperation. Furthermore, we thank Dr. C. de Kovel for helping to estimate the distance to the most recent common founder, Marcel R. Nelen, Mark M. Entius, and Richard A. Zewald for their assistance with the molecular genetic analyses, Jackie Senior for editing the manuscript, and Dr. E.G. Weijers, cardiologist at Groene Hart Ziekenhuis, Gouda, The Netherlands, for referring index patient VII:6.

This work was supported by a grant from the Netherlands Heart Foundation (2007B132).

References

- McKenna WJ, Thiene G, Nava A, Fontaliran F, Blomstrom-Lundqvist C, Fontaine G, Camerini F: Diagnosis of arrhythmogenic right ventricular dysplasia/cardiomyopathy – Task Force of the Working Group Myocardial and Pericardial Disease of the European Society of Cardiology and of the Scientific Council on Cardiomyopathies of the International Society and Federation of Cardiology. *Br Heart J* 1994;71:215–218.
- Marcus FI, McKenna WJ, Sherrill D, Basso C, Bauce B, Bluemke DA, Calkins H, Corrado D, Cox MG, Daubert JP, Fontaine G, Gear K, Hauer R, Nava A, Picard MH, Protonotarios N, Saffitz JE, Sanborn DM, Steinberg JS, Tandri H, Thiene G, Towbin JA, Tsatsopoulou A, Wichter T, Zareba W: Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria. *Circulation* 2010;121:1533–1541.
- Marcus FI, Fontaine GH, Guiraudon G, Frank R, Laurenceau JL, Malergue C, Grossogeat Y: Right ventricular dysplasia: a report of 24 adult cases. *Circulation* 1982;65:384–398.
- Basso C, Corrado D, Marcus FI, Nava A, Thiene G: Arrhythmogenic right ventricular cardiomyopathy. *Lancet* 2009;373:1289–1300.
- Gerull B, Heuser A, Wichter T, Paul M, Basson CT, McDermott DA, Lerman BB, Markowitz SM, Ellinor PT, MacRae CA, Peters S, Grossmann KS, Drenckhahn J, Michely B, Sasse-Klaassen S, Birchmeier W, Dietz R, Breithardt G, Schulze-Bahr E, Thierfelder L: Mutations in the desmosomal protein plakophilin-2 are common in arrhythmogenic right ventricular cardiomyopathy. *Nat Genet* 2004;36:1162–1164.
- Dalal D, Molin LH, Piccini J, Tichnell C, James C, Bomma C, Prakasa K, Towbin JA, Marcus FI, Spevak PJ, Bluemke DA, Abraham T, Russell SD, Calkins H, Judge DP: Clinical features of arrhythmogenic right ventricular dysplasia/cardiomyopathy associated with mutations in plakophilin-2. *Circulation* 2006;113:1641–1649.
- van Tintelen JP, Entius MM, Bhuiyan ZA, Jongbloed R, Wiesfeld AC, Wilde AA, van der Smagt J, Boven LG, Mannens MM, van Langen IM, Hofstra RM, Otterspoor LC, Doevendans PA, Rodriguez LM, van Gelder IC, Hauer RN: Plakophilin-2 mutations are the major determinant of familial arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circulation* 2006;113:1650–1658.
- Syrris P, Ward D, Evans A, Asimaki A, Gandjbakhch E, Sen-Chowdhry S, McKenna WJ: Arrhythmogenic right ventricular dysplasia/cardiomyopathy associated with mutations in the desmosomal gene desmocollin-2. *Am J Hum Genet* 2006;79:978–984.
- Heuser A, Plovie ER, Ellinor PT, Grossmann KS, Shin JT, Wichter T, Basson CT, Lerman BB, Sasse-Klaassen S, Thierfelder L, MacRae CA, Gerull B: Mutant desmocollin-2 causes arrhythmogenic right ventricular cardiomyopathy. *Am J Hum Genet* 2006;79:1081–1088.
- Pilichou K, Nava A, Basso C, Beffagna G, Bauce B, Lorenzon A, Frigo G, Vettori A, Valente M, Towbin J, Thiene G, Danieli GA, Rampazzo A: Mutations in desmoglein-2 gene are associated with arrhythmogenic right ventricular cardiomyopathy. *Circulation* 2006;113:1171–1179.
- Rampazzo A, Nava A, Malacrida S, Beffagna G, Bauce B, Rossi V, Zimbello R, Simonati B, Basso C, Thiene G, Towbin JA, Danieli GA: Mutation in human desmoplakin domain binding to plakoglobin causes a dominant form of arrhythmogenic right ventricular cardiomyopathy. *Am J Hum Genet* 2002;71:1200–1206.
- Asimaki A, Syrris P, Wichter T, Matthias P, Saffitz JE, McKenna WJ: A novel dominant mutation in plakoglobin causes arrhythmogenic right ventricular cardiomyopathy. *Am J Hum Genet* 2007;81:964–973.
- Tsatsopoulou AA, Protonotarios NI, McKenna WJ: Arrhythmogenic right ventricular dysplasia, a cell adhesion cardiomyopathy: insights into disease pathogenesis from preliminary genotype-phenotype assessment. *Heart* 2006;92:1720–1723.
- van Tintelen JP, Hofstra RM, Wiesfeld AC, van den Berg MP, Hauer RN, Jongbloed JD: Molecular genetics of arrhythmogenic right ventricular cardiomyopathy: emerging horizon? *Curr Opin Cardiol* 2007;22:185–192.
- Merner ND, Hodgkinson KA, Haywood AF, Connors S, French VM, Drenckhahn JD, Kupprion C, Ramadanova K, Thierfelder L, McKenna W, Gallagher B, Morris-Larkin L, Bassett AS, Parfrey PS, Young TL: Arrhythmogenic right ventricular cardiomyopathy type 5 is a fully penetrant, lethal arrhythmic disorder caused by a missense mutation in the TMEM43 gene. *Am J Hum Genet* 2008;82:809–821.
- van Tintelen JP, van Gelder IC, Asimaki A, Suurmeijer AJ, Wiesfeld AC, Jongbloed JD, van den Wijngaard A, Kuks JB, Spaendonck-Zwarts KY, Notermans N, Boven L, van den Heuvel F, Veenstra-Knol HE, Saffitz JE, Hofstra RM, van den Berg MP: Severe cardiac phenotype with right ventricular predominance in a large cohort of patients with a single missense mutation in the DES gene. *Heart Rhythm* 2009;6:1574–1583.
- Klauke B, Kossmann S, Gaertner A, Brand K, Stork I, Brodehl A, Dieding M, Walhorn V, Anselmetti D, Gerdes D, Bohms B, Schulz U, Zu Knyphausen E, Vorgerd M, Gummert J, Milting H: De novo desmin-mutation N116S is associated with arrhythmogenic right ventricular cardiomyopathy. *Hum Mol Genet* 2010;19:4595–607.
- Taylor M, Graw S, Sinagra G, Barnes C, Slavov D, Brun F, Pinamonti B, Salcedo EE, Sauer W, Pyxaras S, Anderson B, Simon B, Bogomolovas J, Labeit S, Granzier H, Messtroni L: Genetic variation in titin in arrhythmogenic right ventricular cardiomyopathy-overlap syndromes. *Circulation* 2011;124:876–885.
- Xu T, Yang Z, Vatta M, Rampazzo A, Beffagna G, Pillichou K, Scherer S, Saffitz J, Kravitz J, Zareba W, Danieli G, Lorenzon A, Nava A, Bauce B, Thiene G, Basso C, Calkins H, Gear K, Marcus F, Towbin J: Compound and digenic heterozygosity contributes to arrhythmogenic right ventricular cardiomyopathy. *J Am Coll Cardiol* 2010;55:587–597.

- 20 Fressart V, Duthoit G, Donal E, Probst V, Deharo JC, Chevalier P, Klug D, Dubourg O, Delacretaz E, Cosnay P, Scanu P, Extramiana F, Keller D, Hidden-Lucet F, Simon F, Bessirard V, Roux-Buisson N, Hebert JL, Azarine A, Casset-Senon D, Rouzet F, Lecarpentier Y, Fontaine G, Coirault C, Frank R, Hainque B, Charron P: Desmosomal gene analysis in arrhythmogenic right ventricular dysplasia/cardiomyopathy: spectrum of mutations and clinical impact in practice. *Europace* 2010; 12:861–868.
- 21 Cox MG, van der Zwaag PA, van der Werf C, van der Smagt JJ, Noorman M, Bhuiyan ZA, Wiesfeld AC, Volders PG, van Langen IM, Atsma DE, Dooijes D, van den Wijngaard A, Houweling AC, Jongbloed JD, Jordaens L, Cramer MJ, Doevendans PA, de Bakker JM, Wilde AA, van Tintelen JP, Hauer RN: Arrhythmogenic right ventricular dysplasia/cardiomyopathy: pathogenic desmosome mutations in index-patients predict outcome of family screening – Dutch Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Genotype-Phenotype Follow-Up Study. *Circulation* 2011;123:2690–2700.
- 22 Kaplan SR, Gard JJ, Protonotarios N, Tsatsopoulou A, Spiliopoulou C, Anastasakis A, Squarcioni CP, McKenna WJ, Thiene G, Basso C, Brousse N, Fontaine G, Saffitz JE: Remodeling of myocyte gap junctions in arrhythmogenic right ventricular cardiomyopathy due to a deletion in plakoglobin (Naxos disease). *Heart Rhythm* 2004;1:3–11.
- 23 Basso C, Czarnowska E, Della Barbera M, Bauce B, Beffagna G, Wlodarska EK, Pilichou K, Ramondo A, Lorenzon A, Wozniak O, Corrado D, Daliento L, Danieli GA, Valente M, Nava A, Thiene G, Rampazzo A: Ultrastructural evidence of intercalated disc remodelling in arrhythmogenic right ventricular cardiomyopathy: an electron microscopy investigation on endomyocardial biopsies. *Eur Heart J* 2006;27:1847–1854.
- 24 Garcia-Gras E, Lombardi R, Giocondo MJ, Willerson JT, Schneider MD, Khoury DS, Marian AJ: Suppression of canonical Wnt/beta-catenin signaling by nuclear plakoglobin recapitulates phenotype of arrhythmogenic right ventricular cardiomyopathy. *J Clin Invest* 2006;116:2012–2021.
- 25 Asimaki A, Tandri H, Huang H, Halushka MK, Gautam S, Basso C, Thiene G, Tsatsopoulou A, Protonotarios N, McKenna WJ, Calkins H, Saffitz JE: A new diagnostic test for arrhythmogenic right ventricular cardiomyopathy. *N Engl J Med* 2009;360:1075–1084.
- 26 Syrris P, Ward D, Asimaki A, Sen-Chowdhry S, Ebrahim HY, Evans A, Hitomi N, Norman M, Pantazis A, Shaw AL, Elliott PM, McKenna WJ: Clinical expression of plakophilin-2 mutations in familial arrhythmogenic right ventricular cardiomyopathy. *Circulation* 2006;113:356–364.
- 27 den Haan AD, Tan B, Zikusoka M, Ibanez Llado L, Jain R, Daly A, Tichnell C, James C, Amat-Alarcon N, Abraham T, Russell S, Bluemke D, Calkins H, Dalal D, Judge D: Comprehensive desmosome mutation analysis in North Americans with arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Cardiovasc Genet* 2009;2:428–435.
- 28 van der Zwaag PA, Jongbloed JD, van den Berg MP, van der Smagt JJ, Jongbloed R, Bikker H, Hofstra RM, van Tintelen JP: A genetic variants database for arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Hum Mutat* 2009;30:1278–1283.
- 29 Christensen AH, Benn M, Tybjaerg-Hansen A, Haunso S, Svendsen JH: Missense variants in plakophilin-2 in arrhythmogenic right ventricular cardiomyopathy patients – disease-causing or innocent bystanders? *Cardiology* 2010;115:148–154.
- 30 Lykke-Andersen J: mRNA quality control: marking the message for life or death. *Curr Biol* 2001;11:R88–R91.
- 31 Houwen RH, Baharloo S, Blankenship K, Raeymaekers P, Juyn J, Sandkuijl LA, Freimer NB: Genome screening by searching for shared segments: mapping a gene for benign recurrent intrahepatic cholestasis. *Nat Genet* 1994;8:380–386.
- 32 Bhuiyan ZA, Jongbloed JD, van der Smagt J, Lombardi PM, Wiesfeld AC, Nelen M, Schouten M, Jongbloed R, Cox MG, van Wolferen M, Rodriguez LM, van Gelder IC, Bikker H, Suurmeijer AJ, van den Berg MP, Mannens MM, Hauer RN, Wilde AA, van Tintelen JP: Desmoglein-2 and desmocollin-2 mutations in Dutch arrhythmogenic right ventricular dysplasia/cardiomyopathy patients: results from a multicenter study. *Circ Cardiovasc Genet* 2009;2:418–427.
- 33 van der Zwaag PA, van Tintelen JP, Gerbens F, Jongbloed JD, Boven LG, van der Smagt JJ, van der Roest WP, van Langen IM, Bikker H, Hauer RN, van den Berg MP, Hofstra RM, te Meerman GJ: Haplotype sharing test maps genes for familial cardiomyopathies. *Clin Genet* 2011;79:459–467.
- 34 van der Zwaag PA, Cox MG, van der Werf C, Wiesfeld AC, Jongbloed JD, Dooijes D, Bikker H, Jongbloed R, Suurmeijer AJ, van den Berg MP, Hofstra RM, Hauer RN, Wilde AA, van Tintelen JP: Recurrent and founder mutations in The Netherlands: plakophilin-2 pArg79X mutation causing arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Neth Heart J* 2010;18:583–591.
- 35 Chen X, Bonne S, Hatzfeld M, van Roy F, Green KJ: Protein binding and functional characterization of plakophilin 2: evidence for its diverse roles in desmosomes and beta-catenin signaling. *J Biol Chem* 2002;277:10512–10522.
- 36 Awad MM, Dalal D, Tichnell C, James C, Tucker A, Abraham T, Spevak PJ, Calkins H, Judge DP: Recessive arrhythmogenic right ventricular dysplasia due to novel cryptic splice mutation in PKP2. *Hum Mutat* 2006; 27:1157.