University of Groningen

Food web assembly at the landscape scale

Schrama, Maarten; Jouta, Jeltje; Berg, Matty P.; Olff, Han

Published in:
Ecosystems

DOI:
[10.1007/s10021-013-9636-5](https://doi.org/10.1007/s10021-013-9636-5)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Schrama, M., Jouta, J., Berg, M. P., & Olff, H. (2013). Food web assembly at the landscape scale: Using stable isotopes to reveal changes in trophic structure during succession. *Ecosystems*, 16(4), 627-638. <https://doi.org/10.1007/s10021-013-9636-5>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Electronic appendix. **Food Web Assembly at the Landscape Scale: Using Stable Isotopes to Reveal Changes in Trophic Structure during Succession**

Table 1. General Characteristics for Each of the Sampling Sites

Age succ. stage (y)	Elevation (cm above NAP ±SE)		Flooding freq. (yearly average 1998-2008 ±SE)*		Sediment layer thickness (cm ±SE)		Vegetation height (cm ±SE)	
0	115.2	2.3	184.2	5.8	0.0	0.0	1.2	0.3
10	116.2	2.0	178.3	5.9	4.9	0.3	5.9	0.7
25	116.0	3.2	178.3	5.9	7.2	0.2	10.2	1.3
35	118.4	1.9	164.8	6.5	13.5	0.6	8.1	1.7
45	123.4	1.9	136.9	6.5	15.3	0.2	9.4	0.8
55	125.0	2.1	127.7	7.0	14.4	0.3	19.7	3.8
100	124.4	1.5	131.8	6.8	16.2	0.8	26.1	2.0

* Flooding data were taken from an online archive with freely available measurements, which can be downloaded from http://live.waterbase.nl/waterbase_wns.cfm?taal=en

Appendix 1: Photo's from Each of the Sites, July 2008

Site 1: 0 yrs of succession

Site 2: 10 yrs of succession

Site 3: 25 yrs of succession

Site 4: 35 yrs of succession

Site 5: 45 yrs of succession

Site 6: 55 yrs of succession

Site 7: 100 yrs of succession

Table 1. The Amount of Subsamples per Site for All the Species Collected

Group	Species	0	10	25	35	45	55	100
Primary production								
Terrestrial plants	Artemisia maritima					5		5
	Atriplex portulacoides		5			5		5
	Elytrigia atherica							5
	Festuca rubra		5			5		5
	Limonium vulgare	5	5			5		
	Puccinellia maritima	5	5			5		
	Salicornia europaea	5	5					
	Soil organic matter		2			5		5
	Terrestrial organic matter		5			5		6
Marine prim. prod	Bacillariophyceae (Diatoms)	5	5			5		4
	Particulate organic matter		5			4		3
	Fucus vesiculosus	5	4			5		4
Herbivores								
	Bledius sp.	5	1					
	Cassida vittata	4	4	3	4			
	Auchenorrhyncha spp.							1
	Chrysomelidae sp.		1	1	1	3	1	1
	Elateridae sp.	1	1	1			2	
	Curculionidea sp.		2	1	3	2	4	
Carnivores								
	Bembidion minimum		3	2	3	5	2	5
	Clubiona stagnatilis	5	1	2	1	2	5	1
	Coccinella sedecumpunctata	5		3		2		1
	Dyscherius globus		3	5	4	2		2
	Ichneumonoidea spp.	1	1	1				1
	Erigonidae spp.	4	5	3	5	2	3	3
	Pardosa pubeckensis	5	4	5	4	5	5	4
	Pogonus chalceus			2		1		
	Salda littoralis		1	4	4	3	3	1
	Tytthaspis sedecumpunctata		2		2	4	5	1
Detritivores								
	Helophorus brevipalpis							4
	Isotoma riparia		1				2	
	Platynothrus sp.			1				
	Symplecta stictica		2	4				
	Nemotelus sp.		3	2	1	4	3	
	Ochthebius marinus		4	4	3	3	1	
	Orchestia gammarellus	4	5	4	5	5	5	5
	Ovatella myosotis		4	5	5	5	5	5
	Phyllocia moscorum							1
	Fucellia maritima	4						
	Enchytraea sp.	5						

Every subsample is composed of at least 5 individuals. Not all species were identified up to the species

level, those were given a 'sp.' behind the genus name. When more than one species was collected per taxa, 'spp.' is given behind the genus name.