University of Groningen

Structure determination of the exopolysaccharide of *Lactobacillus fermentum* TDS030603-A revision

Gerwig, Gerrit J.; Dobruchowska, Justyna M.; Shi, Tala; Urashima, Tadasu; Fukuda, Kenji; Kamerling, Johann P.

Published in:
Carbohydrate Research

DOI:
[10.1016/j.carres.2013.04.026](https://doi.org/10.1016/j.carres.2013.04.026)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gerwig, G. J., Dobruchowska, J. M., Shi, T., Urashima, T., Fukuda, K., & Kamerling, J. P. (2013). Structure determination of the exopolysaccharide of *Lactobacillus fermentum* TDS030603-A revision. *Carbohydrate Research*, 378, 84-90. <https://doi.org/10.1016/j.carres.2013.04.026>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Structure determination of the exopolysaccharide of *Lactobacillus fermentum* TDS030603—A revision

Gerrit J. Gerwig^{a,b,*}, Justyna M. Dobruchowska^b, Tala Shi^c, Tadasu Urashima^c, Kenji Fukuda^c,
Johannis P. Kamerling^{a,b}

^a NMR Spectroscopy, Bijvoet Center, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

^b Microbial Physiology, Groningen Biomolecular Sciences and Biotechnology Institute (GBB), University of Groningen, Nijenborgh 7, 8747 AG Groningen, The Netherlands

^c Graduate School of Animal & Food Hygiene, Obihiro University of Agriculture & Veterinary Medicine, Inada cho, Obihiro, Hokkaido 080-8555, Japan

ARTICLE INFO

Article history:

Received 13 March 2013

Received in revised form 17 April 2013

Accepted 18 April 2013

Available online 28 April 2013

In memory of Professor Malcolm Perry and
Professor Lennart Kenne

Keywords:

Lactobacillus fermentum

Exopolysaccharide

Lactic acid bacteria

Structural analysis

NMR spectroscopy

ABSTRACT

The highly viscous neutral exopolysaccharide, produced in large amounts by *Lactobacillus fermentum* TDS030603 in MRS broth and in a chemically defined medium supplemented with glucose, was investigated by GLC(-EIMS) monosaccharide and methylation analysis, periodate oxidation, MALDI-TOF mass spectrometry, and 1D/2D NMR spectroscopy. The GLC(-EIMS) analyses showed that the EPS contained D-glucose and D-galactose in an averaged molar ratio of 2.6:1.0, consisting of terminal D-Glcp, 3-substituted D-Glcp, 2,3-disubstituted D-Glcp, and 6-substituted D-Galp in an averaged molar ratio of 1.3:1.0:1.1:1.1, with a trace of terminal D-Galp (0.1). Combined with the NMR data of the EPS (previously reported constituent analysis: terminal Glcp, 3-substituted Glcp, 2,3-disubstituted Glcp, and 6-substituted Galp in an averaged molar ratio of 1.2:1.1:1.0:1.1) and oligosaccharides isolated after partial acid hydrolysis, a branched tetrasaccharide repeating unit, with some heterogeneity in the side chain, having the following structure is proposed:

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Exopolysaccharides (EPSs) produced by dairy lactic acid bacteria, carrying the GRAS (generally recognized as safe) status, are of increasing interest to the food industry, being often used to improve the texture and consistency of fermented dairy products.¹ In addition to these features, some of the EPSs have been reported to have beneficial health properties such as anti-tumor, anti-ulcer, and immune-stimulating activities.^{2–4}

In earlier studies, it has been found that the neutral exopolysaccharide produced in large amounts by *Lactobacillus fermentum*

strain TDS030603 yields a high degree of ropiness, being a serious candidate for use as an additive to increase the viscosity of food. The strain originated from a collection of lactic acid bacterial strains, isolated from traditional fermented milk products at the Obihiro University of Agriculture and Veterinary Medicine. Besides structural analysis studies of the EPS⁵ and studies on the effects of carbohydrate source on its physicochemical properties,⁶ also attention was paid to the characterization and expression analysis of the EPS gene cluster.⁷ The structural determination data, reported six years ago,⁵ were interpreted as belonging to one major and one minor EPS, both built up from a Glc-/Gal-containing tetrasaccharide repeating unit (two α and two β anomeric residues). The major EPS should contain terminal Glcp, 3-substituted Glcp, 2,3-disubstituted Glcp, and 6-substituted Galp. The minor EPS

* Corresponding author. Tel.: +31 302539926.

E-mail address: g.gerwig@rug.nl (G.J. Gerwig).

should contain terminal Galp, 3-substituted Glcp, 2,3-disubstituted Glcp, and 6-substituted Galp. For the major EPS, two possible structures for the backbone were proposed, built up from two Glc and one Gal residues with one Glc residue as side chain connected to one of the Glc backbone residues. For the minor EPS the same backbone possibilities were proposed, but now with one Gal residue as side chain connected to one of the Glc backbone residues.

Here, we report on a revised structure of the EPS of *Lb. fermentum* strain TDS030603, showing that the earlier idea of one major and one minor EPS was not correct. In fact, a detailed study, making use of monosaccharide and methylation analysis, partial acid hydrolysis, periodate oxidation, MALDI-TOF mass spectrometry,

and 1D/2D NMR spectroscopy, indicated one branched EPS structure (tetrasaccharide repeating unit with a disaccharide backbone), being slightly heterogeneous in its side chain.

2. Results and discussion

2.1. Preparation of EPS from *Lb. fermentum* TDS030603

The EPSs **EPS-MRS** and **EPS-CDM_{Glc}** produced by *Lb. fermentum* TDS030603 in MRS broth and in chemically defined medium supplemented with glucose, respectively, were isolated via absolute

Table 1

Linkage analysis data of *Lb. fermentum* TDS030603 **EPS5'-MRS**, **EPS5'-CDM_{Glc}**, and EPS using whey-mediated broth (**EPS-WMB**⁵)

Partially methylated alditol acetate	R_t^a	Structural feature	Molar ratio ^b %		
			EPS5'-MRS	EPS5'-CDM_{Glc}	EPS-WMB
1,5-Di-O-acetyl-2,3,4,6-tetra-O-methyl-hexitol-1-d	1.00	Glcp(1→	29	29	25
1,5-Di-O-acetyl-2,3,4,6-tetra-O-methyl-hexitol-1-d	1.03	Galp(1→	2	2	6
1,3,5-Tri-O-acetyl-2,4,6-tri-O-methyl-hexitol-1-d	1.16	→3)Glcp(1→	23	20	22
1,5,6-Tri-O-acetyl-2,3,4-tri-O-methyl-hexitol-1-d	1.24	→6)Galp(1→	22	24	25
1,2,3,5-Tetra-O-acetyl-4,6-di-O-methyl-hexitol-1-d	1.28	→2,3)Glcp(1→	24	25	22

^a R_t , GLC retention times relative to 1,5-di-O-acetyl-2,3,4,6-tetra-O-methyl-glucitol-1-d (1.00) on EC-1.

^b Calculated on the basis of FID responses (peak areas), taking into account similar molar adjustment factors.

Figure 1. 500-MHz 1D ¹H NMR spectra of native **EPS-MRS** (A) and **EPS-CDM_{Glc}** (B) produced by *Lb. fermentum* TDS030603 in MRS broth and in chemically defined medium supplemented with glucose, respectively, recorded in D₂O at 334 K. The HOD signal at 4.40 ppm was suppressed.

ethanol precipitation, followed by subsequent anion-exchange and size-exclusion chromatography.⁶

2.2. Composition of EPS-MRS and EPS-CDM_{Glc}

Monosaccharide analysis, including absolute configuration determination, of both **EPS-MRS** and **EPS-CDM_{Glc}** revealed the presence of D-glucose (Glc) and D-galactose (Gal) in a molar ratio of 2.6:1.0, in agreement with earlier data [Glc:Gal = 2.5:1 (whey-mediated broth, **EPS-WMB**),⁵ 2.6:1 (**EPS-MRS**),⁶ 2.6–2.8:1 (**EPS-CDM_{Glc}**, **EPS-CDM_{Gal}**, **EPS-CDM_{Lac}**, and **EPS-CDM_{Suc}**).⁶

As the native EPSs did not dissolve in dimethyl sulfoxide, the partially hydrolyzed (0.5 M TFA; 5 min, 100 °C) EPSs **EPS5'-MRS** and **EPS5'-CDM_{Glc}** were used for methylation analysis. Terminal Glcp, 3-substituted Glcp, 2,3-disubstituted Glcp, and 6-substituted Galp residues were found in roughly similar molar ratios as those reported earlier for **EPS-WMB**⁵ (Table 1), suggesting highly similar/identical branched polysaccharide structures.

2.3. NMR spectroscopy of EPS-MRS and EPS-CDM_{Glc}

The native EPSs **EPS-MRS** and **EPS-CDM_{Glc}** showed high viscosity by dissolving them in D₂O, thus only low-concentration samples could be used for NMR measurements. Due to a carbohydrate signal at δ 4.73, which is in the close neighborhood of the HOD signal at 300 K, all NMR spectra were recorded at

334 K (HOD signal at 4.40 ppm). As is evident from Figure 1, the 500-MHz ¹H NMR spectra of both native EPS probes are identical, and highly similar to those published earlier for **EPS-MRS**, **EPS-WMB**, **EPS-CDM_{Glc}**, **EPS-CDM_{Gal}**, **EPS-CDM_{Lac}**, and **EPS-CDM_{Suc}**, recorded at 343 K.^{5,6} In the anomeric regions four H-1 signals are present (Fig. 1), denoted **A** at δ 5.674, **B** at δ 5.325, **C** at δ 4.985, and **D** at δ 4.731, respectively, according to decreasing chemical shift values. The δ values suggest α -configurations for **A**, **B**, and **C**, and β -configuration for **D**. The signal at δ 4.516 is not stemming from an anomeric proton (HSQC, δ_C 70.1; see below).

As the **EPS5'-MRS** and **EPS5'-CDM_{Glc}** samples, used for methylation analysis, had a higher solubility and a lower viscosity in D₂O, also 1D ¹H NMR spectra of these probes were recorded at 334 K (Fig. 2). Both EPSs gave identical spectra, highly similar to the spectra of the native samples (Fig. 1). The four anomeric signals **A–D** had the same chemical shifts as mentioned above. Both **EPS5'-MRS** and **EPS5'-CDM_{Glc}** were further investigated by 2D NMR spectroscopy. TOCSY with different mixing times (20, 50, 75, 100, and 200 ms) and HSQC experiments were performed to obtain the complete assignment of the ¹H and ¹³C chemical shifts of the constituent residues **A–D** (Table 2), whereas NOESY measurements were carried out to generate monosaccharide sequence data (Table 3). The combined ¹H and ¹³C anomeric δ values support the occurrence of α -configurations for **A**, **B**, and **C**, and β -configuration for **D**. In Figure 3, the 1D ¹H NMR, the HSQC, and the TOCSY (200 ms) spectra, as well as the relevant part of the NOESY spectrum of **EPS5'-MRS** are depicted.

Figure 2. 500-MHz 1D ¹H NMR spectra of **EPS5'-MRS** (A) and **EPS5'-CDM_{Glc}** (B), recorded in D₂O at 334 K. The HOD signal at 4.40 ppm was suppressed. Both samples were obtained from the corresponding native EPSs after treatment with 0.5 M TFA for 5 min at 100 °C.

Table 2¹H and ¹³C NMR chemical shifts^a of monosaccharide residues **A–D** in **EPSS'-CDM_{GLC}** and **EPSS'-MRS**, recorded in D₂O at 334 K

Residue	H-1 C-1	H-2 C-2	H-3 C-3	H-4 C-4	H-5 C-5	H-6a C-6	H-6b
A	5.674	3.94	4.19	3.63	4.13	3.86	3.76
-(1→2,3)-α-D-Glcp-(1→	96.4	73.5^b	79.9	69.8	72.7	61.8	
B	5.325	3.89	4.02	3.96	4.516	3.91	3.71
-(1→6)-α-D-Galp-(1→	95.0	69.8	70.4	72.1	70.1	68.0	
C	4.985	3.57	3.72	3.44	3.74	3.86	3.77
α-D-Glcp-(1→	98.6	72.6	74.9	71.0	74.0	61.8	
D	4.731	3.47	3.74	3.69	3.54	3.94	3.76
-(1→3)-β-D-Glcp-(1→	103.7	73.2	83.1	72.4	77.0	61.8	

^a In ppm relative to the signal of internal acetone at δ 2.225 for ¹H and at δ 31.08 for ¹³C.^b Ring carbons involved in linkages are indicated in gray.**Table 3**Intra- and inter-residue NOE connectivities observed for **EPSS'-MRS** (see Figs. 3 and 4)

NOE	Type	Feature ^a
A H-1– A H-2	Intra	α-D-Glcp unit
A H-1– B H-1	Inter	Close in space
A H-1– D H-3	Inter	→2,3)-α-D-Glcp-(1→3)-β-D-Glcp-(1→
A H-1– D H-4	Inter	Close in space
B H-1– B H-2	Intra	α-D-Galp unit
B H-1– A H-1	Inter	Close in space
B H-1– A H-2	Inter	→6)-α-D-Galp-(1→2,3)-α-D-Glcp-(1→
C H-1– C H-2	Intra	α-D-Glcp unit
C H-1– B H-6a/6b	Inter	α-D-Glcp-(1→6)-α-D-Galp-(1→
D H-1– D H-3	Intra	β-D-Glcp unit
D H-1– D H-5	Intra	β-D-Glcp unit
D H-1– A H-3	Inter	→3)-β-D-Glcp-(1→3,2)-α-D-Glcp-(1→

^a Anomeric configurations and linkages in bold reflect the observed NOE cross-peaks.

Comparison of the TOCSY spectra of **EPSS'-MRS** with increasing mixing times (only the final spectrum of 200 ms is shown in Fig. 3) allowed the assignment of the sequential order of the chemical shifts belonging to the same spin system.

The TOCSY **A** H-1 track (δ 5.674) showed the scalar coupling network H-2,3,4,5,6a. The H-6b value was found via the H-6a signal in the HSQC spectrum. Combined with the downfield shifts for **A** C-2 (δ 73.5; Me α-D-Glcp, δ_{C-2} 72.2⁸) and **A** C-3 (δ 79.9; Me α-D-Glcp, δ_{C-3} 74.1⁸), deduced from the HSQC spectrum, residue **A** corresponds with the →2,3)-α-D-Glcp-(1→ unit, observed in the methylation analysis. The relatively small downfield shift for **A** C-2 is due to the fact that also **A** C-3 is substituted, causing an additional small upfield shift for **A** C-2 (in agreement with a CASPER simulation⁹).

The TOCSY **B** H-1 track (δ 5.325) revealed cross-peaks with **B** H-2,3,4. The cross-peaks with **B** H-6a,6b were present on the **B** H-5 track at δ 4.516, which also contained the cross-peak with **B** H-4. The magnetization transfer on the H-1 track went only as far as **B** H-4, suggesting that **B** is the α-D-Galp residue. The relatively large ¹H downfield shift of **B** H-5 (δ 4.516) is remarkable and could be due to the substitution at C-6 and/or a peculiar tertiary structure of the polysaccharide fragment containing this **B** residue. For residue **B**, the ¹H and ¹³C (**B** C-6, δ 68.0; Me α-D-Galp, δ_{C-6} 62.2⁸) NMR data are in agreement with a →6)-α-D-Galp-(1→ unit.

The TOCSY **C** H-1 track (δ 4.985) revealed cross-peaks with **C** H-2,3,4,5,6a, yielding a set of chemical shifts, typical for a terminal α-D-Glcp-(1→6)-residue.¹⁰

Finally, the set of cross-peaks with **D** H-2,3,4,5,6a,6b, as seen on the TOCSY **D** H-1 track (δ 4.731), matched that of a →3)-β-D-Glcp-(1→ unit.¹¹ The 3-substitution is supported by the **D** C-3 signal at δ 83.1 (Me β-D-Glcp, δ_{C-3} 76.8⁸).

To deduce information about the sequence of the monosaccharide residues in the EPS, a 2D NOESY spectrum was recorded

(Fig. 3), and the observed intra- and inter-residue cross-peaks are listed in Table 3. The intra-residue cross-peaks between **A** H-1 and **A** H-2, **B** H-1 and **B** H-2, and **C** H-1 and **C** H-2 confirmed the α-configurations for the two D-Glcp residues **A** and **C** and the D-Galp residue **B**, whereas the β-configuration for the **D** D-Glcp residue was supported by the intra-residue cross-peaks between **D** H-1 and **D** H-3 and between **D** H-1 and **D** H-5. On the **A** H-1 track an inter-residue cross-peak with **D** H-3 indicated an **A**(1→3)**D** linkage [→3)-α-D-Glcp-(1→3)-β-D-Glcp]. The observed cross-peak with **D** H-4 reflects the proximity in space of this proton to **A** H-1, due to the **A**(1→3)**D** linkage. On the **D** H-1 track an inter-residue cross-peak with **A** H-3 showed a **D**(1→3)**A** linkage [→3)-β-D-Glcp-(1→3)-α-D-Glcp]. On the **B** H-1 track an inter-residue cross-peak with **A** H-2 supported a **B**(1→2)**A** linkage [→6)-α-D-Galp-(1→2)-α-D-Glcp], and on the **C** H-1 track inter-residue cross-peaks with **B** H-6a and 6b corresponded with a **C**(1→6)**B** linkage [α-D-Glcp-(1→6)-α-D-Galp]. The inter-residue cross-peaks between **A** H-1 and **B** H-1 reflect the proximity in space of the 2,3-disubstituted α-D-Glcp **A** and 6-substituted α-D-Galp **B** residues. A structural interpretation of a part of the EPS is shown in Figure 4, including the repeating unit and the observed NOE connectivities.

Integration of the anomeric signals in the 1D ¹H NMR spectrum gave the peak area ratio of **A**:**B**:**C**:**D** = 1.0:1.1:1.2:1.1, in agreement with a branched repeating tetrasaccharide unit. The trace of terminal galactose found by methylation analysis is too low to be detected by NMR spectroscopy. But, it probably indicates a minor heterogeneity in the EPS, missing the terminal α-D-Glcp- unit in the side chain.

2.4. Preparation and isolation of oligosaccharides

Partial acid hydrolysis of native **EPS-MRS** and **EPS-CDM_{GLC}** (0.5 M TFA; 20 min, 100 °C) yielded in each case a complex mixture of oligosaccharides, which was fractionated on Bio-Gel P-2. Twenty 10-min fractions were collected and checked by MALDI-TOF-MS, demonstrating that each fraction contained several oligosaccharides, overall ranging from DP37 to DP2, and differing in one hexose unit. Several oligosaccharide fractions from the **EPS-MRS** fractionation were investigated by 1D ¹H NMR spectroscopy, all showing the same spectra (data not shown), being identical with the ¹H NMR spectra of **EPS-MRS** and **EPS-CDM_{GLC}** (Fig. 1) and **EPSS'-MRS** and **EPSS'-CDM_{GLC}** (Fig. 2). Additionally, 2D TOCSY and HSQC analysis of the DP17–DP37 fraction revealed identical spectra to those recorded for **EPSS'-MRS** and **EPSS'-CDM_{GLC}** (data not shown). Monosaccharide analysis of the DP2 fraction showed the presence of Gal and Glc in the molar ratio 1: 1, and ¹H NMR analysis (Fig. 5) indicated a Glcp(α1→6)Galp disaccharide (Table 4, ¹H and ¹³C chemical shifts obtained from TOCSY and HSQC measurements), in accordance with the proposed side chain. Also here, α-D-Galp H-5 resonates at a relatively downfield position (δ 4.259; C-5, δ 69.4) (see above).

Figure 3. 1D ^1H NMR and 2D HSQC, TOCSY (mixing time, 200 ms), and NOESY (mixing time, 200 ms) spectra of **EP55-MRS**, recorded in D_2O at 334 K. In the 2D spectra, relevant cross-peaks are included (see text). HSQC: A1 refers to a cross-peak between **A** H-1 and **A** C-1, etc. TOCSY: A1 corresponds to the diagonal peak of residue **A** H-1; A2 refers to a cross-peak between **A** H-1 and **A** H-2, etc. NOESY: A1 corresponds to the diagonal peak of residue **A** H-1; A1-D3 (encircled) refers to an inter-residue cross-peak between **A** H-1 and **D** H-3; A2 refers to an intra-residue cross-peak between **A** H-1 and **A** H-2, etc.

2.5. Periodate oxidation

Monosaccharide analysis of the polymeric material, obtained after periodate oxidation and subsequent NaBH₄ reduction of **EPS5'-MRS**, showed the only presence of glucose.

3. Conclusion

Taking together the monosaccharide analysis (on average D-Glc: D-Gal = 2.6: 1.0) and the methylation analysis (on average Glcp(1 $\rightarrow$: $\rightarrow$ 3)Glcp(1 $\rightarrow$: $\rightarrow$ 6)Galp(1 $\rightarrow$: $\rightarrow$ 2,3)Glcp(1 $\rightarrow$: Galp(1 $\rightarrow$: = 1.2:1.0:1.1:1.1:0.2) data of this study and the earlier studies,^{5,6} combined with the detailed 1D/2D NMR analyses of the EPSs and the partial acid hydrolysate, a branched tetrasaccharide repeating unit for the EPS from *Lactobacillus fermentum* TDS030603 is indicated, independent on the used culture media, as follows:

The finding of small amounts of terminal Gal residues supports heterogeneity in the side chain. The reproducible finding of the Gal:Glc molar ratio, being lower than 3:1, indicated that for a few percentage not only the terminal Glc residue in the side chain is absent, but also the whole Glc-Gal side chain. A remarkable feature of the structure is that the non-branched D-GlcP residues in the backbone have β -configuration and the branched ones α -configuration. Looking back at the earlier reported structures, it can be ascertained that in the 1D NMR analysis the peaks at δ 5.674 (here **A** H-1) and δ 4.516 (here **B** H-5) were interpreted in the wrong way.^{5,6} The proposed revised EPS structure was simulated in CASPER (www.casper.organ.su.se/casper) for reference ^1H and ^{13}C chemical shift values, and the calculated data were in close agreement with the measured chemical shifts.

Recently, the composition of an EPS produced by *Lb. fermentum* F6 was reported, also containing glucose and galactose, however, in a molar ratio of 4:3.¹²

4. Experimental

4.1. Production, isolation, and purification of exopolysaccharide from *Lb. fermentum* TDS030603

Lb. fermentum TDS030603 was pre-cultured aerobically in Man-Rogosa-Sharpe (MRS) medium for 24 h at 30 °C. Then, the cells were harvested by centrifugation and washed thoroughly with sterilized phosphate buffered saline. The washed cell suspension was inoculated into 1 L of either MRS or a chemically defined medium,⁶ to which 1% (w/v) glucose was added as a sole carbon source (CDM_{Glc}), yielding optical densities of 0.2 at 600 nm. After aerobic cultivation for 48 h at 30 °C, in each case the cells were removed by centrifugation (17,000g, 1 h, 4 °C). **EPS-MRS** or **EPS-CDM_{Glc}**, secreted into the supernatant, was precipitated by the addition of an equal volume of ice-cold ethanol, collected by centrifugation (17,000g, 30 min, 4 °C), and dissolved in 30 mL pure water. The EPS solutions were dialyzed overnight against pure water at 4 °C, and then lyophilized. The lyophilized EPSs were dissolved in 10 mL 50 mM Tris-HCl (pH 8.7) and purified by a batch method using a 20-mL slurry of DEAE A-50 equilibrated with the same buf-

Figure 4. Schematic presentation of the observed ^1H intra- (blue) and inter-residue (red) NOE connectivities (see Table 3) in a part of the *Lb. fermentum* TDS030603 EPS structure, including the repeating unit.

Figure 5. 500-MHz 1D ^1H NMR spectrum of the EPS-derived disaccharide $\alpha\text{-D-Glcp-(1}\rightarrow\text{6)-D-Galp}$, recorded in D_2O at 334 K. The HOD signal at 4.40 ppm was suppressed.

Table 4

^1H and ^{13}C NMR chemical shifts^a of monosaccharide residues in the EPS-derived disaccharide $\alpha\text{-D-Glcp-(1}\rightarrow\text{6)-D-Galp}$, recorded in D_2O at 334 K

Residue	H-1 C-1	H-2 C-2	H-3 C-3	H-4 C-4	H-5 C-5	H-6a C-6	H-6b
$\alpha\text{-D-Glcp-(1}\rightarrow\text{6)-}$	4.943	3.57	3.72	3.42	3.74	3.86	3.76
$-(1\rightarrow\text{6)-}\alpha\text{-D-Galp}$	99.1	72.2	73.9	70.4	72.6	61.4	
	5.254	3.81	3.84	4.03	4.259	3.89	3.72
$-(1\rightarrow\text{6)-}\beta\text{-D-Galp}$	93.3	69.3	69.8	70.3	69.4	67.5 ^b	
	4.587	3.48	3.65	3.97	3.89	3.77	3.70
	97.3	72.7	73.5	69.6	73.9	67.5	

^a In ppm relative to the signal of internal acetone at δ 2.225 for ^1H and at δ 31.08 for ^{13}C .

^b Ring carbons involved in linkages are indicated in gray.

fer. The non-adsorbed fractions were collected, thoroughly dialyzed against pure water, and lyophilized. **EPS-MRS** was further purified on a column of Toyopearl HW-55F (2.6 × 100 cm, 15 mL/h, pure water), dialyzed against pure water, and lyophilized. Both EPSs were stored in a desiccator until use.

4.2. Chemical analyses

Monosaccharide analyses, absolute configuration determinations, methylation analyses of partially hydrolyzed (0.5 M TFA; 5 min, 100 °C) polysaccharide samples, and periodate oxidation of **EPS-MRS** were carried out as described previously.^{13–17}

4.3. Mild partial acid hydrolysis of EPS-MRS

Polysaccharide samples (5 mg/mL) were treated with 0.5 M trifluoroacetic acid (TFA) for 5 min at 100 °C, and then the solutions were concentrated to dryness under a stream of nitrogen. The resulting products were dissolved in water and lyophilized.

4.4. Preparation of oligosaccharide fragments from EPS-MRS

A polysaccharide sample (~20 mg/2 mL) was treated with 0.5 M TFA for 20 min at 100 °C. After evaporation of the solution to dryness under a stream of nitrogen, the generated mixture of oligosaccharides was dissolved in water and lyophilized. The residue, dissolved in 2 mL water, was fractionated on a Bio-Gel P-2 column (94 cm × 1 cm), eluted with 10 mM ammonium bicarbonate at a flow rate of 13 mL/h. Fractions of 10 min were collected and checked for carbohydrate by an orcinol/H₂SO₄ spot test and for oligosaccharide mass distribution by matrix-assisted laser-desorption ionization time-of-flight mass spectrometry (MALDI-TOF-MS).

4.5. MALDI-TOF mass spectrometry

MALDI-TOF-MS, in positive-ion mode, of oligosaccharide materials was performed on an AXIMATM mass spectrometer (Shimadzu Kratos Inc., Manchester, UK), equipped with a nitrogen laser (337 nm, 3 ns pulse width). Samples were prepared by mixing on the target 0.5 µL oligosaccharide solution with 0.5 µL 2,5-dihydroxybenzoic acid (10 mg/mL) in 50% aqueous acetonitrile as matrix solution.

4.6. NMR spectroscopy

Resolution-enhanced 1D/2D NMR spectra were recorded in D₂O on a Bruker DRX-500 spectrometer (Bijvoet Center, NMR Spectroscopy, Utrecht University) at a probe temperature of 334 K. Before analysis, samples were exchanged twice in D₂O (99.9 atom% D, Cambridge Isotope Laboratories, Inc., Andover, MA) with intermediate lyophilization, and then dissolved in 0.6 mL D₂O. Suppression of the HOD signal was achieved by applying a WEFT pulse

sequence for 1D experiments and by a pre-saturation of 1 s during the relaxation delay for 2D experiments. The 2D TOCSY spectra were recorded using an MLEV-17 mixing sequence with spin-lock times of 20–200 ms. The 2D NOESY spectra were recorded using standard Bruker XWINNMR software with mixing time of 200 ms. The carrier frequency was set at the downfield edge of the spectrum in order to minimize TOCSY transfer during spin-locking. Natural abundance 2D ¹³C–¹H HSQC experiments (¹H frequency 500.0821 MHz, ¹³C frequency 125.7552 MHz) were recorded without decoupling during acquisition of the ¹H FID. Resolution enhancement of the spectra was performed by a Lorentzian-to-Gaussian transformation for 1D spectra or by multiplication with a squared-bell function phase shifted by $\pi/(2.3)$ for 2D spectra, and when necessary, a fifth order polynomial baseline correction was performed. Chemical shifts (δ) are expressed in ppm by reference to internal acetone (δ 2.225 for ¹H and δ 31.08 for ¹³C).

Acknowledgements

We thank Professor Rolf Boelens (NMR Spectroscopy, Bijvoet Center, Utrecht University, The Netherlands) for providing us with measuring time on the 500-MHz NMR instrument.

References

- De Vuyst, L.; Degeest, B. *FEMS Microbiol. Rev.* **1999**, *23*, 153–177.
- Kitazawa, H.; Nomura, M.; Itoh, T.; Yamaguchi, T. *J. Dairy Sci.* **1991**, *74*, 2082–2088.
- Nagaoka, M.; Hashimoto, S.; Watanabe, T.; Yokokura, T.; Mori, Y. *Biol. Pharm. Bull.* **1994**, *17*, 1012–1017.
- Chabot, S.; Yu, H.-L.; De Léséleuc, L.; Cloutier, D.; Van Calsteren, M.-R.; Lessard, M.; Roy, D.; Lacroix, M.; Oth, D. *Lait* **2001**, *81*, 683–697.
- Leo, F.; Hashida, S.; Kumagai, D.; Uchida, K.; Motoshima, H.; Arai, I.; Asakuma, S.; Fukuda, K.; Urashima, T. *J. Appl. Glycosci.* **2007**, *54*, 223–229.
- Fukuda, K.; Shi, T.; Nagami, K.; Leo, F.; Nakamura, T.; Yasuda, K.; Senda, A.; Motoshima, H.; Urashima, T. *Carbohydr. Polym.* **2010**, *79*, 1040–1045.
- Dan, T.; Fukuda, K.; Sugai-Bannai, M.; Takakuwa, N.; Motoshima, H.; Urashima, T. *Biosci. Biotechnol. Biochem.* **2009**, *73*, 2656–2664.
- Bock, K.; Thøgersen, H. *Annu. Rep. NMR Spectroscopy* **1982**, *13*, 1–57.
- <http://www.casper.org.au/se/casper>.
- Dobrurowska, J. M.; Gerwig, G. J.; Kralj, S.; Grijpstra, P.; Leemhuis, H.; Dijkhuizen, L.; Kamerling, J. P. *Glycobiology* **2012**, *22*, 517–528.
- Hreggvidsson, G. O.; Dobrurowska, J. M.; Fridjonsson, O. H.; Jonsson, J. O.; Gerwig, G. J.; Aevansson, A.; Kristjansson, J. K.; Curti, D.; Redgwell, R. R.; Hansen, C.-E.; Kamerling, J. P.; Debeche-Boukhit, T. *Glycobiology* **2011**, *21*, 304–328.
- Zhang, Y.; Li, S.; Zhang, C.; Luo, Y.; Zhang, H.; Yang, Z. *African J. Biotechnol.* **2011**, *10*, 2080–2091.
- Kamerling, J. P.; Vliegthart, J. F. G. *Mass Spectrometry In Clinical Biochemistry—Principles, Methods, Applications*; Lawson, A. M., Ed.; Walter de Gruyter: Berlin, 1989; Vol. 1, pp 175–263.
- Dobrurowska, J. M.; Gerwig, G. J.; Babuchowski, A.; Kamerling, J. P. *Carbohydr. Res.* **2008**, *343*, 726–745.
- Gerwig, G. J.; Kamerling, J. P.; Vliegthart, J. F. G. *Carbohydr. Res.* **1978**, *62*, 349–357.
- Ciucanu, I.; Kerek, F. *Carbohydr. Res.* **1984**, *131*, 209–217.
- Sanchez-Medina, I.; Gerwig, G. J.; Urshev, Z. L.; Kamerling, J. P. *Carbohydr. Res.* **2007**, *342*, 2430–2439.