University of Groningen

OXIDATIVE PHENOL COUPLING WITH CUPRIC-AMINE COMPLEXES

FERINGA, B; WYNBERG, H

Published in:
Tetrahedron Letters

DOI:
[10.1016/S0040-4039\(01\)83533-4](https://doi.org/10.1016/S0040-4039(01)83533-4)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1977

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
FERINGA, B., & WYNBERG, H. (1977). OXIDATIVE PHENOL COUPLING WITH CUPRIC-AMINE COMPLEXES. *Tetrahedron Letters*, 18(50), 4447-4450. [https://doi.org/10.1016/S0040-4039\(01\)83533-4](https://doi.org/10.1016/S0040-4039(01)83533-4)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

OXIDATIVE PHENOL COUPLING WITH CUPRIC-AMINE COMPLEXES

Ben Feringa and Hans Wynberg*

Department of Organic Chemistry,
The University, Zernikelaan, Groningen, The Netherlands

(Received in UK 22 September 1977; accepted for publication 21 October 1977)

The oxidative coupling of phenolic compounds as a mode of formation of a diversity of biologically important structures has been investigated for a long time¹. Phenol oxidations with modified metal complexes have received considerable attention during the last years².

Although the role of copper ions in enzymic (laccase, tyrosinase) phenol oxidations has been recognized^{1,2}, relatively little attention has been paid to the use of copper complexes in phenol oxidations, either as a synthetic method or in a biomimetic type synthesis of natural products. Oxidations of phenols using copper complexes under aerobic conditions yield quinones, polyethers or carbon-carbon bond cleaved products. Thus Brackman and Havinga, in a study of the mode of action of tyrosinase, reported the use of catalytic amounts of a cupric-morpholine complex in the oxidation of phenol giving *o*-benzoquinone as a primary product followed by substitution and oxidation to di-morpholino-substituted *o*-benzoquinone (Scheme I, A)³.

Rogić and coworkers recently showed that oxidation of phenol using an excess of the pyridine-cupric complex PyCuClO₄Me as oxidant gave *cis*-,*cis*-muconic acid *via* an intermediate *o*-benzoquinone. When an oxygen substituent is already present in the *ortho* position of phenol - as in the case of catechol or *o*-benzoquinone - oxidation to *cis*-,*cis*-muconic acid occurs also under anaerobic conditions (Scheme I, B)⁴.

Furthermore, the formation of polyethers from the cupric-pyridine complex catalyzed oxidation of 2,6-dimethylphenol, in which *o*-quinone formation is blocked, under aerobic conditions is a well-described process (Scheme I, C)⁵.

Exclusion of oxygen in the oxidation of phenol 1 could lead to the elimination of pathways A and B, possibly giving coupled phenolic products (Scheme I, D)⁶.

We now wish to report this supposition to be correct. Using cupric-amine complexes under anaerobic conditions in molar quantities, we have observed the intra- and intermolecular coupling of phenols as a general reaction.

Scheme I: Oxidations of phenol and 2,6-dimethylphenol with copper complexes.

M = morpholine

py = pyridine

The green or blue coloured stock solutions of the cupric-amine complexes were prepared by stirring Cu(OAc)₂·H₂O or Cu(NO₃)₂·3H₂O and amines in a 1 to 3 ratio in either methanol, ethanol, water or alcohol-water mixtures at room temperature under nitrogen for 15 minutes.

A typical oxidative coupling using these complexes was carried out in the following manner. To a solution of the complex formed from 0.72 g (3.0 mmol) Cu(NO₃)₂·3H₂O and 1.09 g (9.0 mmol) dl- α -phenylethylamine⁷ (α -P.E.A.) in 30 ml of CH₃OH, 0.43 g (3.0 mmol) of 2-naphthol (2) dissolved in 10 ml CH₃OH was added and the mixture was stirred at room temperature under nitrogen for 20 hours. The dark-brown product - probably the copper-dinaphthol-amine complex - was decomposed with dil. hydrochloric acid and gave after a normal work-up 1,1'-dinaphthol (3) in 62% yield (m.p. 216-218°C). A similar oxidation of 2 in water as a solvent at 100°C using the cupric complex of pyridine gave 3 in 70% yield⁹.

The oxidations of the following phenols using the cupric- α -P.E.A. complex were also investigated.

The oxidation of *p*-cresol gave "Pummerer's ketone" (4) in 15% yield. The dimer (6) of 2,7-dihydroxynaphthalene (5) could be isolated in 25% yield and dimerization of 3,4,8-trimethyl-tetrahydronaphthol (7)¹⁰ gave 8 in 46% yield.

Oxidation of 2,6-dimethylphenol (1) with the cupric- α -P.E.A. complex under nitrogen gave dimers 9 and 10 as a 7 to 1 mixture in 20% yield. Only traces of polymers were present (Scheme I, C and D). The coupling of 3-hydroxyphenanthrene (11)¹¹ occurred at the hindered 4-position and the racemic 4,4'-biphenanthrol (12) was obtained in 20% yield¹². A biogenetic type synthesis of dehydrogriseofulvin^{1,13} could also be achieved. Oxidation of griseophenone 13 with the cupric-dl- α -P.E.A. complex in methanol, water (1:9) gave the C-O coupled product 14 in 40% yield, identical with an independently prepared sample of the compound¹³.

On the basis of the present results it is clear that a diversity of phenol oxidation reactions can be performed with a cupric-amine complex. Considerable control can be exercised over the nature of the products by considering the structure of the starting phenol and the presence or absence of oxygen.

It is not too far fetched to suggest that this multi-parameter dependence in copper-amine oxidations is reminiscent of the known lack of specificity of many phenol oxidases¹.

REFERENCES AND NOTES

- 1) a. W.I. Taylor and A.R. Battersby, Eds. "Oxidative Coupling of Phenols", Marcel Dekker, New York, N.Y., 1967; b. H. Musso, *Angew. Chem.*, **75**, 965 (1963); c. A.I. Scott, *Q. Rev. Chem. Soc.*, **19**, 1 (1965); d. D.H.R. Barton, Pedler Lecture, *Chem. Br.*, **3**, 330 (1967); e. T. Kametani and K. Fukumoto, *Synthesis*, 657 (1972); f. P.D. McDonald and G.A. Hamilton in: "Oxidation in Organic Chemistry", Part B, W.S. Trahanovsky, Ed., Academic Press, New York, London 1973, Chapter II.
- 2) M.J.S. Dewar and T. Nakaya, *J. Am. Chem. Soc.*, **90**, 7134 (1968), S. Tobinaga and E. Kotani, *ibid*, **94**, 309 (1972), E. Kotani, N. Takeuchi and S. Tobinaga, *Tetrahedron Lett.*, 2735 (1973) M.A. Schwartz, R.A. Holton and S.W. Scott, *J. Am. Chem. Soc.*, **91**, 2800 (1969), M.A. Schwartz, B.F. Rose and B. Vishnuvajjala, *ibid.*, **95**, 612 (1973), M.A. Schwartz, B.F. Rose, R.A. Holton, S.W. Scott and B. Vishnuvajjala, *ibid.*, **99**, 2571 (1977), S.M. Kupchan and A.J. Liepa, *ibid.*, **95**, 4062 (1973). Kaeding reported the oxidation of phenols under drastic conditions, using copper (II) salts in an excess of a molten phenol at 160-220°C; W.W. Kaeding, *J. Org. Chem.*, **28**, 1063 (1963).
- 3) W. Brackman and E. Havinga, *Recl. Trav. Chim. Pays-Bas*, **74**, 937, 1021, 1070, 1100, 1107 (1955).
- 4) M.M. Rogić, T.R. Demmin, W.B. Hammond, *J. Am. Chem. Soc.*, **98**, 7441 (1976), J. Tsuji and H. Takayanagi, *ibid*, **96**, 7349 (1974), J. Tsuji, H. Takayanagi, I. Sakai, *Tetrahedron Lett.*, 1245 (1975), J. Tsuji and H. Takayanagi, *ibid*, 1365 (1976), D.G. Brown, L. Beckmann, C.H. Ashby, G.C. Vogel and J.T. Reinprecht, *Tetrahedron Lett.*, 1363 (1977).
- 5) A.S. Hay, H.S. Blanchard, G.F. Endres and J.W. Eustance, *J. Am. Chem. Soc.*, **81**, 6335 (1959), H. Finkbeiner, A.S. Hay, H.S. Blanchard and G.F. Endres, *J. Org. Chem.*, **31**, 549 (1966), E. Tsuchida, M. Kanedo and H. Nishide, *Makromol. Chem.*, **151**, 221 (1972).
- 6) Endres and coworkers have shown that it is possible to get 3,3',5,5'-tetramethyldiphenquinone as a major product from the oxidation of 2,6-dimethylphenol with molar quantities of cupric pyridine complex under nitrogen (ref. 5).
- 7) Using d- α -phenylethylamine, low but reliable asymmetric induction has been achieved and will be the subject of a separate communication. B. Feringa and H. Wynberg, manuscript in preparation.
- 8) Identical with the product of a FeCl₃ oxidation of 2. The remaining 38% consists mainly of starting material.
- 9) Reaction time, 3 minutes.
- 10) B. Feringa and H. Wynberg, *J. Am. Chem. Soc.*, **98**, 3372 (1976).
- 11) L.F. Fieser, *J. Am. Chem. Soc.*, **51**, 2460 (1929).
- 12) The synthesis and properties of this new compound will be reported on shortly in detail.
- 13) A. Brossi, M. Baumann, M. Gerecke, and E. Kyburz, *Helv. Chim. Acta*, **43**, 1444, 2071 (1960). A.C. Day, J. Nabney and A.I. Scott, *Proc. Chem. Soc.*, 284 (1960), *J. Chem. Soc.*, 4067 (1961), G. Stork and M. Tomasz, *J. Am. Chem. Soc.*, **84**, 310 (1962), D. Taub, C.H. Kuo, H.L. Slaters and N.L. Wendler, *Tetrahedron*, **19**, 1 (1963).
- 14) During the preparation of this manuscript Kametani and coworkers reported the performance of a biomimetic type synthesis of isoquinoline alkaloids using the cuprous chloride, pyridine system in an oxygen atmosphere. T. Kametani, M. Ihara, M. Takemura, Y. Satoh, H. Terasawa, Y. Ohta, K. Fukumoto and K. Takahashi, *J. Am. Chem. Soc.*, **99**, 3805 (1977).
- 15) We thank the Netherlands Organization for Pure Research (Z.W.O.) for a graduate fellowship.