University of Groningen

NOVEL DIAMAGNETIC AND PARAMAGNETIC HYDRIDES OF VANADIUM(II)

HESSEN, B; VANBOLHUIS, F; TEUBEN, JH; PETERSEN, JL; Bolhuis, Fré van; Petersen, Jeffrey L.

Published in:
Journal of the American Chemical Society

DOI:
[10.1021/ja00209a050](https://doi.org/10.1021/ja00209a050)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1988

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

HESSEN, B., VANBOLHUIS, F., TEUBEN, JH., PETERSEN, JL., Bolhuis, F. V., & Petersen, J. L. (1988). NOVEL DIAMAGNETIC AND PARAMAGNETIC HYDRIDES OF VANADIUM(II). *Journal of the American Chemical Society*, 110(1), 295-296. <https://doi.org/10.1021/ja00209a050>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

is not identical with transport experiments A and C can be attributed to incomplete removal of Ag^+ or to leaks in the membrane that may have developed during the exchange processes.

At this time, it is not possible to assign the factors responsible for the enhanced fluxes of olefins through the Ag^+ form membranes unambiguously. If facilitated transport, which requires that the $\text{Ag}(\text{olefin})^+$ complex is mobile within the membrane, is the dominant factor, Noble has shown that the facilitation factor, assuming reaction equilibrium and no mass transport resistance at the decane/Nafion interfaces, should be¹⁸

$$F = 1 + \frac{\alpha K_D}{(1 + K_D)} \quad \text{where } K_D = K_{\text{eq}} C_{\text{olefin}}(x=0) \quad \text{and}$$

$$\alpha = \frac{D_{\text{complex}} C_{\text{Ag}^+}(\text{total})}{D_{\text{olefin}} C_{\text{olefin}}(x=0)}$$

We note that since the solubility of olefins in the aqueous membrane phase is low, the value of K_D in our system, ca. 2×10^{-4} is orders of magnitude lower than the theoretically predicted optimal values.¹⁹ This implies that still greater facilitation factors may be achieved by increasing either the complex formation constant or the solubility of the permeate in the membrane phase. We are presently exploring these possibilities as well as attempting to ascertain the factors responsible for the enhanced fluxes.

Acknowledgment. This research is supported by the NSF under Grant No. CBT-8604518. Acknowledgment is also made to the donors of the Petroleum Research Fund, administered by the ACS (17616-AC3). We thank J. J. Peligrino (NBS, Boulder, CO) and L. L. Burton (DuPont, Wilmington, DE) for providing samples of the Nafion membranes and R. Barkely (CU) for assistance with the gas chromatography measurements. Useful discussions with R. D. Noble (NBS, Boulder, CO) and J. D. Way (SRI International, Menlo Park, CA) are also gratefully acknowledged.

(18) Noble, R. D.; Way, J. D.; Powers, L. A. *Ind. Eng. Chem. Fund.* **1986**, 25, 450.

(19) Kemena, L. L.; Noble, R. D.; Kemp, N. J. *J. Membr. Sci.* **1983**, 15, 259.

Novel Diamagnetic and Paramagnetic Hydrides of Vanadium(II)

Bart Hessen, Fré van Bolhuis, and Jan H. Teuben*

Department of Chemistry, Rijksuniversiteit
Groningen, Nijenborgh 16
9747 AG Groningen, The Netherlands

Jeffrey L. Petersen

Department of Chemistry
West Virginia University
Morgantown, West Virginia 26506
Received August 10, 1987

Apart from V(I) complexes like $[\text{CpVH}(\text{CO})_3]^{-1}$ and $\text{HV}(\text{C}-\text{O})_4(\text{PP})$ (PP = chelating diphosphine),² very few vanadium hydride species are known which are as follows: one V(III) compound, $(\text{C}_5\text{Me}_5)_2\text{VH}$, and its CO adduct³ and two dimeric species, $[\text{CpV}(\mu\text{-H})]_2(\mu\text{-C}_6\text{H}_6)^4$ and $\text{V}_2\text{H}_4\text{Zn}_2(\text{BH}_4)_2(\text{PPh}_2\text{Me})_4$.⁵ Here

(1) Kinney, R. J.; Jones, W. D.; Bergman, R. G. *J. Am. Chem. Soc.* **1978**, 100, 7902.

(2) Many derivatives of this type have been described. See, for example: Puttfarcken, U.; Rehder, D. *J. Organomet. Chem.* **1980**, 185, 219 and references cited therein.

(3) Curtis, C. J.; Smart, J. C.; Robbins, J. L. *Organometallics* **1985**, 4, 1283.

(4) Jonas, K.; Wiskamp, V.; Tsay, Y.-H.; Krüger, C. *J. Am. Chem. Soc.* **1983**, 105, 5480.

Figure 1. Molecular structure of $[\text{CpV}(\mu\text{-H})\text{dmpe}]_2$ (**1**). Selected structural parameters: $\text{V}-\text{V}' = 2.701$ (1) Å, $\text{V}-\text{H} = 1.64$ (4) Å, $\text{V}-\text{H}' = 1.72$ (4) Å, $\angle \text{H}-\text{V}-\text{H}' = 73$. (2)°, $\angle \text{V}-\text{H}-\text{V}' = 107$. (2)°, $\angle \text{P}(1)-\text{V}-\text{P}(2) = 78.40$ (3)°, $\angle \text{CT}-\text{V}-\text{P}(1) = 115.41$ (2)°, $\angle \text{CT}-\text{V}-\text{P}(2) = 115.69$ (2)°, $\angle \text{CT}-\text{V}-\text{H} = 119$. (2)°. Prime denotes symmetry related atom; CT is Cp centroid.

we wish to report two new vanadium(II) hydride species, diamagnetic $[\text{CpV}(\mu\text{-H})\text{dmpe}]_2$ (**1**) (dmpe = 1,2-bis(dimethylphosphino)ethane) and paramagnetic $[\text{CpVH}(\text{dmpe})]_2(\mu\text{-dmpe})$ (**2**), produced in the reaction of the V(II) alkyl complex $\text{CpV}(\text{n-Pr})\text{dmpe}$ (**3**) with molecular hydrogen.

When molecular hydrogen (1 atm) is admitted to a dark brown solution of **3** in toluene at 0 °C, the solution turns red in several hours and a small amount of grey precipitate is formed. A Toepler pump experiment indicated that 1 mol of H_2 per mol of V was consumed and 1 mol of propane per mol of V was produced. After filtration, red crystals could be obtained from the liquid by cooling. This compound is diamagnetic (by NMR⁷) and was shown by X-ray analysis⁸ to be the dimeric $[\text{CpV}(\mu\text{-H})\text{dmpe}]_2$ (**1**, Figure 1), formed according to eq 1.

The molecule is centrosymmetric and consists of a planar V_2H_2 unit with two $\eta^5\text{-Cp}$ and two $\eta^2\text{-dmpe}$ ligands in a trans arrangement. A $\text{V}-\text{V}'$ distance of 2.701 (1) Å is found. EHMO calculations on $\text{CpV}(\eta^2\text{-BH}_4)(\text{PH}_3)_2$ have indicated that the d^3 metal center, in such a coordination geometry, has a preference for the low-spin ($S = 1/2$) state.⁶ Thus **1** can be considered to consist of two low-spin d^3 centers forming a single metal-metal bond. This is in accordance with both the observed intermetal

(5) Bansemer, R. L.; Huffman, J. C.; Caulton, K. G. *J. Am. Chem. Soc.* **1983**, 105, 6163.

(6) Hessen, B.; Lemmen, T. H.; Luttikhedde, H. J. G.; Teuben, J. H.; Huffman, J. C.; Caulton, K. G.; Petersen, J. L.; Jägger, S. *Organometallics* **1987**, 6, 2354.

(7) ¹H NMR (300 MHz, THF-*d*₆, 20 °C) of **1**: 4.13 ppm (s, 10 H, Cp), 2.49 ppm (br m, 4 H, P-CH₂), 2.05 ppm (br m, 4 H, P-CH₂), 1.40 ppm (s, 12 H, P-Me), 0.64 ppm (s, 12 H, P-Me), -15.9 ppm (very br, $\Delta\nu_{1/2} = 630$ Hz, 2 H, V-H).

(8) **1** crystallizes in the monoclinic space group $P2_1/n$, $a = 10.128$ (2) Å, $b = 8.511$ (5) Å, $c = 15.458$ (2) Å, $\beta = 100.48$ (1)° (130 K), $Z = 2$. Reflections (2056) with $1^\circ \leq \theta \leq 27^\circ$ were considered observed. All atoms were located from the Fourier difference map. The V-bound hydrogen atoms were refined isotropically, and the other hydrogen atoms were constrained at 0.95 Å from their corresponding C atoms in the final refinement cycles: $R = 0.039$, $R_w = 0.042$ ($w = 1$).

Figure 2. Molecular structure of $[\text{CpV}(\text{H})\text{dmpe}]_2(\mu\text{-dmpe})$ (**2**). Selected structural parameters: $\text{V-P}(1) = 2.369$ (1) Å, $\text{V-P}(2) = \text{V-P}(3) = 2.378$ (1) Å, $\angle\text{P}(1)\text{-V-P}(2) = 95.00$ (4)°, $\angle\text{P}(1)\text{-V-P}(3) = 117.40$ (5)°, $\angle\text{P}(2)\text{-V-P}(3) = 79.02$ (4)°.

distance (which is too long for a triple bond, characteristically 2.2–2.4 Å⁹) and the diamagnetism of **1**. The V–H distances (1.64 (4) and 1.72 (4) Å) and the nonbonding separation of the H atoms of 2.01 (5) Å are similar to those found in $[\text{CpV}(\mu\text{-H})]_2(\mu\text{-C}_6\text{H}_6)$,⁴ but the latter compound contains a puckered V_2H_2 ring with multiple metal–metal bond character ($\text{V-V} = 2.425$ (1) Å).

1 was isolated in 30% yield, but it is not the only product formed. From the mother liquor another compound can be obtained as orange-brown crystals from pentane or diethyl ether in 10–20% yield. This product is paramagnetic (by NMR), while the IR spectrum indicates the presence of a terminal hydride, with $\nu_{\text{V-H}} = 1560$ cm⁻¹ ($\nu_{\text{V-D}} = 1105$ cm⁻¹ in the D_2 reaction product, $\nu_{\text{V-H}}/\nu_{\text{V-D}} = 1.412$).

X-ray diffraction¹⁰ revealed that the compound is the dimeric $[\text{CpV}(\text{H})\text{dmpe}]_2(\mu\text{-dmpe})$ (**2**, Figure 2). Unfortunately, the positions of the hydride ligands could not be determined unequivocally from the Fourier difference map. However, a substantial asymmetry in the P–V–P angles can be seen ($\text{P}(2)\text{-V-P}(3) = 79.02$ (4)°, $\text{P}(1)\text{-V-P}(2) = 95.00$ (4)°, $\text{P}(1)\text{-V-P}(3) = 117.40$ (5)°), leaving room for the hydride ligand in the sector of the metal between P(1) and P(3).

As no metal–metal contacts are present in the structure, the vanadium atoms should behave like two isolated 17-electron low-spin ($S = 1/2$) d^3 centers. Indeed **2** is ESR active and in solution exhibits a complicated hyperfine splitting pattern with $g = 2.0003$. Replacing the hydride by a deuteride simplifies the solution ESR spectra considerably.¹¹ A comparison of these two ESR spectra indicates that the unpaired electron on each vanadium center in **2** interacts with one hydride ligand, $a(^1\text{H}) = 15.5$ G.¹² Simulation of the spectrum of **2-d**₂ confirmed additional interactions of the unpaired electron with the vanadium nucleus, $a(^{51}\text{V}) = 57.4$ G, and with three nonequivalent phosphorus nuclei, $a(^{31}\text{P}) = 30.5, 33.5$, and 36.0 G. The extensive delocalization of the unpaired electron, resultant in low spin-orbit coupling, is in accordance with the relatively large g value.

The mechanism of formation of **2** is unclear. Apparently, parallel with the reaction sequence in eq 1, some disproportionation occurs according to eq 2, where $[\text{CpVH}]$ may be identified with the poorly soluble gray powder that is also observed in the reaction mixture.

Once formed, the hydrido-bridged dimer **1** is fairly unreactive. For instance, it cannot be broken up into monomeric units by Lewis bases like THF or PMe_3 . Pure **1** does not catalyze the hydrogenation and isomerization of 1-hexene at ambient conditions, whereas **3**, dissolved in 1-hexene under 1 atm of H_2 , does.⁶ Apparently it is necessary to trap the monomeric hydrido species immediately after hydrogenolysis of the V–C bond.

Acknowledgment. We thank Prof. K. G. Caulton for stimulating discussions and H. J. Heeres for practical assistance. This investigation was supported by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for the Advancement of Pure Research (ZWO).

Registry No. **1**, 111635-25-7; **2**, 111717-72-7; **3**, 110174-52-2; 1-hexene, 592-41-6.

Supplementary Material Available: Observed and calculated ESR spectra of **2-d**₂, tables of crystallographic data and positional and thermal parameters, and lists of interatomic distances and angles for **1** and **2** (13 pages); listings of observed and calculated structure factors for **1** and **2** (24 pages). Ordering information is given on any current masthead page.

Stereochemistry of the Thermal Isomerization of Bicyclo[3.2.0]hept-2-ene to Bicyclo[2.2.1]hept-2-ene

John E. Baldwin* and Kevin D. Belfield

Department of Chemistry, Syracuse University
Syracuse, New York 13244
Received September 22, 1987

Sigmatropic [1,3] CHD-carbon migrations of unsubstituted bicyclic hydrocarbons constrained to isomerize suprafacially may provide stereochemical information sensitive to reaction mechanistic course and dynamics. Totally stereoselective inversion is consistent with an allowed, orbital symmetry controlled process,¹ while participation of both inversion and retention modes implicates competitive paths,² either parallel concerted processes with opposite stereochemistry³ or the partitioning of a short-lived biradical intermediate. To know the stereochemical features of CHD-carbon shifts as a function of bicyclic system could provide grounds for distinguishing among these subtle but mechanistically significant alternatives.

Such stereochemical information has been gained for only one system: bicyclo[2.1.1]hex-2-ene labeled with deuterium at C5 gives 6-deuteriobicyclo[3.1.0]hex-2-ene with high but not complete stereoselectivity favoring inversion. At 197 °C in isooctane solution the retention component is 6.9%.²

We have followed the stereochemical course of the bicyclo[3.2.0]hept-2-ene (**1**) to bicyclo[2.2.1]hept-2-ene (**2**) isomerization.⁴ Stereochemical assessment of this isomerization is experimentally challenging since the product **2** reacts further to give cyclopentadiene and ethene. At 276 °C, the maximum concentration of **2** to be expected is only 1–2%.^{4,5}

(9) (a) Cotton, F. A.; Millar, M. *J. Am. Chem. Soc.* **1977**, *99*, 7886. (b) Cotton, F. A.; Walton, R. A. *Multiple Bonds between Metal Atoms*; Wiley: New York, 1982.

(10) **2** crystallizes in the monoclinic space group $P2_1/n$, $a = 8.429$ (2) Å, $b = 24.053$ (4) Å, $c = 9.468$ (5) Å, $\beta = 110.16$ (4)° (130 K), $Z = 2$. Reflections (2548) with $1^\circ \leq \theta \leq 26^\circ$ were considered observed. Nonmetal bound hydrogen atoms were located from the Fourier difference map and constrained at 0.95 Å from their corresponding carbon atoms in the final refinement cycles with use of isotropic fixed thermal parameters: $R = 0.042$, $R_w = 0.055$ ($w = 1$).

(11) In general $a(^2\text{H})$ is 1–2 G, which remains unresolved in the spectrum.

(12) The only $a(^1\text{H})$ observed so far in a vanadium system is 14.6 G in the reaction mixture of Cp_2V with HSnEt_3 : Razuvaev, G. A.; Abakumov, G. A.; Gladyshev, E. N.; Fokeev, A. P.; Cherkasov, V. S. *Dokl. Akad. Nauk SSSR* **1982**, *266*, 135.

(1) Woodward, R. B.; Hoffmann, R. *J. Am. Chem. Soc.* **1965**, *87*, 2511–2513. Woodward, R. B.; Hoffmann, R. *The Conservation of Orbital Symmetry*; Verlag Chemie: Weinheim/Bergs., 1971.

(2) Newman-Evans, R. H.; Carpenter, B. K. *J. Am. Chem. Soc.* **1984**, *106*, 7994–7995. For the stereochemical aspects of the isomerizations shown by the isomeric 5-methylbicyclo[2.1.1]hex-2-enes, see: Roth, W. R.; Friedrich, A. *Tetrahedron Lett.* **1969**, 2607–2610.

(3) Berson, J. A.; Salem, L. *J. Am. Chem. Soc.* **1972**, *94*, 8917–8918. Fernandez, A. *J. Chem. Phys.* **1985**, *82*, 3123–3126.

(4) Cocks, A. T.; Frey, H. M. *J. Chem. Soc. A* **1971**, 2564–2566.