University of Groningen

Clinical and Biological Implications of Mutational Spectrum in Acute Myeloid Leukemia of FAB Subtypes M0 and M1

Cheng, Zhiheng; Dai, Yifeng; Pang, Yifan; Jiao, Yang; Zhao, Hongmian; Wu, Sun; Zhang, Lingxiu; Zhang, Yuan; Wang, Xiufeng; Wang, Lihua

Published in:
Cellular physiology and biochemistry

DOI:
[10.1159/000491065](https://doi.org/10.1159/000491065)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Cheng, Z., Dai, Y., Pang, Y., Jiao, Y., Zhao, H., Wu, S., Zhang, L., Zhang, Y., Wang, X., Wang, L., Ma, D., Qin, T., Hu, N., Zhang, Y., Hu, K., Zhang, Q., Shi, J., & Fu, L. (2018). Clinical and Biological Implications of Mutational Spectrum in Acute Myeloid Leukemia of FAB Subtypes M0 and M1. *Cellular physiology and biochemistry*, 47(5), 1853-1861. <https://doi.org/10.1159/000491065>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Original Paper

Clinical and Biological Implications of Mutational Spectrum in Acute Myeloid Leukemia of FAB Subtypes M0 and M1

Zhiheng Cheng^{a,b} Yifeng Dai^{c,d} Yifan Pang^e Yang Jiao^f Hongmian Zhao^g
Sun Wu^h Lingxiu Zhang^h Yuan Zhang^h Xiufeng Wang^h Lihua Wang^h
Dong Ma^h Tong Qin^g Ning Hu^g Yijie Zhangⁱ Kai Hu^b Qingyi Zhang^{h,j}
Jinlong Shi^{a,k,l} Lin Fu^{b,g,m}

^aTranslational Medicine Center, Huaihe Hospital of Henan University, Kaifeng, ^bDepartment of Hematology and Lymphoma Research Center, Peking University, Third Hospital, Beijing, ^cLaboratory of Environmental Medicine and Developmental Toxicology, Shantou University Medical College, Shantou, China, ^dImmunoendocrinology, Division of Medical Biology, Department of Pathology and Medical Biology, University Medical Center Groningen, Groningen, Netherlands, ^eDepartment of Medicine, William Beaumont Hospital, Royal Oak, USA, ^fLife Sciences Institute and Innovation Center for Cell Signaling Network, Zhejiang University, Hangzhou, ^gDepartment of Hematology, Huaihe Hospital of Henan University, Kaifeng, ^hDepartment of Hematology, The First Affiliated Hospital of Xinxiang Medical University, Weihui, ⁱDepartment of Respiratory, Huaihe Hospital of Henan University, Kaifeng, ^jDepartment of Hematology of Air Force PLA General Hospital, ^kDepartment of Biomedical Engineering, Chinese PLA General Hospital, Beijing, ^lDepartment of Medical Big Data, Chinese PLA General Hospital, Beijing, ^mDepartment of Hematology, The Affiliated Hospital of Xuzhou Medical University, Xuzhou, China

Key Words

Acute myeloid leukemia • M0 and M1 • Next generation sequencing • Mutational spectrum • Prognosis

Abstract

Background/Aims: Acute myeloid leukemia (AML) of French-American-British (FAB) subtypes M0 and M1 are both poorly differentiated AML, but their mutational spectrum and molecular characteristics remain unknown. This study aimed to explore the mutational spectrum and prognostic factors of AML-M0 and M1. **Methods:** Sixty-five AML patients derived from The Cancer Genome Atlas (TCGA) database were enrolled in this study. Whole-genome sequencing was performed to depict the mutational spectrum of each patient. Clinical characteristics at diagnosis, including peripheral blood (PB) white blood cell counts (WBC), blast percentages in PB and bone marrow (BM), FAB subtypes and the frequencies of known recurrent genetic mutations were described. Survival was estimated using the Kaplan-Meier methods and log-rank test. Univariate and multivariate Cox proportional hazard models were constructed

K. Hu, Q. Zhang, J. Shi, L. Fu contributed equally to this work.

Lin Fu, MD. PhD.

Department of Hematology and Lymphoma Research Center, Peking University, Third Hospital Beijing, 100191 (China)
Tel. +86-10-82267650, Fax +86-10-82267650, E-Mail fulin022@126.com

for event-free survival (EFS) and overall survival (OS), using a limited backward elimination procedure. **Results:** Forty-six patients had more than five recurrent genetic mutations. *FLT3* had the highest mutation frequency (n=20, 31%), followed by *NPM1* (n=18, 28%), *DNMT3A* (n=16, 25%), *IDH1* (n=14, 22%), *IDH2* (n=12, 18%), *RUNX1* (n=11, 17%) and *TET2* (n=7, 11%). Univariate analysis showed that age ≥ 60 years and *TP53* mutations had adverse effect on EFS ($P=0.015$, $P=0.036$, respectively) and OS ($P=0.003$, $P=0.004$, respectively), WBC count $\geq 50 \times 10^9/L$ and *FLT3-ITD* negatively affected EFS ($P=0.003$, $P=0.034$, respectively), whereas *NPM1* mutations had favorable effect on OS ($P=0.035$) and allogeneic hematopoietic stem cell transplantation (allo-HSCT) on EFS and OS (all $P < 0.001$). Multivariate analysis suggested that allo-HSCT and *NPM1* mutations were independent favorable prognostic factors for EFS and OS (all $P < 0.05$), WBC count $\geq 50 \times 10^9/L$ was an independent risk factor for EFS ($P=0.002$) and *TP53* mutations for OS ($P=0.043$). **Conclusions:** Our study provided new insights into the mutational spectrum and molecular signatures of AML-M0 and M1. We proposed that *FLT3-ITD*, *NPM1* and *TP53* be identified as markers for risk stratification of AML-M0 and M1. Patients with AML-M0 and M1 would likely benefit from allo-HSCT.

Introduction

© 2018 The Author(s)
Published by S. Karger AG, Basel

Acute myeloid leukemia (AML) is a heterogeneous malignancy characterized by clonal expansion and differentiation arrest of myeloid progenitors in the bone marrow and peripheral blood; historically AML had poor prognosis [1]. Optimizing treatment based on accurate diagnosis and prognostic evaluation in individual patients is particularly important due to disease heterogeneity [2]. Recently, next generation sequencing (NGS) has shown great potential in AML diagnosis and risk stratification because of its massive parallel sequencing ability and high throughput multiplexing capacity [3]. NGS helped characterizing several recurrent somatic mutations in AML, drawing the details of its mutational spectrum [4]. The growing list of mutations involve prognosticators such as *NPM1*, *FLT3-ITD*, *CEBPA*, *DNMT3A*, *IDH1* and *IDH2*, as well as genes implicated in leukemogenesis, such as *EZH2*, *U2AF1*, *SMC1A* and *SMC3* [5]. A recent study analyzed 1,540 AML patients by cytogenetic profiling and targeted resequencing of 111 myeloid cancer genes, the patterns of co-occurrence and mutual exclusivities of genetic changes segregated AML patients into 11 nonoverlapping classes, each with a distinct clinical phenotype and outcome [6]. Another study analyzed the genomes of 200 adult AML patients by NGS, and mutations were divided into nine categories. Almost all AML patients had one or more mutations that fell into the nine categories, and a complex interplay of genetic alterations was found [5].

Several decades ago, in order to provide objectivity in the diagnosis of AML that would facilitate comparisons between series of cases, the French-American-British (FAB) Cooperative Group developed a classification system based on conventional morphologic and cytochemical characteristics and divided AML into FAB subtypes (M0-M7) [7], with AML-M0 and M1 being the poorly differentiated subtypes. Although advances in identification of prognostic genetic alterations have facilitated detailed risk stratification [8], currently no research has addressed the mutational spectrum of AML-M0 and M1. It's unclear whether they differ in mutational spectrum and how genetic signatures influence their prognosis. We intended to describe the clinical and molecular prognostic factors for the development of optimal and individualized therapy for AML-M0 and M1 patients.

Materials and Methods

Patients

Sixty-five AML patients derived from The Cancer Genome Atlas (TCGA) database (<https://cancergenome.nih.gov/>) were enrolled in this study [5], including 19 AML-M0 and 46 AML-M1 patients. Poor-risk patients each underwent allogeneic hematopoietic stem cell transplantation (allo-HSCT) if

there was no contraindication and a matched donor was available. Many intermediate risk patients also underwent allo-HSCT. The number of patients receiving allo-HSCT was 37 and the rest 28 had chemotherapy only. Whole-genome sequencing was performed to depict the mutational spectrum of each patient. Clinical characteristics at diagnosis, including peripheral blood (PB) white blood cell counts (WBC), blast percentages in PB and bone marrow (BM), French-American-British (FAB) subtypes and the frequencies of known recurrent genetic mutations were described. Detailed descriptions of clinical and molecular characteristics were publicly accessible from the TCGA website. Event-free survival (EFS) and overall survival (OS) were the primary endpoints of this study. EFS was defined as the time from diagnosis to the first event including relapse, death, absence of complete remission or the last follow up. OS was defined as the time from diagnosis to death from any cause or the last follow-up. All patients provided informed consent, and the study protocol was approved by the Washington University Human Studies Committee.

Statistical Analysis

The clinical and molecular characteristics of patients were summarized using descriptive statistics. Data sets were described with median and/or range. Survival was estimated using the Kaplan-Meier method and the log-rank test. Univariate Cox proportional hazards models were used to identify clinical and molecular variables associated with survival. Multivariate proportional hazards models were constructed for EFS and OS, using a limited backward elimination procedure. $P < 0.05$ was considered statistically significant for all analyses. All statistical tests were two-sided and were performed by SPSS software 20.0 and GraphPad Prism software 5.0.

Results

Demographic and biological characteristics of the patients

The demographic and biological characteristics of the patients were summarized in Table 1. Median age was 58 (range 18-88) years, with 31 cases older than 60. Thirty-seven cases were men. Nineteen patients were AML-M0 and 46 were AML-M1. The median WBC count at diagnosis was $19.8 \times 10^9/L$, and in 16 cases it was $\geq 50 \times 10^9/L$. Forty-eight patients had BM blast percentage more than 70% and 28 had PB blasts more than 70%. Thirty-four patients had abnormal karyotypes. Sixty patients had intermediate or poor risk AML. Chemotherapy was differed in two patients due to old age and poor functional status. Thirty-seven patients received HSCT, of which 24 cases achieved complete remission. Forty-six patients had more than five recurrent genetic mutations. *FLT3* had the highest mutation frequency (n=20, 31%), followed by *NPM1* (n=18, 28%), *DNMT3A* (n=16, 25%), *IDH1* (n=14, 22%), *IDH2* (n=12, 18%), *RUNX1* (n=11, 17%) and *TET2* (n=7, 11%) (Fig. 1).

Table 1. Clinical and molecular characteristics of the patients
Abbreviations: FAB, French American British; WBC, white blood cell; BM, bone marrow; PB, peripheral blood; HSCT, hematopoietic stem cell transplantation; MUD, matched unrelated donor; Allo, allogeneic; Auto, autologous

Characteristics	Median (range) or N/%
Age (years)	58 (18-88)
<60	34/52.3
≥60	31/47.7
Gender	
Male	37/56.9
Female	28/43.1
Race	
Caucasian	44/67.7
Others	21/32.3
FAB subtypes	
M0	19/29.2
M1	46/70.8
WBC count/ $\times 10^9/L$	19.8 (0.7-297.4)
<50	49/75.4
≥50	16/24.6
BM blasts/%	81 (32-100)
<70	17/26.2
≥70	48/73.8
PB blasts/%	55 (0-98)
<70	37/56.9
≥70	28/43.1
Karyotype	
Normal	30/46.9
Abnormal	34/53.1
Risk	
Good	4/6.2
Intermediate	40/62.5
Poor	20/31.3
Recurrent gene mutations	6 (0-12)
<5	19/29.2
≥5	46/70.8
MLL-PTD	
Positive	4/6.2
Negative	61/93.8
FLT3	
FLT3-ITD	15/23.1
FLT3-TKD	5/7.7
Wild type	45/69.2
NPM1	
W288	18/27.7
Wild type	47/72.3
DNMT3A	
R882	6/9.2
Non-R882 mutations	10/15.4
Wild type	49/75.4
IDH1	
R132	14/21.5
Wild type	51/78.5
IDH2	
R140	9/13.9
R172	3/4.6
Wild type	53/81.5
RUNX1	
Mutation	11/16.9
Wild type	54/83.1
TET2	
Mutation	7/10.8
Wild type	58/89.2
CEBPA	
Single-mutation	6/9.2
Wild type	59/90.8
TP53	
Mutation	5/7.7
Wild type	60/92.3
PTPN11	
Mutation	5/7.7
Wild type	60/92.3
MT-CO2	
Mutation	5/7.7
Wild type	60/92.3
ASXL1	
Mutation	5/7.7
Wild type	60/92.3
NRAS	
Mutation	4/6.2
Wild type	61/93.8
KRAS	
Mutation	4/6.2
Wild type	61/93.8
TTN	
Mutation	4/6.2
Wild type	61/93.8
STAG2	
Mutation	4/6.2
Wild type	61/93.8
HSCT	
MUD	21/56.8
Sib alb	12/32.4
Auto	4/10.8

Fig. 1. The mutational spectrum of the patients. FLT3 had the highest mutation frequency (n=20, 31%), followed by NPM1 (n=18, 28%), DNMT3A (n=16, 25%), IDH1 (n=14, 22%), IDH2 (n=12, 18%), RUNX1 (n=11, 17%) and TET2 (n=7, 11%). In addition, CEBPA, TP53, PTPN11, MT-CO2, ASXL1, NRAS, KRAS, TTN and STAG2 also had more than 5% mutation frequency.

Fig. 2. Kaplan-Meier curves of EFS and OS based on clinical parameters. (A, B) Patients older than 60 years had shorter EFS and OS than those younger than 60 years. (C) Patients with WBC count $\geq 50 \times 10^9/L$ had shorter EFS than those with WBC count $< 50 \times 10^9/L$. (D) WBC count had no effect on OS.

Comparison of EFS and OS between different clinical and molecular characteristic groups

EFS and OS of different age (≥ 60 vs. < 60 years), WBC count (≥ 50 vs. $< 50 \times 10^9/L$), BM blasts ($\geq 70\%$ vs. $< 70\%$), PB blasts ($\geq 70\%$ vs. $< 70\%$), allo-HSCT (yes vs. no), FLT3-ITD (positive vs. negative), and the mutation status of other common AML mutations (NPM1, DNMT3A, IDH1, IDH2, RUNX1, CEBPA, TP53, PTPN11, MT-CO2, ASXL1, NRAS, KRAS, TTN and STAG2, mutated vs. wild type), were compared with the Kaplan-Meier method and the log-rank test, as listed in Table 2. Older patients (age ≥ 60) had shorter EFS and OS ($P=0.013$, $P=0.002$, respectively, Fig. 2A and 2B). WBC count $\geq 50 \times 10^9/L$ negatively affected EFS ($P=0.002$, Fig. 2C). Positive FLT3-ITD was associated with shorter EFS ($P=0.031$, Fig. 3A). Patients with TP53 mutations had shorter EFS and OS ($P=0.028$, $P=0.002$, respectively,

Table 2. Comparison of EFS and OS between different clinical and molecular characteristic groups. Abbreviation: EFS, event-free survival; OS, overall survival; WBC, white blood cell; BM, bone marrow; PB, peripheral blood; Allo-HSCT, allogeneic hematopoietic stem cell transplantation

Variables	EFS		OS	
	χ^2	P-value	χ^2	P-value
Age (≥ 60 vs. < 60 years)	6.181	0.013	9.601	0.002
WBC (≥ 50 vs. $< 50 \times 10^9/L$)	9.407	0.002	2.318	0.128
BM blasts ($\geq 70\%$ vs. $< 70\%$)	0.030	0.863	0.229	0.632
PB blasts ($\geq 70\%$ vs. $< 70\%$)	0.221	0.638	0.358	0.550
FLT3-ITD (positive vs. negative)	4.672	0.031	1.524	0.217
NPM1 (mutated vs. wild type)	1.594	0.207	4.609	0.032
DNMT3A (mutated vs. wild type)	0.003	0.955	0.611	0.434
IDH1 (mutated vs. wild type)	2.953	0.086	3.449	0.063
IDH2 (mutated vs. wild type)	0.632	0.427	0.065	0.798
RUNX1 (mutated vs. wild type)	1.049	0.306	3.779	0.052
TET2 (mutated vs. wild type)	0.070	0.792	0.017	0.897
CEBPA (mutated vs. wild type)	0.444	0.505	0.368	0.544
TP53 (mutated vs. wild type)	4.833	0.028	9.870	0.002
PTPN11 (mutated vs. wild type)	0.001	0.973	0.095	0.757
MT-CO2 (mutated vs. wild type)	0.048	0.827	0.416	0.519
ASXL1 (mutated vs. wild type)	0.002	0.960	0.987	0.321
NRAS (mutated vs. wild type)	0.006	0.939	0.009	0.922
KRAS (mutated vs. wild type)	0.056	0.814	0.179	0.673
TTN (mutated vs. wild type)	0.289	0.591	0.607	0.436
STAG2 (mutated vs. wild type)	0.459	0.498	1.011	0.315
Allo-HSCT (yes vs. no)	14.048	< 0.001	14.656	< 0.001

Fig. 3. Kaplan-Meier curves of EFS and OS based on mutated genes. (A) Patients with FLT3-ITD had shorter EFS than negative group. (B) FLT3-ITD had no effect on OS. (C) NPM1 mutations had no effect on EFS. (D) Patients with NPM1 mutations had longer OS than wild type group. (E, F) Patients with TP53 mutations had shorter EFS and OS than wild type group.

Fig. 4. Kaplan-Meier curves of EFS and OS based on allo-HSCT. (A, B) Patients underwent allo-HSCT had longer EFS and OS than those without allo-HSCT.

Fig. 3E and 3F). Patients with *NPM1* mutations had longer OS ($P=0.032$, Fig. 3D). Furthermore, patients received allo-HSCT had longer EFS and OS ($P<0.001$, $P<0.001$, respectively, Fig. 4A and 4B). Other variables did not demonstrate effect on EFS or OS.

Univariate and multivariate analyses of possible prognostic factors

To further explore the prognostic significance of the aforementioned factors, we did univariate analysis and selected factors that had statistical significance to construct the multivariate COX regression model for EFS and OS. Univariate analysis showed that age ≥ 60 years was an unfavorable factor for EFS

Table 3. Univariate analysis for EFS and OS. Abbreviation: EFS, event-free survival; OS, overall survival; HR, hazard ratio; CI, confidence interval; WBC, white blood cell; BM, bone marrow; PB, peripheral blood; Allo-HSCT, allogeneic hematopoietic stem cell transplantation

Variables	EFS		OS	
	HR (95%CI)	P-value	HR (95%CI)	P-value
Age (≥ 60 vs. <60 years)	0.513 (0.300-0.878)	0.015	0.424 (0.243-0.741)	0.003
WBC (≥ 50 vs. $<50 \times 10^9/L$)	0.389 (0.208-0.728)	0.003	0.630 (0.346-1.148)	0.131
BM blasts ($\geq 70\%$ vs. $<70\%$)	0.949 (0.523-1.721)	0.864	1.156 (0.637-2.098)	0.633
PB blasts ($\geq 70\%$ vs. $<70\%$)	0.879 (0.513-1.506)	0.639	1.179 (0.686-2.026)	0.550
FLT3-ITD (positive vs. negative)	1.967 (1.051-3.684)	0.034	1.471 (0.794-2.725)	0.220
NPM1 (mutated vs. wild type)	0.671 (0.358-1.255)	0.211	0.506 (0.269-0.953)	0.035
DNMT3A (mutated vs. wild type)	1.018 (0.553-1.872)	0.955	1.279 (0.689-2.377)	0.436
IDH1 (mutated vs. wild type)	0.552 (0.276-1.100)	0.091	0.525 (0.263-1.048)	0.068
IDH2 (mutated vs. wild type)	0.758 (0.381-1.509)	0.430	1.095 (0.545-2.198)	0.799
RUNX1 (mutated vs. wild type)	1.412 (0.726-2.746)	0.310	1.935 (0.982-3.812)	0.056
TET2 (mutated vs. wild type)	1.121 (0.479-2.624)	0.792	0.946 (0.404-2.211)	0.897
CEBPA (mutated vs. wild type)	1.365 (0.542-3.440)	0.509	1.330 (0.527-3.356)	0.546
TP53 (mutated vs. wild type)	2.727 (1.070-6.951)	0.036	4.166 (1.583-10.958)	0.004
PTPN11 (mutated vs. wild type)	1.016 (0.404-2.556)	0.973	1.156 (0.459-2.913)	0.758
MT-CO2 (mutated vs. wild type)	1.120 (0.404-3.110)	0.827	1.398 (0.545-3.888)	0.521
ASXL1 (mutated vs. wild type)	1.024 (0.406-2.584)	0.960	1.597 (0.629-4.057)	0.325
NRAS (mutated vs. wild type)	1.047 (0.324-3.378)	0.939	0.943 (0.292-3.049)	0.922
KRAS (mutated vs. wild type)	1.130 (0.407-3.135)	0.814	1.247 (0.447-3.484)	0.673
TET2 (mutated vs. wild type)	1.322 (0.475-3.680)	0.594	1.503 (0.535-4.222)	0.439
STAG2 (mutated vs. wild type)	0.671 (0.209-2.155)	0.503	0.554 (0.173-1.781)	0.322
Allo-HSCT (yes vs. no)	0.365 (0.211-0.631)	<0.001	0.358 (0.208-0.618)	<0.001

and OS ($P=0.015$, $P=0.003$, respectively), as well as *TP53* mutations ($P=0.036$, $P=0.004$ for EFS and OS, respectively), WBC count $\geq 50 \times 10^9/L$ and *FLT3-ITD* negatively affected EFS ($P=0.003$, $P=0.034$, respectively), whereas *NPM1* mutations favorably affected OS ($P=0.035$), and allo-HSCT was a favorable factor for EFS and OS (all $P<0.001$) (Table 3). Multivariate analysis suggested that allo-HSCT was an independent favorable factor for EFS (HR: 0.358, 95% CI: 0.201-0.640, $P=0.001$), the effect was more prominent after adjusting for *NPM1* mutation status ($P=0.025$) and WBC count ($P=0.002$). It was also an independent favorable factor for OS (HR: 0.374, 95% CI: 0.209-0.669, $P=0.001$), with more profound effect after adjusting for *NPM1* ($P=0.002$) and *TP53* mutation status ($P=0.043$) (Table 4).

Table 4. Multivariate analysis for EFS and OS. Abbreviation: EFS, event-free survival; OS, overall survival; HR, hazard ratio; CI, confidence interval; WBC, white blood cell; Allo-HSCT, allogeneic hematopoietic stem cell transplantation

Variables	EFS		OS	
	HR (95%CI)	P-value	HR (95%CI)	P-value
Age (≥ 60 vs. <60 years)	0.704 (0.386-1.286)	0.254	0.692 (0.371-1.290)	0.246
WBC (≥ 50 vs. $<50 \times 10^9/L$)	0.321 (0.156-0.664)	0.002	0.522 (0.240-1.134)	0.101
<i>FLT3-ITD</i> (positive vs. negative)	2.000 (0.949-4.217)	0.069	1.719 (0.742-3.986)	0.207
<i>NPM1</i> (mutated vs. wild type)	0.439 (0.214-0.900)	0.025	0.291 (0.133-0.637)	0.002
<i>TP53</i> (mutated vs. wild type)	2.654 (0.956-7.372)	0.061	2.974 (1.034-8.559)	0.043
Allo-HSCT (yes vs. no)	0.358 (0.201-0.640)	0.001	0.374 (0.209-0.669)	0.001

Discussion

AML is a genetically heterogeneous disease resulting from complex interactions among different leukemogenic pathways, so integrated mutational analysis is highly valuable for evaluation [5, 9]. Formerly, the mutational spectrum of AML-M0 and M1 was unclear. In this study, we found that *FLT3-ITD*, *NPM1*, *DNMT3A*, *IDH1*, *IDH2*, *RUNX1* and *TET2* were mutated in more than 10% of all patients with *FLT3-ITD* exhibiting the highest frequency; *CEBPA*, *TP53*, *PTPN11*, *MT-CO2*, *ASXL1*, *NRAS*, *KRAS*, *TTN* and *STAG2* also had more than 5% mutation frequency. This was different from previous reports which showed that *CEBPA*, *NPM1*, *DNMT3A*, *FLT3-ITD*, *NRAS*, *IDH2* and *WT1* were mutated in more than 10% and *CEBPA* mutations were more frequent in intermediate-risk AML [10, 11]. The reported frequency of *CEBPA* mutations in cytogenetically normal AML (CN-AML) was also higher, about 35% [12]. The discrepancy suggested that poorly differentiated AML might have a distinct mutational spectrum.

In uni- and multivariate analyses, we found that age ≥ 60 years was an adverse factor for EFS and OS, which was consistent with the fact that AML patients younger than 60 years had improved prognosis and approximately 35-40% of them would get cured [13]. WBC count $\geq 50 \times 10^9/L$ was also related to shorter EFS and OS, which was consistent with previous finding that WBC count had a significant impact on complete remission rate, EFS and OS in AML patients [14].

FLT3 is a class III family receptor tyrosine kinase that acts as a cytokine receptor for the *FLT3* ligand. *FLT3* is strongly expressed in hematopoietic stem cells with important roles in cell survival and proliferation [15]. *FLT3-ITD* was among the most frequent mutations observed in AML, it could activate *FLT3* signaling, promoting blast proliferation [16]. Furthermore, *FLT3-ITD* was associated with increased risk of relapse in AML [17]. *NPM1* is involved in numerous cellular functions, such as ribosome biogenesis, DNA repair and regulation of apoptosis. *NPM1* mutations were among the most common genetic changes in AML, especially in CN-AML [18]. In the absence of *FLT3-ITD*, *NPM1* mutations were associated with improved outcomes for CN-AML patients. *NPM1* mutations have been associated with chemosensitivity to intensive chemotherapy in both young and old patients, which might account for improved outcomes [19]. *NPM1* mutations were also associated with other recurrent genetic abnormalities, such as *DNMT3A*, *FLT3-ITD* and *IDH* mutations [20]. The pattern of co-mutations largely shaped clinical outcomes. *TP53* mutations were rare in patients lacking chromosomal deletions, and it conferred an adverse prognosis with documented chemoresistance [21, 22]. *TP53* mutations might be responsible for the poor prognosis of complex karyotype AML [23]. Our results showed that *FLT3-ITD* and *TP53* mutations were associated

with shorter EFS and OS, while *NPM1* mutations were associated with favorable prognosis, consistent with previous results. Furthermore, studies indicated that allo-HSCT could lead to better clinical outcomes for patients with unfavorable-risk cytogenetics in the first complete remission [24]. The favorable effect of allo-HSCT was also replicated in our univariate analysis, and the effect was still exist after adjusting for potential confounding factors (age, WBC, *FLT3-ITD*, *NPM1* and *TP53*).

Several limitations need to be acknowledge. First, due to the limited number of our cases, we didn't stratify data more precisely based on factors that could affect the prognosis. So, our results didn't fully account for the effect of mutational spectrum and clinical data on prognosis. Second, our study was a retrospective study which could suffer from inherited biases as opposed to prospective studies.

Conclusion

In summary, we conducted a TCGA database-derived analysis on the mutational profiles and prognosis of AML-M0 and M1 and compared our findings with previous studies. Our study provided new insights into the clinical and biological implications of mutational spectrum in AML-M0 and M1. *FLT3-ITD*, *NPM1* and *TP53* could be incorporated into AML-M0 and M1 risk stratification and these patients would likely benefit from allo-HSCT.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (81500118, 61501519), the China Postdoctoral Science Foundation funded project (project No.2016M600443), Jiangsu Province Postdoctoral Science Foundation funded project (project No.1701184B) and PLAGH project of Medical Big Data (project No.2016MBD-025).

Disclosure Statement

The authors declare to have no conflict of interests.

References

- Schlenk RF, Döhner H: Genomic applications in the clinic: use in treatment paradigm of acute myeloid leukemia. *Hematology Am Soc Hematol Educ Program* 2013;2013:324-330.
- Cazzola M, Della Porta MG, Travaglino E, Malcovati L: Classification and prognostic evaluation of myelodysplastic syndromes. *Semin Oncol* 2011;38:627-634.
- Ding L, Ley TJ, Larson DE, Miller CA, Koboldt DC, Welch JS, Ritchey JK, Young MA, Lamprecht T, McLellan MD, McMichael JF, Wallis JW, Lu C, Shen D, Harris CC, Dooling DJ, Fulton RS, Fulton LL, Chen K, Schmidt H, Kalicki-Veizer J, Magrini VJ, Cook L, McGrath SD, Vickery TL, Wendl MC, Heath S, Watson MA, Link DC, Tomasson MH, Shannon WD, Payton JE, Kulkarni S, Westervelt P, Walter MJ, Graubert TA, Mardis ER, Wilson RK, DiPersio JF: Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature* 2012;481:506-510.
- Garraway LA: Genomics-driven oncology: framework for an emerging paradigm. *J Clin Oncol* 2013;31:1806-1814.
- Ley TJ, Miller C, Ding L, Raphael BJ, Mungall AJ, Robertson A, Hoadley K, Triche TJ Jr, Laird PW, Baty JD, Fulton LL, Fulton R, Heath SE, Kalicki-Veizer J, Kandoth C, Klco JM, Koboldt DC, Kanchi KL, Kulkarni S, Lamprecht TL et al. *N Engl J Med* 2013;368:2059-2074.

- 6 Papaemmanuil E, Gerstung M, Bullinger L, Gaidzik VI, Paschka P, Roberts ND, Potter NE, Heuser M, Thol F, Bolli N, Gundem G, Van Loo P, Martincorena I, Ganly P, Mudie L, McLaren S, O'Meara S, Raine K, Jones DR, Teague JW, Butler AP, Greaves MF, Ganser A, Döhner K, Schlenk RF, Döhner H, Campbell PJ: Genomic Classification and Prognosis in Acute Myeloid Leukemia. *N Engl J Med* 2016;374:2209-2221.
- 7 Bennett JM, Catovsky D, Daniel MT, Flandrin G, Galton DA, Gralnick HR, Sultan C: Proposed revised criteria for the classification of acute myeloid leukemia. A report of the French-American-British Cooperative Group. *Ann Intern Med* 1985;103:620-625.
- 8 Patel JP, Gönen M, Figueroa ME, Fernandez H, Sun Z, Racevskis J, Van Vlierberghe P, Dolgalev I, Thomas S, Aminova O, Huberman K, Cheng J, Viale A, Socci ND, Heguy A, Cherry A, Vance G, Higgins RR, Ketterling RP, Gallagher RE, Litzow M, van den Brink MR, Lazarus HM, Rowe JM, Luger S, Ferrando A, Paietta E, Tallman MS, Melnick A, Abdel-Wahab O, Levine RL: Prognostic Relevance of Integrated Genetic Profiling in Acute Myeloid Leukemia. *N Engl J Med* 2012;366:1079-1089.
- 9 Estey EH: Acute myeloid leukemia: 2013 update on risk-stratification and management. *Am J Hematol* 2013;88:318-327.
- 10 Naoe T, Kiyoi H: Gene mutations of acute myeloid leukemia in the genome era. *Int J Hematol* 2013;97:165-174.
- 11 Grossmann V, Schnittger S, Kohlmann A, Eder C, Roller A, Dicker F, Schmid C, Wendtner CM, Staib P, Serve H, Kreuzer KA, Kern W, Haferlach T, Haferlach C: A novel hierarchical prognostic model of AML solely based on molecular mutations. *Blood* 2012;120:2963-2972.
- 12 Lin LI, Chen CY, Lin DT, Tsay W, Tang JL, Yeh YC, Shen HL, Su FH, Yao M, Huang SY, Tien HF: Characterization of CEBPA mutations in acute myeloid leukemia: most patients with CEBPA mutations have biallelic mutations and show a distinct immunophenotype of the leukemic cells. *Clin Cancer Res* 2005;11:1372-1379.
- 13 Döhner H, Weisdorf DJ, Bloomfield CD: Acute myeloid leukemia. *N Engl J Med* 2015;373:1136-1152.
- 14 de Jonge HJ, Valk PJ, de Bont ES, Schuringa JJ, Ossenkoppele G, Vellenga E, Huls G: Prognostic impact of white blood cell count in intermediate risk acute myeloid leukemia: relevance of mutated NPM1 and FLT3-ITD. *Haematologica* 2011;96:1310-1317.
- 15 Saultz JN, Garzon R: Acute Myeloid Leukemia: A Concise Review. *J Clin Med* 2016;5:E33.
- 16 Zhu GZ, Yang YL, Zhang YJ, Liu W, Li MP, Zeng WJ, Zhao XL, Chen XP: High Expression of AHSP, EPB42, GYPC and HEMGN Predicts Favorable Prognosis in FLT3-ITD-Negative Acute Myeloid Leukemia. *Cell Physiol Biochem* 2017;42:1973-1984.
- 17 Gale RE, Green C, Allen C, Mead AJ, Burnett AK, Hills RK, Linch DC: The impact of FLT3 internal tandem duplication mutant level, number, size, and interaction with NPM1 mutations in a large cohort of young adult patients with acute myeloid leukemia. *Blood* 2008;111:2776-2784.
- 18 Falini B, Nicoletti I, Martelli M F, Mecucci C: Acute myeloid leukemia carrying cytoplasmic/mutated nucleophosmin (NPMc+ AML): biologic and clinical features. *Blood* 2007;109:874-885.
- 19 Döhner K, Schlenk RF, Habdank M, Scholl C, Rücker FG, Corbacioglu A, Bullinger L, Fröhling S, Döhner H: Mutant nucleophosmin (NPM1) predicts favorable prognosis in younger adults with acute myeloid leukemia and normal cytogenetics: interaction with other gene mutations. *Blood* 2005;106:3740-3746.
- 20 Marcucci G, Metzeler KH, Schwind S, Becker H, Maharry K, Mrózek K, Radmacher MD, Kohlschmidt J, Nicolet D, Whitman SP, Wu YZ, Powell BL, Carter TH, Kolitz JE, Wetzler M, Carroll AJ, Baer MR, Moore JO, Caligiuri MA, Larson RA, Bloomfield CD: Age-related prognostic impact of different types of DNMT3A mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J Clin Oncol* 2012;30:742-750.
- 21 Haferlach C, Dicker F, Herholz H, Schnittger S, Kern W, Haferlach T: Mutations of the TP53 gene in acute myeloid leukemia are strongly associated with a complex aberrant karyotype. *Leukemia* 2008;22:1539-1541.
- 22 Kadia TM, Jain P, Ravandi F, Garcia-Manero G, Andreef M, Takahashi K, Borthakur G, Jabbour E, Konopleva M, Daver NG, Dinardo C, Pierce S, Kanagal-Shamanna R, Patel K, Estrov Z, Cortes J, Kantarjian HM: TP53 mutations in newly diagnosed acute myeloid leukemia: Clinicomolecular characteristics, response to therapy, and outcomes. *Cancer* 2016;122:3484-3491.

- 23 Rücker FG, Schlenk RF, Bullinger L, Kayser S, Teleanu V, Kett H, Haddank M, Kugler CM, Holzmann K, Gaidzik VI, Paschka P, Held G, von Lilienfeld-Toal M, Lübbert M, Fröhling S, Zenz T, Krauter J, Schlegelberger B, Ganser A, Lichter P, Döhner K, Döhner H: TP53 alterations in acute myeloid leukemia with complex karyotype correlate with specific copy number alterations, monosomal karyotype, and dismal outcome. *Blood* 2012;119:2114-2121.
- 24 Schetelig J, Schaich M, Schäfer-Eckart K, Hänel M, Aulitzky WE, Einsele H, Schmitz N, Rösler W, Stelljes M, Baldus CD, Ho AD, Neubauer A, Serve H, Mayer J, Berdel WE, Mohr B, Oelschlägel U, Parmentier S, Röhl C, Kramer M, Platzbecker U, Illmer T, Thiede C, Bornhäuser M, Ehninger G: Hematopoietic cell transplantation in patients with intermediate and high-risk AML - results from the randomized study alliance leukemia (SAL) AML 2003 trial. *Leukemia* 2015;29:1060-1068.