University of Groningen

Differential control of ageing and lifespan by isoforms and splice variants across the mTOR network

Razquin Navas, Patricia; Thedieck, Kathrin

Published in:
Essays in biochemistry

DOI:
[10.1042/EBC20160086](https://doi.org/10.1042/EBC20160086)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Razquin Navas, P., & Thedieck, K. (2017). Differential control of ageing and lifespan by isoforms and splice variants across the mTOR network. *Essays in biochemistry*, 61(3), 349-368.
<https://doi.org/10.1042/EBC20160086>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Differential control of ageing and lifespan by isoforms and splice variants across the mTOR network

Patricia Razquin Navas¹ and Kathrin Thedieck^{1,*}

¹Lab for Metabolic Signalling, European Medical School Oldenburg-Groningen; Department of Pediatrics, Section Systems Medicine of Metabolism & Signalling, University of Groningen, University Medical Center Groningen (UMCG), 9713 AV Groningen, The Netherlands; Department of Neuroscience, School of Medicine and Health Sciences, Carl von Ossietzky University Oldenburg, 26111 Oldenburg, Germany

*to whom correspondence should be addressed:

k.thedieck@umcg.nl ; kathrin.thedieck@uni-oldenburg.de ; kathrin.thedieck@metabolic-signaling.eu

Abstract

Ageing can be defined as the gradual deterioration of physiological functions, increasing the incidence of age-related disorders and the probability of death. Therefore, the term of ageing does not only reflect the lifespan of an organism but also refers to progressive functional impairment and disease.

The nutrient-sensing kinase mTOR (mammalian target of rapamycin) is a major determinant of ageing. mTOR promotes cell growth and controls central metabolic pathways including protein biosynthesis, autophagy, and glucose and lipid homeostasis. The concept that mTOR has a crucial role in ageing is supported by numerous reports on the lifespan prolonging effects of the mTOR inhibitor rapamycin in invertebrate and vertebrate model organisms. Interestingly, rapamycin not only increases lifespan but also delays the appearance of age-related metabolic phenotypes and diseases. Dietary restriction increases lifespan and delays ageing phenotypes as well, and mTOR has been assigned a major role in this process. This may suggest a causal relationship between the lifespan of an organism and its metabolic phenotype.

More than 25 years after mTOR's discovery, a wealth of metabolic and ageing-related effects have been reported. In this review we cover the current view on the contribution of the different elements of the mTOR pathway to lifespan and age-related metabolic impairment. We specifically focus on distinct roles of isoforms and splice variants across the mTOR network. The comprehensive analysis of mouse knockout studies targeting these variants does not support a tight correlation between lifespan prolongation and improved metabolic phenotypes and questions the strict causal relationship between them.

Summary

- mTOR inhibition prolongs lifespan and delays the appearance of age-related metabolic diseases, suggesting that mTOR is a major determinant of ageing and metabolic balance.
- The mTOR network in higher organisms harbours many different isoforms and splice variants, which exert complex effects on lifespan and metabolism.
- A comprehensive analysis of mouse knockout studies targeting these variants does not support a tight causal relationship between an improved metabolic phenotype and lifespan prolongation.

Abbreviation list

4E-BP1	eukaryotic translation initiation factor 4E-binding protein 1
4E-BP2	eukaryotic translation initiation factor 4E-binding protein 2
4E-BP3	eukaryotic translation initiation factor 4E-binding protein 3
aa	amino acids
AGC	protein kinase A/ protein kinase G/ protein kinase C
AMPK	AMP-activated protein kinase
ApoE	apolipoprotein E
Atg13	autophagy related 13
BAD	BCL2 associated agonist of cell death
BD	binding domain
<i>C. elegans</i>	<i>Caenorhabditis elegans</i>
CaMKK β	Ca ²⁺ /calmodulin-dependent protein kinase kinase-beta
Cdc42	cell division cycle 42
CTD	C-terminal domain
CREM	cAMP responsive element modulator
<i>D. melanogaster</i>	<i>Drosophila melanogaster</i>
d4E-BP	eukaryotic translation initiation factor 4E-binding in <i>Drosophila melanogaster</i>
DEPTOR	DEP domain containing mTOR-interacting protein
dS6K	ribosomal S6 kinase in <i>Drosophila melanogaster</i>
dTOR	TOR in <i>Drosophila melanogaster</i>
eEF2K	eukaryotic elongation factor-2 kinase
eIF4B	eukaryotic translation initiation factor 4B
eIF4E	eukaryotic translation initiation factor 4E
EMT	endothelial-mesenchymal transition
Er α	estrogen receptor alpha
FAT	FRAP-ATM-TTRAP domain
FATC	FAT-carboxy terminal domain
FoxO	Forkhead box O
GAP	GTPase-activating protein

GLUT4	glucose transporter type 4
Grb2	growth factor receptor-bound protein 2
Grb10	growth factor receptor-bound protein 10
GSK-3	glycogen synthase kinase 3
GWAS	genome-wide association studies
HEAT	huntingtin-elongation factor 3-regulatory subunit A of PP2A-TOR1 repeats
HM	hydrophobic motif
IGF-1	insulin like growth factor 1
IGF1-R	insulin like growth factor 1 receptor
InR	insulin receptor
IRS1	insulin receptor substrate 1
IRS2	insulin receptor substrate 2
IRS3	insulin receptor substrate 3
IRS4	insulin receptor substrate 4
KD	kinase domain
KRLB	kinase regulatory loop binding
MAPK	mitogen-activated protein kinase
Mdm2	Mdm2 proto-oncogene
mLST8	mammalian lethal with SEC13 protein 8
mSin1	mammalian stress-activated protein kinase interacting protein 1
mTOR	mammalian target of rapamycin
mTORC1	mTOR complex 1
NLS	nuclear localization sequence
NTD	N-terminal domain
PDK1	3-phosphoinositide-dependent kinase-1
PH	pleckstrin homology
PHLPP1	PH domain and leucine-rich repeat protein phosphatase 1
PI3K	phosphatidylinositol 3-kinase
PIP2	phosphatidylinositol-3,4-biphosphate
PIP3	phosphatidylinositol-3,4,5-triphosphate
PKC	protein kinase C
PRAS40	proline-rich Akt substrate of 40 kDa
Protor-1	protein observed with rictor 1/ proline-rich protein 5
Protor-2	protein observed with rictor 1/ proline-rich protein 5-like
PTEN	phosphatase and tensin homolog
Rac-1	ras-related C3 botulinum toxin substrate 1
Rag	ras-related GTP-binding proteins
Raptor	regulatory associated protein of mTORC1
RBD	ras-binding domain
Rheb	ras-homologue-enriched-in-brain
Rictor	rapamycin-insensitive companion of mTOR
RNC	raptor N-terminal conserved
ROR γ	RAR-related orphan receptor gamma
S6	ribosomal protein S6
S6K1	ribosomal protein S6 kinase 1
S6K2	ribosomal protein S6 kinase 2
<i>S. cerevisiae</i>	<i>Saccharomyces cerevisiae</i>

SGK	serum and glucocorticoid-regulated kinase
SHP2	src homology 2 domain-containing phosphatase
SIRT	sirtuins
TBC1D7	Tre2-Bub2-Cdc16 1 domain family member 7
Tel2	telomere maintenance 2
TFE3	transcription factor binding to immunoglobulin heavy constant mu enhancer 3
Tti1	TELO2 interacting protein 1
TOR	target of rapamycin
TOS	TOR signalling sequence
TSC	tuberous sclerosis complex
TTP	tristetraprolin
ULK1	unc-51 like autophagy activating kinase 1
ULK2	unc-51 like autophagy activating kinase 2
YY1	yin yang 1

1. The mTOR pathway in mammals

Target of rapamycin (TOR) was discovered in 1991 in the budding yeast *Saccharomyces cerevisiae* (*S. cerevisiae*) (1) and is structurally and functionally conserved in all eukaryotes, including mammals where it is called mammalian TOR (mTOR). mTOR is a serine/threonine kinase that functions as a master regulator of cellular growth and metabolism. mTOR forms two structurally and functionally distinct complexes, mTOR complex 1 (mTORC1) and mTORC2 (2-4). mTORC1's specific binding partners are the scaffold protein raptor (regulatory associated protein of mTORC1) (5, 6), and the inhibitory protein PRAS40 (proline-rich Akt substrate of 40 kDa) (7-11). mTORC2 is formed by rictor (rapamycin-insensitive companion of mTOR) (12, 13), mSin1 (mammalian stress-activated protein kinase interacting protein 1) (14, 15) and Protor (protein observed with rictor) (7, 16). In addition, mTORC1 and mTORC2 share the binding partners mLST8 (mammalian lethal with SEC13 protein 8) (13, 17), DEPTOR (DEP domain containing mTOR-interacting protein) (18) and the Tti1 (TELO2 interacting protein 1)/ Tel2 (telomere maintenance 2) complex (19). Both mTORC1 and mTORC2 are activated by growth factors (e.g., insulin) and amino acids, and mTORC1 is positively regulated by energy (ATP/AMP ratio) (2) (**Figure 1**).

Insulin activates mTORC1 via a well-described signalling cascade that is initiated either by the InR (insulin receptor) or the IGF1-R (insulin like growth factor 1 receptor) (**Figure 1**). Upon insulin/ IGF-1 (insulin like growth factor 1) binding, the receptors dimerize and trans-phosphorylate their cytoplasmic domain (20). This leads to the recruitment of IRS (insulin receptor substrate), which is phosphorylated at tyrosine residues by the InR and IGF-1R, and in response acts as a scaffold for many proteins (20). Class-I PI3Ks (phosphatidylinositol 3-kinases) bind to IRS, where they are activated, and thus, phosphorylate PIP2 (phosphatidylinositol-3,4-bisphosphate) to form PIP3 (phosphatidylinositol-3,4,5-triphosphate). PIP3 is converted back to PIP2 by the phosphatase PTEN (phosphatase and tensin homolog) (21). Via binding to PIP3, proteins containing a PH (pleckstrin homology) domain are recruited to the membrane. The AGC (protein kinase A/protein kinase G/protein kinase C) kinases PDK1 (3-phosphoinositide-dependent kinase-1) and Akt both contain PH domains (21). Upon PIP3 formation they translocate to the plasma membrane, where PDK1 phosphorylates Akt within its kinase domain (22). Akt downstream of PDK1 activates mTORC1, by phosphorylating and inhibiting PRAS40 (23, 24), and the TSC (tuberous sclerosis complex) complex, formed by TSC1, TSC2 and TBC1D7 (Tre2-Bub2-Cdc16 (TBC) 1 domain family member 7) (25, 26). The TSC complex acts as a GAP (GTPase-activating protein) toward the small GTPase Rheb (ras-homologue-enriched-in-brain) (27). Therefore, TSC complex inhibition by Akt de-represses Rheb, which directly binds and activates mTORC1 (28). For Rheb to act on mTORC1, mTORC1 must be localized to lysosomes. Translocation of mTORC1 from the cytosol to the lysosomal surface occurs upon amino acid stimulation and is mediated by the Rag GTPases (Ras-related GTP-binding proteins) (29-32) (reviewed by (2, 4)). In addition, amino acids suppress lysosomal localisation of the TSC complex, leading to de-repression of Rheb (33-35). Amino acids also activate PI3K via an unknown mechanism (36, 37), and AMPK (AMP-activated protein kinase) in a CaMKK β (Ca²⁺/calmodulin-dependent protein kinase kinase-beta) dependent manner (37) (**Figure 1**). Next to growth factors and amino acids, also the cellular energy status modulates mTORC1 activity (2). When the ATP/AMP ratio is low, AMPK (AMP-activated protein kinase) is allosterically activated by AMP, and inhibits mTORC1 by activating the TSC complex (38), and by directly

phosphorylating raptor (39) (**Figure 1**). mTORC1 affects virtually all metabolic processes to ultimately regulate cellular growth and survival (reviewed by Saxton and Sabatini (4)). We focus here on protein homeostasis which mTORC1 controls by activating translation and by inhibiting autophagy (4). mTORC1 positively regulates translation by activation of S6K (ribosomal protein S6 kinase) (40) and inhibition of 4E-BP (eukaryotic translation initiation factor 4E-binding protein) (41), leading to increased translation capacity (42) and enhanced translation initiation (43), respectively (**Figure 1**). mTORC1 inhibits autophagy by phosphorylating and inhibiting ULK1/2 (unc-51 like autophagy activating kinase 1/2) (44) (**Figure 1**), which phosphorylates FIP200 and thereby enhances autophagosome formation (45-48). AMPK activates ULK1, acting as an mTORC1 antagonist in autophagy (44, 49). In addition, mTORC1 and its substrate S6K reduce insulin sensitivity via negative feedback mechanisms. S6K phosphorylates IRS, leading to IRS degradation (50, 51), and mTORC1 phosphorylates and stabilizes Grb10 (growth factor receptor-bound protein 10), leading to InR inhibition (52, 53). Both events render cells refractory to insulin and consequently reduce PI3K and Akt activity.

mTORC2 also controls central metabolic pathways, including glucose metabolism (54), cell survival (54) and cytoskeletal organization (12, 13). The mechanisms leading to mTORC2 activation are relatively poorly explored. Previous research has established that insulin activates mTORC2 (13) in a PI3K dependent manner (55) at the ribosomes (56) (**Figure 1**). However, this PI3K differs from the PI3K upstream of mTORC1, in that mTORC2 is not regulated by mTORC1-driven negative feedback to PI3K or Akt (57). Different PI3K inputs to the two mTOR complexes could be mediated by distinct isoforms of the PI3K catalytic subunit p110, such as in hippocampal progenitor cells, where p110 α activates both mTOR complexes while p110 β activates mTORC2 only (58). Amino acids activate mTORC2 as well, but the exact mechanism remains so far unknown (36). The best described downstream effector of mTORC2 is Akt, which is phosphorylated by mTORC2 within the so-called hydrophobic motif (HM) (59). Akt downstream of mTORC2 inhibits the transcription factors FoxO1/3A (forkhead box proteins O1 and O3A) (60, 61) (**Figure 1**) which enhance apoptosis (60, 62, 63) and mediate stress responses (64-67). Therefore, mTORC2 promotes cellular survival. mTORC2 also activates the AGC kinase SGK (serum and glucocorticoid-regulated kinase) (55) that induces proliferation, migration and cell survival (68).

Isoforms and splice variants across the mTOR network

Next to the complex wiring of the mTOR network, most of its components occur as different variants, contributing to the network's complexity and versatility. These variants comprise similar proteins originating from different genes, referred to as "isoforms" (**Table 1**), and "splice variants" originating from alternative splicing or alternative translation initiation of the same gene (**Table 2**). Isoforms and splice variants can have overlapping, yet also different biological functions. In the following we focus on the differences between these variants regarding their protein structure, tissue expression pattern, and regulation and function within the mTOR network. Isoforms and splice variants of PI3K and FoxO1/3A have been reviewed extensively earlier (69, 70) and are therefore not covered here.

In mammals, the InR occurs in two different splice variants, named InR-A and InR-B (71) (**Table 2**). InR-A is expressed predominantly in the central nervous system and hematopoietic cells, while InR-B is found in adipose tissue, liver, and muscle (72). Both InR-A and InR-B have

alpha and beta domains, but only InR-B has an extra stretch of 12 amino acids in the alpha domain that changes its binding characteristics toward its ligands (72). IR-A and IR-B can form homo- or hetero-dimers and have similar affinities for insulin. However, InR-A binds IGF-I and IGF-II with higher affinity than InR-B (73, 74). This difference in affinity may allow them to activate the mTOR pathway differently in response to the same ligands.

Downstream of the InR, IRS has four different isoforms originating from different genes (IRS1-4), with IRS3 exclusively present in rodents (75). Therefore, we focus here on the differences between IRS1, IRS2, and IRS4 (**Table 1**). IRS1 and IRS2 are ubiquitously expressed (76) and, thus, the most widely studied isoforms. They are highly similar in their domain structure, but IRS2 contains an extra KRLB (kinase regulatory loop binding) domain that binds to the InR and the IGF1-R (77, 78). The KRLB domain limits the tyrosine phosphorylation on IRS2 by the InR and IGF1-R, and thereby inhibits IRS2 function (79). An *in vitro* study in skeletal muscle cells has shown that IRS1 and IRS2 tyrosines are dephosphorylated at different rates by phosphatases upon insulin or IGF-1 stimulation (80). While IRS2 becomes rapidly dephosphorylated after 3-10 min, IRS1 remains phosphorylated for up to one hour (80). It is unknown if this difference is mediated by distinct phosphatases, or by different dynamic behaviour of the same phosphatase toward IRS1 *versus* IRS2. Such differences in dynamic regulation might contribute to the distinct functional outcomes of IRS1 and IRS2. IRS1 enhances phosphorylation of both Akt1 and Akt2, whereas IRS2 signals mainly through Akt2 (81). IRS1 but not IRS2 enhances actin remodelling and GLUT4 (glucose transporter type 4) translocation (81). IRS1 and IRS2 induce the MAPK pathway with IRS2 having a stronger effect than IRS1 (81, 82). IRS1 and IRS2 seem to have opposite effects on metastasis, as IRS2 promotes metastasis in breast cancer cells (83, 84), whereas IRS1 suppresses metastatic spread in mice. Only little is known about IRS4, whose protein expression is limited to brain, kidney, thymus and liver (85). Recent evidence suggests a function of IRS4 during adenoviral infection, where IRS4 upregulation leads to constitutive Akt activation even in the absence of insulin (86).

The PI3K antagonist PTEN exists in three reported variants, termed PTEN, PTEN-Long (87) and PTEN α (88), originating from alternative translation initiation (**Table 2**). PTEN-Long and PTEN α both originate from the start codon CUG⁵¹³ and have the same apparent molecular weight and predicted number of amino acids, suggesting that these two variants could be the same (87, 88). PTEN is ubiquitously expressed while PTEN α is predominantly expressed in skeletal and cardiac muscle. All variants contain the functional PTEN domains, including the phosphatase domain, the C2 domain and the tail domain. PTEN α and PTEN-Long contain an extra N-terminal domain which may lead to different subcellular localizations (87, 88). While PTEN localizes mainly to the cytosol, PTEN α resides at mitochondria (88), and PTEN-Long is secreted to the microenvironment (87). This difference in localization results in distinct functions, as PTEN is the counterpart and antagonist of PI3K at the plasma membrane, whereas PTEN α targets the mitochondrial complex IV (cytochrome c oxidase) to regulate the cellular energy status (88). PTEN-Long also decreases PI3K signalling in a phosphatase-dependent manner, however, as it is secreted to the microenvironment and able to enter neighbouring cells, it could have its main role in controlling signalling at the tissue level (87).

There are three Akt isoforms in mammals, Akt1 (PKB α), Akt2 (PKB β) and Akt3 (PKB γ) (**Table 1**) which are similar in structure, as they all contain a PH domain, a kinase domain and the HM. Akt3 is also present as a shorter splice variant Akt3-1 (**Table 2**), which lacks the

mTORC2 phosphorylation site within the HM, and is less responsive to growth factor stimulation (89, 90). The three Akt isoforms display unique tissue expression patterns. While Akt1 is ubiquitously present (76), Akt2 resides primarily in brown fat, skeletal muscle and liver (91, 92), and Akt3 is mainly expressed in brain and testis (93). Akt1 and Akt2 also differ in their sub-cellular localization: in rat adipocytes, Akt1 is mainly distributed in the cytosol, whereas Akt2 localizes to GLUT4 containing vesicles (94). The phosphatases PHLPP1 (PH domain and leucine-rich repeat protein phosphatase1) and PHLPP2 (95) differently act on the mTORC2-substrate site within the HM of the three Akt isoforms (95): PHLPP1 specifically dephosphorylates Akt2 and Akt3, while PHLPP2 dephosphorylates Akt1 and Akt3 (95). The three Akt isoforms also differ regarding their substrate specificity, as only Akt1 and Akt2 phosphorylate TSC2 (95) to promote cell growth, whereas all three isoforms phosphorylate FoxO1/3A (95) to inhibit apoptosis. It is unknown if the substrate specificity of the different Akt isoforms also relates to PRAS40. It is likely as a number of further Akt-isoform specific substrates have been described: Akt1 phosphorylates palladin, an actin-binding protein, to inhibit cell migration (96) while Akt2 phosphorylates Ankrd2/ARPP, a muscle-specific protein that, when phosphorylated by Akt, prevents muscle differentiation (97). Their different substrate specificities grant the Akt isoforms distinct roles in tumour formation. Akt1 specifically promotes cell survival (98, 99) and cell growth (100, 101), whereas Akt2 is more important for glucose metabolism (102, 103), cell migration (100), tumour metastasis (100), and EMT (endothelial-mesenchymal transition) (101). In contrast, Akt3 inhibits migration (104), and has been linked to neuronal growth and development (105, 106). Interestingly, Akt3 overexpression confers resistance to the Akt inhibitor MK2206 (107) suggesting that Akt3 can be pro-tumourigenic in the context of targeted cancer therapies.

Downstream of Akt, TSC2 is present in healthy individuals in several variants originating from alternative splicing of exons 25 and 31 (108-110) (**Table 2**). TSC2 transcripts lacking exons 25 (Uniprot ID: P49815-2, P49815-3, P49815-5, P49815-6 and P49815-7) and/or 31 (Uniprot ID: P49815-4, P49815-5, P49815-6 and P49815-7) occur at higher levels as compared to the full length variant (Uniprot ID: P49815-1) (110). Exon 25 is detectable in lymph node, muscle and thyroid tissue whereas Exon 31 is found in adipose, adrenal, brain, breast, colon, kidney, lung, prostate, skeletal muscle, testes, and thyroid tissue (110). Deletion of exons 25 and 31 does not affect the TSC complex-dependent inhibition of mTORC1 (110). However, it may affect TSC2's properties as a substrate for Akt and AMPK. Akt phosphorylates TSC2 at residues S939, S981 and T1462 (26, 111, 112) and variants missing exon 25 do not contain the target site at S981. The TSC2 residues phosphorylated by Akt serve as a scaffold for 14-3-3 protein binding to and subsequent degradation of TSC2 (112). This suggests that TSC2 containing exon 25 has a higher 14-3-3 affinity and higher turnover than TSC2 variants without that exon (108). AMPK activates TSC2 by phosphorylation at S1387 and T1271 (38). The AMPK target site at T1271 is contained in exon 31 and, therefore, TSC2 variants that lack exon 31 may be less sensitive to AMPK activation. Further studies are required to delineate the biological consequences of the different structures of the TSC2 splice variants upon Akt and AMPK activation. TSC1 variants originating from alternative splicing have so far not been identified experimentally under healthy conditions. However, mutations in either TSC1 or TSC2 that lead to aberrant splicing and generation of multiple splice variants are commonly found in a wide range of pathologies such as cancer and tuberous sclerosis complex (TSC) (113). TSC is a rare genetic disorder caused by mutations in the TSC1 and TSC2 genes, leading to benign tumors in multiple organ systems (110). The TSC Leiden Open Variation Databases (TSC LOVD, www.lovd.nl) displays all variants reported for TSC1 and TSC2, being at the moment

(May 2017), 870 and 2473 respectively. For TBC1D7, the third member of the TSC protein complex, no isoforms or splice variants have been found experimentally. Also Rheb, the target of the TSC complex, does not present any isoform or splice variant.

Components of the mTOR complexes occur in different variants (**Table 2**). mTOR itself exists in two splice variants, mTOR α and mTOR β , of which mTOR β is the shorter version (114). mTOR α contains the protein-protein interaction domains HEAT (huntingtin-elongation factor 3-regulatory subunit A of PP2A-TOR1 repeats), FAT (FRAP-ATM-TTRAP) and FATC (FAT-carboxy terminal domain) and a kinase domain, while mTOR β lacks the HEAT and FAT domains (114). Studies in rats suggest that mTOR α is ubiquitously expressed whereas mTOR β is mainly expressed in lung, heart, stomach, intestine and liver (114). mTOR β can still form the two mTORC1 and mTORC2 complexes, and phosphorylates the mTORC1 substrates S6K1 and 4E-BP1 as well as the mTORC2 substrate Akt. Cells overexpressing mTOR β have a shorter G1 phase, suggesting a role in cell cycle progression (114). This may be due to a stronger association of c-Myc, a transcription factor that controls cell cycle progression, with mTOR β than with mTOR α . Also raptor exists in two splice variants, termed raptor and raptor-v2. Raptor-v2 (**Table 2**) is the shorter variant lacking the HEAT repeats, which are needed for binding to mTORC1 substrates (115). Raptor is ubiquitously expressed but levels are higher in brain, immune cells, the gastrointestinal tract and kidney (76). Raptor-V2 mainly localizes to the pituitary, nasal mucosa and muscle (115). Raptor-v2 binds mTORC1 but cannot bind S6K1, as it lacks the HEAT repeats (116). This suggests that raptor-v2 inhibits mTORC1 by sequestering mTOR away from the functional complex. However, further studies are required to characterize the biological function of raptor-v2.

Regarding the members of mTORC2, no isoform or splice variant for rictor has been so far described. However, five mSin1 splice variants have been experimentally identified, mSin1.1 - mSin1.5 (14) (**Table 2**). No information is available on their expression pattern. Only mSin1.1, mSin1.2 and mSin1.5 (also termed mSin1, mSin1 α and mSin1 β (117)) are found in mTORC2 and mediate Akt phosphorylation. mSin1.1 is the full-length protein, and contains a ras-binding domain (RBD) and a PH domain. mSin1.2 and mSin1.5 lack part of the RBD, and mSin1.5 also lacks the PH domain (14). mSin1.1 and mSin1.2 overexpression increases Akt phosphorylation in response to insulin, whereas mSin1.5 overexpression enhances Akt phosphorylation independently of insulin (14). This suggests that mTORC2 containing mSin1.1 or mSin1.2 is activated by insulin, whereas mTORC2 with mSin1.5 is constitutively active or responds to signals other than insulin. Protor has two isoforms, Protor-1 and Protor-2 (**Table 1**) (7, 16), that are ubiquitously expressed (16). They do not contain any known functional domain, but their amino-terminal sequence is highly conserved and they both bind mTORC2 via rictor (7, 16). Protor-1 also exists in 3 splice variants, Protor-1 α , Protor-1 β and Protor-1 γ , of which only Protor-1 α and Protor-1 β can form part of mTORC2 (16). In kidney, Protor-1 in mTORC2 is required for the phosphorylation of SGK and activation of sodium transport (118). Protor-2 interacts with and activates the RNA binding protein TTP (tristetraprolin) to ensure mRNA turnover under stress conditions (119). Further research is required to better understand the differences in expression, function and regulation between these variants.

Protein variants downstream of mTORC1 have been extensively characterized, and we discuss them here in the context of cell growth, translation and autophagy. Firstly, S6K has two isoforms encoded by different genes, S6K1 and S6K2 (120-122) (**Table 1**). They both have an N-terminal domain (NTD), a C-terminal domain (CTD) and a kinase domain. In

addition, S6K2 has a nuclear localization sequence (NLS) next to its CTD (123). Both S6K1 and S6K2 are ubiquitously expressed (76) and, although they are both activated via phosphorylation by PDK1 (122, 124) and mTORC1 (40, 122), different isoform-specific signalling inputs contribute to their activation (123). S6K1 is inhibited by leucine starvation (125) and activated by the actin-binding protein neurabin (126), phospholipase D1 (127), the Rho family G proteins Rac1 and Cdc42 (cell division cycle 42) (128), and the deacetylases sirtuin 1 (SIRT1) and SIRT2 (129). S6K2 is activated in response to the cytokine IL-3 (interleukin 3) (130), by the MAPK pathway (131), and PKC (protein kinase C) (132). Both S6K1 and S6K2 phosphorylate S6 (ribosomal protein S6), a protein that forms part of the ribosomal 40S subunit (133) and, thus, the distinct activators of S6K1 and S6K2 may be a means to specifically control translation upon different cellular conditions, such as growth factor stimulation, different cell cycle phases or during the immune response. In addition to acting on S6, S6K1 also activates translation by phosphorylating eIF4B (eukaryotic translation initiation factor 4B) (134) and eEF2K (eukaryotic elongation factor-2 kinase) (135). Furthermore, S6K1 promotes cell survival by inhibiting BAD (BCL2 associated agonist of cell death) (136), Mdm2 (Mdm2 proto-oncogene) (137) and GSK-3 (glycogen synthase kinase 3) (138), and induces transcription by activating $E\alpha$ (estrogen receptor alpha) (139) and CREM (cAMP responsive element modulator) (140). S6K2 activates transcription by binding to the transcription factors YY1 (yin yang 1) (141) and ROR γ (RAR-related orphan receptor gamma) (142). Both S6K isoforms exist as several variants (**Table 2**). The S6K1 mRNA yields two differently translated variants originating from different translational start sites, p70-S6K1 and p85-S6K1 (143), the latter containing an N-terminal NLS. There is also a shorter S6K1 with a truncated kinase domain originating from alternative splicing, and termed p31-S6K1 in mice and hS6K1-h6A and hS6K1-h6C in humans (144). The tissue expression of the S6K1 variants is unknown but within the cell p70-S6K1 localizes to the cytoplasm (145), p85-S6K1 to both cytoplasm and nucleus (145, 146) and p31-S6K1 locates to the nucleus (147). Both p70-S6K1 and p85-S6K1 have been shown to be targeted by mTORC1, as rapamycin treatment decreases both phosphorylation of p70-S6K1 at pT389 and of p85-S6K1 at pT412 (147). However, in cells arrested in prometaphase, p85-S6K1 is phosphorylated at pT412 in an mTORC1 independent manner (148) and hence p85-S6K1 might not be regulated by mTORC1 under all circumstances. There is no conclusive data confirming if p31-S6K1 is targeted by mTORC1 (147). Further studies are needed to discriminate the biological functions of each of these variants. Alternative translation also occurs for S6K2, giving rise to two variants, p56-S6K2 and p54-S6K2, that differ by the presence of an NLS in p56-S6K2 in the NTD, but not in p54-S6K2 (122). Hence, they reside in different compartments, p56-S6K2 in the membranous fraction and p54-S6K2 in the soluble fraction (122). Further work is required to establish the exact sub-cellular localization and distinct functions of the S6K2 variants.

4E-BP, another direct mTORC1 substrate, exists in three isoforms (4E-BP1-3) (**Table 1**). All 4E-BP isoforms share an eIF4E (eukaryotic translation initiation factor 4E) binding domain and a TOS (TOR signalling sequence) motif. In addition, 4E-BP1 and 2 contain a RAIP motif, named after its amino acid sequence. Whereas 4E-BP2 is ubiquitously expressed (149), 4E-BP1 and 4E-BP3 are primarily found in the pancreas and skeletal muscle (149, 150). In addition, 4E-BP1 is expressed in adipose tissue (149), and 4E-BP3 in the kidney (150). mTORC1 inhibits 4E-BP1 and 2 by phosphorylation at their RAIP motifs. Phosphorylated 4E-BP1 and 2 are released from eIF4E, and consequently eIF4E can bind the 5' cap of mRNAs to initiate translation (43, 151). 4E-BP3 does not contain a RAIP motif and therefore is not inhibited by mTORC1, although mTORC1 phosphorylates 4E-BP3 at other sites (151).

Interestingly, prolonged mTORC1 inhibition enhances 4E-BP3 expression at the transcriptional level, and this is mediated by the transcription factor TFE3 (transcription factor binding to immunoglobulin heavy constant mu enhancer 3) (152). In addition to inhibiting eIF4E binding to the cap, 4E-BP3 regulates eIF4E at the nucleus to regulate nuclear mRNA export (153).

mTORC1 regulates autophagy by phosphorylating ULK. ULK has 4 isoforms, ULK1-4, but only ULK1 and ULK2 are inhibited by mTORC1 (47) (**Table 1**). ULK1 and ULK2 are ubiquitously expressed (76) and both induce autophagy (154, 155). ULK1 has a stronger affinity than ULK2 towards the other members of the autophagy-inducing complex, Atg13 (autophagy related 13) and FIP200 (48). This suggests different functions of the ULK1/Atg13/FIP200 and ULK2/Atg13/FIP200 complexes during autophagy, but further studies are needed to test this hypothesis.

2. mTOR and ageing

Ageing can be defined as the gradual deterioration of the physiological functions necessary for survival (156). This concept relates both to the lifespan of an individual and to the manifestation of age-related disorders, such as obesity, diabetes and myopathy. In other words, increased longevity can reflect either an increase in lifespan or a reduction of age-related diseases. The role of mTOR in the ageing process has been a topic of research over the last decades, and we give here an overview of the key findings in invertebrate and vertebrate model organisms and humans.

TOR and lifespan in invertebrates

The process of ageing has been widely studied in model organisms such as the budding yeast *Saccharomyces cerevisiae* (*S. cerevisiae*), the nematode *Caenorhabditis elegans* (*C. elegans*), and the fruit fly *Drosophila melanogaster* (*D. melanogaster*). Much research on ageing has been performed in these organisms due to their short lifespan, their easy manipulation and the availability of powerful genetic tools. Studies in *S. cerevisiae* have shown that inhibition of TORC1 with rapamycin increases the chronological life span (duration of time that cells in stationary phase remain viable) (157). In addition, the longevity phenotype induced by dietary restriction was found to be TOR dependent (157). By generating yeast deletion collections, several long-lived mutants were identified. Among them were strains with mutations in genes of the TOR signalling axis (157) and the TOR substrate Sch9/S6K (158), as well as transcription factors that upregulate genes encoding amino acid biosynthetic enzymes and amino acid permeases (157).

Studies in *C. elegans* provide evidence that rapamycin (159) and dietary restriction (160) increase the lifespan of multicellular organisms as well. CeTORC1, TORC1 in *C. elegans*, inhibition by mutation or inhibition of let-363/CeTOR (161), DAF-15/raptor (162), raga-1/Rag GTPases (159, 163), or RHEB-1/Rheb (164) causes longevity phenotypes. Also mutations in CeTOR's upstream regulators DAF-2/InR (165) and AGE-1/PI3K (166) dramatically extend the lifespan of *C. elegans*. In contrast to the increased longevity of worms in which let-363/CeTOR or daf-15/raptor are targeted by RNAi or mutation, RICT-1/rictor deficiency causes short-lived worms (167), suggesting that the two complexes have opposite roles in the lifespan regulation of *C. elegans*.

Further evidence on the evolutionary conservation of TOR's role in lifespan and ageing arose from the confirmation of these phenotypes in *D. melanogaster*. Rapamycin (168) or dietary

restriction (169) increase lifespan also in flies. Inhibition of the insulin pathway by mutations in the *InR* gene (170) or in the *InR* substrate *CHICO* (171) results in increased longevity. Additionally, overexpression of *Tsc1/TSC1* or *Tsc2/TSC2*, or dominant-negative forms of *dTOR* (*D. melanogaster* TOR) and its substrate *dS6K/S6K*, cause lifespan extension (172). Furthermore, the TORC1 substrate *d4E-BP/4E-BP* has a pivotal role delaying fly ageing (173). Hence, inhibition of the signalling axis converging on TOR and its substrates prolongs lifespan in non-vertebrate model organisms.

mTOR and ageing in mice

The before mentioned studies provide strong evidence that signalling through insulin and TOR restricts the lifespan of invertebrates. Evidence that this mechanism is also conserved in mammals came from mouse studies in which rapamycin or dietary restriction increased lifespan (174, 175). However, dissecting the mechanisms underlying mTOR's role in mammalian ageing proved to be even more challenging than in invertebrates. One reason is the higher complexity of the mammalian insulin-mTOR axis with several isoforms and often various splice variants at almost all levels of the pathway. As detailed earlier, tissue and subcellular distribution varies greatly for the different variants, leading to tissue-specific biological outcomes of mTOR. It is therefore not surprising that knockout of the same protein in different tissues leads to divergent phenotypes (176).

Several knockout studies in mice suggest functions of components of the insulin-mTOR pathway in ageing (**Table 3**). Lifespan extension has been observed in a female mouse model with heterozygous whole body double knockout of the mTOR and *mLST8* genes, *Mtor* and *Mlst8* (177), or with a homozygous whole body knockout of *Irs1* (178). Also a homozygous whole body knockout of *Rps6kb1*, the *S6K1* gene, extends lifespan in mice (179). Lifespan extension has been also reported for heterozygous knockouts of *Irs2* in the whole body, or in brain only (180), but the reproducibility of this phenotype has been questioned (181). In agreement with the studies in *C. elegans* (167), mTORC2 specific knockouts enhance ageing also in mice, as a whole body knockout of *riCTOR* decreases lifespan (182).

Lifespan studies in mice are relatively rare due to their comparatively long lifespan. In contrast, studies that focus on the role of the mTOR pathway in age-related, metabolic phenotypes and diseases are much more common (**Table 3**). Adipose tissue-specific knockouts of the *InR* gene *Insr* (183) or the raptor gene *Rptr* (184) result in mice with substantially less fat that are protected against obesity and hypercholesterolemia. In addition, knockout of either *Tsc1* or *Tsc2* enhances tumour formation (185, 186). This suggests that inhibition of the insulin-mTORC1 axis protects higher organisms against age-related metabolic and tumour disorders. However, this concept is challenged by the fact that most other knockout models of the insulin-mTOR pathway in mice develop phenotypes that positively link with age-related disease and could hence be considered as phenotypes of accelerated ageing (**Table 3**). Such phenotypes encompass for example impaired glucose tolerance, insulin resistance, obesity and myopathy, which correlate with increased age in mice and men (187-189). Impaired glucose tolerance has been observed in whole body knockouts of *Irs4* (190), and *Akt2* (102), in tissue specific knockouts of the *Insr* in muscle (191), beta cells (192) and brain (193), or of the *PDK1* gene, *Pdprk1*, in the liver (194). Insulin resistance occurs in mice lacking *Irs2* (195) or *Akt2* (102), and in tissue specific knockouts of the *Insr* in liver (196) or brain (193). Obesity is also observed for whole body *Irs2* knockout mice (195), as well as for specific knockouts of the *Insr* in muscle (191) or brain (193). Myopathy is a characteristic phenotype of mice with muscle specific

knockouts of *Pdprk1* (197), *Mtor* (198), *Rprtr* (199) and in a double knockout model of *Rps6kb1* and *Rps6kb2* (200).

The observation that so many mouse knockout models of the insulin-mTOR pathway develop metabolic phenotypes which can be linked with age-related metabolic impairment may seem at odds with the studies in invertebrates where a large majority of insulin-TOR pathway mutants display a prolonged lifespan. However, metabolic phenotypes in mouse studies must be interpreted with caution regarding their relationship with lifespan and ageing. The reason is that it is not possible to discriminate whether a detrimental metabolic phenotype in a mouse knockout model is due to an age-inhibitory role of the targeted gene, or due to a potentially essential role of this gene in metabolic processes. A well-known example of such a seemingly-contradictory phenotype has been observed for rapamycin in mice. Rapamycin does increase lifespan via mTORC1 inhibition but, when chronically administered, rapamycin causes secondary effects leading to mTORC2 inhibition and substantial impairment of glucose tolerance and insulin action (177). Hence, rapamycin extends lifespan and severely impairs metabolism at the same time, via distinct mechanisms. The fact that lifespan extension can be observed concomitantly with metabolic impairment suggests that metabolic alterations and lifespan are not always strictly causally linked.

The notion that lifespan and metabolic outcomes of genes can be separated is further strengthened when taking a closer look at isoform specific effects of genes in the insulin-mTOR axis on ageing and metabolic phenotypes in mice (**Table 3**). For example, different IRS isoforms govern metabolism and ageing in distinct, often opposite ways. A homozygous whole body knockout of *Irs1* prolongs lifespan of female mice (178), although they develop insulin resistance (201). In contrast a whole body knockout of *Irs2* shortens the lifespan of both male and female mice (178), and leads to a diabetic phenotype (195, 202). *Irs4* knock out mice have a milder phenotype than *Irs1* or *Irs2* knockouts, with regard to insulin sensitivity defects (190), and no effects on lifespan have been reported. Distinct phenotypes are also observed for knockouts of the different Akt isoforms. Mice with heterozygous *Akt1* knockout display a prolonged lifespan (203). In contrast, *Akt2* deficient mice are insulin resistant with elevated plasma triglycerides and diabetes in males (102). An *Akt3* knockout does not seem to have an ageing-related metabolic phenotype (106). Downstream of mTORC1, a whole body knockout of *Rps6kb1* prolongs lifespan (179), whereas a *Rps6kb2* knockout shows no obvious phenotypic abnormalities (133). More extensive characterization of the *Rps6kb2* knockout model would allow for better understanding of S6K2's potential role in aging. Mice with a knockout of *Eif4ebp1*, 4E-BP1's gene, exhibit no difference in lifespan, although they present an increased metabolic rate and a reduction of adipose tissue (204), again questioning the strict causal relationship between beneficial metabolic features and lifespan extension. The role of 4E-BP2 has only been studied by double knockout of *Eif4ebp1* and *Eif4ebp2*, and these mice are obese and insulin resistant (205). However, no information about the lifespan of this model is available and no 4E-BP3 knockout has been so far reported.

mTOR and ageing in humans

Most research on mTOR in humans focuses on age-related diseases such as cancer and diabetes. Regarding ageing itself, studies in long-lived primates, specifically in Rhesus monkeys, have shown that calorie restriction delays disease onset and possibly mortality (206). The effect of rapamycin on primate lifespan has not yet been reported, but rapamycin improves immune function in elderly humans (207). Given that intervention studies on longevity and ageing in higher primates and humans are scarce, candidate gene and genome-

wide association studies (GWAS) are the main tools to understand the relationship between genetic makeup and human lifespan and disease susceptibility. Such studies use cohorts of advanced age and focus on genetic factors that correlate with exceptional longevity and healthy ageing.

GWAS studies in long-lived humans have so far yielded only very few gene associations that correlate with ageing. Indeed, only associations of ApoE (apolipoprotein E) and FoxO3a genes have been replicated in several studies (208). It is surprising that components of the insulin-mTOR pathway have not been identified in these studies. Possibly the GWAS study does not have enough power to detect such correlations, as this approach only allows the detection of common genetic variations. Development of new techniques such as whole-genome sequencing permit the detection of rare potentially functional genetic variants and raise much hope for the detection of further genetic correlations with lifespan and age-related processes. In a recent study, whole-genome sequencing was used to analyse human healthy ageing, defined as disease-free ageing without medical intervention (209). However, in this study no major genetic contributors to healthy ageing could be identified.

In contrast to these unbiased approaches, analyses of specific sets of candidates in the insulin-mTOR signalling pathway to unravel their potential role in ageing has yielded more success. A study that included 122 Japanese “semisupercentenarians” (older than 105 years) found polymorphic variations of the InR and IRS1 genes that are more frequent than in the control group (210). In addition, a polymorphism in Akt1 that significantly associates with lifespan has been found in three independent Caucasian cohorts (211). In the Leiden Longevity Study, gene expression analysis of nonagenarians shows that expression of 4E-BP1 and PRAS40, two negative effectors of the mTORC1 pathway, is higher in the aged group compared to the middle-aged control group (212). Moreover, raptor is expressed to a lower level in middle-aged members of the longevity families as compared to similarly aged controls (212). Finally, low insulin signalling has been associated with improved old-age survival in women (213). Of note, these targeted studies do not include isoforms or splice variants of the different members of the mTOR signalling pathway and, therefore, some relevant candidates may have been overlooked.

3. Discussion

mTOR signalling is widely recognized as a key element in ageing and age-related metabolic conditions in a wide range of organisms from yeast to rodents (157, 159, 168, 174). In invertebrates, inhibition of the insulin- TOR pathway by mutations or RNA interference extends lifespan, suggesting a positive link between TOR activity and ageing progression. When studying the same genes in higher organisms such as mammals, the relationship between mTOR and ageing becomes more complex. Although there are some clear examples of inhibition of the mTOR pathway that lead to lifespan extension, most knockouts result in the development of metabolic conditions that could be rather be considered as a sign of accelerated ageing. A limitation is that most of these mouse studies only analyse metabolic parameters and not lifespan, and conclusions cannot be drawn from metabolic phenotypes on shortened or prolonged lifespan. The concomitant occurrence of prolonged lifespan and detrimental metabolic phenotypes, or beneficial metabolic features with no lifespan effect in

some of these models challenge the idea of a strict and direct relationship between metabolic alterations in knockout models of genes of the insulin-mTOR axis and lifespan.

This review emphasizes the need to identify and characterize the isoforms and splice variants within the mTOR pathway to achieve a better understanding of the contribution of these different elements to metabolism and ageing, and the interrelationship of both. Although many of these isoforms have been identified long ago, they have been considered as proteins with overlapping function for a long period. This is reflected by the fact that isoforms with the number 1 in their name are often much better studied than their counterparts with higher numbers. Hence, the extent of knowledge on these variants often relates to their arbitrary numbering in databases.

For splice variants we lack even more knowledge as they are ignored by most experimental studies even though the majority of the genes that code for mTOR pathway components have been predicted to produce several splice variants. Furthermore, RNA splicing is required for longevity downstream of dietary restriction and the CeTORC1 pathway in *C. elegans* (214). The gap in our knowledge becomes even more apparent when considering that additional, abnormal splice variants occur in many genetic diseases and cancers (215). A recent global study on the interactomes of splice variants have shown that splice variants share only half of their interaction partners and have distinct tissue expression and, therefore, should be considered as distinct proteins (216). Thus, further work is required to experimentally identify and functionally characterize both natural-occurring and disease-causing variants in the mTOR pathway, and to better understand the relationship of these genes and their splice products with metabolic regulation, ageing and lifespan, and age-related diseases.

Acknowledgements

We thank Dr. Mark Nellist for his expert opinion on TSC2 variants. We thank Alexander Martin Heberle, Laura Corbett, Antje Thien, Marti Cadena Sandoval and Ulrike Bosch for critical reading of this manuscript. The authors sincerely apologize to those whose work could not be cited in this review because of space limitations.

Declarations of interest

The authors declare no competing interests

Funding information

K.T. and P.R.N. are supported by the BMBF e:Med project GlioPATH (01ZX1402B). K.T. is supported by BMBF e:Med MAPTor-NET (031A426B), a Rosalind-Franklin-Fellowship of the University of Groningen, the Ubbo Emmius Fund, Stichting Michelle (calls 2015 and 2017) and the Deutsche Forschungsgemeinschaft (TH 1358/3-1).

Author contribution statement

P.R.N and K.T. wrote the manuscript.

Figure/Table legends

Figure 1. The mTOR pathway. Insulin binds and activates the insulin receptor (InR) which recruits phosphatidylinositol 3-kinase (PI3K) via the insulin receptor substrate (IRS). Once recruited to the membrane, PI3K phosphorylates phospholipid phosphatidylinositol-3,4-bisphosphate (PIP2) to form phosphatidylinositol,3,4,5-triphosphate (PIP3). PIP3 is converted back to PIP2 by PTEN (phosphatase and tensin homolog). PIP3 serves as a membrane anchor for the 3-phosphatidylinositol-dependent kinase 1 (PDK1) and Akt. PDK1 activates Akt, which in turn phosphorylates and inhibits the tuberous sclerosis (TSC) complex and the proline-rich Akt substrate of 40 kDa (PRAS40). Once the TSC complex is inhibited, ras-homologue enriched in brain (Rheb) can exert its activating action on the mammalian target of rapamycin (mTOR) complex 1 (mTORC1). mTORC1 negatively regulates the eukaryotic translation initiation factor 4E-binding protein (4E-BP) and unc-51 like autophagy activating kinase 1/2 (ULK1/2) and positively regulates the ribosomal protein S6 kinase (S6K). Both mTORC1 and S6K contribute to negative feedback mechanisms at the level of the InR and IRS. Insulin also activates mTORC2 in a PI3K dependent manner. mTORC2 activates Akt, which inhibits the forkhead box O transcription factors FoxO1/3A. Amino acids have several activating inputs on the network, at the level of mTORC1 via the Rag GTPases (Ras-related GTP-binding proteins) and at the level of PI3K. A high AMP/ATP ratio leads to allosteric AMP-activated protein kinase (AMPK) activation that inhibits mTORC1 by activating the TSC complex and by directly inhibiting mTORC1. AMPK also phosphorylates and activates ULK1/2. Only the functional mTORC1 and mTORC2 outputs discussed in this review are shown. More extensive overviews of the processes downstream of the mTOR complexes are provided for example by Saxton and Sabatini (4) and Ben-Sahra and Manning (3).

Table 1. Isoforms across the mTOR network. This table highlights the differences among the various isoforms of proteins in the insulin-mTOR axis. Differences are categorized according to different structure, regulation, expression pattern and biological function. References are indicated in brackets. The colour code refers to the different signalling modules of the mTOR network as indicated in Figure 1. *Abbreviations: autophagy related 13 (ATG13), BCL2 associated agonist of cell death (BAD), cell division cycle 42 (Cdc42), epithelial to mesenchymal transition (EMT), estrogen receptor alpha (Erα), eukaryotic elongation factor-2 kinase (eEF2K), eukaryotic translation initiation factor 4B (eIF4B), eukaryotic translation initiation factor 4E-binding protein (4E-BP), forkhead box O (FoxO), glucose transporter type 4 (GLUT4), glycogen synthase kinase 3 (GSK-3), growth factor receptor-bound protein 2 (Grb2), interleukin 3 (IL-3), insulin receptor substrate (IRS), Mdm2 proto-oncogene (Mdm2), mitogen activated protein kinase (MAPK), PH domain and leucine-rich repeat protein phosphatase (PHLPP), protein kinase 3 (PKC), protein observed with rictor (Protor), RAR-related orphan receptor gamma (RORγ), ras-related C3 botulinum toxin substrate 1 (Rac1), ribosomal S6 kinase (S6K), serum and glucocorticoid-regulated kinase (SGK), sirtuins (SIRT), tuberous sclerosis complex 1 (TSC1), tuberous sclerosis complex 2 (TSC2), unc-51 like autophagy activating kinase (ULK), yin yang 2 (YY2).*

Table 2. Splice variants across the mTOR network This table indicates the predicted splice variants in human and mouse (source: uniprot.org) and the experimentally validated splice variants including functional differences. References are indicated in brackets. The colour code refers to the different signalling modules of the mTOR network as indicated in Figure 1.

Abbreviations: 3-phosphoinositide-dependent kinase-1 (PDK1), amino acids (aa), DEP domain containing mTOR-interacting protein (DEPTOR), insulin-like growth factor 1 (IGF-1), insulin receptor (InR), mammalian stress-activated protein kinase interacting protein 1 (mSin1), mammalian target of rapamycin (mTOR), mTOR complex 2 (mTORC2), phosphatase and tensin homolog (PTEN), phosphatidylinositol 3-kinase (PI3K), phosphatidylinositol-3,4-bisphosphate (PIP2), phosphatidylinositol-3,4,5-triphosphate (PIP3), proline-rich Akt substrate of 40 kDa (PRAS40), protein observed with rictor (Protor), rapamycin-insensitive companion of mTOR (rictor), regulatory associated protein of mTOR (raptor), ribosomal S6 kinase 1 (S6K1), ribosomal S6 kinase 2 (S6K2), TBC1 domain family member 7 (TBC1D7), tuberous sclerosis complex 1 (TSC1), tuberous sclerosis complex 2 (TSC2).

Table 3. Knockout phenotypes for the insulin-mTOR pathway in mice. This table highlights the phenotypes of knockout mouse models that display delayed or accelerated ageing phenotypes. Each group is subdivided depending on whether the phenotype relates to the lifespan of the mice or to their metabolic profile. For each mouse model, the target gene is indicated. Whole body knockout was performed if not indicated otherwise. Reference are indicated in brackets. The colour code refers to the different signalling modules of the mTOR network as indicated in Figure 1. As Mlst8 and Mtor are part of both mTOR complexes, the knockout models are indicated for mTORC1 and mTORC2. A more detailed comparison of the phenotypes with knockouts of the complex-specific components Rptor and Rictor is provided in the text. *Abbreviations: 3-phosphoinositide-dependent kinase-1 (Ppdk1), eukaryotic translation initiation factor 4E-binding protein 1 (Eif4ebp1), insulin receptor (Insr), insulin receptor substrate 1 (Irs1), insulin receptor substrate 2 (Irs2), insulin receptor substrate 3 (Irs3), mammalian lethal SEC13 protein 8 (Mlst8), mammalian target of rapamycin (Mtor), rapamycin-insensitive companion of mTOR (rictor), regulatory associated protein of mTORC1 (Rptr), ribosomal S6 kinase 1 (Rps6kb1), tuberous sclerosis complex protein 1 (Tsc1), tuberous sclerosis complex protein 1 (Tsc2).*

References

1. Heitman J, Movva NR, Hall MN. Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast. *Science*. 1991;253(5022):905-9.
2. Gonzalez A, Hall MN. Nutrient sensing and TOR signaling in yeast and mammals. *EMBO J*. 2017.
3. Ben-Sahra I, Manning BD. mTORC1 signaling and the metabolic control of cell growth. *Curr Opin Cell Biol*. 2017;45:72-82.
4. Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. *Cell*. 2017;168(6):960-76.
5. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, et al. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell*. 2002;110(2):163-75.
6. Hara K, Maruki Y, Long X, Yoshino K, Oshiro N, Hidayat S, et al. Raptor, a binding partner of target of rapamycin (TOR), mediates TOR action. *Cell*. 2002;110(2):177-89.
7. Thedieck K, Polak P, Kim ML, Molle KD, Cohen A, Jenö P, et al. PRAS40 and PRR5-like protein are new mTOR interactors that regulate apoptosis. *PLoS One*. 2007;2(11):e1217.
8. Fonseca BD, Smith EM, Lee VH, MacKintosh C, Proud CG. PRAS40 is a target for mammalian target of rapamycin complex 1 and is required for signaling downstream of this complex. *J Biol Chem*. 2007;282(34):24514-24.
9. Oshiro N, Takahashi R, Yoshino K, Tanimura K, Nakashima A, Eguchi S, et al. The proline-rich Akt substrate of 40 kDa (PRAS40) is a physiological substrate of mammalian target of rapamycin complex 1. *J Biol Chem*. 2007;282(28):20329-39.
10. Sancak Y, Thoreen CC, Peterson TR, Lindquist RA, Kang SA, Spooner E, et al. PRAS40 is an insulin-regulated inhibitor of the mTORC1 protein kinase. *Mol Cell*. 2007;25(6):903-15.
11. Vander Haar E, Lee SI, Bandhakavi S, Griffin TJ, Kim DH. Insulin signalling to mTOR mediated by the Akt/PKB substrate PRAS40. *Nat Cell Biol*. 2007;9(3):316-23.
12. Sarbassov DD, Ali SM, Kim DH, Guertin DA, Latek RR, Erdjument-Bromage H, et al. Rictor, a novel binding partner of mTOR, defines a rapamycin-insensitive and raptor-independent pathway that regulates the cytoskeleton. *Curr Biol*. 2004;14(14):1296-302.
13. Jacinto E, Loewith R, Schmidt A, Lin S, Ruegg MA, Hall A, et al. Mammalian TOR complex 2 controls the actin cytoskeleton and is rapamycin insensitive. *Nat Cell Biol*. 2004;6(11):1122-8.
14. Frias MA, Thoreen CC, Jaffe JD, Schroder W, Sculley T, Carr SA, et al. mSin1 is necessary for Akt/PKB phosphorylation, and its isoforms define three distinct mTORC2s. *Curr Biol*. 2006;16(18):1865-70.
15. Jacinto E, Facchinetti V, Liu D, Soto N, Wei S, Jung SY, et al. SIN1/MIP1 maintains rictor-mTOR complex integrity and regulates Akt phosphorylation and substrate specificity. *Cell*. 2006;127(1):125-37.
16. Pearce LR, Huang X, Boudeau J, Pawlowski R, Wullschlegel S, Deak M, et al. Identification of Protor as a novel Rictor-binding component of mTOR complex-2. *Biochem J*. 2007;405(3):513-22.
17. Kim DH, Sarbassov DD, Ali SM, Latek RR, Guntur KV, Erdjument-Bromage H, et al. GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol Cell*. 2003;11(4):895-904.
18. Peterson TR, Laplante M, Thoreen CC, Sancak Y, Kang SA, Kuehl WM, et al. DEPTOR is an mTOR inhibitor frequently overexpressed in multiple myeloma cells and required for their survival. *Cell*. 2009;137(5):873-86.
19. Kaizuka T, Hara T, Oshiro N, Kikkawa U, Yonezawa K, Takehana K, et al. Tti1 and Tel2 are critical factors in mammalian target of rapamycin complex assembly. *J Biol Chem*. 2010;285(26):20109-16.
20. Vigneri R, Goldfine ID, Frittitta L. Insulin, insulin receptors, and cancer. *J Endocrinol Invest*. 2016.
21. Dibble CC, Cantley LC. Regulation of mTORC1 by PI3K signaling. *Trends Cell Biol*. 2015;25(9):545-55.

22. Alessi DR, Andjelkovic M, Caudwell B, Cron P, Morrice N, Cohen P, et al. Mechanism of activation of protein kinase B by insulin and IGF-1. *EMBO J.* 1996;15(23):6541-51.
23. Kovacina KS, Park GY, Bae SS, Guzzetta AW, Schaefer E, Birnbaum MJ, et al. Identification of a proline-rich Akt substrate as a 14-3-3 binding partner. *J Biol Chem.* 2003;278(12):10189-94.
24. Nascimento EB, Snel M, Guigas B, van der Zon GC, Kriek J, Maassen JA, et al. Phosphorylation of PRAS40 on Thr246 by PKB/AKT facilitates efficient phosphorylation of Ser183 by mTORC1. *Cell Signal.* 2010;22(6):961-7.
25. Dibble CC, Elis W, Menon S, Qin W, Klekota J, Asara JM, et al. TBC1D7 is a third subunit of the TSC1-TSC2 complex upstream of mTORC1. *Mol Cell.* 2012;47(4):535-46.
26. Inoki K, Li Y, Zhu T, Wu J, Guan KL. TSC2 is phosphorylated and inhibited by Akt and suppresses mTOR signalling. *Nat Cell Biol.* 2002;4(9):648-57.
27. Inoki K, Li Y, Xu T, Guan KL. Rheb GTPase is a direct target of TSC2 GAP activity and regulates mTOR signaling. *Genes Dev.* 2003;17(15):1829-34.
28. Long X, Lin Y, Ortiz-Vega S, Yonezawa K, Avruch J. Rheb binds and regulates the mTOR kinase. *Curr Biol.* 2005;15(8):702-13.
29. Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L, et al. The Rag GTPases bind raptor and mediate amino acid signaling to mTORC1. *Science.* 2008;320(5882):1496-501.
30. Sancak Y, Sabatini DM. Rag proteins regulate amino-acid-induced mTORC1 signalling. *Biochem Soc Trans.* 2009;37(Pt 1):289-90.
31. Sancak Y, Bar-Peled L, Zoncu R, Markhard AL, Nada S, Sabatini DM. Ragulator-Rag complex targets mTORC1 to the lysosomal surface and is necessary for its activation by amino acids. *Cell.* 2010;141(2):290-303.
32. Kim E, Goraksha-Hicks P, Li L, Neufeld TP, Guan KL. Regulation of TORC1 by Rag GTPases in nutrient response. *Nat Cell Biol.* 2008;10(8):935-45.
33. Demetriades C, Doumpas N, Teleman AA. Regulation of TORC1 in response to amino acid starvation via lysosomal recruitment of TSC2. *Cell.* 2014;156(4):786-99.
34. Carroll B, Maetzel D, Maddocks OD, Otten G, Ratcliff M, Smith GR, et al. Control of TSC2-Rheb signaling axis by arginine regulates mTORC1 activity. *Elife.* 2016;5.
35. Demetriades C, Plescher M, Teleman AA. Lysosomal recruitment of TSC2 is a universal response to cellular stress. *Nat Commun.* 2016;7:10662.
36. Tato I, Bartrons R, Ventura F, Rosa JL. Amino acids activate mammalian target of rapamycin complex 2 (mTORC2) via PI3K/Akt signaling. *J Biol Chem.* 2011;286(8):6128-42.
37. Dalle Pezze P, Ruf S, Sonntag AG, Langelaar-Makkinje M, Hall P, Heberle AM, et al. A systems study reveals concurrent activation of AMPK and mTOR by amino acids. *Nature Communications.* 2016;7:13254.
38. Inoki K, Zhu T, Guan KL. TSC2 mediates cellular energy response to control cell growth and survival. *Cell.* 2003;115(5):577-90.
39. Gwinn DM, Shackelford DB, Egan DF, Mihaylova MM, Mery A, Vasquez DS, et al. AMPK phosphorylation of raptor mediates a metabolic checkpoint. *Mol Cell.* 2008;30(2):214-26.
40. Burnett PE, Barrow RK, Cohen NA, Snyder SH, Sabatini DM. RAFT1 phosphorylation of the translational regulators p70 S6 kinase and 4E-BP1. *Proc Natl Acad Sci U S A.* 1998;95(4):1432-7.
41. Brunn GJ, Hudson CC, Sekulic A, Williams JM, Hosoi H, Houghton PJ, et al. Phosphorylation of the translational repressor PHAS-I by the mammalian target of rapamycin. *Science.* 1997;277(5322):99-101.
42. Holz MK, Ballif BA, Gygi SP, Blenis J. mTOR and S6K1 mediate assembly of the translation preinitiation complex through dynamic protein interchange and ordered phosphorylation events. *Cell.* 2005;123(4):569-80.
43. Pause A, Belsham GJ, Gingras AC, Donze O, Lin TA, Lawrence JC, Jr., et al. Insulin-dependent stimulation of protein synthesis by phosphorylation of a regulator of 5'-cap function. *Nature.* 1994;371(6500):762-7.
44. Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nat Cell Biol.* 2011;13(2):132-41.

45. Carroll B, Korolchuk VI, Sarkar S. Amino acids and autophagy: cross-talk and co-operation to control cellular homeostasis. *Amino Acids*. 2015;47(10):2065-88.
46. Ganley IG, Lam du H, Wang J, Ding X, Chen S, Jiang X. ULK1.ATG13.FIP200 complex mediates mTOR signaling and is essential for autophagy. *J Biol Chem*. 2009;284(18):12297-305.
47. Hosokawa N, Hara T, Kaizuka T, Kishi C, Takamura A, Miura Y, et al. Nutrient-dependent mTORC1 association with the ULK1-Atg13-FIP200 complex required for autophagy. *Mol Biol Cell*. 2009;20(7):1981-91.
48. Jung CH, Jun CB, Ro SH, Kim YM, Otto NM, Cao J, et al. ULK-Atg13-FIP200 complexes mediate mTOR signaling to the autophagy machinery. *Mol Biol Cell*. 2009;20(7):1992-2003.
49. Egan DF, Shackelford DB, Mihaylova MM, Gelino S, Kohnz RA, Mair W, et al. Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects energy sensing to mitophagy. *Science*. 2011;331(6016):456-61.
50. Tzatsos A, Kandror KV. Nutrients suppress phosphatidylinositol 3-kinase/Akt signaling via raptor-dependent mTOR-mediated insulin receptor substrate 1 phosphorylation. *Mol Cell Biol*. 2006;26(1):63-76.
51. Um SH, Frigerio F, Watanabe M, Picard F, Joaquin M, Sticker M, et al. Absence of S6K1 protects against age- and diet-induced obesity while enhancing insulin sensitivity. *Nature*. 2004;431(7005):200-5.
52. Yu Y, Yoon SO, Poulogiannis G, Yang Q, Ma XM, Villen J, et al. Phosphoproteomic analysis identifies Grb10 as an mTORC1 substrate that negatively regulates insulin signaling. *Science*. 2011;332(6035):1322-6.
53. Hsu PP, Kang SA, Rameseder J, Zhang Y, Ottina KA, Lim D, et al. The mTOR-regulated phosphoproteome reveals a mechanism of mTORC1-mediated inhibition of growth factor signaling. *Science*. 2011;332(6035):1317-22.
54. Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, et al. Ablation in mice of the mTORC components raptor, rictor, or mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not S6K1. *Dev Cell*. 2006;11(6):859-71.
55. Garcia-Martinez JM, Alessi DR. mTOR complex 2 (mTORC2) controls hydrophobic motif phosphorylation and activation of serum- and glucocorticoid-induced protein kinase 1 (SGK1). *Biochem J*. 2008;416(3):375-85.
56. Zinzalla V, Stracka D, Oppliger W, Hall MN. Activation of mTORC2 by association with the ribosome. *Cell*. 2011;144(5):757-68.
57. Dalle Pezze P, Sonntag AG, Thien A, Prentzell MT, Godel M, Fischer S, et al. A dynamic network model of mTOR signaling reveals TSC-independent mTORC2 regulation. *Sci Signal*. 2012;5(217):ra25.
58. Wahane SD, Hellbach N, Prentzell MT, Weise SC, Vezzali R, Kreutz C, et al. PI3K-p110-alpha-subtype signalling mediates survival, proliferation and neurogenesis of cortical progenitor cells via activation of mTORC2. *J Neurochem*. 2014;130(2):255-67.
59. Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science*. 2005;307(5712):1098-101.
60. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, Hu LS, et al. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell*. 1999;96(6):857-68.
61. Guo S, Rena G, Cichy S, He X, Cohen P, Unterman T. Phosphorylation of serine 256 by protein kinase B disrupts transactivation by FKHR and mediates effects of insulin on insulin-like growth factor-binding protein-1 promoter activity through a conserved insulin response sequence. *J Biol Chem*. 1999;274(24):17184-92.
62. Dijkers PF, Medema RH, Lammers JW, Koenderman L, Coffey PJ. Expression of the pro-apoptotic Bcl-2 family member Bim is regulated by the forkhead transcription factor FKHR-L1. *Curr Biol*. 2000;10(19):1201-4.
63. Modur V, Nagarajan R, Evers BM, Milbrandt J. FOXO proteins regulate tumor necrosis factor-related apoptosis inducing ligand expression. Implications for PTEN mutation in prostate cancer. *J Biol Chem*. 2002;277(49):47928-37.

64. Kops GJ, Dansen TB, Polderman PE, Saarloos I, Wirtz KW, Coffey PJ, et al. Forkhead transcription factor FOXO3a protects quiescent cells from oxidative stress. *Nature*. 2002;419(6904):316-21.
65. Nemoto S, Finkel T. Redox regulation of forkhead proteins through a p66shc-dependent signaling pathway. *Science*. 2002;295(5564):2450-2.
66. Ramaswamy S, Nakamura N, Sansal I, Bergeron L, Sellers WR. A novel mechanism of gene regulation and tumor suppression by the transcription factor FKHR. *Cancer Cell*. 2002;2(1):81-91.
67. Tran H, Brunet A, Grenier JM, Datta SR, Fornace AJ, Jr., DiStefano PS, et al. DNA repair pathway stimulated by the forkhead transcription factor FOXO3a through the Gadd45 protein. *Science*. 2002;296(5567):530-4.
68. Bruhn MA, Pearson RB, Hannan RD, Sheppard KE. Second AKT: the rise of SGK in cancer signalling. *Growth Factors*. 2010;28(6):394-408.
69. Vanhaesebroeck B, Whitehead MA, Pineiro R. Molecules in medicine mini-review: isoforms of PI3K in biology and disease. *J Mol Med (Berl)*. 2016;94(1):5-11.
70. Martins R, Lithgow GJ, Link W. Long live FOXO: unraveling the role of FOXO proteins in aging and longevity. *Aging Cell*. 2016;15(2):196-207.
71. Mosthaf L, Grako K, Dull TJ, Coussens L, Ullrich A, McClain DA. Functionally distinct insulin receptors generated by tissue-specific alternative splicing. *EMBO J*. 1990;9(8):2409-13.
72. Moller DE, Yokota A, Caro JF, Flier JS. Tissue-specific expression of two alternatively spliced insulin receptor mRNAs in man. *Mol Endocrinol*. 1989;3(8):1263-9.
73. Frasca F, Pandini G, Scalia P, Sciacca L, Mineo R, Costantino A, et al. Insulin receptor isoform A, a newly recognized, high-affinity insulin-like growth factor II receptor in fetal and cancer cells. *Mol Cell Biol*. 1999;19(5):3278-88.
74. Yamaguchi Y, Flier JS, Benecke H, Ransil BJ, Moller DE. Ligand-binding properties of the two isoforms of the human insulin receptor. *Endocrinology*. 1993;132(3):1132-8.
75. Sciacchitano S, Taylor SI. Cloning, tissue expression, and chromosomal localization of the mouse IRS-3 gene. *Endocrinology*. 1997;138(11):4931-40.
76. Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Proteomics. Tissue-based map of the human proteome. *Science*. 2015;347(6220):1260419.
77. He W, Craparo A, Zhu Y, O'Neill TJ, Wang LM, Pierce JH, et al. Interaction of insulin receptor substrate-2 (IRS-2) with the insulin and insulin-like growth factor I receptors. Evidence for two distinct phosphotyrosine-dependent interaction domains within IRS-2. *J Biol Chem*. 1996;271(20):11641-5.
78. Sawka-Verhelle D, Tartare-Deckert S, White MF, Van Obberghen E. Insulin receptor substrate-2 binds to the insulin receptor through its phosphotyrosine-binding domain and through a newly identified domain comprising amino acids 591-786. *J Biol Chem*. 1996;271(11):5980-3.
79. Wu J, Tseng YD, Xu CF, Neubert TA, White MF, Hubbard SR. Structural and biochemical characterization of the KRLB region in insulin receptor substrate-2. *Nat Struct Mol Biol*. 2008;15(3):251-8.
80. Ogihara T, Shin BC, Anai M, Katagiri H, Inukai K, Funaki M, et al. Insulin receptor substrate (IRS)-2 is dephosphorylated more rapidly than IRS-1 via its association with phosphatidylinositol 3-kinase in skeletal muscle cells. *J Biol Chem*. 1997;272(19):12868-73.
81. Huang C, Thirone AC, Huang X, Klip A. Differential contribution of insulin receptor substrates 1 versus 2 to insulin signaling and glucose uptake in L6 myotubes. *J Biol Chem*. 2005;280(19):19426-35.
82. Valverde AM, Lorenzo M, Pons S, White MF, Benito M. Insulin receptor substrate (IRS) proteins IRS-1 and IRS-2 differential signaling in the insulin/insulin-like growth factor-I pathways in fetal brown adipocytes. *Mol Endocrinol*. 1998;12(5):688-97.
83. Nagle JA, Ma Z, Byrne MA, White MF, Shaw LM. Involvement of insulin receptor substrate 2 in mammary tumor metastasis. *Mol Cell Biol*. 2004;24(22):9726-35.
84. Ma Z, Gibson SL, Byrne MA, Zhang J, White MF, Shaw LM. Suppression of insulin receptor substrate 1 (IRS-1) promotes mammary tumor metastasis. *Mol Cell Biol*. 2006;26(24):9338-51.

85. Lavan BE, Fantin VR, Chang ET, Lane WS, Keller SR, Lienhard GE. A novel 160-kDa phosphotyrosine protein in insulin-treated embryonic kidney cells is a new member of the insulin receptor substrate family. *J Biol Chem*. 1997;272(34):21403-7.
86. Shimwell NJ, Martin A, Bruton RK, Blackford AN, Sedgwick GG, Gallimore PH, et al. Adenovirus 5 E1A is responsible for increased expression of insulin receptor substrate 4 in established adenovirus 5-transformed cell lines and interacts with IRS components activating the PI3 kinase/Akt signalling pathway. *Oncogene*. 2009;28(5):686-97.
87. Hopkins BD, Fine B, Steinbach N, Dendy M, Rapp Z, Shaw J, et al. A secreted PTEN phosphatase that enters cells to alter signaling and survival. *Science*. 2013;341(6144):399-402.
88. Liang H, He S, Yang J, Jia X, Wang P, Chen X, et al. PTENalpha, a PTEN isoform translated through alternative initiation, regulates mitochondrial function and energy metabolism. *Cell Metab*. 2014;19(5):836-48.
89. Taniyama Y, Ito M, Sato K, Kuester C, Veit K, Tremp G, et al. Akt3 overexpression in the heart results in progression from adaptive to maladaptive hypertrophy. *J Mol Cell Cardiol*. 2005;38(2):375-85.
90. Brodbeck D, Hill MM, Hemmings BA. Two splice variants of protein kinase B gamma have different regulatory capacity depending on the presence or absence of the regulatory phosphorylation site serine 472 in the carboxyl-terminal hydrophobic domain. *J Biol Chem*. 2001;276(31):29550-8.
91. Altomare DA, Guo K, Cheng JQ, Sonoda G, Walsh K, Testa JR. Cloning, chromosomal localization and expression analysis of the mouse Akt2 oncogene. *Oncogene*. 1995;11(6):1055-60.
92. Altomare DA, Lyons GE, Mitsuuchi Y, Cheng JQ, Testa JR. Akt2 mRNA is highly expressed in embryonic brown fat and the AKT2 kinase is activated by insulin. *Oncogene*. 1998;16(18):2407-11.
93. Brodbeck D, Cron P, Hemmings BA. A human protein kinase Bgamma with regulatory phosphorylation sites in the activation loop and in the C-terminal hydrophobic domain. *J Biol Chem*. 1999;274(14):9133-6.
94. Calera MR, Martinez C, Liu H, Jack AK, Birnbaum MJ, Pilch PF. Insulin increases the association of Akt-2 with Glut4-containing vesicles. *J Biol Chem*. 1998;273(13):7201-4.
95. Brognard J, Sierrecki E, Gao T, Newton AC. PHLPP and a second isoform, PHLPP2, differentially attenuate the amplitude of Akt signaling by regulating distinct Akt isoforms. *Mol Cell*. 2007;25(6):917-31.
96. Chin YR, Toker A. The actin-bundling protein palladin is an Akt1-specific substrate that regulates breast cancer cell migration. *Mol Cell*. 2010;38(3):333-44.
97. Cenni V, Bavelloni A, Beretti F, Tagliavini F, Manzoli L, Lattanzi G, et al. Ankrd2/ARPP is a novel Akt2 specific substrate and regulates myogenic differentiation upon cellular exposure to H(2)O(2). *Mol Biol Cell*. 2011;22(16):2946-56.
98. Chen WS, Xu PZ, Gottlob K, Chen ML, Sokol K, Shiyanova T, et al. Growth retardation and increased apoptosis in mice with homozygous disruption of the Akt1 gene. *Genes Dev*. 2001;15(17):2203-8.
99. Cho H, Thorvaldsen JL, Chu Q, Feng F, Birnbaum MJ. Akt1/PKBalpha is required for normal growth but dispensable for maintenance of glucose homeostasis in mice. *J Biol Chem*. 2001;276(42):38349-52.
100. Dillon RL, Marcotte R, Hennessy BT, Woodgett JR, Mills GB, Muller WJ. Akt1 and akt2 play distinct roles in the initiation and metastatic phases of mammary tumor progression. *Cancer Res*. 2009;69(12):5057-64.
101. Maroulakou IG, Oemler W, Naber SP, Tschlis PN. Akt1 ablation inhibits, whereas Akt2 ablation accelerates, the development of mammary adenocarcinomas in mouse mammary tumor virus (MMTV)-ErbB2/neu and MMTV-polyoma middle T transgenic mice. *Cancer Res*. 2007;67(1):167-77.
102. Cho H, Mu J, Kim JK, Thorvaldsen JL, Chu Q, Crenshaw EB, 3rd, et al. Insulin resistance and a diabetes mellitus-like syndrome in mice lacking the protein kinase Akt2 (PKB beta). *Science*. 2001;292(5522):1728-31.

103. Garofalo RS, Orena SJ, Rafidi K, Torchia AJ, Stock JL, Hildebrandt AL, et al. Severe diabetes, age-dependent loss of adipose tissue, and mild growth deficiency in mice lacking Akt2/PKB beta. *J Clin Invest*. 2003;112(2):197-208.
104. Grottke A, Ewald F, Lange T, Norz D, Herzberger C, Bach J, et al. Downregulation of AKT3 Increases Migration and Metastasis in Triple Negative Breast Cancer Cells by Upregulating S100A4. *PLoS One*. 2016;11(1):e0146370.
105. Diez H, Garrido JJ, Wandosell F. Specific roles of Akt iso forms in apoptosis and axon growth regulation in neurons. *PloS one*. 2012;7(4):e32715.
106. Tschopp O, Yang ZZ, Brodbeck D, Dummmler BA, Hemmings-Mieszczak M, Watanabe T, et al. Essential role of protein kinase B gamma (PKB gamma/Akt3) in postnatal brain development but not in glucose homeostasis. *Development*. 2005;132(13):2943-54.
107. Stottrup C, Tsang T, Chin YR. Upregulation of AKT3 Confers Resistance to the AKT Inhibitor MK2206 in Breast Cancer. *Mol Cancer Ther*. 2016;15(8):1964-74.
108. Xu L, Sterner C, Maheshwar MM, Wilson PJ, Nellist M, Short PM, et al. Alternative splicing of the tuberous sclerosis 2 (TSC2) gene in human and mouse tissues. *Genomics*. 1995;27(3):475-80.
109. Xiao GH, Jin F, Yeung RS. Identification of tuberous sclerosis 2 messenger RNA splice variants that are conserved and differentially expressed in rat and human tissues. *Cell Growth Differ*. 1995;6(9):1185-91.
110. Ekong R, Nellist M, Hoogeveen-Westerveld M, Wentink M, Panzer J, Sparagana S, et al. Variants Within TSC2 Exons 25 and 31 Are Very Unlikely to Cause Clinically Diagnosable Tuberous Sclerosis. *Hum Mutat*. 2016;37(4):364-70.
111. Manning BD, Tee AR, Logsdon MN, Blenis J, Cantley LC. Identification of the tuberous sclerosis complex-2 tumor suppressor gene product tuberlin as a target of the phosphoinositide 3-kinase/akt pathway. *Mol Cell*. 2002;10(1):151-62.
112. Cai SL, Tee AR, Short JD, Bergeron JM, Kim J, Shen J, et al. Activity of TSC2 is inhibited by AKT-mediated phosphorylation and membrane partitioning. *J Cell Biol*. 2006;173(2):279-89.
113. Mayer K, Ballhausen W, Rott HD. Mutation screening of the entire coding regions of the TSC1 and the TSC2 gene with the protein truncation test (PTT) identifies frequent splicing defects. *Hum Mutat*. 1999;14(5):401-11.
114. Panasyuk G, Nemazanyy I, Zhyvoloup A, Filonenko V, Davies D, Robson M, et al. mTORbeta splicing isoform promotes cell proliferation and tumorigenesis. *J Biol Chem*. 2009;284(45):30807-14.
115. Sun C, Southard C, Di Rienzo A. Characterization of a novel splicing variant in the RAPTOR gene. *Mutat Res*. 2009;662(1-2):88-92.
116. Nojima H, Tokunaga C, Eguchi S, Oshiro N, Hidayat S, Yoshino K, et al. The mammalian target of rapamycin (mTOR) partner, raptor, binds the mTOR substrates p70 S6 kinase and 4E-BP1 through their TOR signaling (TOS) motif. *J Biol Chem*. 2003;278(18):15461-4.
117. Schroder WA, Buck M, Cloonan N, Hancock JF, Suhrbier A, Sculley T, et al. Human Sin1 contains Ras-binding and pleckstrin homology domains and suppresses Ras signalling. *Cell Signal*. 2007;19(6):1279-89.
118. Pearce LR, Sommer EM, Sakamoto K, Wullschlegel S, Alessi DR. Protor-1 is required for efficient mTORC2-mediated activation of SGK1 in the kidney. *Biochem J*. 2011;436(1):169-79.
119. Holmes B, Artinian N, Anderson L, Martin J, Masri J, Cloninger C, et al. Protor-2 interacts with tristetraprolin to regulate mRNA stability during stress. *Cell Signal*. 2012;24(1):309-15.
120. Banerjee P, Ahmad MF, Grove JR, Kozlosky C, Price DJ, Avruch J. Molecular structure of a major insulin/mitogen-activated 70-kDa S6 protein kinase. *Proc Natl Acad Sci U S A*. 1990;87(21):8550-4.
121. Kozma SC, Ferrari S, Bassand P, Siegmann M, Totty N, Thomas G. Cloning of the mitogen-activated S6 kinase from rat liver reveals an enzyme of the second messenger subfamily. *Proc Natl Acad Sci U S A*. 1990;87(19):7365-9.
122. Lee-Fruman KK, Kuo CJ, Lippincott J, Terada N, Blenis J. Characterization of S6K2, a novel kinase homologous to S6K1. *Oncogene*. 1999;18(36):5108-14.
123. Tavares MR, Pavan IC, Amaral CL, Meneguello L, Luchessi AD, Simabuco FM. The S6K protein family in health and disease. *Life Sci*. 2015;131:1-10.

124. Pullen N, Dennis PB, Andjelkovic M, Dufner A, Kozma SC, Hemmings BA, et al. Phosphorylation and activation of p70s6k by PDK1. *Science*. 1998;279(5351):707-10.
125. Talvas J, Obled A, Fafournoux P, Mordier S. Regulation of protein synthesis by leucine starvation involves distinct mechanisms in mouse C2C12 myoblasts and myotubes. *J Nutr*. 2006;136(6):1466-71.
126. Burnett PE, Blackshaw S, Lai MM, Qureshi IA, Burnett AF, Sabatini DM, et al. Neurabin is a synaptic protein linking p70 S6 kinase and the neuronal cytoskeleton. *Proc Natl Acad Sci U S A*. 1998;95(14):8351-6.
127. Oh CH, Park SY, Han JS. Phospholipase D1 is required for lipopolysaccharide-induced tumor necrosis factor- α expression and production through S6K1/JNK/c-Jun pathway in Raw 264.7 cells. *Cytokine*. 2014;66(1):69-77.
128. Chou MM, Blenis J. The 70 kDa S6 kinase complexes with and is activated by the Rho family G proteins Cdc42 and Rac1. *Cell*. 1996;85(4):573-83.
129. Hong S, Zhao B, Lombard DB, Fingar DC, Inoki K. Cross-talk between sirtuin and mammalian target of rapamycin complex 1 (mTORC1) signaling in the regulation of S6 kinase 1 (S6K1) phosphorylation. *J Biol Chem*. 2014;289(19):13132-41.
130. Cruz R, Hedden L, Boyer D, Kharas MG, Fruman DA, Lee-Fruman KK. S6 kinase 2 potentiates interleukin-3-driven cell proliferation. *J Leukoc Biol*. 2005;78(6):1378-85.
131. Pardo OE, Arcaro A, Salerno G, Tetley TD, Valovka T, Gout I, et al. Novel cross talk between MEK and S6K2 in FGF-2 induced proliferation of SCLC cells. *Oncogene*. 2001;20(52):7658-67.
132. Valovka T, Verdier F, Cramer R, Zhyvoloup A, Fenton T, Rebholz H, et al. Protein kinase C phosphorylates ribosomal protein S6 kinase beta11 and regulates its subcellular localization. *Mol Cell Biol*. 2003;23(3):852-63.
133. Pende M, Um SH, Mieulet V, Sticker M, Goss VL, Mestan J, et al. S6K1(-/-)/S6K2(-/-) mice exhibit perinatal lethality and rapamycin-sensitive 5'-terminal oligopyrimidine mRNA translation and reveal a mitogen-activated protein kinase-dependent S6 kinase pathway. *Mol Cell Biol*. 2004;24(8):3112-24.
134. Dennis MD, Jefferson LS, Kimball SR. Role of p70S6K1-mediated phosphorylation of eIF4B and PDCD4 proteins in the regulation of protein synthesis. *J Biol Chem*. 2012;287(51):42890-9.
135. Wang X, Li W, Williams M, Terada N, Alessi DR, Proud CG. Regulation of elongation factor 2 kinase by p90(RSK1) and p70 S6 kinase. *EMBO J*. 2001;20(16):4370-9.
136. Harada H, Andersen JS, Mann M, Terada N, Korsmeyer SJ. p70S6 kinase signals cell survival as well as growth, inactivating the pro-apoptotic molecule BAD. *Proc Natl Acad Sci U S A*. 2001;98(17):9666-70.
137. Lai KP, Leong WF, Chau JF, Jia D, Zeng L, Liu H, et al. S6K1 is a multifaceted regulator of Mdm2 that connects nutrient status and DNA damage response. *EMBO J*. 2010;29(17):2994-3006.
138. Zhang HH, Lipovsky AI, Dibble CC, Sahin M, Manning BD. S6K1 regulates GSK3 under conditions of mTOR-dependent feedback inhibition of Akt. *Mol Cell*. 2006;24(2):185-97.
139. Yamnik RL, Digilova A, Davis DC, Brodt ZN, Murphy CJ, Holz MK. S6 kinase 1 regulates estrogen receptor alpha in control of breast cancer cell proliferation. *J Biol Chem*. 2009;284(10):6361-9.
140. de Groot RP, Ballou LM, Sassone-Corsi P. Positive regulation of the cAMP-responsive activator CREM by the p70 S6 kinase: an alternative route to mitogen-induced gene expression. *Cell*. 1994;79(1):81-91.
141. Ismail HM, Myronova O, Tsuchiya Y, Niewiarowski A, Tsaneva I, Gout I. Identification of the general transcription factor Yin Yang 1 as a novel and specific binding partner for S6 kinase 2. *Cell Signal*. 2013;25(5):1054-63.
142. Kurebayashi Y, Nagai S, Ikejiri A, Ohtani M, Ichiyama K, Baba Y, et al. PI3K-Akt-mTORC1-S6K1/2 axis controls Th17 differentiation by regulating Gfi1 expression and nuclear translocation of RORgamma. *Cell Rep*. 2012;1(4):360-73.
143. Grove JR, Banerjee P, Balasubramanyam A, Coffey PJ, Price DJ, Avruch J, et al. Cloning and expression of two human p70 S6 kinase polypeptides differing only at their amino termini. *Mol Cell Biol*. 1991;11(11):5541-50.

144. Ben-Hur V, Denichenko P, Siegfried Z, Maimon A, Krainer A, Davidson B, et al. S6K1 alternative splicing modulates its oncogenic activity and regulates mTORC1. *Cell Rep.* 2013;3(1):103-15.
145. Coffey PJ, Woodgett JR. Differential subcellular localisation of two isoforms of p70 S6 protein kinase. *Biochem Biophys Res Commun.* 1994;198(2):780-6.
146. Reinhard C, Fernandez A, Lamb NJ, Thomas G. Nuclear localization of p85s6k: functional requirement for entry into S phase. *EMBO J.* 1994;13(7):1557-65.
147. Rosner M, Hengstschlager M. Nucleocytoplasmic localization of p70 S6K1, but not of its isoforms p85 and p31, is regulated by TSC2/mTOR. *Oncogene.* 2011;30(44):4509-22.
148. Ruf S, Heberle AM, Langelaar-Makkinje M, Gelino S, Wilkinson D, Gerbeth C, et al. PLK1 (polo like kinase 1) inhibits MTOR complex 1 and promotes autophagy. *Autophagy.* 2017:0.
149. Tsukiyama-Kohara K, Vidal SM, Gingras AC, Glover TW, Hanash SM, Heng H, et al. Tissue distribution, genomic structure, and chromosome mapping of mouse and human eukaryotic initiation factor 4E-binding proteins 1 and 2. *Genomics.* 1996;38(3):353-63.
150. Poulin F, Gingras AC, Olsen H, Chevalier S, Sonenberg N. 4E-BP3, a new member of the eukaryotic initiation factor 4E-binding protein family. *J Biol Chem.* 1998;273(22):14002-7.
151. Tee AR, Proud CG. Caspase cleavage of initiation factor 4E-binding protein 1 yields a dominant inhibitor of cap-dependent translation and reveals a novel regulatory motif. *Mol Cell Biol.* 2002;22(6):1674-83.
152. Tsukumo Y, Alain T, Fonseca BD, Nadon R, Sonenberg N. Translation control during prolonged mTORC1 inhibition mediated by 4E-BP3. *Nat Commun.* 2016;7:11776.
153. Chen CC, Lee JC, Chang MC. 4E-BP3 regulates eIF4E-mediated nuclear mRNA export and interacts with replication protein A2. *FEBS Lett.* 2012;586(16):2260-6.
154. Chan EY, Kir S, Tooze SA. siRNA screening of the kinome identifies ULK1 as a multidomain modulator of autophagy. *J Biol Chem.* 2007;282(35):25464-74.
155. Hara T, Takamura A, Kishi C, Iemura S, Natsume T, Guan JL, et al. FIP200, a ULK-interacting protein, is required for autophagosome formation in mammalian cells. *J Cell Biol.* 2008;181(3):497-510.
156. Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell.* 2013;153(6):1194-217.
157. Powers RW, 3rd, Kaeberlein M, Caldwell SD, Kennedy BK, Fields S. Extension of chronological life span in yeast by decreased TOR pathway signaling. *Genes Dev.* 2006;20(2):174-84.
158. Swinnen E, Ghillebert R, Wilms T, Winderickx J. Molecular mechanisms linking the evolutionary conserved TORC1-Sch9 nutrient signalling branch to lifespan regulation in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 2014;14(1):17-32.
159. Robida-Stubbs S, Glover-Cutter K, Lamming DW, Mizunuma M, Narasimhan SD, Neumann-Haefelin E, et al. TOR signaling and rapamycin influence longevity by regulating SKN-1/Nrf and DAF-16/FoxO. *Cell Metab.* 2012;15(5):713-24.
160. Greer EL, Brunet A. Different dietary restriction regimens extend lifespan by both independent and overlapping genetic pathways in *C. elegans*. *Aging Cell.* 2009;8(2):113-27.
161. Vellai T, Takacs-Vellai K, Zhang Y, Kovacs AL, Orosz L, Muller F. Genetics: influence of TOR kinase on lifespan in *C. elegans*. *Nature.* 2003;426(6967):620.
162. Jia K, Chen D, Riddle DL. The TOR pathway interacts with the insulin signaling pathway to regulate *C. elegans* larval development, metabolism and life span. *Development.* 2004;131(16):3897-906.
163. Schreiber MA, Pierce-Shimomura JT, Chan S, Parry D, McIntire SL. Manipulation of behavioral decline in *Caenorhabditis elegans* with the Rag GTPase raga-1. *PLoS Genet.* 2010;6(5):e1000972.
164. Honjoh S, Yamamoto T, Uno M, Nishida E. Signalling through RHEB-1 mediates intermittent fasting-induced longevity in *C. elegans*. *Nature.* 2009;457(7230):726-30.
165. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice as long as wild type. *Nature.* 1993;366(6454):461-4.
166. Friedman DB, Johnson TE. A mutation in the age-1 gene in *Caenorhabditis elegans* lengthens life and reduces hermaphrodite fertility. *Genetics.* 1988;118(1):75-86.

167. Soukas AA, Kane EA, Carr CE, Melo JA, Ruvkun G. Rictor/TORC2 regulates fat metabolism, feeding, growth, and life span in *Caenorhabditis elegans*. *Genes Dev.* 2009;23(4):496-511.
168. Bjedov I, Toivonen JM, Kerr F, Slack C, Jacobson J, Foley A, et al. Mechanisms of life span extension by rapamycin in the fruit fly *Drosophila melanogaster*. *Cell Metab.* 2010;11(1):35-46.
169. Pletcher SD, Macdonald SJ, Marguerie R, Certa U, Stearns SC, Goldstein DB, et al. Genome-wide transcript profiles in aging and calorically restricted *Drosophila melanogaster*. *Curr Biol.* 2002;12(9):712-23.
170. Tatar M, Kopelman A, Epstein D, Tu MP, Yin CM, Garofalo RS. A mutant *Drosophila* insulin receptor homolog that extends life-span and impairs neuroendocrine function. *Science.* 2001;292(5514):107-10.
171. Clancy DJ, Gems D, Harshman LG, Oldham S, Stocker H, Hafen E, et al. Extension of life-span by loss of CHICO, a *Drosophila* insulin receptor substrate protein. *Science.* 2001;292(5514):104-6.
172. Kapahi P, Zid BM, Harper T, Koslover D, Sapin V, Benzer S. Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr Biol.* 2004;14(10):885-90.
173. Zid BM, Rogers AN, Katewa SD, Vargas MA, Kolipinski MC, Lu TA, et al. 4E-BP extends lifespan upon dietary restriction by enhancing mitochondrial activity in *Drosophila*. *Cell.* 2009;139(1):149-60.
174. Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, et al. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature.* 2009;460(7253):392-5.
175. Weindruch R, Walford RL. Dietary restriction in mice beginning at 1 year of age: effect on life-span and spontaneous cancer incidence. *Science.* 1982;215(4538):1415-8.
176. Cornu M, Albert V, Hall MN. mTOR in aging, metabolism, and cancer. *Curr Opin Genet Dev.* 2013;23(1):53-62.
177. Lamming DW, Ye L, Katajisto P, Goncalves MD, Saitoh M, Stevens DM, et al. Rapamycin-induced insulin resistance is mediated by mTORC2 loss and uncoupled from longevity. *Science.* 2012;335(6076):1638-43.
178. Selman C, Lingard S, Choudhury AI, Batterham RL, Claret M, Clements M, et al. Evidence for lifespan extension and delayed age-related biomarkers in insulin receptor substrate 1 null mice. *FASEB J.* 2008;22(3):807-18.
179. Selman C, Tullet JM, Wieser D, Irvine E, Lingard SJ, Choudhury AI, et al. Ribosomal protein S6 kinase 1 signaling regulates mammalian life span. *Science.* 2009;326(5949):140-4.
180. Taguchi A, Wartschow LM, White MF. Brain IRS2 signaling coordinates life span and nutrient homeostasis. *Science.* 2007;317(5836):369-72.
181. Selman C, Lingard S, Gems D, Partridge L, Withers DJ. Comment on "Brain IRS2 signaling coordinates life span and nutrient homeostasis". *Science.* 2008;320(5879):1012; author reply
182. Lamming DW, Mihaylova MM, Katajisto P, Baar EL, Yilmaz OH, Hutchins A, et al. Depletion of Rictor, an essential protein component of mTORC2, decreases male lifespan. *Aging Cell.* 2014;13(5):911-7.
183. Bluher M, Michael MD, Peroni OD, Ueki K, Carter N, Kahn BB, et al. Adipose tissue selective insulin receptor knockout protects against obesity and obesity-related glucose intolerance. *Dev Cell.* 2002;3(1):25-38.
184. Polak P, Cybulski N, Feige JN, Auwerx J, Ruegg MA, Hall MN. Adipose-specific knockout of raptor results in lean mice with enhanced mitochondrial respiration. *Cell Metab.* 2008;8(5):399-410.
185. Kwiatkowski DJ, Zhang H, Bandura JL, Heiberger KM, Glogauer M, el-Hashemite N, et al. A mouse model of TSC1 reveals sex-dependent lethality from liver hemangiomas, and up-regulation of p70S6 kinase activity in Tsc1 null cells. *Hum Mol Genet.* 2002;11(5):525-34.
186. Kobayashi T, Minowa O, Kuno J, Mitani H, Hino O, Noda T. Renal carcinogenesis, hepatic hemangiomatosis, and embryonic lethality caused by a germ-line Tsc2 mutation in mice. *Cancer Res.* 1999;59(6):1206-11.
187. Brewer RA, Gibbs VK, Smith DL, Jr. Targeting glucose metabolism for healthy aging. *Nutr Healthy Aging.* 2016;4(1):31-46.
188. Boles A, Kandimalla R, Reddy PH. Dynamics of Diabetes and Obesity: Epidemiological Perspective. *Biochim Biophys Acta.* 2017.

189. Gomes MJ, Martinez PF, Pagan LU, Damatto RL, Cezar MD, Lima AR, et al. Skeletal muscle aging: influence of oxidative stress and physical exercise. *Oncotarget*. 2017.
190. Fantin VR, Wang Q, Lienhard GE, Keller SR. Mice lacking insulin receptor substrate 4 exhibit mild defects in growth, reproduction, and glucose homeostasis. *Am J Physiol Endocrinol Metab*. 2000;278(1):E127-33.
191. Bruning JC, Michael MD, Winnay JN, Hayashi T, Horsch D, Accili D, et al. A muscle-specific insulin receptor knockout exhibits features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Mol Cell*. 1998;2(5):559-69.
192. Kulkarni RN, Bruning JC, Winnay JN, Postic C, Magnuson MA, Kahn CR. Tissue-specific knockout of the insulin receptor in pancreatic beta cells creates an insulin secretory defect similar to that in type 2 diabetes. *Cell*. 1999;96(3):329-39.
193. Bruning JC, Gautam D, Burks DJ, Gillette J, Schubert M, Orban PC, et al. Role of brain insulin receptor in control of body weight and reproduction. *Science*. 2000;289(5487):2122-5.
194. Mora A, Lipina C, Tronche F, Sutherland C, Alessi DR. Deficiency of PDK1 in liver results in glucose intolerance, impairment of insulin-regulated gene expression and liver failure. *Biochem J*. 2005;385(Pt 3):639-48.
195. Kubota N, Tobe K, Terauchi Y, Eto K, Yamauchi T, Suzuki R, et al. Disruption of insulin receptor substrate 2 causes type 2 diabetes because of liver insulin resistance and lack of compensatory beta-cell hyperplasia. *Diabetes*. 2000;49(11):1880-9.
196. Michael MD, Kulkarni RN, Postic C, Previs SF, Shulman GI, Magnuson MA, et al. Loss of insulin signaling in hepatocytes leads to severe insulin resistance and progressive hepatic dysfunction. *Mol Cell*. 2000;6(1):87-97.
197. Mora A, Davies AM, Bertrand L, Sharif I, Budas GR, Jovanovic S, et al. Deficiency of PDK1 in cardiac muscle results in heart failure and increased sensitivity to hypoxia. *EMBO J*. 2003;22(18):4666-76.
198. Risson V, Mazelin L, Roceri M, Sanchez H, Moncollin V, Corneloup C, et al. Muscle inactivation of mTOR causes metabolic and dystrophin defects leading to severe myopathy. *J Cell Biol*. 2009;187(6):859-74.
199. Bentzinger CF, Romanino K, Cloetta D, Lin S, Mascarenhas JB, Oliveri F, et al. Skeletal muscle-specific ablation of raptor, but not of rictor, causes metabolic changes and results in muscle dystrophy. *Cell metabolism*. 2008;8(5):411-24.
200. Aguilar V, Alliouachene S, Sotiropoulos A, Sobering A, Athea Y, Djouadi F, et al. S6 kinase deletion suppresses muscle growth adaptations to nutrient availability by activating AMP kinase. *Cell Metab*. 2007;5(6):476-87.
201. Araki E, Lipes MA, Patti ME, Bruning JC, Haag B, 3rd, Johnson RS, et al. Alternative pathway of insulin signalling in mice with targeted disruption of the IRS-1 gene. *Nature*. 1994;372(6502):186-90.
202. Withers DJ, Gutierrez JS, Towery H, Burks DJ, Ren JM, Previs S, et al. Disruption of IRS-2 causes type 2 diabetes in mice. *Nature*. 1998;391(6670):900-4.
203. Nojima A, Yamashita M, Yoshida Y, Shimizu I, Ichimiya H, Kamimura N, et al. Haploinsufficiency of akt1 prolongs the lifespan of mice. *PLoS One*. 2013;8(7):e69178.
204. Tsukiyama-Kohara K, Poulin F, Kohara M, DeMaria CT, Cheng A, Wu Z, et al. Adipose tissue reduction in mice lacking the translational inhibitor 4E-BP1. *Nat Med*. 2001;7(10):1128-32.
205. Le Bacquer O, Petroulakis E, Paglialunga S, Poulin F, Richard D, Cianflone K, et al. Elevated sensitivity to diet-induced obesity and insulin resistance in mice lacking 4E-BP1 and 4E-BP2. *J Clin Invest*. 2007;117(2):387-96.
206. Mattison JA, Colman RJ, Beasley TM, Allison DB, Kemnitz JW, Roth GS, et al. Caloric restriction improves health and survival of rhesus monkeys. *Nat Commun*. 2017;8:14063.
207. Mannick JB, Del Giudice G, Lattanzi M, Valiante NM, Praestgaard J, Huang B, et al. mTOR inhibition improves immune function in the elderly. *Sci Transl Med*. 2014;6(268):268ra179.
208. Murabito JM, Yuan R, Lunetta KL. The search for longevity and healthy aging genes: insights from epidemiological studies and samples of long-lived individuals. *J Gerontol A Biol Sci Med Sci*. 2012;67(5):470-9.

209. Erikson GA, Bodian DL, Rueda M, Molparia B, Scott ER, Scott-Van Zeeland AA, et al. Whole-Genome Sequencing of a Healthy Aging Cohort. *Cell*. 2016;165(4):1002-11.
210. Kojima T, Kamei H, Aizu T, Arai Y, Takayama M, Nakazawa S, et al. Association analysis between longevity in the Japanese population and polymorphic variants of genes involved in insulin and insulin-like growth factor 1 signaling pathways. *Exp Gerontol*. 2004;39(11-12):1595-8.
211. Pawlikowska L, Hu D, Huntsman S, Sung A, Chu C, Chen J, et al. Association of common genetic variation in the insulin/IGF1 signaling pathway with human longevity. *Aging Cell*. 2009;8(4):460-72.
212. Passtoors WM, Beekman M, Deelen J, van der Breggen R, Maier AB, Guigas B, et al. Gene expression analysis of mTOR pathway: association with human longevity. *Aging Cell*. 2013;12(1):24-31.
213. van Heemst D, Beekman M, Mooijaart SP, Heijmans BT, Brandt BW, Zwaan BJ, et al. Reduced insulin/IGF-1 signalling and human longevity. *Aging Cell*. 2005;4(2):79-85.
214. Heintz C, Doktor TK, Lanjuin A, Escoubas CC, Zhang Y, Weir HJ, et al. Splicing factor 1 modulates dietary restriction and TORC1 pathway longevity in *C. elegans*. *Nature*. 2017;541(7635):102-6.
215. Sveen A, Kilpinen S, Ruusulehto A, Lothe RA, Skotheim RI. Aberrant RNA splicing in cancer; expression changes and driver mutations of splicing factor genes. *Oncogene*. 2016;35(19):2413-27.
216. Yang X, Coulombe-Huntington J, Kang S, Sheynkman GM, Hao T, Richardson A, et al. Widespread Expansion of Protein Interaction Capabilities by Alternative Splicing. *Cell*. 2016;164(4):805-17.
217. Mohanty S, Spinas GA, Maedler K, Zuellig RA, Lehmann R, Donath MY, et al. Overexpression of IRS2 in isolated pancreatic islets causes proliferation and protects human beta-cells from hyperglycemia-induced apoptosis. *Exp Cell Res*. 2005;303(1):68-78.
218. Niessen M. On the role of IRS2 in the regulation of functional beta-cell mass. *Arch Physiol Biochem*. 2006;112(2):65-73.
219. Nakatani K, Sakaue H, Thompson DA, Weigel RJ, Roth RA. Identification of a human Akt3 (protein kinase B gamma) which contains the regulatory serine phosphorylation site. *Biochem Biophys Res Commun*. 1999;257(3):906-10.
220. Lin TA, Lawrence JC, Jr. Control of the translational regulators PHAS-I and PHAS-II by insulin and cAMP in 3T3-L1 adipocytes. *J Biol Chem*. 1996;271(47):30199-204.
221. Gingras AC, Sonenberg N. Adenovirus infection inactivates the translational inhibitors 4E-BP1 and 4E-BP2. *Virology*. 1997;237(1):182-6.
222. Accili D, Drago J, Lee EJ, Johnson MD, Cool MH, Salvatore P, et al. Early neonatal death in mice homozygous for a null allele of the insulin receptor gene. *Nat Genet*. 1996;12(1):106-9.
223. Hagiwara A, Cornu M, Cybulski N, Polak P, Betz C, Trapani F, et al. Hepatic mTORC2 activates glycolysis and lipogenesis through Akt, glucokinase, and SREBP1c. *Cell Metab*. 2012;15(5):725-38.

Figure 1.

Table 1.

Isoforms across the mTOR network					
	Name UniProt ID	Regulatory domains	Isoform regulation	Expression pattern	Isoform functions
Signalling modules	insulin	IRS1 P35568	PH, PTB, PI3K-BD, Grb2-BD, SHP2-BD	Dephosphorylated 60 min after insulin stimulation (80) Strong Grb2 association (82)	Ubiquitous (76) Signals to Akt1, Akt2 and MAPK pathway (81, 82) Actin remodelling(81) GLUT4 translocation(81)
		IRS2 Q9Y4H2	PH, PTB, KRLB, PI3K-BD, Grb2-BD, SHP2-BD	Dephosphorylated 10 min after insulin stimulation (80) Weak Grb2 association (82)	Ubiquitous (76) Signals to Akt2 and MAPK pathway (81) Metastasis (83, 84) Maintenance of β -cell mass (217, 218)
		IRS4 O14654	PH, PTB, PI3K-BD, Grb2-BD	Upregulated upon viral infection(86)	Brain, kidney, thymus, liver (85) Upon viral infection, leads to constitutive Akt activation (86)
	Akt	Akt1 P31749	PH, KD, HM	Localizes at the cytosol (94) Dephosphorylated by PHLPP2 (95)	Ubiquitous (76) Phosphorylates TSC2 (95), FoxOs (95) and palladin (96) Cell survival (98, 99) Cell growth (100, 101)
		Akt2 P31751	PH, KD, HM	Localizes in GLUT4 containing vesicles (94) Dephosphorylated by PHLPP1 (95)	Brown fat, skeletal muscle and liver (91, 92) Phosphorylates TSC2 (95), FoxOs (95) and Ankrd2/ARPP (97) Glucose metabolism(102, 103) Cell migration/ metastasis(100, 101) EMT(100, 101)
		Akt3 Q9Y243	PH, KD, HM	Dephosphorylated by PHLPP1 and PLHPP2 (95)	Brain and testes (93, 219) Phosphorylates FoxOs (95) Akt3 inhibition induces migration (104) Neuronal growth (105, 106)
	S6K	S6K1 P23443	NTD, KD, CTD	Activated by neurabin, phospholipase D1, Rac1, Cdc42, SIRT1/SIRT2 (123) Inhibited by leucine starvation (125)	Ubiquitous (76) Activates S6 (133) Targets BAD, Mdm2 GSK-3, eIF4B, eEF2K, Er α and CREM (123) Induces cell survival, translation initiation and elongation and transcription (123)
		S6K2 Q9UBS0	NLS, NTD, KD, CTD	Activated by IL-3, MAPK pathway and PKC (123) Not inhibited by leucine starvation (125)	Ubiquitous (76) Activates S6 (133) Regulates YY2 and ROR γ (123) Induces cell survival, translation and transcription (123)
	4E-BP	4E-BP1 Q13541	RAIP, eIF4E-BD, TOS motif	Inhibited by mTORC1 (151)	Adipose tissue, pancreas, skeletal muscle (149) Redundant function with 4E-BP2 (220, 221) Inhibits cap-dependent translation (43)
		4E-BP2 Q13542	RAIP, eIF4E-BD, TOS motif	Inhibited by mTORC1 (151)	Ubiquitous (149) Redundant function with 4E-BP1 (220, 221) Inhibits cap-dependent translation (43)
		4E-BP3 O60516	eIF4E-BD, TOS motif	Not inhibited by mTORC1 (151) Transcription regulated by mTORC1 (152)	Skeletal muscle, heart, kidney, pancreas (150) Inhibits cap-dependent translation (153) Regulation of eIF4E at the nucleus to regulate mRNA nuclear export (153)
	ULK	ULK1 O75385	KD, Pro/Ser region, CTD	Inhibited by mTOR (48) Strong affinity to Atg13 and FIP200 (48)	Ubiquitous (76) Induces autophagosome formation (154)
		ULK2 Q8IYT8	KD, Pro/Ser region, CTD	Inhibited by mTOR (48) Weak affinity to Atg13 and FIP200 (48)	Ubiquitous (76) Induces autophagosome formation (155)
	mTORC2	Protor-1 P85299	No functional domain known	Binds mTORC2 via rictor (16)	Ubiquitous (76) Promotes SGK phosphorylation (118)
		Protor-2 Q6MZQ0	No functional domain known	Binds mTORC2 via rictor (16)	Spleen and intestine (76) Regulates mRNA stability during stress (119)

Protein domains: pleckstrin homologue (PH), phosphotyrosine-binding (PTB), phosphatidylinositol 3-kinase binding domain (PI3K-BD), binding domain (BD), growth factor receptor-bound protein 2 binding domain (Grb2-BD), src homology 2 domain-containing phosphatase binding domain (SHP2-BD), kinase regulatory loop binding (KRLB), kinase domain (KD), hydrophobic motif (HM), nuclear localization sequence (NLS), N-terminal domain (NTD), C-terminal domain (CTD), eukaryotic translation initiation factor 4E binding domain (eIF4E-BD), TOR signalling sequence (TOS), Proline/Serine (Pro/Ser).

Table 2.

Variants across the mTOR network arising from alternative splicing or alternative translation initiation

UniProt predicted variants			Name UniProt ID	Regulatory domains	Functional differences (citations)
insulin	InR	Mouse: 1 Human: 2	InR-A P06213-2	α domain + β domain	High affinity for insulin, IGF-1 and IGF-2 (73, 74)
			InR-B P06213-1	α domain + β domain (+12 aa)	High affinity for insulin, low for IGF-1 (73, 74)
	PTEN	Mouse: 1 Human: 3	PTEN P60484-1	Phosphatase domain, C2 domain, tail	Counterpart of PI3K. Converts PIP3 into PIP2 (21)
			PTEN α^T P60484-2	NTD, phosphatase domain, C2 domain, tail	Induction of cytochrome c oxidase activity (88)
			PTEN-Long T P60484-2	NTD, phosphatase domain, C2 domain, tail	Counterpart of PI3K. Secreted in the microenvironment (87)
Akt	Akt1	Mouse: 1 Human: 2	*	*	*
	Akt2	Mouse: 1 Human: 2	*	*	*
	Akt3	Mouse: 2 Human: 2	Akt3 Q9Y243-1	PH, KD, HM	Activated by mTORC2 and PDK1 (90)
			Akt3-1 Q9Y243-2	Lacks mTORC2 target site	Activated by PDK1. Less responsive to growth factors (90)
	TSC1	Mouse: 4 Human: 2	*	*	*
	TSC2	Mouse: 7 Human: 8	TSC2 P49815-1	Full protein	Akt target sites pS939, pS981 and T1462 (108) AMPK target sites pS1387 and pT1271 (38)
			P49815-2, P49815-3, P49815-5, P49815-6 P49815-7	Lacks exon 25	Akt target sites pS939 and pT1462 (108)
			P49815-4, P49815-5, P49815-6 P49815-7	Lacks exon 31	AMPK target site pS1387 (38)
	TBC1D7	Mouse: 2 Human: 4	*	*	*
	mTORC1	mTOR	Mouse: 2 Human: 1	mTOR α P42345-1	HEAT, FAT, KD, FATC
mTOR β^L				KD, FATC	Strong binding to c-Myc that allows control of cell cycle (114)
DEPTOR		Mouse: 3 Human: 2	*	*	*
raptor		Mouse: 5 Human: 3	raptor Q8N122-1	RNC, HEAT, WD40 repeats	Forms part of mTORC1. Mediates binding with mTORC1 substrates
			raptor-v2 Q8N122-3	RNC, WD40 repeats	Forms part of mTORC1. Cannot bind substrates (115)
PRAS40		Mouse: 1 Human: 3	*	*	*
S6K1		Mouse: 2 Human: 5	p70-S6K1 T P23443-2	NTD, KD, CTD	Targeted by mTORC1 (147)
			p85-S6K1 T P23443-1	1xNLS, NTD, KD, CTD	Contradiction if targeted by mTORC1 (147, 148)
			p35-S6K1 L	1xNLS, NTD, partial KD	Not known if targeted by mTORC1 (147)
S6K2		Mouse: 1 Human: 2	p54-S6K2 T Q9UBS0-1	1xNLS, NTD, KD, CTD	Resides in soluble fraction of cells (122)
	p56-S6K2 T,L		2xNLS, NTD, KD, CTD	Resides in particulate fraction of cells (122)	
mTORC2	riCTOR	Mouse: 2 Human: 3	*	*	*
	mSin1	Mouse: 3 Human: 6	mSin1.1 Q9BPZ7-1	RBD, PH	Forms part of mTORC2 (14)
			mSin1.2 Q9BPZ7-2	Partial RBD, PH	Forms part of mTORC2 (14)
			mSin1.3 Q9BPZ7-3	Partial RBD, PH	Does not form part of mTORC2 (14)
			mSin1.4	Partial RBD, PH	Does not form part of mTORC2 (14)

		Q9BPZ7-4		
		mSin1.5 ^L	Partial RBD	Forms part of mTORC2 (14)
Protector-1	Mouse: 2 Human: 5	Protector-1α P85299-1	Full protein.	Forms part of mTORC2 (16)
		Protector-1β P85299-3	Shorter variant	Forms part of mTORC2 (16)
		Protector-1γ P85299-4	Shorter variant	Does not form part of mTORC2 (16)
Protector-2	Mouse: 1 Human: 4	*	*	*

Protein domains: N-terminal domain (NTD), pleckstrin homologue (PH), kinase domain (KD), hydrophobic motif (HM), huntingtin-elongation factor 3-regulatory subunit A of PP2A-TOR1 repeats (HEAT repeats), FRAP-ATF-TTRAP (FAT), FRAP-ATM-TTRAP domain (FATC), raptor N-terminal conserved (RNC), C-terminal domain (CTD), nuclear localization signal (NLS), ras-binding domain (RBD). The variants arise from alternative splicing unless marked otherwise. Variants marked ^T originate from alternative translation initiation. *Asterisks designate splice variants that have been predicted, but have not yet been experimentally confirmed. ^L Refer to primary literature as this variant is not listed in UniProt.

Table 3.

Phenotypes of knockout mouse models

	Delayed ageing phenotype		Accelerated ageing phenotype	
	Lifespan	Metabolic profile	Lifespan	Metabolic profile
Insulin	<i>Irs1</i> ^(-/-) (178) <i>Irs2</i> ^{(+/-)?} (180, 181) brain <i>Irs2</i> ^{(+/-)?} (180, 181)	adipose <i>Insr</i> ^(flox/flox) ^L (183)	<i>Insr</i> ^(-/-) ^I (222)	muscle <i>Insr</i> ^(flox/flox) ^{G,O} (191) pancreas <i>Insr</i> ^(flox/flox) ^G (192) brain <i>Insr</i> ^(flox/flox) ^{G,I} (193) liver <i>Insr</i> ^(flox/flox) ^I (196) <i>Irs2</i> ^(-/-) ^{I,O} (202) <i>Irs4</i> ^(-/-) ^G (190)
Akt	<i>Akt1</i> ^(+/-) (203)			liver <i>Pdpr1</i> ^(-/-) ^G (194) muscle <i>Pdpr1</i> ^(-/-) ^M (197) <i>Akt2</i> ^(-/-) ^G (102) <i>Tsc1</i> ^(+/-) ^T (185) <i>Tsc2</i> ^(+/-) ^T (186)
mTORC1	<i>Mtor</i> ^(+/-) <i>Mlst8</i> ^(+/-) (177) <i>Rps6kb1</i> ^(-/-) (179)	adipose <i>Rptor</i> ^(-/-) ^L (184) <i>Eif4ebp1</i> ^(-/-) ^L (204)		muscle <i>Mtor</i> ^(flox/flox) ^M (198) muscle <i>Rptor</i> ^(flox/flox) ^M (199)
mTORC2	<i>Mtor</i> ^(+/-) <i>Mlst8</i> ^(+/-) (177)		<i>Rictor</i> ^(+/-) (182) <i>Rictor</i> ^(-/-) (182)	muscle <i>Mtor</i> ^(flox/flox) ^M (198) liver <i>Rictor</i> ^(flox/flox) ^{G,I} (223)

^L Less adipose tissue, ^G Impaired glucose tolerance, ^I Insulin resistance, ^O Obesity, ^M Myopathy, ^T Appearance of tumours, [?] Phenotype not reproduced