

University of Groningen

Low but contrasting neutral genetic differentiation shaped by winter temperature in European great tits

Lemoine, Melissa; Lucek, Kay; Perrier, Charles; Saladin, Verena; Adriaensen, Frank; Barba, Emilio; Belda, Eduardo J.; Charmantier, Anne; Cichon, Mariusz; Eeva, Tapio

Published in:
Biological Journal of the Linnean Society

DOI:
[10.1111/bij.12745](https://doi.org/10.1111/bij.12745)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lemoine, M., Lucek, K., Perrier, C., Saladin, V., Adriaensen, F., Barba, E., Belda, E. J., Charmantier, A., Cichon, M., Eeva, T., Gregoire, A., Hinde, C. A., Johnsen, A., Komdeur, J., Mand, R., Matthysen, E., Norte, A. C., Pitala, N., Sheldon, B. C., ... Richner, H. (2016). Low but contrasting neutral genetic differentiation shaped by winter temperature in European great tits. *Biological Journal of the Linnean Society*, 118(3), 668-685. <https://doi.org/10.1111/bij.12745>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Low but contrasting neutral genetic differentiation shaped by winter temperature in European great tits

MÉLISSA LEMOINE^{1†}, KAY LUCEK^{2,3}, CHARLES PERRIER⁴, VERENA SALADIN⁵, FRANK ADRIAENSEN⁶, EMILIO BARBA⁷, EDUARDO J. BELDA⁸, ANNE CHARMANTIER⁴, MARIUSZ CICHON⁹, TAPIO EEVA¹⁰, ARNAUD GRÉGOIRE⁴, CAMILLA A. HINDE¹¹, ARILD JOHNSEN¹², JAN KOMDEUR¹³, RAIVO MÄND¹⁴, ERIK MATTHYSEN⁶, ANA CLÁUDIA NORTE¹⁵, NATALIA PITALA¹⁶, BEN C. SHELDON¹⁷, TORE SLAGSVOLD¹⁸, JOOST M. TINBERGEN¹⁹, JÁNOS TÖRÖK²⁰, RICHARD UBELS¹⁹, KEES VAN OERS²¹, MARCEL E. VISSER²¹, BLANDINE DOLIGEZ^{22,23*} and HEINZ RICHNER^{5*}

¹*Institute for Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190, CH-8057, Zurich, Switzerland*

²*Aquatic Ecology and Evolution, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, CH-3012, Bern, Switzerland*

³*Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK*

⁴*Centre d'Ecologie Fonctionnelle et Evolutive, Unité Mixte de Recherche 5175, 1919 Route de Mende, FR-34293, Montpellier Cedex 5, France*

⁵*Division of Evolutionary Ecology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, CH-3012, Bern, Switzerland*

⁶*Evolutionary Ecology Group, Department of Biology, University of Antwerp, BE-2020, Antwerp, Belgium*

⁷*Cavanilles' Institute of Biodiversity and Evolutionary Biology, University of Valencia, C/Catedrático José Beltrán 2, 46980, Paterna, Spain*

⁸*Instituto de Investigación para la Gestión Integrada de Zonas Costeras-IGIC, U.P.V., C/Paraninfo n° 1, ES-46730 Gandia, Valencia, Spain*

⁹*Institute of Environmental Sciences, Jagiellonian University, Gronostajowa 7, 30-387, Kraków, Poland*

¹⁰*Department of Biology, University of Turku, FI-20014, Turku, Finland*

¹¹*Behavioural Ecology Group, Department of Animal Sciences, Wageningen University, 6700, AH Wageningen, The Netherlands*

¹²*Natural History Museum, University of Oslo, PO Box 1172 Blindern, NO-0318, Oslo, Norway*

¹³*Behavioural Ecology and Self-organization, Centre for Ecological and Evolutionary studies, University of Groningen, PO Box 11103, NL-9747, AG Groningen, The Netherlands*

¹⁴*Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu, Vanemuise 46, Tartu, EE-51014, Estonia*

¹⁵*Department of Life Sciences, Largo Marquês de Pombal, Faculty of Sciences and Technology, Marine and Environmental Sciences Centre (MARE), University of Coimbra, 3004-517, Coimbra, Portugal*

¹⁶*Department of Biological and Environmental Science, University of Jyväskylä, PO Box 35, FI-40014, Finland*

¹⁷*Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford, South Parks Road, Oxford, OX1 3PS, England*

¹⁸*Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of*

†Corresponding author. E-mail: melissa.lemoine@gmail.com

*These authors contributed equally to this work.

Co-authors 5-25 are in alphabetical order.

Oslo, PO Box 1066 Blindern, NO-0316, Oslo, Norway

¹⁹Animal Ecology Group, Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 11103, NL-9747, AG Groningen, The Netherlands

²⁰Behavioural Ecology Group, Department of Systematic Zoology and Ecology, Eötvös Loránd University, Pázmány Péter sétány 1/c, 1117, Budapest, Hungary

²¹Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), PO Box 50, 6700, AB Wageningen, The Netherlands

²²Department of Biometry and Evolutionary Biology, CNRS, University of Lyon, UMR 5558, F-69622, Villeurbanne Cedex, France

²³Animal Ecology/Department of Ecology and Genetics, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18d, SE-752 36, Uppsala, Sweden

Received 19 August 2015; revised 30 November 2015; accepted for publication 1 December 2015

Gene flow is usually thought to reduce genetic divergence and impede local adaptation by homogenising gene pools between populations. However, evidence for local adaptation and phenotypic differentiation in highly mobile species, experiencing high levels of gene flow, is emerging. Assessing population genetic structure at different spatial scales is thus a crucial step towards understanding mechanisms underlying intraspecific differentiation and diversification. Here, we studied the population genetic structure of a highly mobile species – the great tit *Parus major* – at different spatial scales. We analysed 884 individuals from 30 sites across Europe including 10 close-by sites (< 50 km), using 22 microsatellite markers. Overall we found a low but significant genetic differentiation among sites ($F_{ST} = 0.008$). Genetic differentiation was higher, and genetic diversity lower, in south-western Europe. These regional differences were statistically best explained by winter temperature. Overall, our results suggest that great tits form a single patchy metapopulation across Europe, in which genetic differentiation is independent of geographical distance and gene flow may be regulated by environmental factors via movements related to winter severity. This might have important implications for the evolutionary trajectories of sub-populations, especially in the context of climate change, and calls for future investigations of local differences in costs and benefits of philopatry at large scales. © 2016 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2016, **118**, 668–685.

KEYWORDS: *F*-statistics – isolation-by-distance – latitude – microsatellites – *Parus major* – Population genetic structure – winter severity.

INTRODUCTION

Gene flow is generally thought to impede local adaptation by introducing locally maladapted genotypes into populations exchanging individuals. Consequently, microevolutionary processes at small scales are predicted to be rare in highly mobile organisms with high gene flow over large spatial scales, due to spatial genetic homogenisation. However, evidence for genetic differentiation and local adaptation at small scales despite high levels of gene flow at large scales has recently started to accumulate in different taxa (e.g. mammals: Musiani *et al.*, 2007; marine invertebrates: Sanford & Kelly, 2011; birds: Charmantier *et al.*, 2016; fish: Junge *et al.*, 2011; trees: Savolainen, Pyhäjärvi & Knürr, 2007). This evidence suggests that dispersal is not a diffusion-like movement process and that ultimately gene flow may vary in space.

Spatial variation in gene flow is probably common, especially in relation to environmental factors in

highly mobile species. High mobility and long-distance dispersal facilitate spatial spread and the colonization of new habitats (Nathan *et al.*, 2003). As a consequence, highly mobile species are likely to experience a large set of environmental conditions that may shape locally adaptive processes. In addition, high mobility combined with the ability to cross physical barriers such as seas or mountains may minimize the influence of geographical factors. Increased mobility may also reduce the impact of historical factors on gene flow by homogenising gene pools, increasing local population size and counteracting genetic drift (Slatkin, 1987). In this case, environmental factors may become the main force shaping gene flow (e.g. Pilot *et al.*, 2006). Assessing gene flow between populations at small and large spatial scales in highly mobile species and the links between gene flow and environmental factors is crucial to understand the ecological mechanisms leading to intraspecific differentiation and diversification. When dispersal movements and immigration rate

do not provide reliable estimates of gene flow, such as in highly mobile species, a population genetic approach may help investigating patterns of gene flow at different spatial scales (Nathan *et al.*, 2003).

The great tit *Parus major*, a widespread passerine bird across Eurasia (Snow & Perrins, 1998), is a particularly interesting biological model to address such questions. This species is considered to be an 'evolutionary winner', given its ability to colonize and rapidly adapt to new habitats. Its rapid spread across Europe since the last glaciation period (Kvist *et al.*, 2003; Pavlova *et al.*, 2006) suggests high dispersal ability and gene flow among sub-populations (Caswell, Lensink & Neubert, 2003; Pilot *et al.*, 2006 but see Peterson & Denno, 1998). Conversely, long-term monitoring studies provide evidence for small-scale local adaptation (Garant *et al.*, 2005; Postma & van Noordwijk, 2005) with a considerable fraction of individuals dispersing over short distances (e.g. Verhulst, Perrins & Riddington, 1997). Thus, although great tits are considered highly mobile and forming a homogeneous population across Europe, microevolutionary processes linked with limited gene flow occur at the small scale, and with it the detection of subtle fine-scale genetic structures (Björklund, Ruiz & Senar, 2010; Van Bers *et al.*, 2012; Garroway *et al.*, 2013). These conflicting observations call for investigating genetic differentiation using microsatellite markers at different spatial scales in this species. Indeed microsatellite markers generating multilocus diploid genotypes provide an ideal resolution to study recent or ongoing microevolutionary processes occurring both at small and large scales (e.g. Wang, 2010).

Moreover, the environmental heterogeneity over the species' range combined with its colonisation history provides excellent conditions to study the influence of environmental factors on population genetic structure in this species. Indeed, phylogeographic studies based on mitochondrial DNA (mtDNA) suggest that other tit species colonized Europe from different glacial refugia, each harbouring distinct mitochondrial lineages and forming secondary contact zones within Europe (Kvist *et al.*, 2004; Päckert, Martens & Sun, 2010; Pentzold *et al.*, 2013). In contrast, all western-European great tits share a common haplotype, suggesting that they originated from a single glacial refugium located in southern Europe (Kvist *et al.*, 2007, 1999; Pavlova *et al.*, 2006; Supporting Information, Figure S1 and Table S1). Genetic differentiation in great tits estimated with microsatellites that evolve faster than mtDNA and are more powerful to detect recent and local microevolutionary processes among populations, therefore, are less likely to result from past genetic discontinuities across different glacial refugia as is

the case for many other species (e.g. Kvist *et al.*, 1999; Hewitt, 2000).

Using 22 microsatellite markers, we investigated population genetic diversity and structure, as well as the scale of genetic differentiation, in great tits by sampling 30 sites across Europe including 10 close-by (i.e. up to 50 km) sites. We expected the genetic differentiation to be correlated with the geographical distance either at small or large scales: the studied geographical scale should allow us to determine at which scale isolation-by-distance would occur in great tits. In addition, a signal of historical range expansion from the South to the North should result in decreased genetic diversity with increasing latitude. In a second step, we explored the influence of environmental factors on the observed genetic diversity and differentiation patterns, focusing on factors that can be expected to affect individual movement. In particular, temperature may strongly shape genetic differentiation among populations by acting on both dispersal movements (e.g. Parn *et al.*, 2012) and establishment success (i.e. survival and reproductive success after settlement) of long-distance immigrants (e.g. Van Doorslaer *et al.*, 2009). Three different patterns may thus be predicted in relation to temperature. First, because temperature can be positively correlated with survival and population density (Garant *et al.*, 2004; Ahola *et al.*, 2009; Parn *et al.*, 2012) that increase dispersal propensities (Forsman & Monkkonen, 2003; Matthysen, 2005), genetic diversity could increase and genetic differentiation decrease with increasing temperature. Second, a negative relationship between temperature and dispersal propensities may be expected in the case of partial migration (e.g. Nilsson *et al.*, 2006). In this case, temperature should relate to environmental conditions during winter, triggering partial migration and favouring dispersal in general or the establishment of migrants in non-natal breeding areas. Genetic diversity should consequently decrease while genetic differentiation should increase with temperature (e.g. Miller *et al.*, 2012). Third, if the establishment success of immigrants is linked to adaptation to temperature, we predicted that genetic differentiation should increase with the difference of temperature between sites.

MATERIAL AND METHODS

SPECIES DESCRIPTION, SAMPLING AND GENOTYPING

The great tit is a hole-nesting passerine bird that readily breeds in nest boxes, providing easy access to breeding pairs. In this study, all individuals from all but one site (FL.TU, see Supporting Information, Table S2) were breeding adults caught in nest boxes

during the nestling period. Thirty woodland sites across Europe were sampled between 2005 and 2010 (Fig. 1, Supporting Information, Table S2), 10 of which were within a range of 50 km on the island of Gotland (57°10'N, 18°20'E). Overall, our studied populations fell along a south-west–north-east gradient (Fig. 1). Either blood or feather samples were obtained. Most sites were sampled once, except when the sample size was too low for statistical analysis (in 10 sites). In this case, samples of two consecutive years were pooled. The number of sampled individuals per site ranged from 18 to 47 with an average of 29.

DNA was extracted with magnetic beads (Magne-Sil Blue, Promega AG, Dübendorf, Switzerland) and genotyped at 22 microsatellite loci (Supporting Information, Table S3, Saladin & Richner, 2012). These 22 microsatellite markers were developed using individuals from CH.BE, a site in the geographical centre of our sampling scheme. For details on the PCR protocols and allele scoring procedure, see Saladin &

Richner (2012). Twelve individuals with missing alleles or atypical profiles at different loci were excluded from all analyses. None of the individuals shared the same multilocus genotype indicating that none of the individuals was sampled twice. Overall, 884 individuals were analysed. Allelic dropout, scoring errors and null alleles were checked for each locus per site with MICRO-CHECKER (Van Oosterhout *et al.*, 2004). Among all loci, no evidence for allelic dropout was detected and only one locus in one sampling site showed scoring errors. Moreover, null alleles were randomly distributed, and present at only 19 (i.e. 2.9%) locus \times site combinations. Genotypic linkage disequilibrium and departure from Hardy–Weinberg equilibrium (HWE) were tested with probability tests per locus per site. In addition, departure from HWE for the overall population, i.e. across loci and sites, was tested using a multisample score test. All tests were performed using GENEPOP on the web (Rousset, 2008). *P*-values for multiple tests were corrected with a sequential Bonferroni procedure (Rice, 1989).




Figure 1. Location of the 30 sampling sites across Europe. The inset shows the 10 sampling sites on the island of Gotland, Sweden. The dashed line shows the 47° latitude. IBD analysis treats all populations into a single quantity assuming that all local populations have similar characteristics. In contrast, the DPR analysis extracts the elements of individual local population from the information on an entire metapopulation and identifies five groups differing in relative strengths of gene flow and genetic drift patterns (i.e. different patterns of genetic differentiation and IBD summarised by different colours, see Table 1 for details).

GENETIC DIVERSITY AND DIFFERENTIATION AMONG SITES

To assess genetic diversity at each sampling site, both the observed and unbiased expected heterozygosity (H_O and H_E) were calculated using GENALEX v6 (Peakall & Smouse, 2006). In addition, the mean allelic richness per site (A_R) based on 18 individuals, corresponding to the smallest number of individuals sampled in a given site, was estimated with FSTAT v2.9.3 (Goudet, 1995). Genetic differentiation among sites was quantified using pairwise and global F_{ST} calculated in FSTAT with 10 000 permutations to assess significance. Because F_{ST} estimates may be strongly affected by the polymorphism of the markers used (Meirmans & Hedrick, 2011), standardized estimators G'_{ST} and D were calculated with GENODIVE 2.0b27 (Meirmans & Van Tienderen, 2004).

To test for a spatial pattern of genetic differentiation among sites, two methods were used: (i) a principal coordinate analysis (PCoA), based on codominant genotypic distance among sites with a standardized covariance matrix, using GENALEX 6.5; and (ii) a neighbour-joining (NJ) phenogram based on Nei's genetic distance between sites, using PHYLIP v3.68 (Felsenstein, 1989). The presence of genetic clusters was also tested using two methods. First, an individual-based Bayesian cluster analysis was implemented in STRUCTURE v2.2 (Pritchard, Stephens & Donnelly, 2000). Ten runs of an admixture model with correlated allele frequencies among sites and LOCPRIOR were performed for each value of putative population number (K) between 1 and 40 with a burn-in of 50 000 iterations followed by 100 000 iterations in the Markov chain. The most likely number of genetically different populations was determined from the posterior probability of the data for a given K and the ΔK (Evanno, Regnaut & Goudet, 2005). To test for a potential bias due to the inclusion of 10 close-by sites from Gotland, the PCoA and STRUCTURE analyses were run once using individuals from all 30 sites and once using individuals from 21 sites including only a single site from Gotland (SE.OG). As the results did not qualitatively differ (Supporting Information, Figs S2–S6 and Table S4), we presented only the results based on 30 sites. In addition, assignment probabilities of individuals to their original site (P_A) were calculated using a discriminant analysis of principal components (DAPC – Jombart, Devillard & Balloux, 2010) in R 3.0.1 (R Core Team, 2013). Second, the clustering of sites into groups was investigated by a K-means clustering using an analysis of molecular variance (AMOVA) with 40 independent Markov chains with 50 000 iterations each assuming 2–15 clusters with GENODIVE. The most likely number of clusters was determined from the smallest

bayesian information criterium (BIC). Furthermore, genetic differentiation was quantified between groups and among sampling sites within groups using an AMOVA with 10 000 permutations to assess significance using GENODIVE. Additionally, within-group global F_{ST} values were calculated and compared with 10 000 permutations using FSTAT.

To test for the presence of isolation-by-distance patterns, a decomposed pairwise regression analysis (DPR) was conducted in R to account for potential between-site differences in the gene flow-drift equilibrium (Koizumi, Yamamoto & Maekawa, 2006). Briefly, DPR first detects outlier sites based on the distribution of residuals from the overall regression between genetic and geographical distances. In a second step, genetic distances are regressed against geographical distances for each site against all other non-outlier sites to obtain a regression intercept and slope per site. The intercept and slope of the decomposed regressions measure genetic differentiation to other populations and isolation-by-distance (IBD) respectively for each site (see Koizumi *et al.*, 2006 for details).

TESTING FOR THE INFLUENCE OF ENVIRONMENTAL FACTORS ON DIFFERENCES AMONG SITES

To investigate potential mechanisms underlying differences in genetic diversity, the relationships between indices of genetic diversity per site and the following environmental factors, which may be expected to influence individuals' movements, were tested: (i) geographical location (latitude and longitude); (ii) vegetation type (deciduous or coniferous trees; excluding SP.MU and ES.KI, where birds were sampled in orange tree plantations or mixed areas); (iii) temperature and (iv) minimal distance to the sea. Latitude, longitude and minimal distance to the sea were obtained using GOOGLE EARTH v5.2.1. Using the position along a south-west–north-east axis as a geographical location did not affect the results, and thus only results including latitude and longitude are reported. Temperatures were obtained from the European photovoltaic geographical information system (Huld *et al.*, 2006). The measures based on temperature were: (i) average daily temperature per month; (ii) temperature variance per year; (iii) difference between the most extreme annual temperatures; and (iv) average temperature during autumn–winter (September–February) and spring–summer (March–August). In addition to the indices of genetic diversity per site, we calculated an estimate of effective population size (N_e) with the linkage disequilibrium method using a threshold of 0.05 for the exclusion of rare alleles in N_e ESTIMATOR v2 (Do *et al.*,

2014) and the kinship coefficient of Loiselle *et al.* (1995) averaged per site with GENODIVE. The relationships between genetic diversities (A_R and H_E), assignment probabilities, kinship coefficients, effective population sizes and environmental factors were tested using linear models since all indices were normally distributed (residuals were checked for normality and homoscedasticity). Because the environmental factors were correlated with each other (correlation coefficients ranging from 0.31 to 0.86, all $P < 0.001$, results not shown), Akaike's information criterion (AIC) values of models including each factor separately were compared in order to identify the environmental factor(s) that best explained the data using the package AICmodavg (Mazerolle, 2015) in R. The best models included the model with the smallest AIC and all models with a difference in AIC (ΔAIC) to this model of < 2 (Burnham, Anderson & Huyvaert, 2011). Once the best models were identified, the significance of the effects retained was assessed with an F test.

In a second step, the influence of the following environmental factors on genetic differentiation among sampling sites was tested: (i) geographical distance between sites; (ii) mean geographical location of sites; (iii) absolute difference in average daily temperature between sites; and (iv) mean of the average daily temperature of sites. Because previous analyses showed that genetic diversity was best explained by temperatures in autumn–winter (see Results section), only the difference in average autumn–winter temperatures between sites (hereafter called autumn–winter temperature difference) and the mean of the average daily temperature in autumn–winter of sites (hereafter called mean autumn–winter temperature) were tested in the analyses of genetic differentiation. Similarly, only the latitude was retained here to characterize geographical location for analyses on genetic differentiation since site latitude and longitude were correlated in our study (i.e. sites were distributed along a south–west–north–east axis). The difference between values for the two sites in pairwise comparisons provides a measure of the environmental contrast between sites, whereas the mean value gives a measure of the position of the pair of sites in each pairwise comparison along the environmental gradient considered (geographical position or winter severity). The genetic differentiation between sites was calculated for each pair of sites and summarized in a pairwise matrix; the same approach was used for the differences and mean values of the environmental factors between sites. Correlations between levels of pairwise genetic differentiation based on either F_{ST} , G''_{ST} or D and pairwise differences in environmental factors were investigated with Mantel tests (or

partial Mantel tests when more than two matrices were compared) with 10 000 permutations using the package vegan (Oksanen *et al.*, 2011) in R. By homogenising the genetic composition of connected populations, gene flow should reduce both the mean level and the variability of genetic differentiation between populations (Hutchison, Templeton & R., 1999). Consequently, a factor affecting gene flow should be correlated with both the level of genetic differentiation and the absolute values of residuals of the linear regression between the factor and the level of genetic differentiation (hereafter called residual pairwise F_{ST} , G''_{ST} or D respectively) (Hutchison *et al.*, 1999). Therefore, the correlation between matrices of environmental factors and their residual pairwise genetic differentiation was also tested.

RESULTS

GENETIC DIVERSITY AND EQUILIBRIUM

No evidence for linkage disequilibrium at any locus in any site or departure from HWE was found after correction for multi-comparisons. Pooling all sites, a significant deviation from HWE was observed (score test: $P < 0.001$), suggesting the existence of subpopulations. The number of alleles per locus ranged from 4 to 41 with an average of 16 alleles across loci. Mean allelic richness per site ranged from 6.32 to 7.66 (Supporting Information, Table S2). Expected heterozygosity varied between 0.60 and 0.68 and the number of effective alleles between 3.94 and 4.92 (Supporting Information, Table S2). F_{IS} per site ranged from -0.049 to 0.047 (Supporting Information, Table S2), but no F_{IS} value differed significantly from zero after correcting for multiple tests, as expected under within-site HWE.

GENETIC DIFFERENTIATION AMONG SAMPLING SITES

Genetic differentiation among sampling sites across Europe was low, but significant (global $F_{ST} = 0.008$, $G''_{ST} = 0.024$, $D = 0.016$, all $P < 0.001$). Pairwise F_{ST} ranged from -0.004 to 0.040 (Supporting Information, Table S3). Out of 435 pairwise F_{ST} comparisons, 147 (i.e. 33.8%) were significantly different from zero after sequential Bonferroni correction. Interestingly, the majority of significant comparisons (134 out of 147, i.e. 91.1%) involved six (out of seven) sampling sites located in the south-western part of Europe, i.e. below $47^\circ N$ (CH.BE, FR.MO, SP.MU, SP.FR, SP.MA and PO.CO), indicating different levels of genetic differentiation between northern and southern sites (Fig. 2). FR.RO was the only site located in the southern region for which pairwise F_{ST} values were non-significant. Results of both the PCoA analysis




Figure 2. Heatmap of the pairwise F_{ST} values between all sites. Sites are ordinated by pairwise F_{ST} values. Black bars highlight the sites located below the 47° latitude (i.e. south-western sites).

and NJ phenogram based on Nei's genetic distance were congruent with the observed pairwise F_{ST} pattern for six out of the seven southern sites (Fig. 3). The PCoA accounted for 62% of the total genetic variation on the first three axes (26.5, 18 and 17.5% respectively). Independently of the axes considered, PO.CO, SP.MU, SP.MA, SP.FR, FR.MO were identified as being rather distinct from all other sites (i.e. outside the 50% and close to the 95% limit of the confidence interval; Fig. 3A, B). These south-western sites were also differentiated from each other, except SP.MA and SP.FR, which also showed lower pairwise F_{ST} values. Only CH.BE, which had relatively low F_{ST} values, was not identified as a differentiated site by the PCoA and the NJ phenogram analyses. Furthermore the central cluster was randomly distributed on each PCoA axis, in particular with no clumping of the 10 close-by sampling sites located on Gotland (Fig. 3A, B), which was confirmed on the NJ phenogram. In fact, populations on Gotland showed similar levels of differentiation among themselves as among the other sites from northern Europe (Fig. 3C). Depending on the method used, some of the northern sites appeared differentiated from the central cluster (e.g. SE.LO, Figs 2B, 3A; SE.SA, Fig. 3A–C; or NE.LA, Fig. 3A) suggesting that they could be distinct from the central cluster yet less differentiated than the south-western sites. Overall, the results indicate that: (i) genetic differentiation

among sampling sites was low (Figs 2, 3); (ii) many sites (including close-by ones) presented similar and low levels of genetic differentiation without spatial structure (e.g. a centred star-like pattern; Fig. 3C); and (iii) at least five southern sites were differentiated from the central cluster and differentiated from each other, except SP.MA and SP.FR (Figs 2, 3).

STRUCTURE identified three genetic clusters ($K = 3$) following the Evanno correction (Supporting Information, Figs S4, S5). Two of these clusters were mainly associated with the four Iberian sites, where the Portuguese site (PO.CO) was further distinct from all Spanish sites (SP.MU, SP.MA and SP.FR), however no individual was fully assigned to either cluster (Supporting Information, Fig. S6). All other sites were predominantly assigned to a third cluster except for CH.BE, which showed evidence for introgression from south-western Europe. Concordantly, the AMOVA based K-means clustering identified two groups: one comprising the four Iberian sites and CH.BE and a second including all other sites (all northern sites and the two sites in France). The AMOVA using south-western (i.e. below 47° latitude: PO.CO, SP.MU, SP.MA, SP.FR, FR.MO, FR.RO, CH.BE) and northern (above 47° latitude) sites as grouping variable suggested low but significant genetic differentiation between these groups ($F_{\text{group-total}} = 0.002$, $P < 0.001$) and among sites within groups ($F_{\text{sites-group}} = 0.008$, $P < 0.001$).




Figure 3. (A and B) Principal coordinate analysis (PCoA) contrasting axes 1 vs. 2 (A) and 1 vs. 3 (B), and (C) NJ phenogram based on Nei's genetic distance, with bootstrap values of specific clusters. (A and B) On the PCoA plots, the smallest and largest ellipses represent the 50 and 95% confidence intervals respectively; black dots represent the five sites identified as satellites, grey dots potential other satellites and white dots non-differentiated sites. (C) On the NJ phenogram, the three grey circles indicate identified clusters and the five sites identified as satellites are indicated in bold.

In addition, the differentiation was higher within southern sites than other sites (global $F_{ST} = 0.016$ and 0.005 , $G''_{ST} = 0.052$ and 0.014 , $D = 0.034$ and 0.009 , for southern sites and other sites, respectively; $P < 0.001$). Excluding CH.BE, FR.MO and FR.RO, did not change qualitatively the results of the hierarchical AMOVA and the level of differentiation, suggesting that the observed clustering was mainly driven by the four Iberian sites, which are more differentiated than the other south-western sites. Interestingly, the weak overall differentiation among the northern sites did not result from differentiation between specific sampling sites since 19 sites had to be excluded one after the other

(starting from the sites with the highest mean pairwise F_{ST} value and going downwards) for the overall differentiation to become non-significant (results not detailed). Moreover, differentiation among the close-by sites on Gotland (with distance ranging from 3 to 50 km) was not lower than among other northern sites (global $F_{ST} = 0.006$ and 0.004 respectively, $P = 0.646$; Fig. 5A).

Finally, the DPR analysis identified FR.MO (the only urban site) as an outlier, as the model excluding this site had a lower AIC (-94.78) and higher R^2 (0.17) values, although other models (either comprising all sites or with additional outliers) could not be excluded ($\Delta AIC < 1.28$). Overall, the DPR divided

sampling sites into five groups (Table 1 and see Fig. 1 for location): (1) two southern sites (FR.MO and FR.RO) showed a significant atypical negative IBD pattern and significant differentiation from other sites; (2) the four Iberian sites (SP.MU, SP.MA, SP.FR and PO.CO) and CH.BE showed no significant IBD but significant differentiation from other sites; (3) ten northern sites in Fennoscandia showed both significant differentiation from other sites and an IBD pattern; (4) nine northern sites from different locations showed no differentiation from other sites but significant IBD; and (5) four central sites (UK.WY, UK.CA, BE.CE, BE.BO) showed no differentiation and no IBD. Interestingly, all but two close-by sites on Gotland showed both significant differentiation from other sites and an IBD pattern.

EXPLORING THE INFLUENCE OF ENVIRONMENTAL FACTORS ON GENETIC DIFFERENCES AMONG SITES

Models including latitude, longitude, and the variance and difference in daily temperature were retained for none of the five indices (allelic richness A_R , expected heterozygosity H_E , assignment probability P_A , kinship

coefficient and effective population size N_e , $\Delta AIC > 2$ in all cases; Supporting Information, Table S5). Conversely, models with average daily temperature for months September to January, and consequently average autumn–winter temperature, were among the models best explaining the data for P_A , A_R , H_E ($\Delta AIC < 2$ in all cases; Supporting Information, Table S5). For kinship coefficient, models with average daily temperature for months August, September and December were among the models best explaining the data ($\Delta AIC < 2$) but not the model with average autumn–winter temperature despite a relative low AIC ($\Delta AIC < 2.5$). The model including vegetation type was the only best model in explaining the data for the effective population size. Allelic richness decreased ($F_{1,28} = 6.90$, $P = 0.014$, $R^2 = 0.20$) while assignment probabilities and kinship coefficients increased ($F_{1,28} = 10.57$, $P = 0.003$, $R^2 = 0.27$; $F_{1,28} = 17.04$, $P < 0.001$, $R^2 = 0.36$ respectively) with increasing average autumn–winter temperature (Fig. 4). Expected heterozygosity and effective population size were not correlated with average autumn–winter temperature ($F_{1,28} = 0.81$, $P = 0.38$; $F_{1,24} = 0.56$, $P = 0.46$ respectively, Fig. 4). Effective population size was




Figure 4. Relationships between average autumn–winter temperature and (A) latitude, and (B–F) different population indices: assignment probability (B), allelic richness (C), mean pairwise kinship (D), unbiased expected heterozygosity (E), effective population size (F).

similar for coniferous and deciduous forests ($F_{1,22} = 0.03$, $P = 0.87$). Models with other factors were retained for part of the indices only: temperatures in summer months (July–August) for A_R and H_E , temperature in February for H_E , average spring-summer temperature for H_E , vegetation type for A_R and distance to the sea for H_E (Supporting Information, Table S5). However, allelic richness was similar in coniferous and deciduous forests ($F_{1,26} = 0.08$, $P =$

0.77), and expected heterozygosity was not correlated with spring-summer temperature or distance to the sea ($F_{1,28} < 2.5$, $P > 0.12$). Based on these results, only the average autumn–winter temperature was retained among temperature measures for the analyses of genetic differentiation.

All pairwise genetic differentiation indices increased with geographical distance between sites, autumn–winter temperature difference between sites

Table 1. Decomposed pairwise regression (DPR) of the genetic differentiation with geographic distance for each sampling site

Site	Intercept ± SE (10 ⁻²)		P	Slope ± SE (10 ⁻⁶)		P	R ²	Genetic differentiation pattern
FR.MO	2.57	0.26	0.000	-3.87	1.71	0.032	0.159	Negative IBD, differentiated sites
FR.RO	0.95	0.27	0.002	-3.75	1.77	0.044	0.147	
SP.MU	1.79	0.21	0.000	0.07	1.06	0.950	0.000	No IBD, differentiated sites
PO.CO	1.70	0.22	0.000	0.34	0.97	0.729	0.005	
SP.MA	1.28	0.23	0.000	-0.24	1.09	0.828	0.002	
SP.FR	1.21	0.21	0.000	-0.58	0.99	0.561	0.013	
CH.BE	0.61	0.22	0.010	1.73	1.80	0.344	0.035	
SE.SA	0.92	0.14	0.000	2.65	1.13	0.027	0.175	IBD, differentiated sites
SE.LO	0.87	0.12	0.000	4.35	1.01	0.000	0.418	
SE.BO	0.52	0.12	0.000	2.86	1.02	0.010	0.232	
FI.TU	0.50	0.17	0.006	2.36	1.11	0.043	0.148	
NO.DA	0.44	0.13	0.002	2.59	1.02	0.017	0.199	
SE.OG	0.42	0.11	0.001	3.50	0.92	0.001	0.355	
SE.JA	0.35	0.11	0.005	5.31	0.93	0.000	0.558	
SE.GE	0.33	0.12	0.009	2.51	0.97	0.016	0.203	
SE.SI	0.30	0.13	0.035	3.34	1.08	0.005	0.269	
SE.BI	0.26	0.11	0.021	2.61	0.86	0.005	0.264	
NE.LA	0.43	0.24	0.081	6.43	2.27	0.009	0.235	IBD, undifferentiated sites
FI.KO	0.20	0.17	0.251	2.38	0.94	0.018	0.197	
SE.ET	0.10	0.12	0.407	3.23	0.99	0.003	0.292	
NE.HO	0.07	0.21	0.754	4.33	2.02	0.042	0.150	
SE.DT	0.01	0.13	0.944	4.13	1.06	0.001	0.369	
NE.WE	0.00	0.17	0.994	3.94	1.55	0.018	0.199	
ES.KI	0.00	0.14	0.997	2.14	0.97	0.036	0.159	
PL.PU	-0.16	0.25	0.520	4.30	1.98	0.039	0.154	
HU.PI	-0.21	0.29	0.490	5.21	2.24	0.028	0.172	
UK.WY	0.45	0.25	0.089	0.80	1.96	0.687	0.006	
BE.BO	0.38	0.22	0.098	2.83	2.03	0.175	0.070	
BE.CE	0.27	0.17	0.127	2.76	1.57	0.090	0.106	
UK.CA	0.22	0.23	0.331	2.84	1.83	0.132	0.085	
All	0.45	0.06	0.000	3.21	0.40	0.000	0.130	IBD, differentiated sites

Intercepts indicate the level of differentiation of sites, and slopes indicate isolation-by-distance (IBD). FR.MO was identified as an outlier site and was therefore excluded to calculate the pairwise regressions of other sites. Significant values are indicated in bold.

and mean autumn–winter temperature of the two sites in pairwise comparisons, and decreased with mean latitude of the two sites (Table 2; Fig. 5). Each environmental factor explained 36–57% of the variation in pairwise genetic differentiation. Furthermore, both mean autumn–winter temperature and latitude, but not geographical distance or autumn–winter temperature difference, were correlated with their respective residual pairwise genetic differentiation (Table 2). This suggests that genetic differentiation is mainly driven by site characteristics (latitude, mean autumn–winter temperature) rather than environmental contrast between sites. Mean autumn–winter temperature remained significantly correlated with genetic differentiation after correcting for latitude (partial Mantel test: $r_M = 0.31$, $P = 0.019$), whereas mean latitude was not correlated with genetic differentiation anymore after correcting for mean autumn–winter temperature (partial Mantel test: $r_M = -0.03$, $P = 0.534$). This suggests that mean autumn–winter temperature was the best predictor of genetic differentiation among the tested environmental effects.

DISCUSSION

BIOLOGICAL RELEVANCE OF THE OBSERVED GENETIC DIFFERENTIATION

The low but significant global genetic differentiation based on microsatellite markers suggests extensive gene flow among great tit populations across Europe. Nevertheless, the overall deviation from HWE, the absence of inbreeding within sites (as revealed by heterozygosity) and the overall population differentiation support a Wahlund effect, i.e. a substructure

among sites. Individual-based clustering methods failed to characterise discrete genetic groups, yet found some indication for substructure among south-western sites. This is consistent with the high proportion of the genetic variance (> 98%) observed within populations (e.g. Latch *et al.*, 2006; Chen *et al.*, 2007). We are nevertheless confident about the validity of the significant global genetic differentiation given the relatively large sample sizes and because none of the analyses suggested a bias in both global and pairwise genetic differentiation due to variation in sample size among sites or being associated by specific loci and sites.

In general, a significant IBD supports the biological relevance of low genetic differentiation among populations (e.g. F_{ST} values around 0.003), especially in species characterised by large population sizes and high gene flow such as birds (e.g. Prochazka *et al.*, 2011) or marine fishes (e.g. Purcell *et al.*, 2006). But low genetic differentiation even in absence of IBD may also reflect heterogeneity in gene flow affecting ongoing microevolutionary processes in highly mobile organisms. This is illustrated by the case of a physically isolated island population of great tits, where immigrants from the mainland can be easily identified (Postma & van Noordwijk, 2005). In this population, direct (i.e. observed movements of individuals) and indirect (i.e. genetic, based on microsatellite markers) measures of gene flow were compared. The genetic differentiation between resident and immigrant individuals was low but significant ($F_{ST} = 0.007$; Postma *et al.*, 2009). Consistent with a higher immigration rate in the western part (43%) compared to the eastern part (13%) of the study island, a low but significant genetic differentiation was found between the two parts ($F_{ST} = 0.011$;

Table 2. Effects of environmental factors on the genetic differentiation between sampling sites across Europe and its variation based on Mantel tests (r_M)

Response variable:	F_{ST}		Residuals on F_{ST}				Residuals on G''_{ST}				Residuals on D			
	r_M	P	r_M	P	r_M	P	r_M	P	r_M	P	r_M	P	r_M	P
Mean autumn–winter temperature	0.57	< 0.001	0.17	0.022	0.57	< 0.001	0.16	0.025	0.57	< 0.001	0.15	0.023		
Latitude	-0.50	0.002	-0.23	0.010	-0.50	0.002	-0.22	0.010	-0.51	0.001	-0.22	0.007		
Geographic distance	0.36	< 0.001	0.06	0.205	0.37	0.001	0.07	0.188	0.37	< 0.001	0.07	0.166		
Difference in autumn–winter temperature	0.39	0.002	0.10	0.125	0.39	0.002	0.10	0.120	0.39	0.002	0.10	0.114		

Genetic distance was measured as pairwise F_{ST} , G''_{ST} and D and their variation was investigated using the residuals of a linear regression between each environmental factor and the respective pairwise genetic distances. See text for details. Significant correlations are indicated in bold.




Figure 5. Relationship between pairwise F_{ST} values and (A) geographic distance between sites and (B) mean autumn–winter temperature of the two sites in pairwise comparisons. White dots: pairwise F_{ST} values between northern sites; grey dots: pairwise F_{ST} values between one northern and one south-western site; black dots: pairwise F_{ST} values between south-western sites.

Postma *et al.*, 2009). Because mainland individuals lay larger clutches, immigration was shown to impede local adaptation in the western but not the eastern part of the island (Postma & van Noordwijk, 2005). Using similar microsatellite markers in the present study, we also found comparable levels of genetic differentiation between populations, supporting the biological implications of our findings. Lastly, using a restricted set of microsatellite markers, we retrieved a comparable level of genetic differentiation between two sites (NE.HO and UK.WY; $F_{ST} = 0.005$) as has been observed with several thousand SNP markers for the same sites (Van Bers *et al.*, 2012; $F_{ST} = 0.010$). The slightly higher level of genetic differentiation in their study could be due to the inclusion of some highly divergent outlier loci. Another study also using the same SNP set further identified cryptic genetic differentiation within the UK.WY

site, which was similarly driven by few (< 1%) markers (Garroway *et al.*, 2013). Thus our microsatellite data set seems to be suitable to accurately calculate population genetic estimates that resemble average genome-wide patterns (i.e. Van Bers *et al.*, 2012), whereas few genomic regions may exist that underlie patterns of local adaptation (Van Bers *et al.*, 2012; Garroway *et al.*, 2013).

Our analyses revealed higher genetic differentiation in south-western compared to northern European sites. This finding suggests decreased gene flow between south-western and northern Europe as well as within south-western Europe. Subsequent generalisations towards other southern European populations need to be done with caution since our sampling design focused only on south-western populations. A similar pattern was reported for different passerine species as well as for plants and mammals (Hewitt, 2000; Kvist *et al.*, 2004; Prochazka *et al.*, 2011; Pentzold *et al.*, 2013) and is generally interpreted as the result of post-glacial recolonization. In the present case, the higher divergence of southern populations compared to northern ones could be due to the fact that both groups may have derived from different glacial refugia (Hewitt, 2000). Such a scenario has been suggested for other tit species, for which distinct glacial refugia may have existed in the Mediterranean region (Kvist *et al.*, 2004) and across Europe (Pentzold *et al.*, 2013). However, for several reasons, the genetic differentiation observed in great tits using microsatellite markers seems unlikely to result from the occurrence of one or several genetic lineages that have recolonized northern Europe from distinct refugia. First, the presence of several glacial refugia would have led to the existence, at least in south-western populations, of genetic variations specific to the multiple refugia causing a higher genetic diversity within the Iberian Peninsula (Prochazka *et al.*, 2011; Pentzold *et al.*, 2013). In contrast, the Iberian Peninsula harboured a level of allelic richness at microsatellite markers that was comparable with all other sites (7.26 and 7.32 alleles respectively). Similarly, phylogenetic studies showed in great tits a homogeneous mitochondrial diversity from northern to southern Europe (Supporting Information, Fig. S1), which is consistent with a colonisation from a single refugium and the absence of strong geographical barriers to dispersal (Kvist *et al.*, 1999, 2003; Pavlova *et al.*, 2006). Second, a rapid post-glacial range expansion from a single refugium is likely to result in lower genetic diversity within the colonized range as opposed to the ancestral refugium (Pavlova *et al.*, 2006; Antoniazza *et al.*, 2014). In contrast, Iberian populations had a slightly lower allelic richness per site compared with all other sites (6.71 ± 0.29 and

7.11 ± 0.29 alleles respectively, Supporting Information, Table S2). Interestingly haplotype diversity was lower in all south-western populations than in the north-eastern populations in coal tits (Pentzold *et al.*, 2013) suggesting that a lower genetic diversity in southern regions could have arisen long time ago. However such a pattern was not detected with mitochondrial DNA in great tits (Pavlova *et al.*, 2006). Therefore, the observed patterns of genetic differentiation at microsatellite loci among great tit populations are unlikely to result from post-glacial recolonization processes from one or several refugia but rather represent other historical and/or recent processes. Further studies using genetic modelling approaches combined with increased genomic coverage are, however, necessary to elucidate the factors underlying the pattern observed here.

HOW COULD GENE FLOW BE SHAPED BY TEMPERATURE?

Latitude and (autumn–winter) temperature were significantly correlated with both the level of genetic differentiation among populations and its level of variation in contrast with the geographical distance and the difference in temperature that explained only the level of genetic differentiation among populations. Moreover, only temperature was significantly associated with the level of genetic differentiation after taking into account latitude. Finally, temperature but not latitude explained the decrease of genetic diversity from the South to the North. The effect of temperature on different components of the genetic variation suggests a strong relationship between temperature and neutral genetic structure among great tit populations. We cannot exclude, however, that temperature is correlated with additional environmental factors such as photoperiod or irradiance cues (De Frenne *et al.*, 2013) and then the correlation between temperature and genetic differentiation is a by-product of the effect of environmental factors on genetic variation that we did not measure here. Nonetheless the relationship between temperature and neutral genetic structure suggests that genetic differentiation, and hence gene flow, may be related to winter local movements and partial migration (Nowakowski & Vähätalo, 2003; Nilsson, Alerstam & Nilsson, 2008). This finding could also be associated with winter severity: food availability may be especially restricted in northern Europe (Newton, 2012 but see Nilsson *et al.*, 2008; Nowakowski & Vähätalo, 2003) when insect abundances are lowest and great tits become mainly granivorous (Vel'ky, Kanuch & Kristin, 2011). Great tits are considered to be resident in southern and western Europe, but partial migrants in northern

Europe, as shown in particular by captures at migratory passage sites in the autumn and spring (Gosler, 2002; Nowakowski & Vähätalo, 2003; Poluda, 2011). Part of the birds (especially juveniles) may move during winter over short to long distances (up to >1000 km; Nilsson *et al.*, 2008; Nowakowski & Vähätalo, 2003). In spring, these migrants may either stay on the wintering grounds or return to their natal region to breed more or less close to their natal site (Gosler, 2002; Nowakowski & Vähätalo, 2003; Nilsson *et al.*, 2008). Partial migration could therefore generate on average longer dispersal distances, associated with higher variance, in the northern compared to southern European populations (see Orell *et al.*, 1999). Although part of the immigrant individuals (often around 50% of local breeders in monitored populations) may originate from the surroundings of study areas (e.g. Verhulst *et al.*, 1997), differences in immunological, behavioural and/or life-history traits between potential immigrants (i.e. not previously captured in the population) and locally born individuals (e.g. Snoeijs *et al.*, 2004; Postma & van Noordwijk, 2005) may support the existence of long-distance immigration in great tits. Because obtaining additional information on the origin of immigrant individuals in the field is highly challenging, this hypothesis, however, remains difficult to test.

Interestingly, similar genetic structures across Europe have been found in other small passerine species, i.e. for the bluethroat (*Luscinia svecica*; Johnsen *et al.*, 2006) and the pied flycatcher (*Ficedula hypoleuca*; Lehtonen *et al.*, 2009). In the latter case, no large-scale differentiation was observed in north-eastern Europe but small-scale differentiation was found in southern Europe. Because the pied flycatcher is an obligatory migratory species, wintering in sub-Saharan Africa, the lower genetic differentiation of northern sites cannot be explained by differences in winter movements linked to winter severity. Nevertheless, lower philopatry and local recruitment rates, and thus higher dispersal rates, have been suggested in northern compared to southern sites for several migratory species, including the pied flycatcher (Lehtonen *et al.*, 2009) and the barn swallow (Balbontin *et al.*, 2009). In these species dispersal may be linked to other environmental factors such as e.g. habitat stability, fragmentation or elevation. Both here and in the study by Lehtonen *et al.* (2009), southern populations were sampled in specific habitats, including high elevation sites (great tits: SP.MA, SP.FR and CH.BE > 500 m.a.s.l.; pied flycatchers: Lehtonen *et al.*, 2009), urban environment (FR.MO) or plantations (SP.MU), in contrast with northern sites located mainly in temperate lowland forests. In southern Europe, stable habitat hetero-

geneity, niche specialisation or high temperature may promote local adaptation (e.g. Husby, Visser & Kruuk, 2011). This could increase local genetic differentiation and select against dispersal to a higher degree than in the northern regions (Van Doorslaer *et al.*, 2009), where the availability of large and/or homogeneous habitat patches may reduce dispersal costs (Travis & Dytham, 1999) in both migratory and sedentary species. Individuals of the southern populations may therefore be less prone to accept breeding in new sites, leading to lower gene flow. Consequently, intraspecific differentiation might be more likely than neutral differentiation in southern sites (e.g. Johnsen *et al.*, 2006; Lehtonen *et al.*, 2012).

CONCLUSION

Non-random dispersal and genetic structure in great tits have previously been investigated at small scales, providing evidence for local adaptation (i.e. within a few km; Garant *et al.*, 2005; Postma & Noordwijk, 2005; Postma *et al.*, 2009; Garroway *et al.*, 2013). Here, we compared populations across Europe and found low but significant genetic differentiation among populations. This differentiation was unrelated to geographical distance between sites but was influenced by geographic location and environmental factors, in particular autumn-winter temperature. This finding might have important implications for the evolutionary trajectories of great tit populations and other species showing similar patterns. The northern populations may represent a single large population in which gene flow drives demographic and evolutionary processes. In this case, habitat choice and assortative mating may play a central role in local adaptation processes (e.g. Postma & van Noordwijk, 2005). In contrast, the southern populations may be more isolated and experience stronger genetic drift and/or higher selective pressures (e.g. Lehtonen *et al.*, 2012). Studying potentially ongoing intraspecific diversification may be particularly relevant in these populations.

The association between genetic differentiation and winter severity may have further implications in the context of climate change. If the increase of winter temperatures favours increased philopatry in northern populations (e.g. Van Vliet, Musters & Ter Keurs, 2009), the latter may reach a gene flow-drift equilibrium. As a consequence, increased genetic differentiation and IBD could arise, favouring neutral genetic differentiation and/or local adaptation. Conversely, southern populations may become extinct if genetic adaptation or phenotypic plasticity fail to allow to adapt sufficiently fast (Visser, 2008; Boeye

et al., 2013). Alternatively, an increase of philopatry among northern populations, induced by warmer winters could intensify competition especially during the breeding season, leading to a population decline (Kokko, 2011 but see Stenseth *et al.*, 2015). And southern populations may persist if climate change combined with habitat fragmentation select for less emigration but larger dispersal distances (Boeye *et al.*, 2013; Fronhofer *et al.*, 2014). If global warming results in population extinction, proportionally more genetic diversity would be lost in the South than in the North of Europe. Because most studies on great tits have been conducted in north-central Europe, further work is needed to assess both the large-scale variation of philopatry, its relation to local and regional winter partial migration movements and its consequence in terms of gene flow between populations.

ACKNOWLEDGEMENTS

The study was funded by the Swiss National Science Foundation (3100A0-102017 to HR, P2BEP3_152103 to KL and PMPDP3_151361/161858 to ML). We thank L. Gustafsson and J. Forsman who kindly allowed us to use their nest boxes. We gratefully thank E. Bezault, L. Cornetti and two anonymous reviewers for valuable advice on genetic analyses and helpful comments on the manuscript. All samples were collected under licenses of national authorities and financially supported by the Academy of Finland grant (to NP and 265859 to TE), the Netherlands Organisation for Scientific Research (NWO-VICI 86503003 to JK and NWO-VICI to MV), the Netherlands Genomics Initiative (Horizon grant to KvO), the Hungarian Scientific Research Fund (OTKA 75618 to JT), the Estonian Ministry of Education and Science (IUT 34-8 to RM), the OSU-OREME, the Spanish Ministry of Education and Science (CGL2010-21933-CO-02 to EB), the Portuguese Foundation for Science and Technology (SFRH/BD/13594/2003 to ACN) and the Agence Nationale de la Recherche (ANR-08-JCJC-0041-01 to BD).

REFERENCES

- Ahola MP, Laaksonen T, Eeva T, Lehikoinen E. 2009. Great tits lay increasingly smaller clutches than selected for: a study of climate- and density-related changes in reproductive traits. *Journal of Animal Ecology* **78**: 1298–1306.
- Antoniazza S, Kanitz R, Neuenschwander S, Burri R, Gaigher A, Roulin A, Goudet J. 2014. Natural selection in a postglacial range expansion: the case of the colour cline in the European barn owl. *Molecular Ecology* **23**: 5508–5523.

- Balbontin J, Moller AP, Hermosell IG, Marzal A, Reviriego M, de Lope F. 2009. Geographic patterns of natal dispersal in barn swallows *Hirundo rustica* from Denmark and Spain. *Behavioral Ecology and Sociobiology* **63**: 1197–1205.
- Björklund M, Ruiz I, Senar JC. 2010. Genetic differentiation in the urban habitat: the great tits (*Parus major*) of the parks of Barcelona city. *Biological Journal of the Linnean Society* **99**: 9–19.
- Boeye J, Travis JMJ, Stoks R, Bonte D. 2013. More rapid climate change promotes evolutionary rescue through selection for increased dispersal distance. *Evolutionary Applications* **6**: 353–364.
- Burnham KP, Anderson DR, Huyvaert KP. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* **65**: 23–35.
- Caswell H, Lensink R, Neubert MG. 2003. Demography and dispersal: life table response experiments for invasion speed. *Ecology* **8**: 1968–1978.
- Charmantier A, Doutrelant C, Dubuc-Messier G, Fargevieille A, Szulkin M. 2016. Mediterranean blue tits as a case study of local adaptation. *Evolutionary Applications* **9**: 135–152.
- Chen C, Durand E, Forbes F, Francois O. 2007. Bayesian clustering algorithms ascertaining spatial population structure: a new computer program and a comparison study. *Molecular Ecology Notes* **7**: 747–756.
- De Frenne P, Graae BJ, Rodriguez-Sanchez F, Kolb A, Chabrierie O, Decocq G, De Kort H, De Schrijver A, Diekmann M, Eriksson O, Gruwez R, Hermy M, Lenoir J, Plue J, Coomes DA, Verheyen K. 2013. Latitudinal gradients as natural laboratories to infer species' responses to temperature. *Journal of Ecology* **101**: 784–795.
- Do C, Waples RS, Peel D, Macbeth GM, Tillett BJ, Ovenden JR. 2014. NEESTIMATOR v2: re-implementation of software for the estimation of contemporary effective population size (N_e) from genetic data. *Molecular Ecology Resources* **14**: 209–214.
- Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology* **14**: 2611–2620.
- Felsenstein J. 1989. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics* **5**: 164–166.
- Forsman JT, Monkkonen M. 2003. The role of climate in limiting European resident bird populations. *Journal of Biogeography* **30**: 55–70.
- Fronhofer EA, Stelz JM, Lutz E, Poethke HJ, Bonte D. 2014. Spatially correlated extinctions select for less emigration but larger dispersal distances in the spider mite *Tetranychus urticae*. *Evolution* **68**: 1838–1844.
- Garant D, Kruuk LEB, McCleery RH, Sheldon BC. 2004. Evolution in a changing environment: a case study with great tit fledging mass. *American Naturalist* **164**: E115–E129.
- Garant D, Kruuk LEB, Wilkin TA, McCleery RH, Sheldon BC. 2005. Evolution driven by differential dispersal within a wild bird population. *Nature* **433**: 60–65.
- Garroway CJ, Radersma R, Sepil I, Santure AW, De Cauwer I, Slate J, Sheldon BC. 2013. Fine-scale genetic structure in a wild bird population: the role of limited dispersal and environmentally based selection as causal factors. *Evolution* **67**: 3488–3500.
- Gosler AG. 2002. Great tit, *Parus major*. In: Wernham C, Siriwardena G, Toms M, Marchant JH, Clark JA, Baillie S, eds. *The migration atlas: movements of the birds of Britain and Ireland*. London: T & AD Poyser, 602–605.
- Goudet J. 1995. FSTAT (Version 1.2): a computer program to calculate F-statistics. *Journal of Heredity* **86**: 485–486.
- Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.
- Huld TA, Suri M, Dunlop ED, Micale F. 2006. Estimating average daytime and daily temperature profiles within Europe. *Environmental Modelling & Software* **21**: 1650–1661.
- Husby A, Visser ME, Kruuk LEB. 2011. Speeding up microevolution: the effects of increasing temperature on selection and genetic variance in a wild bird population. *PLoS Biology* **9**: 9.
- Hutchison DW, Templeton AR. 1999. Correlation of pairwise genetic and geographic distance measures: inferring the relative influences of gene flow and drift on the distribution of genetic variability. *Evolution* **53**: 1898–1914.
- Johnsen A, Andersson S, Garcia Fernandez J, Kempnaers B, Pavel V, Questiau S, Raess M, Rindal E, Lifjeld JT. 2006. Molecular and phenotypic divergence in the bluethroat (*Luscinia svecica*) subspecies complex. *Molecular Ecology* **15**: 4033–4047.
- Jombart T, Devillard S, Balloux F. 2010. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics* **11**: 94.
- Junge C, Vøllestad LA, Barson NJ, Haugen TO, Otero J, Sætre G-P, Leder EH, Primmer CR. 2011. Strong gene flow and lack of stable population structure in the face of rapid adaptation to local temperature in a spring-spawning salmonid, the European grayling (*Thymallus thymallus*). *Heredity* **106**: 460–471.
- Koizumi I, Yamamoto S, Maekawa K. 2006. Decomposed pairwise regression analysis of genetic and geographic distances reveals a metapopulation structure of stream-dwelling Dolly Varden charr. *Molecular Ecology* **15**: 3175–3189.
- Kokko H. 2011. Directions in modelling partial migration: how adaptation can cause a population decline and why the rules of territory acquisition matter. *Oikos* **120**: 1826–1837.
- Kvist L, Ruokonen M, Lumme J, Orell M. 1999. The colonization history and present-day population structure of the European great tit (*Parus major major*). *Heredity* **82**: 495–502.
- Kvist L, Martens J, Higuchi H, Nazarenko AA, Valchuk OP, Orell M. 2003. Evolution and genetic structure of the great tit (*Parus major*) complex. *Proceedings of Royal Society of London Serie B: Biological Sciences* **270**: 1447–1454.

- Kvist L, Viiri K, Dias PC, Rytönen S, Orell M. 2004.** Glacial history and colonization of Europe by the blue tit *Parus caeruleus*. *Journal of Avian Biology* **35**: 352–359.
- Kvist L, Arbabi T, Päckert M, Orell M, Martens J. 2007.** Population differentiation in the marginal populations of the great tit (*Paridae: Parus major*). *Biological Journal of the Linnean Society* **90**: 201–210.
- Latch EK, Dharmarajan G, Glaubitz JC, Rhodes OE. 2006.** Relative performance of Bayesian clustering software for inferring population substructure and individual assignment at low levels of population differentiation. *Conservation Genetics* **7**: 295–302.
- Lehtonen P, Laaksonen T, Artemyev AV, Belskii E, Both C, Bureš S, Bushuev AV, Krams I, Moreno J, Mägi M, Nord A, Potti J, Ravussin P-A, Sirkiä P, Sætre G-P, Primmer CR. 2009.** Geographic patterns of genetic differentiation and plumage colour variation are different in the pied flycatcher (*Ficedula hypoleuca*). *Molecular Ecology* **18**: 4463–4476.
- Lehtonen P, Laaksonen T, Artemyev AV, Belskii E, Berg PR, Both C, Buggiotti L, Bureš S, Burgess M, Bushuev AV, Krams I, Moreno J, Mägi M, Nord A, Potti J, Ravussin P-A, Sirkiä P, Sætre G-P, Winkel W, Primmer CR. 2012.** Candidate genes for colour and vision exhibit signals of selection across the pied flycatcher (*Ficedula hypoleuca*) breeding range. *Heredity* **108**: 431–440.
- Lemoine M, Lucek K, Perrier C, Saladin V, Adriaenssens F, Barba E, Belda EJ, Charmantier A, Cichoń M, Eeva T, Grégoire A, Hinde CA, Johnsen A, Komdeur J, Mänd R, Matthysen E, Norte AC, Pitala N, Sheldon BC, Slagsvold T, Tinbergen JM, Török J, Ubels R, Oers KN, Visser ME, Doligez B, Richner H. 2016.** Data from: Low but contrasting neutral genetic differentiation shaped by winter temperature in European great tits. *Dryad Digital Repository* doi:10.5061/dryad.7md3q
- Loiselle BA, Sork VL, Nason J, Graham C. 1995.** Spatial genetic-structure of a tropical understory shrub, *Psychotria officinalis* (Rubiaceae). *American Journal of Botany* **82**: 1420–1425.
- Matthysen E. 2005.** Density-dependent dispersal in birds and mammals. *Ecography* **28**: 403–416.
- Mazerolle MJ. 2015.** AICcmodavg: Model selection and multi-model inference based on (Q)AIC(c). R package version 2.0-3.
- Meirmans PG, Hedrick PW. 2011.** Assessing population structure: F-ST and related measures. *Molecular Ecology Resources* **11**: 5–18.
- Meirmans PG, Van Tienderen PH. 2004.** GenoType and GenoDive: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Notes* **4**: 792–794.
- Miller MP, Mullins TD, Parrish JW Jr, Walters JR, Haig SM. 2012.** Variation in migratory behavior influences regional genetic diversity and structure among american kestrel populations (*Falco sparverius*) in North America. *Journal of Heredity* **103**: 503–514.
- Musiani M, Leonard JA, Cluff D, Gates C, Mariani S, Paquet PC, Vilà C, Wayne RK. 2007.** Differentiation of tundra/taiga and boreal coniferous forest wolves: genetics, coat colour and association with migratory caribou. *Molecular Ecology* **16**: 4149–4170.
- Nathan R, Perry G, Cronin JT, Strand AE, Cain ML. 2003.** Methods for estimating long-distance dispersal. *Oikos* **103**: 261–273.
- Newton I. 2012.** Obligate and facultative migration in birds: ecological aspects. *Journal of Ornithology* **153**: 171–180.
- Nilsson ALK, Linstrom A, Jonzen N, Nilsson SG, Karlsson L. 2006.** The effect of climate change on partial migration - the blue tit paradox. *Global Change Biology* **12**: 2014–2022.
- Nilsson ALK, Alerstam T, Nilsson JA. 2008.** Diffuse, short and slow migration among blue tits. *Journal of Ornithology* **149**: 365–373.
- Nowakowski JK, Vähätalo A. 2003.** Is the great tit *Parus major* an irruptive migrant in North-east Europe? *Ardea* **91**: 231–244.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, O'Hara RB, Simpson GL, Solymos PH, Stevens MH, Wagner H. 2011.** vegan: Community Ecology Package R package version 1.17-6: <http://CRAN.R-project.org/package=vegan>.
- Orell M, Lahti K, Koivula K, Rytönen S, Welling P. 1999.** Immigration and gene flow in a northern willow tit (*Parus montanus*) population. *Journal of Evolutionary Biology* **12**: 283–295.
- Päckert M, Martens J, Sun Y-H. 2010.** Phylogeny of long-tailed tits and allies inferred from mitochondrial and nuclear markers (Aves: Passeriformes, Aegithalidae). *Molecular Phylogenetics and Evolution* **55**: 952–967.
- Parn H, Ringsby TH, Jensen H, Saether B-E. 2012.** Spatial heterogeneity in the effects of climate and density-dependence on dispersal in a house sparrow metapopulation. *Proceedings of the Royal Society B-Biological Sciences* **279**: 144–152.
- Pavlova A, Rohwer S, Drovetski SV, Zink RM. 2006.** Different post-pleistocene histories of Eurasian parids. *Journal of Heredity* **97**: 389–402.
- Peakall R, Smouse PE. 2006.** GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* **6**: 288–295.
- Pentzold S, Tritsch C, Martens J, Tietze DT, Giacalone G, Lo Valvo M, Nazarenko AA, Kvist L, Päckert M. 2013.** Where is the line? Phylogeography and secondary contact of western Palearctic coal tits (*Periparus ater*: Aves, Passeriformes, Paridae). *Zoologischer Anzeiger* **252**: 367–382.
- Peterson MA, Denno RF. 1998.** The influence of dispersal and diet breadth on patterns of genetic isolation by distance in phytophagous insects. *The American Naturalist* **152**: 428–446.
- Pilot M, Jędrezejewski W, Branicki W, Sidorovich VE, Jędrezejewska B, Stachura K, Funk SM. 2006.** Ecological factors influence population genetic structure of European grey wolves. *Molecular Ecology* **15**: 4533–4553.
- Poluda AM. 2011.** Spatial-temporal characteristics of migrations of the great tits *parus major* (Aves, Passeriformes, Paridae) in Ukraine. *Vestnik Zoologii* **45**: 343–357.

- Postma E, van Noordwijk AJ. 2005.** Gene flow maintains a large genetic difference in clutch size at a small spatial scale. *Nature* **433**: 65–68.
- Postma E, Den Tex R-J, Van noordwijk AJ, Mateman AC 2009.** Neutral markers mirror small-scale quantitative genetic differentiation in an avian island population. *Biological Journal of the Linnean Society* **97**: 867–875.
- Pritchard JK, Stephens M, Donnelly P. 2000.** Inference of population structure using multilocus genotype data. *Genetics* **155**: 945–959.
- Prochazka P, Stokke BG, Jensen H, Fainova D, Bellinvia E, Fossoy F, Vikan JR, Bryja J, Soler M. 2011.** Low genetic differentiation among reed warbler *Acrocephalus scirpaceus* populations across Europe. *Journal of Avian Biology* **42**: 103–113.
- Purcell JFH, Cowen RK, Hughes CR, Williams DA. 2006.** Weak genetic structure indicates strong dispersal limits: a tale of two coral reef fish. *Proceedings of the Royal Society B-Biological Sciences* **273**: 1483–1490.
- R Core Team. 2013. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.
- Rice W. 1989.** Analyzing tables of statistical tests. *Evolution* **43**: 223–225.
- Rousset F. 2008.** GENEPOP '007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources* **8**: 103–106.
- Saladin V, Richner H. 2012.** A set of 48 microsatellite loci for the great tit *Parus major* including 15 novel markers. *Molecular Ecology Resources* **12**: 185–189.
- Sanford E, Kelly MW. 2011.** Local adaptation in marine invertebrates. *Annual Review of Marine Science* **3**: 509–535.
- Savolainen O, Pyhäjärvi T, Knürr T. 2007.** Gene flow and local adaptation in Trees. *Annual Review of Ecology, Evolution and Systematics* **38**: 595–619.
- Slatkin M. 1987.** Gene flow and the geographic structure of natural populations. *Science* **236**: 787–792.
- Snoeijs T, Van de Castele T, Adriaensen F, Matthysen E, Eens M. 2004.** A strong association between immune responsiveness and natal dispersal in a songbird. *Proceedings of Royal Society of London Serie B: Biological sciences* **271**: S199–S201.
- Snow DW, Perrins CM. 1998.** *The birds of the Western Palearctic*. Oxford, New York: Oxford University Press.
- Stenseth NC, Durant JM, Fowler MS, Matthysen E, Adriaensen F, Jonzen N, Chan K-S, Liu H, De Laet J, Sheldon BC, Visser ME, Dhondt AA 2015.** Testing for effects of climate change on competitive relationships and coexistence between two bird species. *Proceedings of the Royal Society B-Biological Sciences* **282**: 20141958.
- Travis JMJ, Dytham C. 1999.** Habitat persistence, habitat availability and the evolution of dispersal. *Proceedings of Royal Society of London Serie B: Biological sciences* **266**: 723–728.
- Van Bers NEM, Santure AW, Van Oers K, De Cauwer I, Dibbitts BW, Mateman C, Crooijmans RPMA, Sheldon BC, Visser ME, Groenen MAM, Slate J. 2012.** The design and cross-population application of a genome-wide SNP chip for the great tit *Parus major*. *Molecular Ecology Resources* **12**: 753–770.
- Van Doorslaer W, Vanoverbeke J, Duvivier C, Rousseaux S, Jansen M, Jansen B, Feuchtmayr H, Atkinson D, Moss B, Stoks R, De Meester L. 2009.** Local adaptation to higher temperatures reduces immigration success of genotypes from a warmer region in the water flea *Daphnia*. *Global Change Biology* **15**: 3046–3055.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* **4**: 535–538.
- Van Vliet J, Musters CJM, Ter Keurs W. 2009.** Changes in migration behaviour of blackbirds *Turdus merula* from the Netherlands. *Bird Study* **56**: 276–281.
- Vel'ky M, Kanuch P, Kristin A 2011.** Food composition of wintering great tits (*Parus major*): habitat and seasonal aspects. *Folia Zoologica* **60**: 228–236.
- Verhulst S, Perrins CM, Riddington R. 1997.** Natal dispersal of great tits in a patchy environment. *Ecology* **78**: 864–872.
- Visser ME. 2008.** Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proceedings of the Royal Society B-Biological Sciences* **275**: 649–659.
- Wang IJ. 2010.** Recognizing the temporal distinctions between landscape genetics and phylogeography. *Molecular Ecology* **19**: 2605–2608.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1. Haplotype network based on 103 available sequences from GenBank of the mitochondrial control region (578 bp) for 15 sites in Europe performed with the software POPART (Leigh and Bryant, 2015).

Figure S2. Principal coordinate analysis (PCoA) contrasting axes 1 vs. 2 when (a) 30 populations, and (b) 21 populations are included.

Figure S3. Principal coordinate analysis (PCoA) contrasting axes 1 vs. 3 when (a) 30 populations, (a) and (b) 21 populations are included.

Figure S4. (a) Mean (\pm SD) of estimated posterior likelihood, and (b) estimation of ΔK over 10 STRUCTURE runs for successive K values when 30 (i.e. all) populations are included in the analysis.

Figure S5. (a) Mean (\pm SD) of estimated posterior likelihood, and (b) estimation of ΔK over 10 STRUCTURE runs for successive K values when 21 (i.e. only 1 population from Gotland is included) populations are included in the analysis.

Figure S6. Assignment plots for $K = 3$ based on a sampling including (a) 30 populations, and (b) 21 populations (i.e. only one out of 10 populations from Gotland).

Table S1. Origin of the 103 sequences from GenBank of the mitochondrial control region (578 bp).

Table S2. Description of sampling sites and average genetic diversity indices per site.

Table S3. Characteristics of microsatellite loci developed on individuals from CH.BE: Locus name, repeat type and motif, species for which a locus has been described initially, as well as intra site variation in the number of alleles found, the smallest allele size and number of sites with null alleles at a specific locus.

Table S4. Probability assignments of STRUCTURE to cluster 1 and 2 by sites for $K = 3$ when 21 and 30 sites are included in the analysis. Populations are indicated in Figure 1.

Table S5. Comparison of models testing the effect of environmental factors on indices of genetic diversity per site and other parameters (A_R : allelic richness, H_E : unbiased expected heterozygosity, P_A : assignment probability, Kinship and N_e : effective population size).

SHARED DATA

Data deposited in the Dryad digital repository (Lemoine *et al.*, 2016).