University of Groningen

Geriatric assessment of patients treated for cutaneous head and neck malignancies in a tertiary referral center

de Vries, Julius; Heirman, Anne N.; Bras, Linda; Plaat, Boudewijn E.C.; Rácz, Eموke; van Kester, Marloes S.; Festen, Suzanne; de Bock, Geertruida H.; van der Laan, Bernard F.A.M.; Halmos, Gyorgy B.

Published in:
European Journal of Surgical Oncology

DOI:
[10.1016/j.ejso.2019.08.008](https://doi.org/10.1016/j.ejso.2019.08.008)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

de Vries, J., Heirman, A. N., Bras, L., Plaat, B. E. C., Rácz, E., van Kester, M. S., Festen, S., de Bock, G. H., van der Laan, B. F. A. M., & Halmos, G. B. (2020). Geriatric assessment of patients treated for cutaneous head and neck malignancies in a tertiary referral center: Predictors of postoperative complications. *European Journal of Surgical Oncology*, 46(1), 123-130.
<https://doi.org/10.1016/j.ejso.2019.08.008>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title

Geriatric assessment of patients treated for cutaneous head and neck malignancies in a tertiary referral center: predictors of postoperative complications

Authors

Julius de Vries, MD ^a

Anne N. Heirman, BSc ^a

Linda Bras, MD ^a

Boudewijn E.C. Plaat, MD, PhD ^a

Emoke Rácz, MD, PhD ^b

Marloes S. van Kester, MD, PhD ^b

Suzanne Festen, MD ^c

Geertruida H. de Bock, PhD ^d

Bernard F.A.M. van der Laan, MD, PhD ^a

Gyorgy B. Halmos, MD, PhD ^a

Author affiliations

^a Department of Otorhinolaryngology, Head and Neck Surgery, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

^b Department of Dermatology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

^c Department of Geriatric Medicine, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

^d Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands.

Corresponding author

Julius de Vries, MD

PO box 30.001 | 9700RB | Groningen, The Netherlands

E-mail: j.de.vries01@umcg.nl

Telephone: +31 (0)50 361 73 32

Fax: +31 (0)50 361 19 72

Funding: This article has no funding source.

Conflict of interest: The authors have no conflict of interest to declare.

Abstract

Introduction: As cutaneous head and neck malignancies are highly prevalent especially in older patients, the risk of surgical complications is substantial in this potentially vulnerable population. The objective of this study was to evaluate the value of geriatric assessment of this population with respect to postoperative complications.

Methods: Patients were prospectively included in OncoLifeS, a databiobank. Before surgery, patients underwent a geriatric assessment including multiple validated screening tools for frailty, comorbidity, polypharmacy, nutrition, functional status, social support, cognition and psychological status. Postoperatively, complications (Clavien-Dindo $\geq$ grade II) were registered. Uni- and multivariable logistic regression analyses were performed yielding odds ratios (ORs) and 95% confidence intervals (95%CI).

Results: 151 patients undergoing surgery for cutaneous head and neck malignancies were included in this study (mean age 78.9 years, 73.5% male). In a multivariable analysis, frailty measured by the Geriatric 8 (G8) (OR=6.34; 95%CI:1.73-23.25) was the strongest independent predictor of postoperative complications, among other predictors such as major treatment intensity (OR=2.73; 95%CI:1.19-6.26) and general anesthesia (OR=4.74; 95%CI:1.02-22.17), adjusted for age and sex.

Conclusion: Frailty, measured by G8, is the strongest predictor of postoperative complications in patients undergoing surgery for cutaneous head and neck malignancies in addition to treatment intensity and type of anesthesia. Geriatric screening on multiple domains is recommended for patients with cutaneous malignancies undergoing head and neck surgery is recommended, as this population includes old patients and frequently suffers postoperative complications.

Key words

Geriatric screening, frailty, skin malignancy, head and neck surgery, postoperative complications.

64 Abbreviations

65 95%CI = 95% Confidence Interval

66 ACE-27 = Adult Comorbidity Evaluation 27

67 ADL = Activities of Daily Living

68 BCC = Basal Cell Carcinoma

69 BMI = Body Mass Index

70 CGA = Comprehensive Geriatric Assessment

71 CM = Cutaneous Melanoma

72 G8 = Geriatric 8

73 GDS-15 = Geriatric Depression Scale 15

74 GFI = Groningen Frailty Indicator

75 IADL = Instrumental Activities of Daily Living

76 MCC = Merkel Cell Carcinoma

77 MMSE = Mini Mental State Examination

78 MUST = Malnutrition Universal Screening Tool

79 NL = Netherlands

80 NMSC = Non-melanoma Skin Cancer

81 OR = Odds Ratio

82 SCC = Squamous Cell Carcinoma

83 SD = Standard Deviation

84 SES = Socioeconomic Status

85 TUG = Timed Up and Go

86 UMCG = University Medical Center Groningen

87 Introduction

88 Skin cancer is the most common type of cancer worldwide.¹ In the United States, the incidence of non-
 89 melanoma skin cancer (NMSC) and cutaneous melanoma (CM) is estimated to be at least over 5.5 million
 90 annually.^{2,3} The incidence of both NMSC and CM are dramatically on the rise,⁴⁻⁶ with especially the proportion
 91 of older patients increasing.⁷ This results from the expanding older population in general and also due to older
 92 patients' higher cumulative sun exposure. Possibly, associated diseases,^{8,9} use of immunosuppressive
 93 medications,¹⁰ or exposure to prior radiation therapy,¹¹ contribute to this as well.

94 Cutaneous malignancies of the head and neck occur more frequently^{12,13} and are at higher risk for
 95 metastasis than other subsites.¹⁴ The cornerstone of treatment in most of the cases is surgery, ranging from a
 96 straightforward local excision to extended resections with neck dissections and even complex reconstructive
 97 surgery. If radical surgery is beyond possibilities, because of expected functional or cosmetic impairments or
 98 foreseen complications in older patients, radiotherapy is an effective treatment modality both as primary
 99 therapy or as an adjuvant therapy.¹⁵ With surgery remaining the primary choice of treatment, the risk of
 100 postoperative complications is substantial in this elderly and possibly vulnerable population, like previously
 101 described after head and neck oncological surgery.¹⁶

102 Comprehensive Geriatric Assessment (CGA) by a geriatrician or specialized nurse is the gold standard
 103 to expose vulnerabilities in older patients, which may be treated to prevent perioperative complications.¹⁷ CGA
 104 focuses on multiple geriatric domains such as comorbidities, polypharmacy, nutritional status, functional
 105 status, social support and psychological status.¹⁸ Because of its time consuming nature, screening tools such as
 106 the Groningen Frailty Indicator (GFI) and the Geriatric 8 (G8) have been developed to detect vulnerable
 107 patients who may benefit from a CGA.^{19,20}

108 The role of geriatric screening is established in many oncological patient populations, but not in
 109 cutaneous malignancies, even though this population is relatively old. Therefore, in the present study, we
 110 evaluated the role of geriatric assessment and frailty screening with respect to postoperative complications in
 111 surgically treated patients for cutaneous head and neck malignancies in a tertiary center.

Materials and methods

Study design

The present cohort study included patients who were enrolled in OncoLifeS, a prospective oncological databiobank at the University Medical Center Groningen. Study protocol was approved by the OncoLifeS scientific board.

Study population

Between October 2014 and October 2018 all consecutive patients referred for a cutaneous malignancy to the Department of Otorhinolaryngology, Head and Neck Surgery were included, regardless of age. Treatment strategies were according to national guidelines and discussed within the multidisciplinary head and neck tumor board and melanoma board, if applicable. If curative treatment was not possible or if patients received other primary treatment than surgery, patients were excluded from this study.

Data collection

Patient, tumor- and treatment characteristics were obtained from the electronic medical record and OncoLifeS database. Tumor stage was defined according to the seventh edition of the Union for International Cancer Control TNM Classification.²¹ At the first day of consultation, patients underwent a geriatric assessment at the outpatient clinic of our department, including the following geriatric domains: comorbidities, polypharmacy, nutritional status, functional status, social support, cognition and psychological status. Comorbidities were graded using the Adult Comorbidity Evaluation (ACE-27) as *none, mild, moderate or severe*.²² Polypharmacy was defined as the prescription of five or more medications on a daily basis.²³ Nutritional status was assessed using the Malnutrition Universal Screening Tool (MUST).²⁴ Functional status consisted of Activities of Daily Living (Katz-ADL), Instrumental Activities of Daily Living (IADL), Timed Up & Go (TUG) and history of falls.^{25–27} Social support was based on patient reported questionnaires. Socioeconomic status (SES) scores are publicly available scores, based on income, employment rate and educational status of postal code areas.²⁸ Cognition was assessed by the Mini Mental State Examination (MMSE) and presence of risk factors for delirium.^{29,30} Psychological status was scored using the Geriatric Depression Scale (GDS-15).³¹ Furthermore, two frailty screening instruments were completed including the Groningen Frailty Indicator (GFI) and the Geriatric 8

(G8).^{19,20} Postoperative complications occurring within 30 days after surgery were assessed from medical files using the Clavien-Dindo classification.³²

Statistical analysis

Patient characteristics were presented as mean $\pm$ standard deviation, median (range) or value (percentage). Univariable logistic regression analyses were performed to identify factors associated with postoperative complications. Analyses yielded odds ratios (ORs) with 95% confidence intervals (95% CIs). For multivariable logistic regression analysis with step backward method, variables with $p < 0.10$ were included. When collinearity was present between variables using Pearson and Spearman correlation coefficients, only clinically most relevant variables were selected. For variables eligible for multivariable analysis, missing values were imputed using multiple imputation. The multivariable model was fitted using a stepwise selection of predictors. All statistical analysis was performed with SPSS Statistics 23.0 software (IBM, Armonk, New York, United States of America). P-value < 0.05 was considered statistically significant.

Results

Study selection

Between October 2014 and October 2018, 197 patients with cutaneous head and neck malignancies were included in the OncoLifeS databiobank. After exclusion of patients treated with other primary treatment modalities than surgery and patients with no curative treatment options, a total of 151 patients remained eligible for analysis (Figure 1). There were no significant age and sex differences after exclusion.

Patient characteristics

Patient characteristics are presented in Table 1. The mean age of the patients was 78.9 years, ranging from 46.6 to 96.7 years. In this tertiary referral center, less than half of patients were referred with a primary tumor (49.7%), and others with residual tumor after recent treatment (29.8%) or recurrent tumor (20.5%). Most frequent histopathological subtypes of malignancies were squamous cell carcinoma (SCC; 59.6%), basal cell carcinoma (BCC; 18.5%), cutaneous melanoma (CM; 11.3%) and Merkel cell carcinoma (MCC; 6.0%).

Univariable analysis of predictors for postoperative complications

Occurrence of postoperative complications is listed in Table 2. Forty patients (26.5%) experienced complications grade II and higher according to the Clavien-Dindo classification. Factors associated with postoperative complications are shown in Table 3. Age was not a significant predictor (OR 0.98; 95%CI 0.94-1.02). Tumor characteristics, such as advanced tumor stage (OR 6.53; 95%CI 1.86-22.99) and large tumor diameter (OR 3.89; 95%CI 1.12-13.51) significantly predicted postoperative complications. Treatment characteristics, including locoregional surgery (OR 4.38; 95%CI 1.98-9.68), major treatment intensity (OR 3.46; 95%CI 1.62-7.39) and general anesthesia (OR 7.70; 95%CI 1.75-33.81), were also significantly related to postoperative complications.

Among the individual domains of geriatric assessment, only polypharmacy (OR 2.36; 95%CI 1.11-5.07) predicted postoperative complications respectively significantly (Table 3). Comorbidities, or impairments in functional status, social support, cognitive status or psychological status alone were not significantly associated with postoperative complications. Of the frailty screeners, the G8 was a strong, significant predictor of complications (OR 5.83; 95%CI 1.68-20.26) and GFI was not (OR 1.43; 95%CI 0.63-3.26).

182 *Independent predictors of postoperative complications*

183 A multivariable model was fitted with eligible variables (Table 4). Within the multivariable model, adjusted for
184 age and sex, major treatment intensity (OR 2.73; 95%CI 1.19-6.26), surgery under general anesthesia (OR 4.74;
185 95%CI 1.02-22.17) and frailty, measured by G8 (OR 6.34; 95%CI 1.73-23.25) were the most significant
186 independent predictors of postoperative complications grade II and higher.

Discussion

Patients with complex cutaneous head and neck malignancies are old and frequently experience postoperative complications. To our knowledge, this is the first study evaluating the value of geriatric assessment in a cohort of patients with cutaneous head and neck malignancies. Key findings show that frailty, measured by G8, is the strongest predictor of postoperative complications. Furthermore, tumor features, such as tumor size and stage, and treatment related predictors, such as treatment intensity and type of anesthesia seem to be related to postoperative complications.

With a mean age of nearly 80 years, the population of patients with cutaneous head and neck malignancies being referred to our tertiary hospital was remarkably aged. However, age did not predict postoperative complications within this population. This corresponds with other dermatological cohorts with head and neck skin malignancies.^{33–35} Pascual et al. showed that complications did not significantly differ between patients younger and older than 80 years, except for hemorrhagic complications.³⁶ This finding is in line with a large prospective cohort of Amici et al., showing more hemorrhagic complications in the elderly as well.³⁷ As significance disappears after correcting for use of anticoagulant medications, the higher amount of hemorrhagic complications is probably related to the increased use of anticoagulants with aging, and not to age itself. Just as age does not predict postoperative complications, it neither affects prognosis of patients with skin cancer.³⁸ Moreover, the majority of patients with a lower life expectancy, defined as age 85 years and older or a Charlson Comorbidity Index of 3 or higher, die of other causes than NMSC.³⁹ Whilst this does not apply directly to our cohort with much more complex cases, it does call the attention to the dilemma of “time to benefit”, referring to a clinical prediction, estimating whether the patient will live long enough to benefit from the treatment.³⁷ It is suggested that a comprehensive approach towards treatment decisions should at least include consideration of comorbidity, functional status and anticipated life expectancy in this specific population.⁴⁰

Complications after surgery of cutaneous head and neck malignancies performed by a dermatologist are usually rare. Percentages of the largest cohorts range between 3 and 6%.^{37,41–43} With 26.5% of patients suffering postoperative complications in our cohort, these outcomes seem much worse. However, our cohort suffers from a negative bias; higher tumor stage, more complex locations, more often lymph node metastasis, and consequently more major surgeries under general anesthesia. Furthermore, referral to a tertiary center may include more residual or recurrent tumor, which was the case in more than half of the patients. Clinical

research on tertiary cohorts of cutaneous head and neck malignancies are rarely reported; therefore comparison is difficult.

Our results show that tumor features such as histopathological type, tumor size and stage, and treatment characteristics, such as treatment intensity, adjuvant neck dissection and type of anesthesia, predict postoperative complications. Many of these variables are closely related to each other. After all, increased tumor size and more aggressive histopathological tumor type lead to more advanced stage, requiring extended surgery, possibly including neck dissection and general anesthesia. As a result, only the strongest predictors were included in multivariable analysis. Treatment intensity, defined as surgery time more than 120 minutes or 3 or more stages of Mohs micrographic surgery, and surgery under general anesthesia were found to be the most important predictors of postoperative complications. Length of surgery and neck dissection has been proven to predict postoperative complications in general head and neck oncological surgery as well.^{16,44–46} Even in case of excision under local anesthesia, length of surgery predicts postoperative complications in skin cancer surgery.³⁷

Frailty, measured by G8, was mostly associated with postoperative complications in this cohort. As far as we know, frailty has never been examined in a cohort undergoing surgery for cutaneous malignancies. Valdatta et al. investigated the FRAIL index in a cohort undergoing *reconstructive surgery* after NMSC excision.⁴⁷ A higher score on the FRAIL index was associated with more moderate to severe complications. Furthermore, Bras et al. included 45 patients with skin malignancies in their cohort of head and neck oncological patients.⁴⁵ The domain *health problems* of the GFI significantly predicted postoperative complications; however, subgroup analysis for patients with skin malignancies was not performed in that study. Interestingly, in our analysis, GFI showed no prognostic value. Comparing these studies is difficult, as there are large differences among frailty screening tools.⁴⁸ Domains that are covered by the G8 are nutritional status, polypharmacy, neuropsychological status and mobility. The G8 has been proven to be a useful tool in liver and colorectal surgery as a predictor of surgical complications.^{49,50} However, the value of G8 remains questionable, as the majority of our patients scored frail on the G8 (73.3%). This is in line with Pottel et al. and Hamaker et al. evaluating the G8 and other screening tools.^{48,51} They found that the G8 is very *sensitive* but not very *specific* with respect to its gold standard, a CGA. Referring all frail patients, based on G8 to a geriatrician for a CGA would be infeasible.

From all individual geriatric domains, polypharmacy and malnutrition were most significantly related with post-operative complications in our population. These domains are both well represented in the G8 as well. Polypharmacy is related to frailty and comorbidities, but also associated with outcome parameters such as postoperative complications, delirium, (chemo)radiation toxicity, increased hospital stay and mortality.²³ Across literature, however, polypharmacy lacks definition and cut-off values range largely, with ≥ 5 being the mostly used.²³ Whether certain specific medications such as anticoagulants were related to postoperative complications, just like in the study of Amici et al., was not possible to investigate using the current dataset.³⁷ Malnutrition is very common and undertreated in elderly.⁵² Evaluation of the nutritional status is therefore important in preoperative screening. Higher risk of malnutrition using MUST is associated with postoperative complications, increased hospital stay and mortality.^{53–55} Often, the body mass index (BMI) is used as an indicator for nutritional status, just as in MUST. However, normal values of 18.5–24.9 kg/mm² are based on mortality risk within a young and healthy population.⁵⁶ For older patients, a BMI <23 kg/mm² is already associated with increased mortality, and may therefore be a better cut-off value for underweight. The 7.9% of patients having risk of malnutrition measured by MUST in our cohort may be an underestimation of the real prevalence of malnutrition. Identification of such deficits is particularly important, as a geriatrician or a dietary consultant may be able to respectively manage polypharmacy or prevent malnutrition, lowering the risk of complications.

Based on our results, it seems that G8 is a very predictive screening tool. However, lack of specificity does not make it possible to adequately select vulnerable patients. Meanwhile, individual geriatric domains such as polypharmacy or malnutrition are too incomprehensive to point out patients at risk for surgical complications. The question arises what would then be an adequate screening strategy for elderly patients with cutaneous malignancies. As a recommendation, a two-step approach may bring a solution to this problem. The first step would be a short geriatric screening by a trained nurse, gathering information on all geriatric domains including comorbidities, polypharmacy, nutritional status, functional status, social support, cognition and psychological status, using short screening instruments. Then, the patients' screening information is discussed within a multidisciplinary team for elderly patients, in which the nurse, a geriatrician, and head and neck surgeon are present. The geriatrician may then already advise on perioperative management, or indicate a CGA and start pre-treatment optimization (second step). In this way, all potentially vulnerable patients have been reviewed prior to treatment, efficiently with respect to limited capacity of geriatric health care.

A strength of this work is the broad range of validated geriatric instruments and screening tools that were used to assess patients at baseline. Besides, many patient, tumor and treatment characteristics were available to adjust for existing differences between patients. Furthermore, patients were prospectively included and the selection of the study population was done carefully with respect to changes through exclusion process.

Limitations of our study may include that it is a single center study in a tertiary care hospital. As a result, the cohort contains a high percentage of complex cases, regarding tumor and treatment characteristics. Furthermore, the population was heterogenic, also in terms of tumor characteristics, like histopathology. However, as we were primarily investigating patient-related factors, this seemed to be less relevant in our study. Lastly, most complications have only temporary effect on the patients' lives. Other outcome parameters, such as health related quality of life may be of more value to this specific population and should be studied.

284 Conclusion

285 Frailty, measured by G8, is the strongest factor associated with postoperative complications in patients
286 undergoing surgery for cutaneous head and neck malignancies, besides treatment related predictors, such as
287 treatment intensity and type of anesthesia. Geriatric screening on multiple domains is recommended in
288 patients with cutaneous head and neck malignancies, as this population includes old patients and frequently
289 suffers postoperative complications.

290 **Acknowledgements**

291 None.

References

1. Leiter U, Eigentler T, Garbe C. Epidemiology of skin cancer. *Adv Exp Med Biol.* 2014;810:120-140.
2. Rogers HW, Weinstock MA, Feldman SR, Coldiron BM. Incidence Estimate of Nonmelanoma Skin Cancer (Keratinocyte Carcinomas) in the US Population, 2012. *JAMA Dermatology.* 2015;151(10):1081-1086. doi:10.1001/jamadermatol.2015.1187
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin.* 2019;69(1):7-34. doi:10.3322/caac.21551
4. Apalla Z, Lallas A, Sotiriou E, Lazaridou E, Ioannides D. Epidemiological trends in skin cancer. *Dermatol Pract Concept.* 2017;7(2):1-6. doi:10.5826/dpc.0702a01
5. Leiter U, Keim U, Eigentler T, Katalinic A, Holleczek B, Martus P, et al. Incidence, Mortality, and Trends of Nonmelanoma Skin Cancer in Germany. *J Invest Dermatol.* 2017;137(9):1860-1867. doi:https://doi.org/10.1016/j.jid.2017.04.020
6. Nikolaou V, Stratigos AJ. Emerging trends in the epidemiology of melanoma. *Br J Dermatol.* 2014;170(1):11-19. doi:10.1111/bjd.12492
7. Garcovich S, Colloca G, Sollena P, Andrea B, Balducci L, Cho WC, et al. Skin Cancer Epidemics in the Elderly as An Emerging Issue in Geriatric Oncology. *Aging Dis.* 2017;8(5):643-661. doi:10.14336/AD.2017.0503
8. Brewer JD, Shanafelt TD, Khezri F, Seda IMS, Zubair AS, Baum CL, et al. Increased incidence and recurrence rates of nonmelanoma skin cancer in patients with non-Hodgkin lymphoma: A Rochester Epidemiology Project population-based study in Minnesota. *J Am Acad Dermatol.* 2015;72(2):302-309. doi:https://doi.org/10.1016/j.jaad.2014.10.028
9. Quesenberry Charles P. J, Warton EM, Engels EA, Asgari MM, Silverberg MJ, Leyden W. HIV Infection Status, Immunodeficiency, and the Incidence of Non-Melanoma Skin Cancer. *JNCI J Natl Cancer Inst.* 2013;105(5):350-360. doi:10.1093/jnci/djs529
10. Engels EA, Pfeiffer RM, Fraumeni JF, Kasiske BL, Israni AK, Snyder JJ, et al. Spectrum of Cancer

- 318 Risk Among US Solid Organ Transplant Recipients. *JAMA*. 2011;306(17):1891-1901.
 319 doi:10.1001/jama.2011.1592
- 320 11. Van Vloten WA, Hermans J, Van Daal WAJ. Radiation-induced skin cancer and radiodermatitis
 321 of the head and neck. *Cancer*. 1987;59(3):411-414. doi:10.1002/1097-
 322 0142(19870201)59:3<411::AID-CNCR2820590310>3.0.CO;2-Z
- 323 12. Katalinic A, Kunze U, Schäfer T. Epidemiology of cutaneous melanoma and non-melanoma
 324 skin cancer in Schleswig-Holstein, Germany: incidence, clinical subtypes, tumour stages and
 325 localization (epidemiology of skin cancer). *Br J Dermatol*. 2003;149(6):1200-1206.
 326 doi:10.1111/j.1365-2133.2003.05554.x
- 327 13. Buettner PG, Raasch BA. Incidence rates of skin cancer in Townsville, Australia. *Int J Cancer*.
 328 1998;78(5):587-593. doi:10.1002/(SICI)1097-0215(19981123)78:5<587::AID-IJC10>3.0.CO;2-E
- 329 14. Brougham NDLS, Dennett ER, Cameron R, Tan ST. The incidence of metastasis from cutaneous
 330 squamous cell carcinoma and the impact of its risk factors. *J Surg Oncol*. 2012;106(7):811-815.
 331 doi:10.1002/jso.23155
- 332 15. Terra JB, Gaster MB, Halmos GB, Roodenburg JL, van der Vegt B, Romeijn TR, et al. Local
 333 control of 151 head and neck cutaneous squamous cell carcinoma after radiotherapy: a
 334 retrospective study on efficacy and prognostic factors. *Clin Otolaryngol*. 2017;42(4):851-855.
 335 doi:10.1111/coa.12707
- 336 16. Peters TTA, van Dijk BAC, Roodenburg JLN, van der Laan BFAM, Halmos GB. Relation Between
 337 Age, Comorbidity, and Complications in Patients Undergoing Major Surgery for Head and Neck
 338 Cancer. *Ann Surg Oncol*. 2014;21(3):963-970. doi:10.1245/s10434-013-3375-x
- 339 17. Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet*.
 340 2013;381(9868):752-762. doi:10.1016/S0140-6736(12)62167-9
- 341 18. Extermann M, Hurria A. Comprehensive geriatric assessment for older patients with cancer. *J*
 342 *Clin Oncol*. 2007;25(14):1824-1831. doi:10.1200/JCO.2007.10.6559
- 343 19. Schuurmans H, Steverink N, Lindenberg S, Frieswijk N, Slaets JPJ. Old or Frail: What Tells Us

- 344 More? *Journals Gerontol Ser A Biol Sci Med Sci*. 2004;59(9):M962-M965.
 345 doi:10.1093/gerona/59.9.M962
- 346 20. Delva F, Bellera CA, Mathoulin-Pélissier S, Rainfray M, Mertens C, Fonck M, et al. Screening
 347 older cancer patients: first evaluation of the G-8 geriatric screening tool. *Ann Oncol*.
 348 2012;23(8):2166-2172. doi:10.1093/annonc/mdr587
- 349 21. Sobin LH, Gospodarowicz MK, Wittekind C. *TNM Classification of Malignant Tumours*. 7th ed.
 350 Oxford, UK: Wiley-Blackwell; 2009.
- 351 22. Piccirillo JF. Importance of comorbidity in head and neck cancer. *Laryngoscope*.
 352 2000;110(4):593-602. doi:10.1097/00005537-200004000-00011
- 353 23. Sharma M, Loh KP, Nightingale G, Mohile SG, Holmes HM. Polypharmacy and potentially
 354 inappropriate medication use in geriatric oncology. *J Geriatr Oncol*. 2016;7(5):346-353.
 355 doi:https://doi.org/10.1016/j.jgo.2016.07.010
- 356 24. Elia M. *The "MUST" Report. Nutritional Screening for Adults: A Multidisciplinary Responsibility.*
 357 *Development and Use of the "Malnutrition Universal Screening Tool" (MUST) for Adults.*
 358 British Association for Parenteral and Enteral Nutrition (BAPEN); 2003.
- 359 25. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of Illness in the Aged: The
 360 Index of ADL: A Standardized Measure of Biological and Psychosocial Function. *JAMA*.
 361 1963;185(12):914-919. doi:10.1001/jama.1963.03060120024016
- 362 26. Lawton MP, Brody EM. Assessment of older people: Self-maintaining and instrumental
 363 activities of daily living. *Gerontologist*. 1969;9(3):179-186. doi:10.1093/geront/9.3_Part_1.179
- 364 27. Podsiadlo D, Richardson S. The Timed "Up & Go": A Test of Basic Functional Mobility for Frail
 365 Elderly Persons. *J Am Geriatr Soc*. 1991;39(2):142-148. doi:10.1111/j.1532-
 366 5415.1991.tb01616.x
- 367 28. Sociaal en Cultureel Planbureau. Statusscores 2016.
 368 [https://www.scp.nl/Onderzoek/Lopend_onderzoek/A_Z_alle_lopende_onderzoeken/Statussc](https://www.scp.nl/Onderzoek/Lopend_onderzoek/A_Z_alle_lopende_onderzoeken/Statusscores)
 369 [ores](https://www.scp.nl/Onderzoek/Lopend_onderzoek/A_Z_alle_lopende_onderzoeken/Statusscores). Accessed January 9, 2019.

- 370 29. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": A practical method for grading the
 371 cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12(3):189-198.
 372 doi:[https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
- 373 30. Oud FMM, de Rooij SEJA, Schuurman T, Duijvelaar KM, van Munster BC. [Predictive value of
 374 the VMS theme "Frail elderly": delirium, falling and mortality in elderly hospital patients]. *Ned*
 375 *Tijdschr Geneesk.* 2015;159:A8491.
- 376 31. Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS): Recent evidence and development of
 377 a shorter version. *Clin Gerontol.* 1986;5(1-2):165-173. doi:10.1300/J018v05n01_09
- 378 32. Dindo D, Demartines N, Clavien P-A. Classification of surgical complications: a new proposal
 379 with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.*
 380 2004;240(2):205-213. doi:10.1097/01.sla.0000133083.54934.ae
- 381 33. Paradela S, Pita-Fernández S, Peña C, Fernández-Jorge B, García-Silva J, Mazaira M, et al.
 382 Complications of ambulatory major dermatological surgery in patients older than 85 years. *J*
 383 *Eur Acad Dermatology Venereol.* 2010;24(10):1207-1213. doi:10.1111/j.1468-
 384 3083.2010.03628.x
- 385 34. Dhiwakar M, Khan NA, McClymont LG. Surgery for head and neck skin tumors in the elderly.
 386 *Head Neck.* 2007;29(9):851-856. doi:10.1002/hed.20605
- 387 35. Bouhassira J, Bosc R, Greta L, Hersant B, Niddam J, Zehou O, et al. Factors associated with
 388 postoperative complications in elderly patients with skin cancer: A retrospective study of 241
 389 patients. *J Geriatr Oncol.* 2016;7(1):10-14. doi:10.1016/j.jgo.2015.11.004
- 390 36. Pascual JC, Belinchón I, Ramos JM. Cutaneous Surgery Complications in Individuals Aged 80
 391 and Older Versus Younger Than 80 After Excision of Nonmelanoma Skin Cancer. *J Am Geriatr*
 392 *Soc.* 2015;63(1):188-190. doi:10.1111/jgs.13226
- 393 37. Amici JM, Rogues AM, Lasheras A, Gachie JP, Guillot P, Beylot C, et al. A prospective study of
 394 the incidence of complications associated with dermatological surgery. *Br J Dermatol.*
 395 2005;153(5):967-971. doi:10.1111/j.1365-2133.2005.06861.x

38. Smith JA, Virk S, Palme CE, Low T-H (Hubert), Ch'ng S, Gupta R, et al. Age is not a predictor of prognosis in metastatic cutaneous squamous cell carcinoma of the head and neck. *ANZ J Surg.* 2018;88(4):E273-E277. doi:10.1111/ans.13757
39. Linos E, Parvataneni R, Stuart SE, Boscardin WJ, Landefeld CS, Chren M-M. Treatment of Nonfatal Conditions at the End of Life: Nonmelanoma Skin CancerNonfatal Conditions at the End of Life. *JAMA Intern Med.* 2013;173(11):1006-1012. doi:10.1001/jamainternmed.2013.639
40. Rogers EM, Connolly KL, Nehal KS, Dusza SW, Rossi AM, Lee E, et al. Comorbidity scores associated with limited life expectancy in the very elderly with nonmelanoma skin cancer. *J Am Acad Dermatol.* 2017;78(6):1119-1124. doi:10.1016/j.jaad.2017.12.048
41. Bordeaux JS, Martires KJ, Goldberg D, Pattee SF, Fu P, Maloney ME. Prospective evaluation of dermatologic surgery complications including patients on multiple antiplatelet and anticoagulant medications. *J Am Acad Dermatol.* 2011;65(3):576-583. doi:https://doi.org/10.1016/j.jaad.2011.02.012
42. Arguello-Guerra L, Vargas-Chandomid E, Diaz-Gonzalez JM, Mendez-Flores S, Ruelas-Villavicencio A, Dominguez-Cherit J. Incidence of complications in dermatological surgery of melanoma and non-melanoma skin cancer in patients with multiple comorbidity and/or antiplatelet-anticoagulants. Five year experience in our Hospital. *Cir Cir.* 2019;86(1):15-23. doi:10.24875/CIRUE.M18000003
43. Dalal AJ, Ingham J, Collard B, Merrick G. Review of outcomes of 500 consecutive cases of non-melanoma skin cancer of the head and neck managed in an oral and maxillofacial surgical unit in a District General Hospital. *Br J Oral Maxillofac Surg.* 2018;56(9):805-809. doi:https://doi.org/10.1016/j.bjoms.2018.08.015
44. Ferrier MB, Spuesens EB, Le Cessie S, de Jong RJ. Comorbidity as a Major Risk Factor for Mortality and Complications in Head and Neck Surgery. *Arch Otolaryngol Neck Surg.* 2005;131(1):27-32. doi:10.1001/archotol.131.1.27

- 422 45. Bras L, Peters TTA, Wedman J, Plaat BEC, Witjes MJH, Leeuwen BL, et al. Predictive value of
 423 the Groningen Frailty Indicator for treatment outcomes in elderly patients after head and
 424 neck, or skin cancer surgery in a retrospective cohort. *Clin Otolaryngol*. 40(5):474-482.
 425 doi:10.1111/coa.12409
- 426 46. Schwam ZG, Sosa JA, Roman S, Judson BL. Complications and mortality following surgery for
 427 oral cavity cancer: analysis of 408 cases. *Laryngoscope*. 2015;125(8):1869-1873.
 428 doi:10.1002/lary.25328
- 429 47. Valdatta L, Perletti G, Maggiulli F, Tamborini F, Pellegatta I, Cherubino M. FRAIL scale as a
 430 predictor of complications and mortality in older patients undergoing reconstructive surgery
 431 for non-melanoma skin cancer. *Oncol Lett*. 2019;17(1):263-269. doi:10.3892/ol.2018.9568
- 432 48. Hamaker ME, Jonker JM, de Rooij SE, Vos AG, Smorenburg CH, van Munster BC. Frailty
 433 screening methods for predicting outcome of a comprehensive geriatric assessment in elderly
 434 patients with cancer: a systematic review. *Lancet Oncol*. 2012;13(10):e437-e444.
 435 doi:https://doi.org/10.1016/S1470-2045(12)70259-0
- 436 49. Kaibori M, Ishizaki M, Matsui K, Iida H, Inoue K, Nagashima F, et al. Geriatric assessment as a
 437 predictor of postoperative complications in elderly patients with hepatocellular carcinoma.
 438 *Langenbeck's Arch Surg*. 2016;401(2):205-214. doi:10.1007/s00423-016-1388-1
- 439 50. Souwer ETD, Verweij NM, van den Bos F, Bastiaannet E, Slangen RME, Steup WH, et al. Risk
 440 stratification for surgical outcomes in older colorectal cancer patients using ISAR-HP and G8
 441 screening tools. *J Geriatr Oncol*. 2018;9(2):110-114.
 442 doi:https://doi.org/10.1016/j.jgo.2017.09.003
- 443 51. Pottel L, Boterberg T, Pottel H, Goethals L, Van Den Noortgate N, Duprez F, et al.
 444 Determination of an adequate screening tool for identification of vulnerable elderly head and
 445 neck cancer patients treated with radio(chemo)therapy. *J Geriatr Oncol*. 2012;3(1):24-32.
 446 doi:10.1016/j.jgo.2011.11.006
- 447 52. Stratton RJ, Hackston A, Longmore D, Dixon R, Price S, Stroud M, et al. Malnutrition in hospital

- outpatients and inpatients: prevalence, concurrent validity and ease of use of the
'malnutrition universal screening tool' ('MUST') for adults. *Br J Nutr.* 2004;92(5):799–808.
doi:10.1079/BJN20041258
53. Stratton RJ, King CL, Stroud MA, Jackson AA, Elia M. 'Malnutrition Universal Screening Tool'
predicts mortality and length of hospital stay in acutely ill elderly. *Br J Nutr.* 2006;95(2):325–
330. doi:10.1079/BJN20051622
54. Henderson S, Moore N, Lee E, Witham MD. Do the malnutrition universal screening tool
(MUST) and Birmingham nutrition risk (BNR) score predict mortality in older hospitalised
patients? *BMC Geriatr.* 2008;8(1):26. doi:10.1186/1471-2318-8-26
55. van der Kroft G, Janssen-Heijnen MLG, van Berlo CLH, Konsten JLM. Evaluation of nutritional
status as an independent predictor of post-operative complications and morbidity after
gastro-intestinal surgery. *Clin Nutr ESPEN.* 2015;10(4):e129-e133.
doi:https://doi.org/10.1016/j.clnesp.2015.05.005
56. Winter JE, Wattanapenpaiboon N, MacInnis RJ, Nowson CA. BMI and all-cause mortality in
older adults: a meta-analysis. *Am J Clin Nutr.* 2014;99(4):875-890.
doi:10.3945/ajcn.113.068122

465 **Figure 1**

466

467 **Figure 1.** Flowchart diagram representing the inclusion of patients into the final cohort of 151 patients who
468 were surgically treated for cutaneous head and neck malignancies. * Cohorts showed no significant differences
469 in age and sex throughout exclusion process.

470 **Table 1**

Variable	Value
	n=151
Age	
Mean $\pm$ SD, y	78.9 $\pm$ 9.0
Median (range), y	78.9 (46.6-96.7)
Categories	
< 70	27 (17.9%)
70-80	55 (36.4%)
80-90	53 (35.1%)
$\geq$ 90	15 (10.6%)
Sex	
Male	111 (73.5%)
Female	40 (26.5%)
Reason for referral	
Primary tumor	75 (49.7%)
Residual tumor	45 (29.8%)
Recurrent tumor	31 (20.5%)
Primary tumor location	
Frontal	9 (6.0%)
Scalp	33 (21.9%)
Temporal	10 (6.6%)
Ear	56 (37.1%)
Cheek	9 (6.0%)
Peri-orbital	7 (4.6%)
Nose	21 (13.9%)
Peri-oral	3 (2.0%)
Neck	3 (2.0%)
Histopathology	
Basal cell carcinoma	28 (18.5%)
Squamous cell carcinoma	90 (59.6%)
Malignant melanoma	17 (11.3%)
Merkel cell carcinoma	9 (6.0%)
Other ^a	7 (4.6%)
Stage of disease	
Stage I	59 (39.1%)
Stage II	53 (35.1%)
Stage III	25 (16.6%)
Stage IV	14 (9.3%)
Immunocompromised ^b	
No	130 (86.1%)
Yes	21 (13.9%)

510

511 **Table 1.** Characteristics of surgically treated patients with cutaneous malignancies of the head and neck area,
512 seen in a tertiary referral head and neck oncology center. ^a Included malignancies were angiosarcoma, atypical
513 fibroxanthoma, malignant adnexal tumor, pleomorphic dermal sarcoma, dermatofibrosarcoma protuberans
514 and adenoid cystic carcinoma. ^b Immunosuppression included patients who have been using long-term
515 immunosuppressive medication e.g. post transplantation, chronic lymphocytic leukemia, Non-Hodgkin's
516 lymphoma, severe rheumatism and Crohn's disease.

517 **Table 2**

Clavien-Dindo	Value	518
	n=151	510
No complications	89 (58.9%)	
Grade I	22 (14.6%)	
Grade II	25 (16.6%)	
Grade III	13 (8.6%)	
Grade IV	2 (1.3%)	
Grade V	0 (0.0%)	

523 **Table 2.** Postoperative complications in patients undergoing surgery for cutaneous head and neck
524 malignancies.

525 **Table 3**

Variable	Value (%) n=151	Univariable analysis Odds ratio (95% CI)	p-value
Patient characteristics			
Age			
Mean $\pm$ SD, y	78.9 $\pm$ 9.0	0.98 (0.94-1.02)	0.27
Median (range), y	78.9 (46.6-96.7)		
Sex			
Male	111 (73.5%)	1	
Female	40 (26.5%)	0.90 (0.39-2.06)	0.80
Immunocompromised ^a			
No	130 (86.1%)	1	
Yes	21 (13.9%)	1.89 (0.72-4.96)	0.20
Tumor characteristics			
Reason for referral			
Primary tumor	75 (49.7%)	1	0.71
Residual tumor	45 (29.8%)	0.95 (0.41-2.25)	0.91
Recurrent tumor	31 (20.5%)	1.40 (0.56-3.51)	0.47
Stage			
Stage I	59 (39.1%)	1	< 0.05
Stage II	53 (35.1%)	1.93 (0.78-4.78)	0.15
Stage III	25 (16.6%)	1.91 (0.63-5.76)	0.25
Stage IV	14 (9.3%)	6.53 (1.86-22.99)	< 0.01
Tumor diameter			
< 20 mm	72 (59.5%)	1	< 0.05
20-40 mm	36 (29.8%)	2.57 (1.04-6.36)	< 0.05
$\geq$ 40mm	13 (10.7%)	3.89 (1.12-13.51)	< 0.05
Invasion depth			
Mean $\pm$ SD, mm	5.2 $\pm$ 3.3	1.13 (0.99-1.29)	0.06
Median (range), mm	4.7 (0.3-19.5)		
Histopathology			
Basal cell carcinoma	28 (18.5%)	1	0.23
Squamous cell carcinoma	90 (59.6%)	3.96 (1.11-14.20)	< 0.05
Malignant melanoma	17 (11.3%)	3.47 (0.71-16.99)	0.13
Merkel cell carcinoma	9 (6.0%)	1.04 (0.10-11.47)	0.97
Other ^b	7 (4.6%)	3.33 (0.44-25.39)	0.25
Treatment characteristics			
Primary treatment			
Local surgery	113 (74.8%)	1	
Locoregional surgery	38 (25.2%)	4.38 (1.98-9.68)	< 0.01
Treatment intensity ^c			
Minor	96 (63.6%)	1	
Major	55 (36.4%)	3.46 (1.62-7.39)	< 0.01
Anesthesia			
Local anesthesia	34 (22.5%)	1	
General anesthesia	117 (77.5%)	7.70 (1.75-33.81)	< 0.01
Reconstructive surgery			
No reconstructive surgery	45 (29.8%)	1	0.10
Intraoperative reconstruction	81 (53.6%)	1.07 (0.45-2.56)	0.88
Subsequent reconstructive surgery	25 (16.6%)	2.75 (0.96-7.92)	0.06
Intoxications			
Smoking			
Never or former	113 (86.3%)	1	
Current	18 (13.7%)	2.03 (0.72-5.74)	0.18
Drinking			
None or mild	117 (88.6%)	1	
Heavy (> 2/day)	15 (11.4%)	2.78 (0.93-8.35)	0.07
Comorbidities			
ACE-27			
None or mild	53 (35.1%)	1	
Moderate or severe	98 (64.9%)	1.61 (0.73-3.55)	0.24
Polypharmacy			
Medication count			

< 5 medications	95 (65.1%)	1	
≥ 5 medications	51 (34.9%)	2.36 (1.11-5.07)	< 0.05
Nutritional status			
MUST			
Low risk	128 (92.1%)	1	
Medium to high risk	11 (7.9%)	3.46 (0.99-12.07)	0.05
Functional status			
ADL			
No restrictions (< 1)	114 (82.6%)	1	
Restrictions (≥ 1)	24 (17.4%)	1.69 (0.65-4.39)	0.28
IADL			
No restrictions (< 1)	100 (69.4%)	1	
Restrictions (≥ 1)	44 (30.6%)	1.07 (0.48-2.38)	0.87
TUG			
Mean ± SD, s	11.4 ± 6.7	1.04 (0.98-1.11)	0.19
Median (range), s	10 (5-70)		
History of falls			
No	124 (91.2%)	1	
Yes	12 (8.8%)	0.96 (0.24-3.76)	0.95
Social support			
Education			
Low level of education	60 (48.8%)	1	0.64
Middle level of education	38 (30.9%)	1.52 (0.61-3.76)	0.37
High level of education	25 (20.3%)	1.04 (0.35-3.10)	0.95
Marital status			
In a relationship	89 (67.9%)	1	0.69
Widow	32 (24.4%)	1.38 (0.60-3.37)	0.47
Single	10 (7.6%)	0.76 (0.15-3.86)	0.74
Social Economic Statusscore (SES)			
Below average (NL)	119 (79.3%)	1	
Above average (NL)	31 (20.7%)	0.99 (0.40-2.44)	0.98
Cognitive status			
MMSE			
Normal cognition (> 24)	108 (76.6%)	1	
Declined cognition (≤ 24)	33 (23.4%)	0.83 (0.34-2.05)	0.69
Risk of delirium			
No	113 (77.4%)	1	
Yes	33 (22.6%)	0.85 (0.35-2.08)	0.72
Psychological status			
GDS-15			
No depression (< 6)	113 (81.3%)	1	
Depression (≥ 6)	26 (18.7%)	1.17 (0.45-3.09)	0.75
Frailty screeners			
G8			
Non-frail (> 14)	39 (26.7%)	1	
Frail (≤ 14)	107 (73.3%)	5.83 (1.68-20.26)	< 0.01
GFI			
Non-frail (< 4)	98 (70.5%)	1	
Frail (≥ 4)	41 (29.5%)	1.43 (0.63-3.26)	0.40

527 **Table 3.** Patient-, tumor- and treatment characteristics and domains of geriatric assessment in a univariable
 528 logistic regression predicting postoperative complications grade II and higher. Abbreviations: CI=Confidence
 529 Interval, SD=Standard Deviation, ACE-27=Adult Comorbidity Evaluation 27, MUST=Malnutrition Universal
 530 Screening Tool, ADL=Activities of Daily Living, IADL=Instrumental Activities of Daily Living, TUG=Timed Up and
 531 Go, NL=Netherlands, MMSE=Mini Mental State Examination, GDS-15=Geriatric Depression Scale 15,
 532 G8=Geriatric 8, GFI=Groningen Frailty Indicator. ^a Immunosuppression included patients who have been using
 533 long-term immunosuppressive medication e.g. post transplantation, chronic lymphocytic leukemia, Non-
 534 Hodgkin's lymphoma, severe rheumatism and Crohn's disease. ^b Included malignancies were angiosarcoma,
 535 atypical fibroxanthoma, malignant adnexal tumor, pleomorphic dermal sarcoma, dermatofibrosarcoma
 536 protuberans and adenoid cystic carcinoma. ^c Defined as surgery > 120 minutes or three or more stages of Mohs
 537 micrographic surgery.

538 **Table 4**

Variable	← No complications	Complications →	Multivariable model ^a	
			Odds ratio (95% CI)	p-value
Treatment intensity ^b			1	
Minor			2.73 (1.19-6.26)	< 0.05
Major				
Anesthesia			1	
Local anesthesia			4.74 (1.02-22.17)	< 0.05
General anesthesia				
Frailty on G8			1	
Non-frail (> 14)			6.34 (1.73-23.25)	< 0.01
Frail (≤ 14)				

539

540 **Table 4.** Multivariable logistic regression model predicting postoperative complications grade II and higher
541 patients receiving in surgery for cutaneous head and neck malignancies. ^a Adjusted for age and sex. ^b Defined as
542 surgery > 120 minutes or three or more stages of Mohs micrographic surgery.