University of Groningen

Speed related changes in muscle activity from normal to very slow walking speeds

den Otter, A.R.; Mulder, T.; Duysens, J.

Published in:
Gait & Posture

DOI:
[10.1016/S0966-6362\(03\)00071-7](https://doi.org/10.1016/S0966-6362(03)00071-7)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2004

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

den Otter, A. R., Mulder, T., & Duysens, J. (2004). Speed related changes in muscle activity from normal to very slow walking speeds. *Gait & Posture*, 19(3), 270 - 278. [https://doi.org/10.1016/S0966-6362\(03\)00071-7](https://doi.org/10.1016/S0966-6362(03)00071-7)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Speed related changes in muscle activity from normal to very slow walking speeds

A.R. den Otter^{a,*}, A.C.H. Geurts^a, T. Mulder^b, J. Duysens^{a,c}

^a SMK-Research, Sint Maartenskliniek, P.O. Box 9011, 6500 GM Nijmegen, The Netherlands

^b Institute of Human Movement Science, University of Groningen, Groningen, The Netherlands

^c Department of Biophysics, University of Nijmegen, Nijmegen, The Netherlands

Received 24 December 2002; received in revised form 23 March 2003; accepted 15 May 2003

Abstract

The study of neuromuscular activity at very slow walking speeds may lead to a better understanding of the mechanisms underlying speed regulation during walking, and may aid the interpretation of gait data in patients who walk slowly. Extreme reductions in walking speed will cause changes in locomotor task demands that may eventually result in modifications of the patterning of muscle activity that underlies walking. The aim of the present study was to investigate patterns of lower limb muscle activity during very slow walking ($< 0.28 \text{ m s}^{-1}$), and to study the neuromuscular gain functions that reflect the phase dependent effects of walking speed on electromyographic (EMG) amplitude. Nine healthy young adults walked at seven different walking speeds (1.39, 0.83, 0.28, 0.22, 0.17, 0.11, and 0.06 m s^{-1}) while EMG was recorded from eight lower extremity muscles. Results showed that the phasing of muscle activity remained relatively stable over walking speeds despite substantial changes in its amplitude. However, between 1.39 and 0.28 m s^{-1} , epochs of Rectus femoris, Biceps femoris and Tibialis anterior activities were found that were typical for specific speed ranges. When walking speed decreased further to almost standing still (0.06 m s^{-1}), negative gain values were found in Peroneus longus during midstance and Rectus femoris in late swing, indicating the emergence of new bursts of activity with decreasing walking speed. It is proposed that these extra activities may be attributed to increased demands on postural stability, and the altered dynamics of the swinging limb at very slow speeds.

© 2003 Elsevier B.V. All rights reserved.

Keywords: Gait; Electromyographic; Walking speed

1. Introduction

During walking, the ability to adjust the speed of progression is an important mechanism that adapts locomotor activity to changes in environmental demands, e.g. to accommodate time pressure or to enhance safety. In general, changes in walking speed require the adjustment of a neuromuscular gain factor, whereas the global timing characteristics of the muscle activity patterns are essentially preserved [1–6]. This strategy for speed regulation, through which a phase specific gain factor modulates relatively invariant basis patterns, simplifies control, and enhances computational efficiency.

The main role of the muscles in the regulation of walking speed is to control the accelerating and decelerating forces

of individual body segments to establish safe forward progression [5]. As a result, the amplitude of muscle activity increases with walking speed because of the need for larger muscular force output. This normal positive relationship may be challenged if walking speed is markedly reduced, due to changes in the underlying locomotor task demands. Extreme reductions in walking speed will prolong substantially the time spent in double support, and one may expect a switch from locomotor to merely postural muscular synergies. Also, the larger horizontal excursions of the centre of mass associated with slow walking may necessitate more explicit muscular efforts to maintain frontal plane balance during walking [7]. This presumed relationship between speed reduction and dynamic instability is substantiated further by recent evidence showing that a decrease in walking speed may cause an increase in potentially destabilising vestibulospinal drive [8], probably due to diminished proprioceptive input [9,10]. Strong reductions in walking speed may also affect the neuromuscular control of the swinging limb. While

* Corresponding author. Tel.: +31-24-365-9367;
fax: +31-24-365-9154.

E-mail address: r.otter@smk-research.nl (A.R. den Otter).

at comfortable and fast speeds, the swinging leg describes a largely ballistic trajectory and moves under passive, gravitational control [11], strongly reduced walking speeds may necessitate a more active mode of control to counteract gravity and guarantee sufficient ‘air time’ for the swinging leg.

Alterations in swing phase control, and the increased emphasis on dynamic balance, may induce modifications in the patterning of lower limb muscle activity that are uniquely related to these very low speed ranges. Within normal ranges of walking speed ($0.75\text{--}1.75\text{ m s}^{-1}$), speed effects on electromyographic (EMG) patterns can be described quite accurately by superimposing a few speed related gain functions on a small number of speed independent basis patterns [6]. It follows that speed induced changes in the timing of muscle activity are established primarily through modulation by muscle specific gain functions of a relatively invariant locomotor rhythm [12]. Given the associated changes in locomotor task demands described above, one may wonder how these same principles are extrapolated to the lowest speed ranges.

The primary aim of this study was to investigate changes in speed related neuromuscular gain, when walking speed was systematically varied from normal speeds to almost standing still. While the lower boundaries of the walking speed continuum are not usually encountered during normal human walking, they still may provide information on the strategies used by the neuromuscular system to induce speed modifications. The present study does not provide a database for systematic comparison of patient data with standardised values, yet the results from the present study may still be of clinical interest. Since reduced walking speed is a key feature of pathological gait [13], clinical gait analysis as well as gait research on pathological populations may profit from the assessment of neuromuscular patterns associated with very slow walking. More specifically, studies on very slow walking may help dissociate between those aberrations that are directly related to pathological changes in neuromuscular coordination, and those that reflect adaptations to speed related changes in locomotor task demands.

2. Methods

2.1. Subjects

Nine healthy volunteers (five females, four males; mean (S.D.) age 22.4 (2.35) years; height 1.81 (6.7) m; body mass 66.6 (7.84) kg) participated. None of the subjects suffered from any neurological or orthopaedic disorder that might interfere with the goals of the study.

2.2. Procedure

Subjects walked on a motor driven treadmill ($2.0 \times 0.7\text{ m}$) at seven different walking speeds (1.39, 0.83, 0.28, 0.22, 0.17, 0.11, and 0.06 m s^{-1}). Two 40-s registrations were

completed at each walking speed in two quasi-randomised series of seven trials. The order of each series was the same for all subjects. Subjects were allowed to walk on the treadmill at a self-selected speed to become accustomed to walking on the treadmill. During the experiment, no particular instructions were given with respect to stride length or cadence.

2.3. Data recording

EMG recordings were made using disposable surface electrodes (MediTrace ECG 1801 Pellet, (Ag/AgCl)) with a diameter of 10–12 mm and a minimum inter-electrode distance of 24 mm (Graphics Controls, Buffalo, NY). The electrodes were placed on the skin to measure activity from eight lower extremity muscles: Biceps femoris (BF), Semitendinosus (ST), Rectus femoris (RF), Vastus medialis (VM), Gastrocnemius medialis (MG), Soleus (SO), Peroneus longus (PL) and Tibialis anterior (TA). Electrodes were placed according to SENIAM conventions [14]. Simultaneously, footswitch data of the heel and toe were collected using custom made footswitches. Signals were fed into a K-lab SPA 20/8 pre-amplifier with a common mode rejection ratio $> 95\text{ dB}$ and a noise level of $< 1\text{ }\mu\text{V rms}$. Incoming signals were high-pass filtered using a third order Butterworth filter (-3 dB point at 20 Hz), and low-pass filtered using a second order Butterworth filter (-3 dB point at 500 Hz). The incoming EMG signals were monitored on-line, to ensure sufficient quality of the data. In case of obviously distorted signals, e.g. due to movement artefacts or poor skin electrode contact, the necessary measures were taken, e.g. renewal of electrodes; skin abrasion, to rectify the problem. The EMG signals as well as the footswitch data were digitised at 2400 Hz and stored on a computer hard disk for offline processing.

2.4. Data processing

Footswitch data were used to determine initial contact (IC) and toe off (TO) of each recorded step. EMG signals were full-wave rectified and low-pass filtered at 25 Hz using custom MATLAB[®] software. After exclusion of steps that contained clearly distorted signals, the EMG data were time normalised with the gait cycle taken as 100%. Earlier studies have shown that walking speed not only affects step length and step duration but also the relative durations of the stance and swing phase [3,4]. Therefore, to justify a point to point comparison between EMG profiles found at different walking speeds, a second normalisation procedure was applied in which the stance and swing phases were normalised separately, to 150 and 100 data points, respectively.

For both types of time-normalised data, the individual strides were averaged for each subject to obtain an ensemble average for every subject at each walking speed. To obtain an accurate estimate of the individual ensemble av-

erages, as many strides as were available for each condition were used to calculate the average profiles (mean number of strides ranged from 15.5 (S.D. = 8.7) at 0.06 m s^{-1} , to 73.5 (S.D. = 4.0) at 1.39 m s^{-1}). For each muscle, these individual averages were then amplitude normalised with the peak amplitude at the fastest walking condition (1.39 m s^{-1}) set to 100%. Finally, these data were averaged over all nine subjects, to obtain a group mean envelope for each of the eight muscles for each of the seven walking speeds.

2.5. Data analysis

To estimate the gain functions that describe the speed related changes in EMG amplitude over the time normalised gait cycle, the walking speeds of the individual subjects were first normalised to body height [6] as follows:

$$\hat{v}_{ni} = \frac{v_n}{\sqrt{g l_{0i}}} \quad (1)$$

where $\hat{v}_{ni}$ —normalised speed of subject i for walking speed condition n ; v_n —treadmill speed for walking speed condition n ; g —acceleration of gravity ($= 9.81 \text{ m s}^{-2}$ on earth); l_{0i} —body height of subject i .

Next, the slope of the presumed linear relationship between walking speed and EMG values was calculated, as

follows:

$$b = \frac{\left(\sum_{i=1}^9 \frac{x_{ijb} - x_{ija}}{9} \right)}{v_b - v_a} \quad \text{for } j = 1 \dots 250 \quad (2)$$

where $b_{b,a}$ = rate of change between walking speed conditions a and b in normalised EMG amplitude per unit change in walking speed ($= \text{m s}^{-1}$); x_{ijb} , x_{ija} = mean EMG value of subject i (1–9) at instant j (1–250) at speed b/a ; $v_b - v_a$ = difference between walking speeds v_b and v_a (m s^{-1}).

This is identical to the slope value found by regressing the amplitude normalised EMG-values on walking speed using a simple least squares criterion (see Fig. 1). This type of analysis was used to compare differences in EMG amplitude between the following pairs of walking speed conditions: (a) 1.39 m s^{-1} vs. 0.83 m s^{-1} , (b) 0.83 m s^{-1} vs. 0.28 m s^{-1} , and (c) 0.28 m s^{-1} vs. 0.06 m s^{-1} .

Regression of the amplitude normalised EMG values (1–250) on walking speed ($1.39, 0.83, 0.28$, and 0.06 m s^{-1}) yielded slope coefficients for each point in the normalised gait cycle. These coefficients quantify the linear dependency of EMG amplitude on walking speed, and thus reflect the gain applied by the neuromuscular system to establish changes in walking speed. This approach is conceptually similar to that used by Hof et al. [6] for the modeling of

Fig. 1. Example of the calculation of the gain factor 'B' for one point on the normalised gait cycle of the RF ($j = 150$) between two walking speed conditions (treadmill speeds: 0.83 m s^{-1} vs. 1.39 m s^{-1}). Depicted are the amplitude normalised EMG values for all nine subjects at both speeds, and the regression line with slope $B = 312.08$.

speed effects on EMG (see also Ref. [15]). In the present study, this approach was chosen to compare speed effects between different pairs of walking speed conditions. It is important to note that this type of analysis reflects both the magnitude of the speed effect as well as its direction: positive slope values indicate an increase in EMG amplitude with increasing walking speed, whereas a negative value indicates an increase in EMG amplitude with decreasing walking speed (i.e. a negative linear relationship).

An important assumption underlying this type of analysis is that all points of the gait cycle at different walking

speeds are appropriately aligned and share a similar time base. Because the relative durations of swing and stance phases are known to vary with walking speed [3], a simple point-to-point comparison between walking speeds based on the normalised step cycle would be inappropriate. For instance, activity found in the late stance phase at lower speeds would be compared to early swing activity at higher speeds if conventional step cycle normalisation would be applied. Therefore, differences in stance–swing distribution were corrected by normalising stance and swing phase separately to 150 and 100 point, respectively.

Fig. 2. Group ensemble average profiles for all eight muscles at all seven speeds employed in the experiment. Profiles are shown in the order of speed, with the treadmill speeds given on the right (m s^{-1}). (Abbreviations: BF, Biceps femoris; ST, Semitendinosus; RF, Rectus femoris; VM, Vastus medialis; MG, Medial gastrocnemius; SO, Soleus; PL, Peroneus longus and TA, Tibialis anterior.)

3. Results

3.1. Effects of walking speed on the group ensemble average

After normalising the step cycle to 100%, the group average profiles were calculated for every muscle at each of the seven walking speeds. In general, the amplitude of muscle activity decreased with decreasing walking speed. When the peak amplitude of the group ensemble average was taken as 100%, peak activity at 0.06 m s^{-1} was found to be decreased to the following percentages: BF = 16.7%, ST = 17.6%, RF = 38.2%, VM = 18.0%, MG = 39.9%, SO = 24.4%, PL = 44.0%, and TA = 31.8%.

For a few instances, EMG bursts were found to be very velocity-specific. In five out of nine subjects, an extra burst of muscle activity was found in the BF at speeds of $\leq 0.28 \text{ m s}^{-1}$. This burst typically occurred around the transition between stance and swing phase, and varied considerably in amplitude between the five subjects. However, in all five cases, this burst occurred selectively at speeds $\leq 0.28 \text{ m s}^{-1}$, was absent at the two fastest speeds, and tended to decrease in amplitude when speed became slower than 0.28 m s^{-1} . Although similar episodes of activity were apparent in ST as well, these bursts were considerably smaller in amplitude than those found in BF. A typical example of this extra burst in the BF, taken from a single subject can be seen in Fig. 3. Note that for this particular subject the amplitude of this extra burst in the BF at 0.28 m s^{-1} exceeds the peak amplitude of this muscle at 1.39 m s^{-1} by 22%. Although a similar phenomenon can be seen in ST, the amplitude of this burst is much smaller than that in the BF and reached only 28% of peak ST activity at 1.39 m s^{-1} , for this subject. Inspection of Fig. 2 further suggests that a large burst emerges in RF during late stance/early swing phase at 1.39 m s^{-1} . Although this burst was never absent, it was present only in a strongly reduced form at speeds $\leq 0.83 \text{ m s}^{-1}$. This speed dependent burst was found in all of our nine subjects. Note that a similar epoch of speed dependent activity is not apparent in VM.

During normal human walking, TA shows two distinct bursts of activity during the course of the gait cycle. The group ensemble averages for this muscle show that its second burst (late swing–early stance phase) is more strongly affected by walking speed than is its first burst (early swing). At very slow walking speeds ($\leq 0.28 \text{ m s}^{-1}$), the second burst is reduced strongly in amplitude so that the overall profile at these speeds has a rather uniphasic appearance.

Although the amplitude of SO and MG activity is strongly reduced at slow speeds, the ensemble average profiles at these speeds still show a clear resemblance with the typical profiles found at normal speeds. In PL, a local change in the shape of its burst becomes apparent at very slow speeds, due to a shift in the peak of activity from late stance to midstance.

3.2. Changes in the neuromuscular gain functions

Figs. 4 and 5 depict the estimated gain functions for all eight muscles, as they were found for three specific comparisons between walking speeds (1.39 m s^{-1} vs. 0.83 m s^{-1} ; 0.83 m s^{-1} vs. 0.28 m s^{-1} ; 0.28 m s^{-1} vs. 0.06 m s^{-1}). Between 1.39 and 0.28 m s^{-1} , gain values are generally positive, indicating a decrease in EMG amplitude with decreasing walking speed. In BF, a short phase with negative slope values was found at late stance/early swing between 0.83 and 0.28 m s^{-1} , whereas between 0.28 and 0.06 m s^{-1} , large positive gain values were found for this phase of the gait cycle. This suggests that, during this particular phase, BF activity increases when walking speed approaches 0.28 m s^{-1} and then again decreases when walking is slowed down further to speeds slower than 0.28 m s^{-1} . As becomes apparent from Fig. 4, a similar relationship between walking speed and EMG amplitude was not found for ST. As could be expected, in RF an episode of large positive gain values can be found in late stance/early swing, indicating that, for speeds between 0.83 and 1.39 m s^{-1} , EMG amplitude is strongly dependent on walking speed for this phase of the gait cycle (see Fig. 4). Note that this epoch of positive slope values is absent for comparisons between the other speeds. Another salient feature of the RF gain func-

Fig. 3. Left panel: Average EMG profiles of BF for one subject. Right panel: Average EMG profiles of ST for the same subject. Note the extra burst in BF that occurs in late stance/early swing phase at slower walking speeds ($\leq 0.28 \text{ m s}^{-1}$). Also note the much smaller amplitude of a similar burst in ST.

Fig. 4. Estimated gain functions for BF, ST, RF and VM for three separate comparisons of walking speeds: Top = 1.39 m s^{-1} vs. 0.83 m s^{-1} ; Middle = 0.83 m s^{-1} vs. 0.28 m s^{-1} ; Bottom = 0.28 m s^{-1} vs. 0.06 m s^{-1} ; Positive slope values indicate a positive linear relationship between walking speed and EMG amplitude (i.e. a decrease in EMG amplitude with decreasing walking speed), whereas negative slope values indicate a negative linear relationship between walking speed and EMG amplitude (i.e. an increase in EMG amplitude with decreasing walking speed). Vertical lines indicate the onset of the swing phase. Note that the gain functions are offset to enhance clarity.

Fig. 5. Estimated gain functions for MG, SO, PL and TA for three separate comparisons of walking speeds: Top = 1.39 m s^{-1} vs. 0.83 m s^{-1} ; Middle = 0.83 m s^{-1} vs. 0.28 m s^{-1} ; Bottom = 0.28 m s^{-1} vs. 0.06 m s^{-1} ; For further explanation, see Fig. 4.

tions is the episode of negative values that is found during late swing between 0.06 and 0.28 m s⁻¹. Apparently, the amplitude of muscle activity increased for this period of the gait cycle when walking speed became increasingly slow.

In PL, a short period with negative gain values can be seen during midstance, reflecting increased activity for this phase at very slow speeds (see Fig. 5). For SO and MG, the gain functions remain relatively constant over all three speed comparisons, indicating that the activity of these muscles continues to be sensitive to speed changes, even at the slowest walking speeds. Note that, within the lowest speed ranges, the PL slope function deviates rather clearly from those found for MG and SO, despite the relative similarity of the slope functions of MG, SO and PL between 0.28 and 1.39 m s⁻¹.

4. Discussion

The results of the present study show that, in general, the amplitude of lower extremity muscle activity tends to increase with the speed of progression. In RF, TA, PL and BF, changes in the pattern of activity were found that could be uniquely related to specific speed ranges. For most muscles, the gain functions that regulate the speed related adaptations in muscle activity appeared to be relatively stable down to speeds as slow as 0.28 m s⁻¹. When speed was decreased to below 0.28 m s⁻¹, negative gain values could be detected in some muscles (PL and RF) that are indicative of an increase in activity with decreasing speed.

It was decided to conduct the experiment on a treadmill rather than on a walkway to exert maximum experimental control over the extremely low walking speeds employed in this study. When walking on a solid surface, extremely slow speeds may be realised by sequencing of more or less 'static' postures. In contrast, on the treadmill, both cadence and step length are imposed on the subject by the moving walking surface. This may have affected our results to some extent but the differences are likely to be very small. Indeed, the limited number of EMG studies that have compared treadmill walking and floor walking failed to show clear differences in the underlying patterns of muscle activity, other than minor changes in the amplitude of muscle activity [16,17]. While these studies did not extend to the very low speed range it is unlikely that major differences would appear.

4.1. Lower leg muscles

In the calf muscles, it was found that the activity during late stance (50–80% of the stance phase) was strongly affected by walking speed. The large positive gain factor values indicate that the amplitude of MG and SO activity during this phase of the gait cycle is strongly dependent on walking speed, a result that is in agreement with earlier work [18]. In contrast, MG and SO activity during 30–50% of the stance phase appears to be relatively stable over walking

speeds. The latter finding is in accordance with an earlier study by Winter [19] who found that the amount of energy absorption by the ankle plantarflexors during weight acceptance was relatively insensitive to differences in walking speed. Furthermore, a recent simulation study showed that MG and SO deliver nearly all of the positive work during late to terminal stance (40–60% of the gait cycle) and that the contribution of these two muscles to forward progression is larger than that of all other muscles taken together [20]. The importance of this mechanism is underscored by the finding that late stance MG and SO activity continues to be affected by walking speed, even within the lowest speed range (0.28–0.06 m s⁻¹). Apparently, even when walking speed is decreased to almost standing still, the plantar flexors still contribute to the production of forward motion. The present result of a continuous scaling of EMG during late stance also demonstrates that detection of leg loading in this phase through calf muscle tendon organs could be a useful mechanism to control swing onset in humans, analogous to that seen in the cat, for the full speed scale [21].

At speeds ≥ 0.28 m s⁻¹, both the global phasing characteristics of PL, SO and MG, as well as their associated gain functions showed strong similarities. However, at speeds < 0.28 m s⁻¹, the PL gain function showed a short episode with negative values, indicating an increase in activity with a decrease in walking speed. It is believed that PL plays a role in the maintenance of frontal plane balance during walking, and especially its activity during the single support phase appears to be important in this respect. An earlier study on the function of PL during walking showed that PL activity during foot flat is sensitive to speed changes, and that its activity in this phase increases when walking speed is decreased [22]. It is known that walking speed affects the medio-lateral distribution of foot loading, and that the pattern of foot loading shifts to the lateral side of the foot when walking speed is decreased [23]. Because the lateral component of the ground reaction force as well as the amount of foot eversion are correlated to the amplitude of PL activity during midstance [24], it can be argued that the negative relationship that was found between walking speed and PL activity for speed < 0.28 m s⁻¹ serves to generate additional foot everting forces to prevent excessive foot inversion during the single support phase.

During normal human walking at regular speeds, TA activity is represented by a biphasic pattern that peaks in early swing (first burst) and in late swing-early stance (second burst). In the present study, it was found that the second burst of TA activity was more sensitive to changes in walking speed than the first burst, eventually resulting in a near absence of the second burst at the slowest speeds. While the first burst of TA activity serves to dorsiflex the foot and achieve foot clearance during early swing, the second burst represents an anticipatory response to the upcoming leg loading during early stance. Because the amount of leg loading increases with walking speed, the amplitude of the second TA burst that anticipates this loading can be expected

to be velocity-dependent. In contrast, foot clearance must be achieved regardless of walking speed, which explains why TA activity during early swing appears to be relatively insensitive to speed changes. The strong speed sensitivity of the second TA burst (end swing-early stance) may be of clinical interest since the absence or reduced amplitude of this burst has been reported in Parkinsonian and hemiparetic persons [25,26]. Because these pathologies are usually associated with strongly reduced gait speeds, the results from the present study suggest that such phenomena could at least partly be explained by a speed effect.

4.2. Upper leg muscles

Between 0.83 and 1.39 m s^{-1} , a substantial increase in RF activity was found at terminal stance–early swing. Although this burst was never absent, not even at the slowest speeds, peak activity in this phase at 0.83 m s^{-1} was only 20.8% of peak RF activity at 1.39 m s^{-1} in the same phase. This phenomenon has been described on numerous occasions in the literature [4,27–30]. This particular epoch of RF activity is assumed to generate an additional hip moment to accelerate the leg forward at higher walking speeds, and is probably related to the deceleration of the shank during initial swing. At walking speeds $\leq 0.28 \text{ m s}^{-1}$, additional RF activity was found during late swing while similar speed dependent effects were absent in VM. The independence of the activities in these two muscles is important to demonstrate that in this case RF activity did not result from vastus cross-talk. Recent evidence has indeed pointed out that end swing RF activity is sometimes due to cross-talk from the vasti [31].

The origin of the negative speed–amplitude relationship found for RF at the end of the swing phase at very slow speeds is likely to be found in its action on the hip. It can be hypothesized that, at very slow walking speeds, the pendulum-like properties of the swinging leg can no longer be utilized for forward progression throughout the full swing phase. Instead, the passive dynamics may force the leg to swing backwards and to prematurely end the swing phase. Additional RF activity may be used to establish extra hip flexion and counteract this problem.

In BF, an extra epoch of activity was found during late stance/early swing when walking speed was decreased to 0.28 m s^{-1} . When speed was lowered to $< 0.28 \text{ m s}^{-1}$, this burst gradually decreased in amplitude. Although this speed dependent component of BF activity has been described on previous occasions in the literature [4,28], it has not received much attention. The observation that this phenomenon did not occur in all of our subjects suggests that, it is possibly related to a neuromuscular strategy to adapt to exceptionally low walking speeds. Since similar bursts of activity were either absent or of considerably smaller amplitude in ST, it can be assumed that this speed dependent phase of BF activity is related to the production of a force aimed at an late stance external rotation of the lower leg as both muscles

are synergists for hip extension and knee flexion but they differ with respect to external rotation.

References

- [1] Milner M, Basmajian JV, Quanbury AO. Multifactorial analysis of walking by electromyography and computer. *Am J Phys Med* 1971;50:235–58.
- [2] Murray MP, Mollinger LA, Gardner GM, Sepic SB. Kinematic and EMG patterns during slow, free, and fast walking. *J Orthop Res* 1984;2:272–80.
- [3] Nilsson J, Thorstensson A, Halbertsma J. Changes in leg movements and muscle activity with speed of locomotion and mode of progression in humans. *Acta Physiol Scand* 1985;123:457–75.
- [4] Shiavi R, Bugle HJ, Limbird T. Electromyographic gait assessment, part 1: adult EMG profiles and walking speed. *J Rehabil Res Dev* 1987;24:13–23.
- [5] Yang JF, Winter DA. Surface EMG profiles during different walking cadences in humans. *Electroenc Clin Neurophys* 1985;60:485–91.
- [6] Hof AL, Elzinga H, Grimmius W, Halbertsma JPK. Speed dependence of averaged EMG profiles in walking. *Gait Posture* 2002;16:78–86.
- [7] Bauby CE, Kuo AD. Active control of lateral balance in human walking. *J Biomech* 2000;33:1433–40.
- [8] Brandt T, Strupp M, Benson J. You are better off running than walking with acute vestibulopathy. *Lancet* 1999;35:746.
- [9] Dietz V, Baaken B, Colombo G. Proprioceptive input overrides vestibulo-spinal drive during human locomotion. *NeuroReport* 2001;12:2743–6.
- [10] Horak FB, Hlavacka F. Somatosensory loss increases vestibulospinal sensitivity. *J Neurophysiol* 2001;86:575–85.
- [11] Mac Mahon TA. Muscles, reflexes and locomotion. Princeton: Princeton University Press, 1984.
- [12] Duysens J, van de Crommert H. Neural control of locomotion Part 1: the central pattern generator from cats to humans. *Gait Posture* 1998;7:131–41.
- [13] Van Emmerik REA, Wagenaar RC. Dynamics of movement coordination and tremor during gait in Parkinson's disease. *Hum Mov Sci* 1996;15:203–35.
- [14] Freriks B, Hermens H, Disselhorst-Klug C, Rau G. The recommendations for sensor and sensor placement procedures for surface electromyography. In: Hermens H, editor. European recommendations for surface electromyography. Enschede: Roessingh Research and Development, 1999:15–53.
- [15] Ramsay JO, Silverman BW. Functional data analysis. New York: Springer, 1997.
- [16] Murray MP, Spurr GB, Sepic SB, Gardner GM, Mollinger LA. Treadmill vs floor walking: kinematics, electromyogram, and heart rate. *J Appl Physiol* 1985;59:87–91.
- [17] Arsenault AB, Winter DA, Marteniuk RG. Treadmill versus walkway locomotion in humans: an EMG study. *Ergonomics* 1986;29:665–76.
- [18] Duysens J, Tax AA, van der Doelen B, Trippel M, Dietz V. Selective adaptation of human soleus or gastrocnemius in reflex responses during walking and running. *Exp Brain Res* 1991;87:193–204.
- [19] Winter DA. Energy generation and absorption at the ankle and knee during fast, natural and slow cadences. *Clin Orthop* 1983;175:147–54.
- [20] Neptune RR, Kautz SA, Zajac FE. Contributions of the individual ankle plantar flexors to support, forward progression and swing initiation during walking. *J Biomech* 2001;34:1387–98.
- [21] Duysens J, Clarac F, Cruse H. Load-regulating mechanisms in gait and posture: comparative aspects. *Physiol Rev* 2000;80:83–133.
- [22] Louwerens JWK, van Linge B, de Klerk LWL, Mulder PGH, Snijders CJ. Peroneus longus and tibialis anterior muscle activity in stance phase. *Acta Orthop Scand* 1995;66:517–23.

- [23] Perttunen J, Komi PV. Effects of walking speed on foot loading patterns. *J Human Move Stud* 2001;40:291–304.
- [24] Matsusaka N. Control of medial–lateral balance in walking. *Acta Orthop Scand* 1986;57:555–9.
- [25] Cioni M, Richards CL, Malouin F, Bedard PJ, Lemieux R. Characteristics of the electromyographic patterns of lower limb muscles during gait in patients with Parkinson's disease when OFF and ON L-dopa treatment. *Ital J Neurol Sci* 1997;18:195–208.
- [26] Burridge JH, Wood DE, Taylor PN, McLellan DL. Indices to describe different muscle activation patterns, identified during treadmill walking, in people with spastic drop foot. *Med Eng Phys* 2001;23:427–34.
- [27] Perry J, 1992. *Gait analysis—normal and pathological gait*. Slack Incorporated, USA, 1992.
- [28] Scott L, Ringwelsky D, Carroll N. Transfer of rectus femoris: effects of transfer site on moment arms about the knee and hip. *J Biomech* 1994;27:1201–11.
- [29] Annaswamy TM, Giddings CJ, Della Croce U, Kerrigan DC. Rectus femoris: its role in normal gait. *Arch Phys Med Rehabil* 1999;80:930–4.
- [30] Nene A, Mayagoitia R, Veltink P. Assessment of rectus femoris function during initial swing phase. *Gait Posture* 1999;9:1–9.
- [31] Nene A, Byrne C, Hermens HJ. Is rectus femoris really a part of quadriceps? *Gait Posture*, in press.