

## University of Groningen

### Tree of the sea

Witbaard, Rob

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1997

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Witbaard, R. (1997). *Tree of the sea: The use of the internal growth lines in the shell of Arctica islandica (Bivalvia, Mollusca) for the retrospective assessment of marine environmental change*. [Thesis fully internal (DIV), University of Groningen]. s.n.

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

---

## REFERENCES

- Aebischer, N.J., J.C. Coulson & J.M. Colebrook, 1990. Parallel long-term trends across four marine trophic levels and weather. —*Nature* 347: 753-755.
- Anger, K., U. Rogal, G. Schriever & C. Valentin, 1977. In situ investigation on the echinoderm *Asterias rubens* as predator of soft bottom communities in the western Baltic Sea. —*Helgoländer Wiss. Meeresunters.* 29(4): 417-532.
- Anon., 1991. Visserij-economisch onderzoek. —*In* Visserijonderzoek jaarverslag 1991, p. 97-105. Uitgave van het Rijksinstituut voor Visserijonderzoek, het Instituut voor Visserijproducten en het Landbouw-Economisch Instituut. de Tunnel, Velsen-noord: 136 pp.
- Anon., 1992a. Report for ACMP on the effects of physical disturbance on the sea floor on the benthos. Review of current research reported at ICES Benthos Ecology Working Group, Bergen 3-9 May 1992. ICES document: 4 pp.
- Anon., 1992b. Vissen naar evenwicht. Structuurnota Zee- en Kustvisserij. — Veenman druk, Wageningen: 97 pp.
- Arntz, W.E., 1972. Über das auftreten des nemertinen *Malacobdella grossa* in *Cyprina islandica* aus der Kieler Bucht. —*Kieler Meeresforsch.* 28: 199-203.
- Arntz, W.E., 1974. A contribution to the feeding ecology of juvenile cod *Gadus morhua* L.) in the western Baltic. —*Rapp. P.-V. Réun. Cons. int. Explor.*, 166: 13-19.
- Arntz, W.E. & W. Weber, 1970. *Cyprina islandica* (L.) Mollusca, Bivalvia als Nahrung von Dorsch und Kliesche in der Kieler Bucht. —*Ber. Dt. Wiss. Komm. Meeresforsch.* 21 H. 1-4: 193-209.
- Asthorsson, O.S., 1990. Ecology of euphausiids *Thysanoessa raschi*, *T. inermis* and *Meanyctiphanes norvegica* in Ísafjord-deep, northwest-Iceland. —*Mar. Biol.* 107: 147-157.
- Austen, M.C., J.B. Buchanan, H.G. Hunt, A.B. Josefson & M.A. Kendall, 1991. Comparison of long-term trends in benthic and pelagic communities of the North Sea. —*J. Mar. Biol. Ass. U.K.* 71: 179-190.
- Babkov, A.I. & A.N. Golikov, 1984. Hydrobiocomplexes of the White Sea. USSR Academy of Sciences, Zoological Institute: 102 pp.
- Basford, D. & A. Eleftheriou, 1988. The benthic environment of the North Sea (56° to 61°N). —*J. mar. Biol. Ass. U.K.* 68 (1): 25-141.
- Basford, D.J., A. Eleftheriou & D. Raffaelli. 1989. The epifauna of the northern North Sea (56°-61°N). —*J. Mar. Biol. Ass. U.K.* 69: 387-407.
- Bayne, B.L., 1971. Oxygen consumption by three species of Lamellibranch molluscs in declining ambient oxygen tension. —*Comp. Biochem. Physiol.* 40A: 955-970.
- Bearse, D.T., 1976. Density and distribution of the ocean quahog (*Arctica islandica*) in Rhode Island waters relative to various environmental factors. Master of science thesis University of Rhode Island: 90pp.
- Becker, G., A. Corten & H. Dooley, 1992. Recent high salinity in the English Channel/southern North Sea. —*ICES hydrography committee. C.M.* 1992/C6: 11pp.

## References

---

- Becker, G.A. & G. Wegner, 1993. North Sea surface temperature means 1981 to 1990. —ICES C.M. 1993/C47: 4pp.
- Berger, R., 1979. Artificial radiocarbon in the troposphere. —Proc. 9th Int. Radiocarbon Conf. Univ. of Calif., Los Angeles: 309-312.
- Bergman, M.J.N., 1992. Long term effects of beamtrawl fishing on the benthic ecosystem in the North Sea: p.25-27. —*In*: Effects of beamtrawl fishery on the bottom fauna III-The 1991 studies; BEON report 16: 27pp.
- Bergman, M.J.N. & M. Hup. 1992. Direct effects of beamtrawling on macrofauna in a sandy sediment in the southern North Sea. —ICES J. Mar. Sci. 49: 5-11.
- Bergman, M.J.N., M. Fonds, W., Lewis, P. van der, Puyl, A, Stam & D. den Uyl, 1990. Direct effects of beamtrawl fishing on benthic fauna in the North Sea - a pilot study: p. 33-57. —*In*: Effects of beamtrawl fishery on the bottom fauna in the North Sea; BEON-rapport 8: 57pp.
- Blacker, R.W., 1957. Benthic animals as indicators of hydrographic conditions and climatic change in Svalbard waters. —Fishery Investigations II. XX 10: 1-49.
- Blom, W. C., 1990. Some remarks about the beamtrawl. p: 26-31. —*In*: Effects of beamtrawl fishery on the bottom fauna in the North Sea; BEON-rapport 8: 57pp.
- Borg, K van der, C.A. Aldersliesten , C.M. Houston, A.F.M. de Jong & N.A. van Zwol, 1987. Accelerator mass spectrometry with  $^{14}\text{C}$  and  $^{10}\text{Be}$  in Utrecht. —Nucl. Instr. Meth. B29: 143-145.
- Brey, T., W.E. Arntz, D. Pauly & H. Rumohr, 1990. *Arctica (Cyprina) islandica* in Kiel Bay (Western Baltic): growth, production and ecological significance. —J. Exp. Mar. Biol. Ecol. 136: 217-235.
- Bridger, J.P., 1970. Some effects of the passage of a trawl over the seabed. —ICES C.M. 1970/B10: 10pp.
- Bridger, J.P., 1972. Some observations on the penetration into the seabed of ticklerchains on a beamtrawl. —ICES C.M. 1972/B7: 6 pp.
- Broecker, W.S., T-H. Peng, G. Ostlund & M. Stuiver, 1985. The distribution of bomb radiocarbon in the ocean. —J. Geophys. Res. 90 (c4): 6953-6970.
- Broom, M.J. & J. Mason, 1978. Growth and spawning in the Pectinid *Chlamys opercularis* in relation to temperature and phytoplankton concentration. —Mar. Biol. 47: 277-285.
- Caddy, J.F., 1968. Underwater observation on scallop (*Placopecten magelanicus*) behavior and drag efficiency. Biological Station st. Andrews. —J. Fish. Res. Brd. of Canada, 25(10): 2123-2141.
- Christensen, A.M., 1970. Feeding biology of the seastar *Astropecten irregularis* Pennant. —Ophelia 8: 1-134.
- Cook, E.K. & L.A. Kariukstis, 1990. Methods of dendrochronology. Applications in the environmental sciences. —Kluwer Academic Publishers, Dordrecht.
- Corten, A. & G. v/d Kamp, 1992. Natural changes in pelagic stocks of the North Sea in the 1980s: p. 402-417. —*In*: Dickson R.R., et al. (eds) Hydrobiological variability in the ICES area, 1980-1989. —ICES Marine. Science. Symposia 195: 514pp.
- Cramer, S. & N. Daan, 1986. Consumption of benthos by North Sea cod and haddock in 1981. —ICES C.M. 1986/G56: 14pp.
- Crawley, M.J., 1993. GLIM for ecologists. —Oxford: 379 pp.

---

## *References*

- Creutzberg, F. 1985. A persistent chlorophyll a maximum coinciding with an enriched benthic zone: p. 97-108. —*In*: P.E. Gibbs (*ed*) Proceedings of the 19th EMBS. Cambridge University Press: 541pp.
- Creutzberg, F., G.C.A. Duineveld & G.J. van Noort, 1987. The effect of different numbers of tickler chains on beam-trawl catches. —*Rapp. P. -v Réun. Cons. int. Explor. Mer*, 43: 159-168.
- Creutzberg, F. & H. Postma, 1979. An experimental approach to the distribution of mud in the southern North Sea. —*Neth. J. Sea Res.* 13 (1): 99-116.
- Creutzberg, F., P. Wapenaar, G. Duineveld & N. Lopez Lopez, 1984. Distribution and density of the benthic fauna in the southern North Sea in relation to bottom characteristics and hydrographic conditions. —*Rapp. P. -v Réun. Cons. int. Explor. Mer*, 183: 101-110.
- Crisp, D.J., 1984. Energy flow measurements: p. 284-372. —*In*: N.A. Holme & A.D. McIntyre (*eds*) 1984. Methods for the study of marine benthos. Blackwell scientific publications, Oxford: 387 pp.
- Christensen, H. & E. Kannevorff, 1985. Sedimenting phytoplankton as major food source for suspension and deposit feeders in the Øresund. —*Ophelia* 24(3): 223-244.
- Cushing, D.H., 1982. Climate and Fisheries. Academic Press. London: 373pp.
- Davies, J.M. & R. Payne, 1984. Supply of organic matter to the sediment in the northern North Sea during a spring phytoplankton bloom. —*Mar. Biol.* 78: 315-324.
- Davis, F.M., 1923. Quantitative studies on the fauna of the sea Bottom. No 1. Preliminary investigation of the Doggerbank. —*Fishery Investigations Series II vol VI no 2*: 54 pp.
- Davis, F.M., 1925. Quantitative studies on the fauna of the Sea bottom. No 2. Results of the investigation in the southern North Sea. 1921-24. —*Fishery Investigations. Series II vol VIII no 4*: 50 pp.
- Dickson, R.R., J. Meincke, S.A. Malmberg & V. Lee, 1988a. The "great salinity anomaly" in the northern North Atlantic. —*Progress in Oceanography* 20:103-151.
- Dickson, R.R., P.M. Kelly, J.M. Colebrook, W.S. Wooster & D.H. Cushing, 1988b. North winds and production in the eastern North Atlantic. —*J. Plankt. Res.* 10(1):151-169.
- Dooley, H.D & G.K. Furnes, 1981. The influence of the wind field on the transport of the northern North Sea: p. 57-71. —*In*: R. SÆtre & M. Mork (*eds*). The Norwegian coastal current. Proceedings from the Norwegian Coastal Current Symposium. Geilo, 1980. University of Bergen: 369pp.
- Dries, R.R. & H. Theede, 1974. Sauerstoffmangelresistenz mariner Bodenvertebraten aus der Westlichen Ostsee. —*Mar. Biol.* 25: 327-333.
- Duineveld, G.C.A. & M.I. Jenness, 1984. Differences in growth rates of the sea urchin *Echinocardium cordatum* as estimated by the parameter  $\omega$  of the von Bertalanffy equation applied to skeletal rings. —*Mar. Ecol. Progr. Ser.* 19: 65-72.
- Duineveld, G.C.A., A. Künitzer, U. Niermann, P.A.W.J. de Wilde & J. Gray, 1991. The macrobenthos of the North Sea. —*Neth. J. Sea Res.* 28 (1/2): 53-65.

## References

---

- Duineveld, G.C.A., P.A.W.J. de Wilde & A. Kok, 1990. A synopsis of the macrobenthic assemblages and benthic ETS activity in the Dutch sector of the North Sea. —Neth. J. Sea Res. 26: 125-138.
- Evans, F. & A. Edwards, 1993. Changes in the zooplankton community off the coast of Northumberland between 1969 and 1988, with notes on changes in the phytoplankton and benthos. —J. Exp. Mar. Biol. Ecol. 172: 11-29.
- Eleftheriou, A. & D.J. Basford, 1989. The macrobenthic infauna of the offshore northern North Sea. —J. Mar. Biol. Ass. U.K. 69: 123-143.
- Ellett, D.J. & J. Blindheim, 1992. Climate and hydrographic variability in the ICES area during the 1980s: p. 11-31. —In: Dickson, R.R. et al. (eds). Hydrobiological variability in the ICES area, 1980-1989. ICES Marine Science Symposia. 195:514pp.
- Emerson, C.W., 1990. Influence of sediment disturbance and water flow on the growth of the soft-shell clam, *Mya arenaria* L. —Can. J. Fish. Aquat. Sci. 47:1655-1663.
- Epstein, S., R. Buchsbaum, H.A. Lowenstein & H.C. Urey, 1953. Revised carbonate-water isotopic temperature scale. —Bull. of the Geol. Soc. of America, G4: 1315-1326.
- Erlenkeuser, H. 1976.  $^{14}\text{C}$  and  $^{13}\text{C}$  isotope concentration in modern marine mussels from sedimentary habitats. —Naturwissenschaften 63: 338.
- Faubel, A., E. Hartwig & H. Thiel, 1983. On the ecology of the benthos of sublittoral sediments, Fladen Ground, North Sea I. Meiofauna standing stock and estimation of production. —Meteor Forschungs Ergebnisse Reihe D 36: 35-48.
- Fonds, M., 1994. Catch composition of 12-m beam trawl and 4-m beam trawl for sole fishing: p. 95-130. —In: Groot, S.J. & H.J. Lindeboom (eds). Environmental impact of bottom gears on benthic fauna in relation to natural resources management and protection of the North Sea: 257pp.
- Fonds, M., 1991. Measurements of catch composition and survival of benthic animals in beam trawl fishery for sole in the southern North Sea: p. 53-67. —In: Effects of beam trawl fishery on the bottom fauna in the North Sea. II- 1990 studies; BEON report 13: 85pp.
- Fonds, M., B.L. Verboom & K. Groeneveld, 1992. Catch composition and survival of fish and benthic invertebrates in commercial beamtrawls for sole and plaice fishery in the southern North Sea: p.17-22. —In: Effects of beamtrawl fishery on the bottom fauna in the North Sea. III-The 1991 studies; BEON-report 16: 27pp.
- Forster, G.R., 1981. A note on the growth of *Arctica islandica*. —J. mar. Biol. Ass. U.K. 61: 817.
- Franz, H.G. & W.W.C. Gieskes, 1984. The unbalance of phytoplankton and copepods in the North Sea. —Rapp. P. -v Réun. Cons. int. Explor. Mer. 183: 218-225.
- Frechette, M., D. Lefavre & C.A. Butman, 1993. Bivalve feeding in the benthic boundary layer: p. 325-370. —In: R.F. Dame (ed). Bivalve Filter Feeders in Estuarine and Coastal Ecosystem processes. —NATO ASI series G: Ecological sciences 33: 579pp.

---

## References

- Fritz, L.W., 1991. Seasonal condition change, morphometrics, growth and sex ratio of the ocean quahog, *Arctica islandica* (Linnaeus, 1767) off New Jersey, (U.S.A). —J. Shellfish Res. 10(1): 79-88.
- Fry, B. & E.B. Sherr, 1988.  $^{13}\text{C}$  measurements as indicators of carbon flow in marine and freshwater ecosystems: p. 197-229. —In: P.W. Rundel, J.R. Ehleringer & K.A. Nagy (eds). Stable isotopes in ecological research. Springer verlag: 525pp.
- Furnes, G.K., 1992. Long-term variability of the in-out flow to the North Sea: p.103-109. —In: Dickson R.R., et al. (eds). Hydrobiological variability in the ICES area, 1980-1989. ICES Marine Science. Symposia 195: 514pp.
- Gaard, E. & H. Mortensen, 1993. Phyto- and zooplankton communities on the Faroe Bank and their relations to the physical and chemical environment. —ICES C.M. 1993/L54: 17pp.
- Gee A. de, M.A. Baars & H.W. van der Veer, 1991. De ecologie van het Friese Front. Waarnemingen aan een biologisch rijke zone in de Noordzee, gelegen tussen de Zuidelijke Bocht en de Oestergronden. —NIOZ Rapport 1991-2. NIOZ, Texel: 96pp.
- Gering, G., 1911. Beiträge zur Kenntnis von *Malacobdella grossa* (Mull). —Z. wiss. zool. 97: 673-720.
- Gibson, R., 1967. Occurrence of the entocommensal rhynchocoelan, *Malacobdella grossa*, in the oval piddock, *Zirfaea crispata* on the yorkshire coast. —J. mar. biol Ass. U.K. 47: 301-317.
- Gibson, R., 1968. Studies on the biology of the entocommensal rhynchocoelan *Malacobdella grossa*. —J. mar. Biol. Ass. U.K. 48: 637-656.
- Gieskes, W.W.C & G.W. Kraay, 1984. Phytoplankton, its pigments and primary production at a central North Sea station in May, July and September 1981. —Neth. J. Sea Res. 18: 51-70.
- Goedecke, E., J. Smed & G. Tomczak, 1967. Monatskarten des Salzgehaltes der Nordsee dargestellt für verschiedene Tiefenhorizonten. —Ergänzungsh. Dt. Hydrogr. Z., Reihe B(4<sup>0</sup>) nr. 9.
- Gosselck, F., F. Doerschel & T. Doerschel, 1987. Further developments of macrozoobenthos in Lübeck Bay following recolonisation in 1980/81. —Int. Rev. Gesamt. Hydrobiol. 72(5): 631-638.
- Graf, G. W. Bengtsson, A. Faubel, L.-A. Meyer-Reil, R. Schulz, H. Theede & H. Thiel, 1984. The importance of the spring phytoplankton bloom for the benthic system of Kiel Bight. —Rapp. P.-v. Réun. Cons. int Explor. Mer. 183: 136-143.
- Grizzle, R.E., R. Langan & W.H. Howell, 1992. Growth responses of suspension-feeding bivalve molluscs to changes in water flow: Differences between siphonate and nonsiphonate taxa. —J. Exp. Mar. Biol. Ecol. 162: 213-228.
- Grizzle R.E. & R.A. Lutz, 1989. A statistical model relating horizontal seston fluxes and bottom sediment characteristics to growth of *Mercenaria mercenaria*. —Mar. Biol. 102: 95-105.
- Grizzle, R.E. & P.J. Morin, 1989. Effect of tidal currents, seston and bottom sediments on growth of *Mercenaria mercenaria*: Results of a field experiment. —Mar. Biol. 102: 85-93.

## References

---

- Groot, S.J. de, 1973. De invloed van trawlen op de zeebodem. —Visserij, 26(7): 401-409.
- Groot, S.J. de, 1984. The impact of bottom trawling on benthic fauna of the North Sea. —Ocean Management, 9: 177-190.
- Groot, S.J. de & H.J. Lindeboom, 1994. Environmental impact of bottom gears on benthic fauna in relation to natural resources management and protection of the North Sea. —NIOZ rapport 1994-1/RIVO-DLO report C026/94: 257pp.
- Grossman, E.L. & T-L. Ku, 1986. Oxygen and carbon isotope fractionation in biogenic aragonite: temperature effects. —Chem. Geol. 59: 59-74.
- Hagmeier, A., 1930. Die Bodenfauna der Ostsee im April 1929 nebst einiger Vergleichen mit April 1925 und Juli 1926. —Ber. Deutsche Wissenschaftl. kommission f. Meeresforsch. neue Folge 88: 156-173.
- Hartwig, E., A. Faubel & H. Thiel, 1983. On the ecology of the benthos of sublittoral sediments, Fladen Ground, North Sea II. Quantitative studies on macrobenthic assemblages. —Meteor Forschungs Ergeb. Reihe D 36: 49-64.
- Heller, J., 1990. Longevity in molluscs. —Malacologia 31(2): 259-295.
- Høisæter, T., 1986. An annotated check-list of marine molluscs of the Norwegian coast and adjacent waters. —Sarsia 71: 73-145.
- Holme, N.A., 1953. The biomass of the bottom fauna in the English channel off Plymouth. —J. Mar. Biol. Ass. U.K. 32: 1-49.
- Holtmann, S.E. & A. Groenewold, 1992. Distribution of the zoobenthos of the Dutch continental shelf: The Oyster Ground, Frisian Front, Vlieland Ground and Terschelling Bank (1991). —NIOZ rapport 1992-8, NIOO-Cemo rapporten en verslagen 1992-6. North Sea Directorate. Rijswijk: 129pp.
- Holtman, S.E. & A. Groenewold, 1994. Distribution of the zoobenthos of the Dutch continental shelf: The western Frisian Front, Brown Bank and Broad Fourteens (1992/1993). —NIOZ rapport 1994-1, NIOO-Cemo rapporten en verslagen 1994-1. North Sea Directorate. Rijswijk: 136pp.
- Hunt, O.D., 1925. The Food of the bottomfauna of the Plymouth fishing grounds. —J. Mar. Biol. Ass. U.K. 13: 560-599.
- Jones, D.S., 1979. The nemertean *Malacobdella grossa* in the ocean quahog, *Arctica islandica* (Bivalvia). —The Nautilus 93(1): 29-30.
- Jones, D.S., 1980. Annual cycle of shell growth increment formation in two continental shelf bivalves and its paleoecologic significance. —Paleobiology 6(3):331-340.
- Jones, D.S., 1983. Sclerochronology: reading the record of molluscan shell. —American Scientist 71: 384-391.
- Jones, D.S., M.A. Arthur & D.J. Allard, 1989. Sclerochronological records of temperature and growth form shells of *Mercenaria mercenaria* from Narragansett Bay, Rhode Island. —Mar. Biol. 102: 225-234.
- Jongman, R.H.G., C.J.F. terBraak, & O.F.R. van Tongeren, 1987. Data analysis in community and landscape ecology. —Pudoc Wageningen 299: pp.
- Josefson, A.B., J.N. Jensen, T. G. Nielsen & B. Rasmussen, 1995. Growth parameters of a benthic suspension feeder along a depth gradient across the pycnocline in the southern Kattegat, Denmark. —Mar. Ecol. Progr. Ser. 125: 107-115.

---

## *References*

- Kalish, J.M., 1991.  $^{13}\text{C}$  and  $^{18}\text{O}$  isotopic disequilibria in fish otoliths: metabolic and kinetic effects. —Mar. Ecol. Prog. Ser. 75: 191-203.
- Kennish M.J., 1980. Shell microgrowth analysis: *Mercenaria mercenaria* as a type example for research in population dynamics: p.255-295. —In: D.C. Rhoads & R. Lutz (eds). Skeletal growth of aquatic organisms. Plenum Press New York and London: 750 pp.
- Kennish, M.J. & R.A. Lutz, 1995. Assessment of the ocean quahog, *Arctica islandica* (Linnaeus, 1767), in the New Jersey fishery. —J. Shellfish Res. 14(1): 45-52.
- Kjørboe, T., F. Møhlenberg & O. Nøhr, 1981. Effect of suspended bottom material on growth and energetics in *Mytilus edulis*. —Mar. Biol. 61: 283-288.
- Krantz, D.E., D.S. Jones & D.F. Williams, 1984. Growth rates of the sea scallop *Placopecten magellanicus* determined from the  $^{18}\text{O}/^{16}\text{O}$  record in shell calcite. —Biol. Bull. 167: 186-199.
- Kraus, M.G., B.F. Beal, S.R. Chapman & L. McMartin, 1992. A comparison of growth rates in *Arctica islandica* (Linnaeus, 1767) between field and laboratory populations. —J. Shellfish Res. 11(2): 289-294.
- Kummel, B., Raup, D., 1965. Handbook of paleontological techniques. —W.H. Freeman and Company, San Francisco and London.
- Künitzer, A., 1990. Zur Verbreitung und populations Dynamik der Bodenfauna der centralen Nordsee. —Dissertation zur erlangung des grades eines Doktors der Naturwissenschaften. Alfred Wegener Institut für Polar und Meeresforschung Bremerhafen BRD: 218 pp.
- Laban, C. & H. Lindeboom, 1991. Penetration depth of beamtrawl gear: p.11-36. —In: Effects of beamtrawl fishery on the bottomfauna in the North Sea. II- The 1990 studies; BEON report 13: 85pp.
- Landers, W.S., 1976. Reproduction and early development of the ocean quahog *Arctica islandica*, in the laboratory. —The Nautilus 90 (2): 88-92.
- Li, H., J.J.M. van Haren & H.M. van Aken, 1989. Cluster analysis as a method to discriminate water masses in shelf seas. —ICES C.M. 1989 /C3: 14 pp.
- Lindley, J.A., J. Roskell, A.J. Warner, N.C. Halliday, H.G. Hunt, A.W.G. John & T.D. Jonas, 1990. Doliolids in the German Bight in 1989: Evidence for exceptional inflow into the North Sea. —J. Mar. Biol. Ass. U.K. 70, 679-682.
- Loosanoff, V.L., 1953. Reproductive cycle in *Cyprina islandica*. —Bull. Mar. Biol. Lab., Woodshole 104: 146-155.
- Lovén, Per Magnus, 1929. Undersökningar över Öresund. Untersuchungen aus dem Öresund XVI. Beiträge zur Kenntnis der *Cyprina islandica* L. im Öresund. —Lunds Universitets Årsskrift. N.F. Avd 2. Bd 26 nr 1. Kungl Fysiografiska sällskapets handlingar N.F. Bd 41. nr 1 Lund Hakans Buchdruckerei.
- Lutz, R.A. & L.W. Fritz, 1990. Growth patterns within the shell of the ocean quahog, *Arctica islandica*. A review and recent observations. —J. Shellfish Res. 8: 463-464.
- Lutz, R.A., R. Mann, J.G. Goodsell & M. Castagna, 1982. Larval and early post-larval development of *Arctica islandica*. —J. mar. Biol Ass. U.K. 62: 745-769.
- Mackie, A.S.Y., P.G. Oliver & E.I.S. Rees, 1995. Benthic biodiversity in the southern Irish Sea. Studies in Marine Biodiversity and systematics from the national museum of Wales. —BIOMÔR reports 1: 263pp.



## References

---

- Mann, R., 1982. The seasonal cycle of gonadal development in *Arctica islandica* from the southern New England Shelf. —Fish. Bull. 80(2): 315-326.
- Mann, R., 1986. *Arctica islandica* (Linné) larvae: Active depth regulators or passive particles. —American Malacological Bulletin special edition no. 3: 51-57.
- Mann, R. & C.C. Wolff, 1983. Swimming behaviour of larvae of the ocean quahog *Arctica islandica* in response to pressure and temperature. —Mar. Ecol. Prog. Ser. 13: 211-218.
- Margetts, A.R. & J.P. Bridger, 1971. The effect of a beam trawl on the sea bed. —ICES C.M. 1971/B8: 9pp.
- McGill, R., J.W. Tukey & W.A. Larsen, 1978. Variations of box plots. —American statistician 32(1): 12-16.
- McIntyre, A.D., 1961. Quantitative differences in the fauna of boreal mud associations. —J. Mar. Biol. Ass. 41: 559-616.
- Meisner, J. & S. Brager, 1990/ The feeding ecology of wintering eiders *Somateria mollissima* and common scoters *Melanitta nigra* on the Baltic Sea coast of Schleswig-Holstein, FRG. —Wader study group bulletin 58:10-12.
- Merrill, A.S., J.L. Camberlin & J.W. Ropes, 1969. Ocean Quahog Fishery. —In: F.E. Firth (ed). The Encyclopedia of Marine resources. Van Norstrand Reinhold Company: 740pp.
- Merrill, A.S. & J.W. Ropes, 1969. The general distribution of the surfclam and ocean quahog. —Proc. Nat. Shellfish Assoc. 59: 40-45.
- Miller, W. III, 1983. Biogenic damage in the small gastropod *Odostomia impressa* (Say). —Tulane studies in Geology and Paleontology 17:105-116.
- Mills, E.L. & R.O. Fournier, 1979. Fish production and the marine ecosystems of the Scotian Shelf, eastern Canada. —Mar. Biol. 45: 101-108.
- Moorsel, G.W.N.M. van, 1993. Long-term recovery of geomorphology and population development of large molluscs after gravel extraction at the Klaverbank (North Sea). —Bureau Waardenburg Culemborg. Report 92-16: 41pp.
- Moorsel, G.W.N.M van & H.W. Waardenburg, 1991. Short-term recovery of geomorphology and macrobenthos of the Klaverbank (North Sea) after gravel extraction. —Bureau Waardenburg, Culemborg: 53pp.
- Møhlenberg, F. & T. Kiørboe, 1981. Growth and energetics in *Spisula subtruncata* (Da Costa) and the effect of suspended bottom material. —Ophelia, 20(1):79-90.
- Møhlenberg, F. & H.U. Riisgård, 1979. Filtration rate, using a new indirect technique, in thirteen species of suspension feeding bivalves. —Mar. Biol. 54: 143-147.
- Mulder, R. & M.W. Manuals, 1982. Description of a drifter system used in experiments in the southern part of the North Sea. —Deutsche Hydrographische Zeitschrift. 35: 203-210.
- Murawski, S.A., J.W. Ropes, & F.M. Serchuk, 1982. Growth of the ocean quahog, *Arctica islandica* in the middle Atlantic Bight. —Fish. Bull. 80 (1): 21-34.
- Murawski, S.A. & F.M. Serchuk, 1979a. Mechanized Shellfish harvesting and its management: the offshore clam fishery of the eastern United States: p.479-700. —In: J.F. Caddy ed. Marine invertebrate fisheries: their assessment and management. John Wiley and Sons, New York: 752pp.

---

## References

- Murawski, S.A. & F.M. Serchuk, 1979b. Shell length-meat weight relationships of ocean quahogs, *Arctica islandica* from the Middle Atlantic Shelf. —Proc. Nat. Shellfish. Ass. 69: 40-46.
- Murawski, S.A., T.R. Azarovitz & D.J. Radosh, 1989. Long-term biological effects of hypoxic water conditions off New Jersey, USA 1976-1989. —ICES CM 1989/E11: 12pp.
- Muus, K., 1973. Settling, growth & mortality of young bivalves in the Øresund. —Ophelia 12: 79-116.
- Naletova, I.A., V.V. Sapozhnikov & M.P. Metreveli, 1994. Production-destruction processes in the White Sea: p. 76-83. —In: V.V. Sapozhnikov (ed). Complex studies of the White Sea ecosystem. Collected papers Moscow: 123pp.
- Nicol, D., 1951. Recent species of the Veneroid pelecypod *Arctica*. —J. Wash. Acad. Sci. 41: 102-106.
- Nielsen, C., 1975. Observations on *Buccinum undatum* L. attacking bivalves and on prey responses with a short review on attack methods of other Prosobranchs. —Ophelia 13: 87-108.
- Nielsen, T.G., B. Løkkegaard, K. Richardson, F.B. Petersen & L. Hansen, 1993. Structure of plankton communities in the Doggerbank area (North Sea) during a stratified situation. —Mar. Ecol. Prog. Ser. 95 (1/2): 115-131.
- Noort, G.J. van & F. Creutzberg, 1981. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 5: Trawl survey October-November 1974. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1981-1: 116 pp.
- Noort, G.J. van, F. Creutzberg & R. Dapper, 1982. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 6: Trawl survey September-November, 1975. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1982-7: 68pp.
- Noort, G.J. van, F. Creutzberg & R. Dapper, 1983. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 7: Trawl survey February-March 1976. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel, 1983-3: 76pp.
- Noort, G.J. van , F. Creutzberg & R. Dapper, 1984. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 8: Trawl survey October 1976. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1984-1: 72 pp.
- Noort, G.J. van, F. Creutzberg & G.C.A. Duineveld, 1986. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 9: Trawl survey May and July 1980. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1986-3: 55pp.
- Noort, G.J. van , F. van Leeuwen & F. Creutzberg, 1979. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 1. Trawl survey April-May 1972. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1979-5: 104pp.
- Noort, G.J. van, F. van Leeuwen & F. Creutzberg, 1979. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea report 2. Trawl survey June-July 1972. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1979-6: 107pp.

## References

---

- Noort, G.J. van, F. van Leeuwen & F. Creutzberg, 1979. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 3: Trawl survey october-november 1972. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel 1979-9: 115 pp.
- Noort, G.J. van, F. van Leeuwen & F. Creutzberg, 1979. "Aurelia"-cruise reports on the benthic fauna of the southern North Sea. Report 4: Trawl survey January-February 1973. —Interne verslagen Nederlands Instituut voor Onderzoek der Zee, Texel. 1979-8: 100pp.
- Nørrevang, A., T. Brattegard, A.B. Josefson, J.A. Sneli & O.S. Tendal, 1994. List of Biofar stations. —Sarsia 79(3): 165-180.
- Nydal, R., K. Lövseth, & S. Gulliksen, 1979. A survey of radiocarbon variation in nature since the test ban treaty. —Proc. 9th Int. Radiocarbon Conf. Univ. of Calif. Los Angeles: 313-323.
- Nydal, R. & K Lövseth, 1983. Tracing bomb  $^{14}\text{C}$  in the atmosphere 1962-1980. —J. Geoph. Res. 88 c6: 3621-3642.
- Nydal, R., S. Gulliksen, K. Lövseth & F.H. Skogseth, 1984. Bomb  $^{14}\text{C}$  in the ocean surface 1966-1981. —Radiocarbon 26(1): 7-45.
- Nyström, J., 1995. Environmental archives disclosed by nuclear microscopy of mussel shells. Acta Univ. Ups. Comprehensive summaries of Uppsalla Dissertation from the faculty of Science and Technology 127: 38 pp.
- O'Reilly, J.E. & D.A. Busch, 1984. Phytoplankton primary production on the northwestern Atlantic shelf. —Rapp. P.-v. Réun. Cons. int Explor. Mer. 183: 255-268.
- Oertzen, J.A. Von, 1972. Cycles and rates of reproduction in six Baltic bivalves of different zoogeographical origin. —Mar. Biol. 14: 143-149.
- Oeschger, R., 1990. Long-term anaerobiosis in sublittoral marine invertebrates from the western Baltic Sea: *Halicryptus spinulosus* (Priapulida), *Astarte borealis* and *Arctica islandica* (Bivalvia). —Mar. Ecol. prog. Ser. 59: 133-143.
- Oeschger, R.E. & K.B. Storey, 1993. Impact of anoxia and hydrogen sulphide on the metabolism of *Arctica islandica* L. (Bivalvia). —J. Exp. Mar. Biol. Ecol., 170: 213-226.
- Palmer, R.E. 1979. A histological and histochemical study of the digestion in the bivalve *Arctica islandica* L. —Biol. Bull. 156: 115-129.
- Pearson, T.H., A.B. Josefson & R. Rosenberg, 1985. Petersen's benthic stations revisited I. Is the Kattegat becoming eutrophic? —J. Exp. Mar. Biol. Ecol. 92: 157-206.
- Petersen, C.G.J., 1913. Valuation of the Sea II. The animal communities of the sea bottom and importance for marine zoogeography. —Report of the Danish Biological Station XXI: 144pp.
- Petersen, C.G.J., 1915. The animal communities in the Skagerak, the Christiania Fjord and the Danish Waters. —Report of the Danish biological station XXIII: 32pp.
- Petersen, C.G.J., 1918. The sea bottom and its production of fish food. —Report of the Danish Biological station XXV: 62pp.
- Rees, H.L. & P.J. Dare, 1993. Sources of mortality and associated life-cycle traits of selected benthic species: a review. —Fisheries Research data report number 33. MAFF, Lowestoft: 36pp.

---

## References

- Reid, P.C., C. Lancelot, W.W.C. Gieskes, E. Hagmeier & G. Weichart, 1990. Phytoplankton of the North Sea and its dynamics: A review. —Neth. J. Sea. Res. 26 (2-4): 295-331.
- Reid, P.C., S.C. Surey-Gent, H.G. Hunt & A.E. Durrant, 1992. *Thalassiothrix longissima*, a possible oceanic indicator species in the North Sea: p.127-143. — In: R.R. Dickson et al. (eds). Hydrobiological variability in the ICES area, 1980-1989. ICES Marine Science Symposia 195: 514pp.
- Riegman, R. & F. Colijn, 1991. Evaluation of measurements and calculation of primary production in the Dogger Bank area (North Sea) in summer 1988. — Mar. Ecol. Prog. Ser. 69: 125-132.
- Romanek, C.S., E.L. Grossman & J.W. Morse, 1992. Carbon isotope fractionation in synthetic aragonite and calcite: Effects of temperature and precipitation rate. — Geochim. Cosmochim. Acta 56: 419-430.
- Ropes, J.W., 1985. Modern methods to age oceanic bivalves. —Nautilus 99(2-3): 53-57.
- Ropes, J.W. 1988. Ocean quahog *Arctica islandica*: p.129-132. —In: J. Penttila & L.M. Dery (eds). Age determination methods for Northwest Atlantic Species. NOAA technical report NMFS 72: 136pp.
- Ropes, J.W., Pyoas, D., 1982. Preliminary age and growth observations of ocean quahogs, *Arctica islandica* Linné from Georges Bank.—ICES C.M. 1982/k:15 Shellfish committee 6 pp.
- Ropes, J.W., D.S. Jones, S.A. Murawski, F.M. Serchuk & A. Jearld Jr., 1984a. Documentation of annual growth lines in ocean quahogs, *Arctica islandica* Linné. —Fishery Bulletin, 82(1): 1-20.
- Ropes, J.W., S.A. Murawski & F.M. Serchuk, 1984b. Size, age, sexual maturity, and sex ratio in ocean quahogs, *Arctica islandica* Linné off Long Island, New York. —Fishery Bulletin 82(2): 253-267.
- Rowell, T.W. & D.R. Chaisson, 1983. Distribution and abundance of the Ocean Quahog (*Arctica islandica*) and stimpson's surf clam (*Spisula polynyma*) resource on the Scotian Shelf. Canadian industry report of fisheries and aquatic sciences 142: 75pp.
- Rowell, T.W., D.R. Chaisson & J.T. Mclane, 1990. Size and age of sexual maturity and annual gametogenic cycle in the ocean quahog, *Arctica islandica* (Linnaeus, 1767), from coastal waters in Nova Scotia, Canada. —J. Shellfish Res., 9(1): 195-203.
- Rumohr, H., T. Brey & S. Ankar, 1987. A compilation of biometric conversion factors for benthic invertebrates of the Baltic. —The Baltic marine biologists publication no. 9: 56 pp.
- Rumohr, H. & P. Krost, 1991. Experimental evidence of damage to benthos by bottom trawling with special reference to *Arctica islandica*. —Meeresforsch. 33: 340-345.
- Saleuddin, A.S.M., 1964. Observations on the habit and functional anatomy of *Cyprina islandica* (L.). —Proc. Malac. Soc. Lond. 36: 149-162.
- Schäfer, W., 1972. Ecology and palaeoecology of marine environments. — Translated by I. Oertel. Edited by G.Y. Graig. The University of Chicago Press: 568pp.

## References

---

- Schmidt, N., 1988. Paleobiological implications of shell repair in recent marine gastropods from the northern gulf of California. —*Historical Biology* 3: 127-139.
- Smith, J.N., R. Nelson & S.E. Campana, 1991. The use of Pb-210/Ra 226 and  $^{228}\text{Th}/^{228}\text{Ra}$  disequilibria in the ageing of otoliths of marine fish. —*In*: P.J. Kershaw & D.S. Woodshead (eds). *Radionuclides in the study of marine processes*. Elsevier applied science: 350-359.
- Schultz, S., 1969. Das Makrobenthos der südlichen Beltsee. —*Beiträge zur Meereskunde* 26: 21-46.
- Schweingruber, F.H. 1989. *Tree rings. Basics and applications of dendro-chronology*. —Kluwer Academic Publishers. Dordrecht: 273 pp.
- Seaward, D.R., 1990. *Distribution of the marine molluscs of north west Europe*. Nature conservatory council. Interpretative branch nature conservancy council. —Shrewbury, England: 114 pp.
- Smetacek, V., 1984. The supply of food to the benthos. —*In*: M.J.R. Fasham (ed). *Flows of energy and material in marine ecosystems; theory and practice*. NATO conference series IV: marine sciences: 733 pp.
- Sokal, R.R. & F.J. Rohlf, 1981. *Biometry*. —W.H. Freeman and Company New York: 859pp.
- Steele, J.H., 1956. Plant production on the Fladen Ground. —*J. Mar. Biol. Ass. U.K.* 35:1-33.
- Steele, J.H., 1974. *The structure of marine ecosystems*. —Harvard Univ. Press, Cambridge, Mass., 128pp.
- Stefánsson, U. & S. Jónsdóttir, 1974. Near-Bottom temperature around Iceland. —*Rit Fiskideildar* 5(2): 3-73.
- Stuiver, M. & H.A. Polach, 1977. Discussion reporting of  $^{14}\text{C}$  data. —*Radiocarbon* 19: 355-363.
- Suess, H.E., 1958. The radioactivity of the atmosphere and hydrosphere. —*Annual review of nuclear science* 8: 243-256.
- Svendsen, E. & A.K. Magnusson, 1992. Climatic variability in the North Sea: p.144-158. —*In*: Dickson R.R., et al. (eds). *Hydrobiological variability in the ICES area, 1980-1989*. ICES Marine Science Symposia 195: 514pp.
- Svendsen, E., R. SÆtre & M. Mork, 1991. Features of the northern North Sea circulation. —*Cont. Shelf Res.* 11: 493-508.
- Tan, F.C., D. Cai & D.L. Roddick, 1988. Oxygen isotope studies on sea scallops, *Placopecten magellanicus*, from Browns Bank, Nova Scotia. Can. —*J. Fish. aquat. Sci.* 45: 1378-1386.
- Tanaka, N, M.C. Monaghan & K.K. Turekian, 1990.  $\Delta^{14}\text{C}$  balance for the Gulf of Maine, Long Island Sound and the northern Middle Atlantic Bight: Evidence for the extent of the Antarctic Intermediate water contribution. —*J. Mar. Res.*, 48: 75-87.
- Taylor, A.C., 1976. Burrowing behaviour and anaerobiosis in the bivalve *Arctica islandica* (L.). —*J. Mar. Biol. Ass. U.K.* 56: 95-109.
- Taylor, A.C. & A.R. Brand, 1975a. Effects of hypoxia and body size on the oxygen consumption of the bivalve *Arctica islandica* (L.) —*J. Exp. Mar. Biol. Ecol.* 19: 187-196.

---

## References

- Taylor, A.C. & A.R. Brand, 1975b. A comparative study of the respiratory responses of the bivalves *Arctica islandica* (L.) and *Mytilus edulis* (L.) to declining oxygen tension. —Proc. R. Soc. Lond. B. 190: 443-456.
- Thompson, I. & D.S. Jones, 1977. The ocean quahog, *Arctica islandica*, "tree" of the north Atlantic shelf. —Abstract Annual Meeting Geological Society of America: p 1199.
- Thompson, I., D.S. Jones & D. Dreibelbis, 1980a. Annual internal growth banding and life history of the ocean quahog, *Arctica islandica* (Mollusca: Bivalvia). —Mar. Biol. 57: 25-34.
- Thompson, I., D.S. Jones, & J.W. Ropes, 1980b. Advanced age for sexual maturity in the ocean quahog *Arctica islandica* (Mollusca: Bivalvia). —Mar. Biol. 57: 35-39.
- Thórarinsdóttir G.G. & S.T. Einarsson, 1994. Distribution, abundance, population structure, meat yield, size of sexual maturity and sex ratio of ocean quahog, *Arctica islandica* in Icelandic waters. —ICES C.M. 1994/k:39: 7pp.
- Thórdardóttir, T., 1973. Successive measurements of primary production and composition of phytoplankton at two stations west of Iceland. —Norw. J. Bot. 20: 257-270.
- Thórdardóttir, T., 1976. Preliminary assessment of the annual production in the shelf area around Iceland. —ICES C.M./L: 32: 4pp.
- Thors, K., 1978. The sea-bed of the southern part of the Faxaflói, Iceland. Jökli 28: 42-51.
- Tijssen, S.B. & F. Wetsteyn, 1983. Temperature, salinity and oxygen graphs and data, obtained near a subsurface drifter in the central North Sea in May, July and September 1981. —Interne verslagen NIOZ, Texel 4: 48pp.
- Tomczak, G. & E. Goedecke, 1964. Die thermische Schichtung der Nordsee und Monatskarten der Temperatur der Nordsee. —Ergänzungsh. Dt. Hydrogr.Z., Reihe B(4)8: 182pp.
- Turekian, K.K., J.K. Cochran, Y. Nozaki, I. Thompson, & D.S. Jones, 1982. Determination of shell deposition rates of *Arctica islandica* from the New York Bight using natural  $^{228}\text{Ra}$  and  $^{228}\text{Th}$  bomb produced  $^{14}\text{C}$ . —Limnol. Oceanogr. 27(4): 737-741.
- Trutsschler, K. & C. Samtleben, 1988. Shell growth of *Astarte elliptica* from Kiel Bay (Western Baltic Sea). —Mar. Ecol. Prog. Ser. 42: 155-162.
- Turrell, W.R., 1992a. New hypotheses concerning the circulation of the northern North Sea and its relation to North Sea fish stock recruitment. —ICES J. Mar. Sci. 49: 107-123.
- Turrell, W.R., 1992b. The East Shetland Atlantic Inflow: p.127-143. —In: Dickson R.R., et al. (eds). Hydrobiological variability in the ICES area, 1980-1989. ICES Marine Science Symposia 195:514pp.
- Vale, K.F. & M.A. Rex., 1988. Repaired shell damage in deep-sea prosobranch gastropods from the western north Atlantic. —Malacologia 28(1/2): 65-79.
- Vermeij, G.J., E.D. Schindel, & E. Zipser, 1981. Predation through geological time: Evidence from gastropod shell repair. —Science 214. 4524: 1024-1026.
- Vermeij, G.J., 1983. Traces and trends of predation with special reference to bivalved animals. —Palaeontology 26(3): 455-465.

## References

---

- Vogel, J.S., J.R. Southon, D.E. Nelson & D.A. Brown, 1984. Performance of catalytically condensed carbon for use in accelerator mass spectrometry. — Nucl. Intsr. Meth. B5: 289-293.
- Wefer, G., 1985. Die Verteilung stabiler Isotope in Kalkschalen mariner Organismen. — Geol. Jb. A82: 3-111.
- Wefer, G. & W.H. Berger, 1991. Isotope paleontology: growth and composition of extant calcareous species. — Mar. Geol. 100: 207-248.
- Weidman, C.R. & G.A. Jones, 1993a. A shell-derived time history of bomb  $^{14}\text{C}$  on Georges Bank and its Labrador Sea implications. — J. geophys. Res., 98(c8): 14577-14588.
- Weidman, C.R. & G.A. Jones, 1993b. A new High-temporal resolution paleoceanographic tool for the northern north Atlantic: The mollusc *Arctica islandica*. — ICES C.M. 1993/K17: 11pp.
- Weidman, C.R., G.A. Jones & K.C. Lohmann, 1994. The long-lived mollusc *Arctica islandica*: A new paleoceanographic tool for the reconstruction of bottom temperatures for the continental shelves of the northern north Atlantic Ocean. — J. geophys. Res. 99: 18305-18314.
- Welleman, H., 1989. Literatuurstudie naar de effecten van de bodemvisserij op de bodem en het bodemleven. — IJmuiden, RIVO MO 89-201: 58 pp.
- Wilde de, P.A.W.J., Berghuis, E.M., Kok, A., 1986. Biomass and activity of benthic fauna on the Fladen Ground (northern North Sea). — Neth. J. Sea. Res. 20 (2/3) 313-323.
- Wildish, D.J. & D.D. Kristmanson, 1985. Control of suspension feeding bivalve production by current speed. — Helgoländer Meeresunters. 39: 237-243.
- Wildish, D.J. & D.D. Kristmanson, 1993. Hydronamic control of bivalve filter feeders: a conceptual view: p.229-325. — In.: R.F. Dame (ed). Bivalve filter feeders in estuarine and coastal ecosystem processes. NATO ASI series G: Ecological sciences 33: 579pp.
- Winter, J.E., 1969. Über den Einfluß der Nahrungskonzentration und anderer Faktoren auf Filtrierleistung und Nahrungsausnutzung der Muscheln *Arctica islandica* und *Modiolus modiolus*. — Mar. Biol. 4: 87-135.
- Winter, J.E., 1970. Filter feeding and food utilization in *Arctica islandica* L. and *Modiolus modiolus* L. at different food concentrations: p.196-206. — In: J.H. Steele (editor) Marine food chains. Oliver and Boyd, Edingburgh: 552pp.
- Winter, J.E., 1978. A review on the knowledge of suspension feeding in Lamellibranchiate bivalves, with special reference to artificial aquaculture systems. — Aquaculture, 13: 1-33
- Witbaard, R., 1995. Ecoprofiel Noordkromp. Een overzicht van de biologie en ecologie van *Arctica islandica*. Uitgevoerd op het NIOZ in opdracht van Rijkswaterstaat in samenwerking met het Rijksinstituut voor kust en Zee. — Rapport NZ 864: 69 pp.
- Witbaard, R., 1996. Growth variations in *Arctica islandica* L. (Mollusca): a reflection of hydrography-related food supply. — ICES journ. Mar. Sci. 53; 981-987.
- Witbaard, R. & G.C.A. Duineveld, 1990. Shell-growth of the bivalve *Arctica islandica* (L.), and its possible use for evaluating the status of the benthos in the subtidal North Sea. — Basteria 54: 63-74.

---

*References*

- Witbaard, R. & R. Klein, 1994. Long-term trends on the effects of the southern North Sea beamtrawl fishery on the bivalve mollusc *Arctica islandica* (L.) (Mollusca, bivalvia). —ICES J. mar. Sci. 51: 99-105.
- Witbaard, R., M.I. Jenness, K. v/d. Borg & G. Ganssen, 1994. Verification of annual growth increments in *Arctica islandica* L. from the North Sea by means of oxygen and carbon isotopes. —Neth. J. Sea Res. 33(1): 91-101.
- Yager, P.L., A.R.M. Nowell & P.A. Jumars, 1993. Enhanced deposition to pits: a local food source for benthos. —J. Mar. Res. 51: 209-236.
- Zatsepin, V.I. & Z.A. Filatova, 1961. The bivalve mollusc *Cyprina islandica* (L.) Its geographic distribution and role in the communities of benthic fauna. —Transactions of the Institute of Oceanology. XLVI: 2-24.
- Zwier, G.J., 1983. De noordkromp en andere verhalen. —Elsevier Amsterdam Brussel: 110pp.
-