

University of Groningen

Precise Measurement of Born Cross Sections for $++\rightarrow^-$ at $\sqrt{s}=3.80\text{--}4.95\text{ GeV}$

BESIII Collaboration; Kalantar-Nayestanaki, N.; Kavatsyuk, M.; Messchendorp, J.

Published in:
Physical Review Letters

DOI:
[10.1103/PhysRevLett.133.081901](https://doi.org/10.1103/PhysRevLett.133.081901)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2024

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
BESIII Collaboration, Kalantar-Nayestanaki, N., Kavatsyuk, M., & Messchendorp, J. (2024). Precise Measurement of Born Cross Sections for $++\rightarrow^-$ at $\sqrt{s}=3.80\text{--}4.95\text{ GeV}$. *Physical Review Letters*, 133(8), Article 081901. <https://doi.org/10.1103/PhysRevLett.133.081901>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Precise Measurement of Born Cross Sections for $e^+e^- \rightarrow D\bar{D}$ at $\sqrt{s} = 3.80\text{--}4.95$ GeV

M. Ablikim *et al.*^{*}
(BESIII Collaboration)

 (Received 6 February 2024; revised 24 June 2024; accepted 29 July 2024; published 21 August 2024)

Using data samples collected with the BESIII detector at the BEPCII collider at center-of-mass energies ranging from 3.80 to 4.95 GeV, corresponding to an integrated luminosity of 20 fb^{-1} , a measurement of Born cross sections for the $e^+e^- \rightarrow D^0\bar{D}^0$ and D^+D^- processes is presented with unprecedented precision. Many clear peaks in the line shape of $e^+e^- \rightarrow D^0\bar{D}^0$ and D^+D^- around the mass range of $G(3900)$, $\psi(4040)$, $\psi(4160)$, $Y(4260)$, and $\psi(4415)$, etc., are foreseen. These results offer crucial experimental insights into the nature of hadron production in the open-charm region.

DOI: [10.1103/PhysRevLett.133.081901](https://doi.org/10.1103/PhysRevLett.133.081901)

The production of hadrons in e^+e^- annihilation above the open-charm threshold is a topic of ongoing theoretical and experimental research. In 1980, a theoretical calculation for the charm cross section in e^+e^- annihilation was first attempted based on a coupled-channel potential model [1]. This calculation presented a prediction of the ΔR value ($\Delta R = \sum_i R_i$, where R_i stands for the ratio of individual hadron cross section to muon cross section in electron-positron collisions and i runs over the two-body channels) with the $D\bar{D}$ final states. According to this prediction, there are five vector charmonium states between 3.773 [$\psi(3773)$, $1D$ state] and 4.95 GeV, namely, the $3S$, $2D$, $4S$, $3D$, and $5S$ states, dominated by the $D\bar{D}$ final states. In experimental studies, besides the three well-established structures observed in the inclusive hadronic cross sections [2], i.e., $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$, many new states, such as $Y(4230)$, $Y(4260)$, $Y(4360)$, and $Y(4660)$, have been reported in the initial state radiation (ISR) processes at the B factories [3–11] or in the direct e^+e^- production at the CLEO [12] and BESIII experiments [13–20]. Among them, the BESIII experiment found that the mass of $Y(4360)$ is around $4.3\text{ GeV}/c^2$ [14], which is different from the value given by the Particle Data Group (PDG) [2]. The overpopulation of structures in this region and the mismatch of the properties between the potential model predictions and experimental measurements have led to various interpretations, such as hybrid states, tetraquark states, or molecular states [21,22]. Although this information enriches our understanding of

these exotic structures, the nature of these states is still not understood.

The studies of charmed meson pairs in e^+e^- annihilation above the open-charm threshold are expected to clarify the current understanding of these states. At present, the available observed cross sections of the $e^+e^- \rightarrow D\bar{D}$ process with limited energy points have been reported by B factories [23,24] using the ISR process and through direct e^+e^- production at the CLEO experiment [25]. Although the interpretations for the possible structure featured in the $D\bar{D}$ final states are performed [26,27], the understanding for the properties of vector charmonium (like) states is still limited except for the $\psi(3770)$ state. A precise measurement, particularly of the exclusive Born cross sections for $e^+e^- \rightarrow D\bar{D}$, is highly desirable to validate the interpretations of the established states and provide insight into the energy region above the open-charm threshold.

In this Letter, we report a precise measurement of Born cross sections for the $e^+e^- \rightarrow D^0\bar{D}^0$ and $e^+e^- \rightarrow D^+D^-$ processes, specifically, at 150 center-of-mass (c.m.) energy points. Many clear structures in the line shape of $e^+e^- \rightarrow D^0\bar{D}^0$ and $e^+e^- \rightarrow D^+D^-$ in the c.m. energy at 3.90, 4.05, 4.20, 4.42 GeV, etc., can be seen. The datasets used in this Letter correspond to a total luminosity of approximately 20 fb^{-1} of e^+e^- collisions, which includes the so-called XYZ data sample [28,29] and the R -scan data sample [30], collected at c.m. energies from 3.80 to 4.95 GeV with the BESIII detector [31] at the BEPCII collider [32].

The BESIII detector's cylindrical core, encapsulating 93% of the 4π solid angle, integrates a helium-based multilayer drift chamber (MDC), a plastic scintillator time-of-flight (TOF) system, and a CsI (TI) electromagnetic calorimeter. These components are all nestled within a superconducting solenoidal magnet that generates a 1.0 T magnetic field. An octagonal flux-return yoke, fortified with steel-interleaved resistive plate chamber muon identifier

^{*}Full author list given at the end of the Letter.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

modules, supports the solenoid. This sophisticated setup achieves a charged-particle momentum resolution of 0.5% at 1 GeV/ c and a dE/dx resolution of 6% for electrons emanating from Bhabha scattering. The TOF system's barrel section boasts a time resolution of 68 ps, in contrast to the end cap section's 110 ps. Notably, the end cap TOF system underwent an enhancement in 2015, adopting multigap resistive plate chamber technology to furnish a time resolution of 60 ps.

In order to achieve a high efficiency for the selection of $e^+e^- \rightarrow D\bar{D}$ events, we employ a single tag technique instead of a full reconstruction. With this technique, we reconstruct only one D^0 (D^+) meson through the $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) mode, while the corresponding antiparticle \bar{D}^0 (D^-) is extracted from the recoil side. Unless otherwise noted, the charge-conjugate mode of the D^0 (D^+) process is included by default. To determine the detection efficiency for $e^+e^- \rightarrow D\bar{D}$, 50 000 simulated events are generated for each energy point using the KKMC generator [33,34] according to the VSS model [35,36], where the ISR effect is included. The D^0 (D^+) meson decays to the $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) modes are simulated with the amplitude sampling [37] via EVTGEN [35,36], and the antimesons are set to decay inclusively according to the known branching fractions provided by PDG [2]. The response of the BESIII detector is modeled with Monte Carlo (MC) simulation using a framework based on GEANT4 [38,39].

Charged tracks are reconstructed in the MDC with points of closest approach to the e^+e^- interaction point that are within 10 cm in the beam direction and 1 cm transverse to the beam direction and within the angular coverage of the MDC $|\cos\theta| < 0.93$, where θ is the polar angle with respect to the symmetry axis of the MDC. Information from the specific ionization energy loss measured in the MDC, combined with the time of flight, is used to determine the particle identification (PID) confidence levels for the pion and kaon hypotheses. Each track is assigned to the particle type with the higher probability. For the D^0 mode, events with at least one negatively charged kaon, one negatively charged pion, and two positively charged pions are kept for further analysis; for the D^+ mode, events with at least one negatively charged kaon and two positively charged pions for the D^+ mode are kept for further analysis. The D^0 (D^+) candidates are reconstructed from the $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) combination by requiring its invariant mass to be within 14(16) MeV/ c^2 around the nominal D^0 (D^+) mass, which corresponds to 3 times the mass resolution.

The antimeson candidates \bar{D}^0 (D^-) are inferred by the mass recoiling against the tagged meson (M_D^{recoil}) via the $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) system:

$$M_D^{\text{recoil}} = \sqrt{(\sqrt{s} - E_D)^2 - |\mathbf{p}_D|^2}, \quad (1)$$




FIG. 1. The two-dimensional distributions of M_D versus M_D^{recoil} at $\sqrt{s} = 4.1992$ GeV. The dashed lines represent the signal region for the D meson in the $D^0\bar{D}^0$ mode (top) and the D^+D^- mode (bottom).

where E_D and \mathbf{p}_D are the energy and momentum of the selected $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) candidate in the c.m. system, respectively, and \sqrt{s} is the c.m. energy [40]. To improve the resolution, a correction is applied to M_D^{recoil} given by $M_D^{\text{recoil}} + M_D - m_D$, where M_D is the invariant mass of the selected $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) candidate and m_D is the nominal D mass [2]. Figure 1 shows the 2D distributions of M_D^{recoil} versus M_D for the $K^-\pi^+\pi^+\pi^-$ ($K^-\pi^+\pi^+$) final states at $\sqrt{s} = 4.1992$ GeV. After applying all the aforementioned selection criteria, the remaining background exhibits a smooth shape in the region of interest based on the sideband study.

The signal yields for the $e^+e^- \rightarrow D\bar{D}$ process at each energy point are extracted by performing an extended maximum likelihood fit to the M_D^{recoil} spectrum in the range from 1.80 to 1.95 GeV/ c^2 . In the fit, the signal shape for the $e^+e^- \rightarrow D\bar{D}$ process is described by the convolution of the MC-simulated shape with a Gaussian function, which accounts for the difference in mass resolution between the data and the MC simulation. The parameters of the Gaussian function are floating for XYZ data points




FIG. 2. Fits to the M_D^{recoil} spectra at $\sqrt{s} = 4.1992$ GeV for the $D^0\bar{D}^0$ mode (top) and the D^+D^- mode (bottom). The dots with error bars represent the data, the blue solid lines indicate the fit results, the red short-dashed lines represent the signal, and the green long-dashed lines correspond to the background.

and some R -scan data points with higher statistics, while they are fixed for the other R -scan data points with low statistics. The fixed parameters are determined from the fits to the neighbor XYZ data points. The background contributions are modeled using a second-order polynomial function. Figure 2 illustrates the results of the fits to the M_D^{recoil} distributions at $\sqrt{s} = 4.1992$ GeV. The signal yields obtained from the fits are summarized in Supplemental Material [41].

The Born cross section for $e^+e^- \rightarrow D\bar{D}$ is calculated by

$$\sigma^B(s) = \frac{N_{\text{obs}}}{2\mathcal{L}(1+\delta)\frac{1}{|1-\Pi|^2}\epsilon\mathcal{B}}, \quad (2)$$

where N_{obs} represents the number of observed signal events, the factor of 2 accounts for the charge-conjugate mode, \mathcal{L} corresponds to the integrated luminosity, $(1+\delta)$ is the ISR correction factor, $(1/|1-\Pi|^2)$ is the vacuum polarization correction factor [45], ϵ denotes the detection efficiency, and \mathcal{B} stands for the branching fractions of $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ mode and $D^+ \rightarrow K^-\pi^+\pi^+$ mode, taken

from the Particle Data Group [2]. The ISR correction factor is obtained through QED calculations [46], where the cross sections measured in this analysis are used as inputs and iterated until convergence. The measured Born cross sections, along with the results from the CLEO-c [25], BABAR [23], and Belle [24] experiments, are shown in Fig. 3 and summarized in Ref. [41], which also includes all the numbers used in the calculation. To evaluate the resonance influence to the measurement of Born cross section, a least- χ^2 method combined with the simultaneous fit of the dressed cross sections $\sigma^{\text{dressed}} = \sigma^B/|1-\Pi|^2$ for the $e^+e^- \rightarrow D^0\bar{D}^0$ and D^+D^- processes, parametrized as the coherent sum of eight relativistic Breit-Wigner (BW) functions:

$$\sigma^{\text{dressed}}(\sqrt{s}) = \left| \sum_{k=1}^8 e^{i\phi_k} \text{BW}_k(\sqrt{s}) \sqrt{\frac{P(\sqrt{s})}{P(M)}} \right|^2, \quad (3)$$

has been attempted, where $\text{BW}(\sqrt{s})$ is given by

$$\text{BW}(\sqrt{s}) = \frac{\sqrt{12\pi}\Gamma_{ee}\mathcal{B}\Gamma}{s - M^2 + iM\Gamma}. \quad (4)$$

Here, the masses M and total widths Γ for seven known resonances $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, $Y(4230)$, $Y(4360)$, $\psi(4415)$, and $Y(4660)$ are fixed at individual PDG values [2], while they are free for another structure regarded as $G(3900)$ around 3.9 GeV [42–44]. For Γ_{ee} , electronic partial widths, and \mathcal{B} , the branching fractions of the decay, they are free for all resonances. The relative phase between different BW functions is denoted by ϕ , which is set to be different in the simultaneous fit for both modes. And $P(\sqrt{s})$ represents the two-body phase space factor. To account for the beam energy spread, $\sigma^{\text{dressed}}(\sqrt{s})$ is convolved with a Gaussian function with the standard deviation $\sigma = (-2.147 + 0.9454\sqrt{s})$ GeV. The influence to the measurement of Born cross section has been incorporated in the uncertainty of the line-shape model below. Note that the parameters for all assumed resonances strongly depend on the chosen fit model and indicating the need for further in-depth research, such as a coupled-channel K -matrix analysis [43]. The details for this fit are listed in Supplemental Material [41].

Systematic uncertainties in the measurement of the cross sections for the $e^+e^- \rightarrow D^0\bar{D}^0$ and D^+D^- processes originate from the luminosity measurement, the efficiencies of tracking and PID, the requirement of the D mass window, the fit of the M_D^{recoil} spectrum, the branching fractions of $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ and $D^+ \rightarrow K^-\pi^+\pi^+$, and the line-shape structures. The uncertainty due to the vacuum polarization is negligible. The integral luminosity is measured to an uncertainty up to 1.0% [28]. The uncertainties due to data-MC differences in BESIII tracking and particle ID efficiencies are both 1% per track [47]. The




FIG. 3. Cross sections with total uncertainties for the $e^+e^- \rightarrow D\bar{D}$ process as a function of c.m. energy from 3.80 to 4.95 GeV compared between this Letter for Born cross section (σ^B) and the previous measurements for observed cross section (σ^{obs}). Clear peaks around the mass range of $G(3900)$, $\psi(4040)$, $\psi(4160)$, $\psi(4415)$, etc. can be seen.

requirement of the D mass window is studied by varying the nominal requirements by $\pm 1\sigma$, resulting in uncertainties of 2.1% and 1.8% in the neutral and charged modes, respectively. The systematic uncertainty arising from the fit of the M_D^{recoil} spectrum includes the fit range and the background shape. Varying the mass range by ± 5 MeV/ c^2 results in an uncertainty of 1.5% for the neutral mode and 1.6% for the charged mode. The uncertainty due to the background shape is estimated to be 1.0% for the neutral mode and 0.9% for the charged mode using alternative fits with a second- or third-order polynomial function. The branching fractions of $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ and $D^+ \rightarrow K^-\pi^+\pi^+$ are quoted with uncertainties of 2.4% for both modes from the PDG [2]. The uncertainty arising from the line-shape model, including the ISR correction factor, is estimated by comparing the $(1 + \delta) \cdot \epsilon$ values with and without the addition of one more resonance in the fit of the cross sections as the input line shape, which introduces an uncertainty of 1.5% for the neutral mode and 1.6% for the charged mode between the nominal and alternative models. Assuming all sources are independent, the total systematic uncertainties on the cross section measurements are determined to be 7.0% for the $e^+e^- \rightarrow D^0\bar{D}^0$ mode and 6.5% for the $e^+e^- \rightarrow D^+D^-$ mode by quadratic sum.

In summary, a measurement of exclusive Born cross sections for the $e^+e^- \rightarrow D^0\bar{D}^0$ and D^+D^- processes is presented at 150 c.m. energy points ranging from 3.80 to 4.95 GeV with unprecedented precision. The result is in qualitative agreement with previous experiments [23–25] and

the prediction of the coupled-channel model [1]. Many clear peaks in the line shape of $e^+e^- \rightarrow D\bar{D}$ around the mass range of $G(3900)$, $\psi(4040)$, $\psi(4160)$, $\psi(4260)$, and $\psi(4415)$, etc., are identified. It implies that there may be some potential contributions from charmonium(like) states. In particular, the possible structure around 3.9 GeV, which was featured and interpreted by $G(3900)$ at the B factories [23,24], has been discussed recently by the theoretical models as the first P -wave $D\bar{D}^*$ molecular resonance [42], threshold enhancement [43], or the final-state interaction [44]. Thus, more detailed study related to a coupled-channel K -matrix analysis is needed to validate this structure.

Our results for describing the cross section provided in Supplemental Material [41] depend on the chosen model, which simplifies the analysis by ignoring the interactions between different decay channels. However, according to the model calculation by the Cornell group [1], strong coupled-channel effects need to be considered, which also is proposed by the recent theoretical works [42–44]. It is out of the scope of this Letter, but a more comprehensive approach based on K -matrix formalism to fit the cross section results of various exclusive channels is expected to test the scenarios [26,27,48–52] of charmonium(like) states above the open-charm threshold. This Letter provides important experimental evidences with unprecedented precision and insights into the nature above the open-charm region, especially for the understanding of the property of charmoniumlike states.

Acknowledgments—The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2020YFA0406300 and No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 12075107, No. 12247101, No. 11635010, No. 11735014, No. 11835012, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 12022510, No. 12025502, No. 12035009, No. 12035013, No. 12061131003, No. 12192260, No. 12192261, No. 12192262, No. 12192263, No. 12192264, No. 12192265, No. 12221005, No. 12225509, and No. 12235017; the 111 Project under Grant No. B20063; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contract No. U1832207; CAS Key Research Program of Frontier Sciences under Contracts No. QYZDJ-SSW-SLH003 and No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; European Union’s Horizon 2020 research and innovation program under Marie Skłodowska-Curie grant agreement under Contract No. 894790; German Research Foundation DFG under Contract No. 455635585, Collaborative Research Center CRC 1044, FOR5327, and GRK 2149; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; National Science Research and Innovation Fund (NSRF) via the Program Management Unit for Human Resources and Institutional Development, Research and Innovation of Thailand under Contract No. B16F640076; Polish National Science Centre under Contract No. 2019/35/O/ST2/02907; The Swedish Research Council; and U.S. Department of Energy under Contract No. DE-FG02-05ER41374.

[1] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T. M. Yan, *Phys. Rev. D* **21**, 203 (1980).
 [2] R. L. Workman *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
 [3] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **95**, 142001 (2005).
 [4] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **98**, 212001 (2007).
 [5] C. Z. Yuan *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 182004 (2007).
 [6] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **99**, 142002 (2007).

[7] G. Pakhlova *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **101**, 172001 (2008).
 [8] J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. D* **86**, 051102(R) (2012).
 [9] Z. Q. Liu *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **110**, 252002 (2013).
 [10] J. P. Lees *et al.* (BABAR Collaboration), *Phys. Rev. D* **89**, 111103(R) (2014).
 [11] X. L. Wang *et al.* (Belle Collaboration), *Phys. Rev. D* **91**, 112007 (2015).
 [12] T. E. Coan *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **96**, 162003 (2006).
 [13] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **114**, 092003 (2015).
 [14] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **118**, 092001 (2017).
 [15] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **118**, 092002 (2017).
 [16] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **124**, 032002 (2020).
 [17] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **104**, L091104 (2021).
 [18] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **11** (2023) 228.
 [19] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **05** (2024) 022.
 [20] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **07** (2024) 258.
 [21] H. X. Chen, W. Chen, X. Liu, and S. L. Zhu, *Phys. Rep.* **639**, 1 (2016).
 [22] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo, and C. Z. Yuan, *Phys. Rep.* **873**, 1 (2020).
 [23] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **76**, 111105(R) (2007).
 [24] G. Pakhlova *et al.* (Belle Collaboration), *Phys. Rev. D* **77**, 011103(R) (2008).
 [25] D. Cronin-Hennessy *et al.* (CLEO Collaboration), *Phys. Rev. D* **80**, 072001 (2009).
 [26] Y. J. Zhang and Q. Zhao, *Phys. Rev. D* **81**, 034011 (2010).
 [27] M. L. Du, U. G. Meißner, and Q. Wang, *Phys. Rev. D* **94**, 096006 (2016).
 [28] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **39**, 093001 (2015).
 [29] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 113002 (2022).
 [30] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **41**, 063001 (2017).
 [31] Y. F. Wang, *Int. J. Mod. Phys. A* **21**, 5371 (2006).
 [32] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
 [33] S. Jadach, B. F. L. Ward, and Z. Was, *Comput. Phys. Commun.* **130**, 260 (2000).
 [34] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001).
 [35] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
 [36] R. G. Ping, *Chin. Phys. C* **32**, 599 (2008).
 [37] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **95**, 072010 (2017).

- [38] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
 [39] J. Allison *et al.*, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
 [40] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **45**, 103001 (2021).
 [41] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.133.081901> for a summary of luminosity, detection efficiency, ISR correction factor, vacuum polarization factor, number of signal events, and Born cross section at each energy point, as well as the fit to the dressed cross section, which includes Refs. [2,42–44].
 [42] Z. Y. Lin, J. Z. Wang, J. B. Cheng, L. Meng, and S. L. Zhu, [arXiv:2403.01727](https://arxiv.org/abs/2403.01727).
 [43] N. Hüskén, R. F. Lebed, R. E. Mitchell, E. S. Swanson, Y. Q. Wang, and C. Z. Yuan, *Phys. Rev. D* **109**, 114010 (2024).
 [44] S. G. Salmikov and A. I. Milstein, *Phys. Rev. D* **109**, 114015 (2024).
 [45] S. Actis *et al.*, *Eur. Phys. J. C* **66**, 585 (2010).
 [46] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001).
 [47] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **101**, 112008 (2020).
 [48] J. Z. Wang, D. Y. Chen, X. Liu, and T. Matsuki, *Phys. Rev. D* **99**, 114003 (2019).
 [49] J. Z. Wang, R. Q. Qian, X. Liu, and T. Matsuki, *Phys. Rev. D* **101**, 034001 (2020).
 [50] F. J. Llanes-Estrada, *Phys. Rev. D* **72**, 031503(R) (2005).
 [51] B. Q. Li and K. T. Chao, *Phys. Rev. D* **79**, 094004 (2009).
 [52] Q. F. Cao, H. R. Qi, G. Y. Tang *et al.*, *Eur. Phys. J. C* **81**, 83 (2010).

M. Ablikim,¹ M. N. Achasov,^{5,b} P. Adlarson,⁷⁵ X. C. Ai,⁸¹ R. Aliberti,³⁶ A. Amoroso,^{74a,74c} M. R. An,⁴⁰ Q. An,^{71,58} Y. Bai,⁵⁷ O. Bakina,³⁷ I. Balossino,^{30a} Y. Ban,^{47,g} V. Batozskaya,^{1,45} K. Begzsuren,³³ N. Berger,³⁶ M. Berlowski,⁴⁵ M. Bertani,^{29a} D. Bettoni,^{30a} F. Bianchi,^{74a,74c} E. Bianco,^{74a,74c} A. Bortone,^{74a,74c} I. Boyko,³⁷ R. A. Briere,⁶ A. Brueggemann,⁶⁸ H. Cai,⁷⁶ X. Cai,^{1,58} A. Calcaterra,^{29a} G. F. Cao,^{1,63} N. Cao,^{1,63} S. A. Cetin,^{62a} J. F. Chang,^{1,58} T. T. Chang,⁷⁷ W. L. Chang,^{1,63} G. R. Che,⁴⁴ G. Chelkov,^{37,a} C. Chen,⁴⁴ Chao Chen,⁵⁵ G. Chen,¹ H. S. Chen,^{1,63} M. L. Chen,^{1,58,63} S. J. Chen,⁴³ S. L. Chen,⁴⁶ S. M. Chen,⁶¹ T. Chen,^{1,63} X. R. Chen,^{32,63} X. T. Chen,^{1,63} Y. B. Chen,^{1,58} Y. Q. Chen,³⁵ Z. J. Chen,^{26,h} W. S. Cheng,^{74c} S. K. Choi,¹¹ X. Chu,⁴⁴ G. Cibinetto,^{30a} S. C. Coen,⁴ F. Cossio,^{74c} J. J. Cui,⁵⁰ H. L. Dai,^{1,58} J. P. Dai,⁷⁹ A. Dbeyssi,¹⁹ R. E. de Boer,⁴ D. Dedovich,³⁷ Z. Y. Deng,¹ A. Denig,³⁶ I. Denysenko,³⁷ M. Destefanis,^{74a,74c} F. De Mori,^{74a,74c} B. Ding,^{66,1} X. X. Ding,^{47,g} Y. Ding,³⁵ Y. Ding,⁴¹ J. Dong,^{1,58} L. Y. Dong,^{1,63} M. Y. Dong,^{1,58,63} X. Dong,⁷⁶ M. C. Du,¹ S. X. Du,⁸¹ Z. H. Duan,⁴³ P. Egorov,^{37,a} Y. H. Fan,⁴⁶ J. Fang,^{1,58} S. S. Fang,^{1,63} W. X. Fang,¹ Y. Fang,¹ R. Farinelli,^{30a} L. Fava,^{74b,74c} F. Feldbauer,⁴ G. Felici,^{29a} C. Q. Feng,^{71,58} J. H. Feng,⁵⁹ K. Fischer,⁶⁹ M. Fritsch,⁴ C. D. Fu,¹ J. L. Fu,⁶³ Y. W. Fu,¹ H. Gao,⁶³ Y. N. Gao,^{47,g} Yang Gao,^{71,58} S. Garbolino,^{74c} I. Garzia,^{30a,30b} P. T. Ge,⁷⁶ Z. W. Ge,⁴³ C. Geng,⁵⁹ E. M. Gersabeck,⁶⁷ A. Gilman,⁶⁹ K. Goetzen,¹⁴ L. Gong,⁴¹ W. X. Gong,^{1,58} W. Gradl,³⁶ S. Gramigna,^{30a,30b} M. Greco,^{74a,74c} M. H. Gu,^{1,58} Y. T. Gu,¹⁶ C. Y. Guan,^{1,63} Z. L. Guan,²³ A. Q. Guo,^{32,63} L. B. Guo,⁴² M. J. Guo,⁵⁰ R. P. Guo,⁴⁹ Y. P. Guo,^{13,f} A. Guskov,^{37,a} T. T. Han,⁵⁰ W. Y. Han,⁴⁰ X. Q. Hao,²⁰ F. A. Harris,⁶⁵ K. K. He,⁵⁵ K. L. He,^{1,63} F. H. H. Heinsius,⁴ C. H. Heinz,³⁶ Y. K. Heng,^{1,58,63} C. Herold,⁶⁰ T. Holtmann,⁴ P. C. Hong,^{13,f} G. Y. Hou,^{1,63} X. T. Hou,^{1,63} Y. R. Hou,⁶³ Z. L. Hou,¹ H. M. Hu,^{1,63} J. F. Hu,^{56,i} T. Hu,^{1,58,63} Y. Hu,¹ G. S. Huang,^{71,58} K. X. Huang,⁵⁹ L. Q. Huang,^{32,63} X. T. Huang,⁵⁰ Y. P. Huang,¹ T. Hussain,⁷³ N. Hüskén,^{28,36} N. in der Wiesche,⁶⁸ M. Irshad,^{71,58} J. Jackson,²⁸ S. Jaeger,⁴ S. Janchiv,³³ J. H. Jeong,¹¹ Q. Ji,¹ Q. P. Ji,²⁰ X. B. Ji,^{1,63} X. L. Ji,^{1,58} Y. Y. Ji,⁵⁰ X. Q. Jia,⁵⁰ Z. K. Jia,^{71,58} H. J. Jiang,⁷⁶ P. C. Jiang,^{47,g} S. S. Jiang,⁴⁰ T. J. Jiang,¹⁷ X. S. Jiang,^{1,58,63} Y. Jiang,⁶³ J. B. Jiao,⁵⁰ Z. Jiao,²⁴ S. Jin,⁴³ Y. Jin,⁶⁶ M. Q. Jing,^{1,63} T. Johansson,⁷⁵ X. Kui,¹ S. Kabana,³⁴ N. Kalantar-Nayestanaki,⁶⁴ X. L. Kang,¹⁰ X. S. Kang,⁴¹ M. Kavatsyuk,⁶⁴ B. C. Ke,⁸¹ A. Khoukaz,⁶⁸ R. Kiuchi,¹ R. Kliemt,¹⁴ O. B. Kolcu,^{62a} B. Kopf,⁴ M. Kuessner,⁴ A. Kupsc,^{45,75} W. Kühn,³⁸ J. J. Lane,⁶⁷ P. Larin,¹⁹ A. Lavanaia,²⁷ L. Lavezzi,^{74a,74c} T. T. Lei,^{71,58} Z. H. Lei,^{71,58} H. Leithoff,³⁶ M. Lellmann,³⁶ T. Lenz,³⁶ C. Li,⁴⁴ C. Li,⁴⁸ C. H. Li,⁴⁰ Cheng Li,^{71,58} D. M. Li,⁸¹ F. Li,^{1,58} G. Li,¹ H. Li,^{71,58} H. B. Li,^{1,63} H. J. Li,²⁰ H. N. Li,^{56,i} Hui Li,⁴⁴ J. R. Li,⁶¹ J. S. Li,⁵⁹ J. W. Li,⁵⁰ K. L. Li,²⁰ Ke Li,¹ L. J. Li,^{1,63} L. K. Li,¹ Lei Li,³ M. H. Li,⁴⁴ P. R. Li,^{39,j,k} Q. X. Li,⁵⁰ S. X. Li,¹³ T. Li,⁵⁰ W. D. Li,^{1,63} W. G. Li,¹ X. H. Li,^{71,58} X. L. Li,⁵⁰ Xiaoyu Li,^{1,63} Y. G. Li,^{47,g} Z. J. Li,⁵⁹ Z. X. Li,¹⁶ C. Liang,⁴³ H. Liang,^{1,63} H. Liang,³⁵ H. Liang,^{71,58} Y. F. Liang,⁵⁴ Y. T. Liang,^{32,63} G. R. Liao,¹⁵ L. Z. Liao,⁵⁰ Y. P. Liao,^{1,63} J. Libby,²⁷ A. Limphirat,⁶⁰ D. X. Lin,^{32,63} T. Lin,¹ B. J. Liu,¹ B. X. Liu,⁷⁶ C. Liu,³⁵ C. X. Liu,¹ F. H. Liu,⁵³ Fang Liu,¹ Feng Liu,⁷ G. M. Liu,^{56,i} H. Liu,^{39,j,k} H. B. Liu,¹⁶ H. M. Liu,^{1,63} Huanhuan Liu,¹ Huihui Liu,²² J. B. Liu,^{71,58} J. L. Liu,⁷² J. Y. Liu,^{1,63} K. Liu,¹ K. Y. Liu,⁴¹ Ke Liu,²³ L. Liu,^{71,58} L. C. Liu,⁴⁴ Lu Liu,⁴⁴ M. H. Liu,^{13,f} P. L. Liu,¹ Q. Liu,⁶³ S. B. Liu,^{71,58} T. Liu,^{13,f} W. K. Liu,⁴⁴ W. M. Liu,^{71,58} X. Liu,^{39,j,k} Y. Liu,^{39,j,k} Y. Liu,⁸¹ Y. B. Liu,⁴⁴ Z. A. Liu,^{1,58,63} Z. Q. Liu,⁵⁰ X. C. Lou,^{1,58,63} F. X. Lu,⁵⁹ H. J. Lu,²⁴ J. G. Lu,^{1,58} X. L. Lu,¹ Y. Lu,⁸ Y. P. Lu,^{1,58} Z. H. Lu,^{1,63} C. L. Luo,⁴² M. X. Luo,⁸⁰ T. Luo,^{13,f} X. L. Luo,^{1,58} X. R. Lyu,⁶³ Y. F. Lyu,⁴⁴ F. C. Ma,⁴¹ H. L. Ma,¹ J. L. Ma,^{1,63} L. L. Ma,⁵⁰ M. M. Ma,^{1,63} Q. M. Ma,¹

R. Q. Ma,^{1,63} R. T. Ma,⁶³ X. Y. Ma,^{1,58} Y. Ma,^{47,g} Y. M. Ma,³² F. E. Maas,¹⁹ M. Maggiora,^{74a,74c} S. Malde,⁶⁹ Q. A. Malik,⁷³
 A. Mangoni,^{29b} Y. J. Mao,^{47,g} Z. P. Mao,¹ S. Marcello,^{74a,74c} Z. X. Meng,⁶⁶ J. G. Messchendorp,^{14,64} G. Mezzadri,^{30a}
 H. Miao,^{1,63} T. J. Min,⁴³ R. E. Mitchell,²⁸ X. H. Mo,^{1,58,63} N. Yu. Muchnoi,^{5,b} J. Muskalla,³⁶ Y. Nefedov,³⁷ F. Nerling,^{19,d}
 I. B. Nikolaev,^{5,b} Z. Ning,^{1,58} S. Nisar,^{12,l} Q. L. Niu,^{39,j,k} W. D. Niu,⁵⁵ Y. Niu,⁵⁰ S. L. Olsen,⁶³ Q. Ouyang,^{1,58,63}
 S. Pacetti,^{29b,29c} X. Pan,⁵⁵ Y. Pan,⁵⁷ A. Pathak,³⁵ P. Patteri,^{29a} Y. P. Pei,^{71,58} M. Pelizaeus,⁴ H. P. Peng,^{71,58} Y. Y. Peng,^{39,j,k}
 K. Peters,^{14,d} J. L. Ping,⁴² R. G. Ping,^{1,63} S. Plura,³⁶ V. Prasad,³⁴ F. Z. Qi,¹ H. Qi,^{71,58} H. R. Qi,⁶¹ M. Qi,⁴³ T. Y. Qi,^{13,f}
 S. Qian,^{1,58} W. B. Qian,⁶³ C. F. Qiao,⁶³ J. J. Qin,⁷² L. Q. Qin,¹⁵ X. P. Qin,^{13,f} X. S. Qin,⁵⁰ Z. H. Qin,^{1,58} J. F. Qiu,¹ S. Q. Qu,⁶¹
 C. F. Redmer,³⁶ K. J. Ren,⁴⁰ A. Rivetti,^{74c} M. Rolo,^{74c} G. Rong,^{1,63} Ch. Rosner,¹⁹ S. N. Ruan,⁴⁴ N. Salone,⁴⁵ A. Sarantsev,^{37,c}
 Y. Schelhaas,³⁶ K. Schoenning,⁷⁵ M. Scodreggio,^{30a,30b} K. Y. Shan,^{13,f} W. Shan,²⁵ X. Y. Shan,^{71,58} J. F. Shangguan,⁵⁵
 L. G. Shao,^{1,63} M. Shao,^{71,58} C. P. Shen,^{13,f} H. F. Shen,^{1,63} W. H. Shen,⁶³ X. Y. Shen,^{1,63} B. A. Shi,⁶³ H. C. Shi,^{71,58} J. L. Shi,¹³
 J. Y. Shi,¹ Q. Q. Shi,⁵⁵ R. S. Shi,^{1,63} X. Shi,^{1,58} J. J. Song,²⁰ T. Z. Song,⁵⁹ W. M. Song,^{35,1} Y. J. Song,¹³ Y. X. Song,^{47,g}
 S. Sosio,^{74a,74c} S. Spataro,^{74a,74c} F. Stieler,³⁶ Y. J. Su,⁶³ G. B. Sun,⁷⁶ G. X. Sun,¹ H. Sun,⁶³ H. K. Sun,¹ J. F. Sun,²⁰ K. Sun,⁶¹
 L. Sun,⁷⁶ S. S. Sun,^{1,63} T. Sun,^{1,63} W. Y. Sun,³⁵ Y. Sun,¹⁰ Y. J. Sun,^{71,58} Y. Z. Sun,¹ Z. T. Sun,⁵⁰ Y. X. Tan,^{71,58} C. J. Tang,⁵⁴
 G. Y. Tang,¹ J. Tang,⁵⁹ Y. A. Tang,⁷⁶ L. Y. Tao,⁷² Q. T. Tao,^{26,h} M. Tat,⁶⁹ J. X. Teng,^{71,58} V. Thoren,⁷⁵ W. H. Tian,⁵⁹
 W. H. Tian,⁵² Y. Tian,^{32,63} Z. F. Tian,⁷⁶ I. Uman,^{62b} S. J. Wang,⁵⁰ B. Wang,¹ B. L. Wang,⁶³ Bo Wang,^{71,58} C. W. Wang,⁴³
 D. Y. Wang,^{47,g} F. Wang,⁷² H. J. Wang,^{39,j,k} H. P. Wang,^{1,63} J. P. Wang,⁵⁰ K. Wang,^{1,58} L. L. Wang,¹ M. Wang,⁵⁰
 Meng Wang,^{1,63} S. Wang,^{13,f} S. Wang,^{39,j,k} T. Wang,^{13,f} T. J. Wang,⁴⁴ W. Wang,⁷² W. Wang,⁵⁹ W. P. Wang,^{71,58} X. Wang,^{47,g}
 X. F. Wang,^{39,j,k} X. J. Wang,⁴⁰ X. L. Wang,^{13,f} Y. Wang,⁶¹ Y. D. Wang,⁴⁶ Y. F. Wang,^{1,58,63} Y. H. Wang,⁴⁸ Y. N. Wang,⁴⁶
 Y. Q. Wang,¹ Yaqian Wang,^{18,1} Yi Wang,⁶¹ Z. Wang,^{1,58} Z. L. Wang,⁷² Z. Y. Wang,^{1,63} Ziyi Wang,⁶³ D. Wei,⁷⁰ D. H. Wei,¹⁵
 F. Weidner,⁶⁸ S. P. Wen,¹ C. W. Wenzel,⁴ U. Wiedner,⁴ G. Wilkinson,⁶⁹ M. Wolke,⁷⁵ L. Wollenberg,⁴ C. Wu,⁴⁰ J. F. Wu,^{1,63}
 L. H. Wu,¹ L. J. Wu,^{1,63} X. Wu,^{13,f} X. H. Wu,³⁵ Y. Wu,⁷¹ Y. H. Wu,⁵⁵ Y. J. Wu,³² Z. Wu,^{1,58} L. Xia,^{71,58} X. M. Xian,⁴⁰
 T. Xiang,^{47,g} D. Xiao,^{39,j,k} G. Y. Xiao,⁴³ S. Y. Xiao,¹ Y. L. Xiao,^{13,f} Z. J. Xiao,⁴² C. Xie,⁴³ X. H. Xie,^{47,g} Y. Xie,⁵⁰
 Y. G. Xie,^{1,58} Y. H. Xie,⁷ Z. P. Xie,^{71,58} T. Y. Xing,^{1,63} C. F. Xu,^{1,63} C. J. Xu,⁵⁹ G. F. Xu,¹ H. Y. Xu,⁶⁶ Q. J. Xu,¹⁷ Q. N. Xu,³¹
 W. Xu,^{1,63} W. L. Xu,⁶⁶ X. P. Xu,⁵⁵ Y. C. Xu,⁷⁸ Z. P. Xu,⁴³ Z. S. Xu,⁶³ F. Yan,^{13,f} L. Yan,^{13,f} W. B. Yan,^{71,58} W. C. Yan,⁸¹
 X. Q. Yan,¹ H. J. Yang,^{51,e} H. L. Yang,³⁵ H. X. Yang,¹ Tao Yang,¹ Y. Yang,^{13,f} Y. F. Yang,⁴⁴ Y. X. Yang,^{1,63} Yifan Yang,^{1,63}
 Z. W. Yang,^{39,j,k} Z. P. Yao,⁵⁰ M. Ye,^{1,58} M. H. Ye,⁹ J. H. Yin,¹ Z. Y. You,⁵⁹ B. X. Yu,^{1,58,63} C. X. Yu,⁴⁴ G. Yu,^{1,63} J. S. Yu,^{26,h}
 T. Yu,⁷² X. D. Yu,^{47,g} C. Z. Yuan,^{1,63} L. Yuan,² S. C. Yuan,¹ X. Q. Yuan,¹ Y. Yuan,^{1,63} Z. Y. Yuan,⁵⁹ C. X. Yue,⁴⁰
 A. A. Zafar,⁷³ F. R. Zeng,⁵⁰ X. Zeng,^{13,f} Y. Zeng,^{26,h} Y. J. Zeng,^{1,63} X. Y. Zhai,³⁵ Y. C. Zhai,⁵⁰ Y. H. Zhan,⁵⁹ A. Q. Zhang,^{1,63}
 B. L. Zhang,^{1,63} B. X. Zhang,¹ D. H. Zhang,⁴⁴ G. Y. Zhang,²⁰ H. Zhang,⁷¹ H. C. Zhang,^{1,58,63} H. H. Zhang,⁵⁹ H. H. Zhang,³⁵
 H. Q. Zhang,^{1,58,63} H. Y. Zhang,^{1,58} J. Zhang,⁸¹ J. J. Zhang,⁵² J. L. Zhang,²¹ J. Q. Zhang,⁴² J. W. Zhang,^{1,58,63} J. X. Zhang,^{39,j,k}
 J. Y. Zhang,¹ J. Z. Zhang,^{1,63} Jianyu Zhang,⁶³ Jiawei Zhang,^{1,63} L. M. Zhang,⁶¹ L. Q. Zhang,⁵⁹ Lei Zhang,⁴³ P. Zhang,^{1,63}
 Q. Y. Zhang,^{40,81} Shuihan Zhang,^{1,63} Shulei Zhang,^{26,h} X. D. Zhang,⁴⁶ X. M. Zhang,¹ X. Y. Zhang,⁵⁰ Xuyan Zhang,⁵⁵
 Y. Zhang,⁶⁹ Y. Zhang,⁷² Y. T. Zhang,⁸¹ Y. H. Zhang,^{1,58} Yan Zhang,^{71,58} Yao Zhang,¹ Z. H. Zhang,¹ Z. L. Zhang,³⁵
 Z. Y. Zhang,⁴⁴ Z. Y. Zhang,⁷⁶ G. Zhao,¹ J. Zhao,⁴⁰ J. Y. Zhao,^{1,63} J. Z. Zhao,^{1,58} Lei Zhao,^{71,58} Ling Zhao,¹ M. G. Zhao,⁴⁴
 S. J. Zhao,⁸¹ Y. B. Zhao,^{1,58} Y. X. Zhao,^{32,63} Z. G. Zhao,^{71,58} A. Zhemchugov,^{37,a} B. Zheng,⁷² J. P. Zheng,^{1,58} W. J. Zheng,^{1,63}
 Y. H. Zheng,⁶³ B. Zhong,⁴² X. Zhong,⁵⁹ H. Zhou,⁵⁰ L. P. Zhou,^{1,63} X. Zhou,⁷⁶ X. K. Zhou,⁷ X. R. Zhou,^{71,58} X. Y. Zhou,⁴⁰
 Y. Z. Zhou,^{13,f} J. Zhu,⁴⁴ K. Zhu,¹ K. J. Zhu,^{1,58,63} L. Zhu,³⁵ L. X. Zhu,⁶³ S. H. Zhu,⁷⁰ S. Q. Zhu,⁴³ T. J. Zhu,^{13,f} W. J. Zhu,^{13,f}
 Y. C. Zhu,^{71,58} Z. A. Zhu,^{1,63} J. H. Zou,¹ and J. Zu^{71,58}

(BESIII Collaboration)

¹Institute of High Energy Physics, Beijing 100049, People's Republic of China

²Beihang University, Beijing 100191, People's Republic of China

³Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China

⁴Bochum Ruhr-University, D-44780 Bochum, Germany

⁵Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia

⁶Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

⁷Central China Normal University, Wuhan 430079, People's Republic of China

⁸Central South University, Changsha 410083, People's Republic of China

⁹China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China

- ¹⁰*China University of Geosciences, Wuhan 430074, People's Republic of China*
- ¹¹*Chung-Ang University, Seoul 06974, Republic of Korea*
- ¹²*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*
- ¹³*Fudan University, Shanghai 200433, People's Republic of China*
- ¹⁴*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*
- ¹⁵*Guangxi Normal University, Guilin 541004, People's Republic of China*
- ¹⁶*Guangxi University, Nanning 530004, People's Republic of China*
- ¹⁷*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*
- ¹⁸*Hebei University, Baoding 071002, People's Republic of China*
- ¹⁹*Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*
- ²⁰*Henan Normal University, Xinxiang 453007, People's Republic of China*
- ²¹*Henan University, Kaifeng 475004, People's Republic of China*
- ²²*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*
- ²³*Henan University of Technology, Zhengzhou 450001, People's Republic of China*
- ²⁴*Huangshan College, Huangshan 245000, People's Republic of China*
- ²⁵*Hunan Normal University, Changsha 410081, People's Republic of China*
- ²⁶*Hunan University, Changsha 410082, People's Republic of China*
- ²⁷*Indian Institute of Technology Madras, Chennai 600036, India*
- ²⁸*Indiana University, Bloomington, Indiana 47405, USA*
- ^{29a}*INFN Laboratori Nazionali di Frascati, INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*
- ^{29b}*INFN Sezione di Perugia, I-06100 Perugia, Italy*
- ^{29c}*University of Perugia, I-06100 Perugia, Italy*
- ^{30a}*INFN Sezione di Ferrara, INFN Sezione di Ferrara, I-44122 Ferrara, Italy*
- ^{30b}*University of Ferrara, I-44122 Ferrara, Italy*
- ³¹*Inner Mongolia University, Hohhot 010021, People's Republic of China*
- ³²*Institute of Modern Physics, Lanzhou 730000, People's Republic of China*
- ³³*Institute of Physics and Technology, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia*
- ³⁴*Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile*
- ³⁵*Jilin University, Changchun 130012, People's Republic of China*
- ³⁶*Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
- ³⁷*Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*
- ³⁸*Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*
- ³⁹*Lanzhou University, Lanzhou 730000, People's Republic of China*
- ⁴⁰*Liaoning Normal University, Dalian 116029, People's Republic of China*
- ⁴¹*Liaoning University, Shenyang 110036, People's Republic of China*
- ⁴²*Nanjing Normal University, Nanjing 210023, People's Republic of China*
- ⁴³*Nanjing University, Nanjing 210093, People's Republic of China*
- ⁴⁴*Nankai University, Tianjin 300071, People's Republic of China*
- ⁴⁵*National Centre for Nuclear Research, Warsaw 02-093, Poland*
- ⁴⁶*North China Electric Power University, Beijing 102206, People's Republic of China*
- ⁴⁷*Peking University, Beijing 100871, People's Republic of China*
- ⁴⁸*Qufu Normal University, Qufu 273165, People's Republic of China*
- ⁴⁹*Shandong Normal University, Jinan 250014, People's Republic of China*
- ⁵⁰*Shandong University, Jinan 250100, People's Republic of China*
- ⁵¹*Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China*
- ⁵²*Shanxi Normal University, Linfen 041004, People's Republic of China*
- ⁵³*Shanxi University, Taiyuan 030006, People's Republic of China*
- ⁵⁴*Sichuan University, Chengdu 610064, People's Republic of China*
- ⁵⁵*Soochow University, Suzhou 215006, People's Republic of China*
- ⁵⁶*South China Normal University, Guangzhou 510006, People's Republic of China*
- ⁵⁷*Southeast University, Nanjing 211100, People's Republic of China*
- ⁵⁸*State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China*
- ⁵⁹*Sun Yat-Sen University, Guangzhou 510275, People's Republic of China*
- ⁶⁰*Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand*
- ⁶¹*Tsinghua University, Beijing 100084, People's Republic of China*
- ^{62a}*Turkish Accelerator Center Particle Factory Group, Istinye University, 34010 Istanbul, Turkey*
- ^{62b}*Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey*
- ⁶³*University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China*
- ⁶⁴*University of Groningen, NL-9747 AA Groningen, The Netherlands*
- ⁶⁵*University of Hawaii, Honolulu, Hawaii 96822, USA*

⁶⁶*University of Jinan, Jinan 250022, People's Republic of China*

⁶⁷*University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom*

⁶⁸*University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany*

⁶⁹*University of Oxford, Keble Road, Oxford OX13RH, United Kingdom*

⁷⁰*University of Science and Technology Liaoning, Anshan 114051, People's Republic of China*

⁷¹*University of Science and Technology of China, Hefei 230026, People's Republic of China*

⁷²*University of South China, Hengyang 421001, People's Republic of China*

⁷³*University of the Punjab, Lahore-54590, Pakistan*

^{74a}*University of Turin and INFN, University of Turin, I-10125 Turin, Italy*

^{74b}*University of Eastern Piedmont, I-15121 Alessandria, Italy*

^{74c}*INFN, I-10125 Turin, Italy*

⁷⁵*Uppsala University, Box 516, SE-75120 Uppsala, Sweden*

⁷⁶*Wuhan University, Wuhan 430072, People's Republic of China*

⁷⁷*Xinyang Normal University, Xinyang 464000, People's Republic of China*

⁷⁸*Yantai University, Yantai 264005, People's Republic of China*

⁷⁹*Yunnan University, Kunming 650500, People's Republic of China*

⁸⁰*Zhejiang University, Hangzhou 310027, People's Republic of China*

⁸¹*Zhengzhou University, Zhengzhou 450001, People's Republic of China*

^aAlso at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

^bAlso at the Novosibirsk State University, Novosibirsk, 630090, Russia.

^cAlso at the NRC "Kurchatov Institute," PNPI, 188300, Gatchina, Russia.

^dAlso at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

^eAlso at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China.

^fAlso at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.

^gAlso at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China.

^hAlso at School of Physics and Electronics, Hunan University, Changsha 410082, China.

ⁱAlso at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.

^jAlso at MOE Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People's Republic of China.

^kAlso at Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province, and Key Laboratory for Quantum Theory and Applications of MoE, Lanzhou University, Lanzhou 730000, People's Republic of China.

^lAlso at the Department of Mathematical Sciences, IBA, Karachi 75270, Pakistan.