

University of Groningen

Macroglial diversity and its effect on myelination

Werkman, Inge

DOI:
[10.33612/diss.113508108](https://doi.org/10.33612/diss.113508108)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Werkman, I. (2020). *Macroglial diversity and its effect on myelination*. Rijksuniversiteit Groningen.
<https://doi.org/10.33612/diss.113508108>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Stellingen

1. The finding that remyelination is more effective in grey matter than white matter tissue in both human MS and rodent remyelination models indicates an evolutionary conserved and robust background¹⁻⁶.
2. The contribution of the tight intertwining of lipid homeostasis and the immune system in astrocytes to MS pathology is currently underestimated⁷.
3. The identification of an immunological phenotype in MS oligodendroglial cells based on immunohistochemistry, while this subclass is not apparent in transcriptomic studies, indicates that single cell proteomic studies might be more valuable compared to transcriptomic studies^{8,9}.
4. To prevent “p-hacking” and the publication of significant -but clinically irrelevant- data, science should convert the archaic probability system to confidence-intervals and include relevant effect sizes¹⁰.
5. Improved understanding of regional heterogeneity in macroglial cells will contribute to MS research as a whole, and may open therapeutic avenues aimed at enhancing remyelination. (this thesis)
6. When studying individual parts of a vacuum cleaner, one will not obtain understanding of the mechanism of suction. Likewise, a pitfall of cell biology research is over-specification, thereby overlooking the bigger picture.
7. Although the study of single cell types is definitely interesting and relevant for the understanding of cell functioning, the study of interactions between different cell types is at least as important for the understanding of pathology.
8. Astrocytes are the real stars of the central nervous system.

“What’s so great about discovery? It’s a violent, penetrative act that scars whatever it explores.” - Ian Malcolm, Jurassic Park

References

1. Gudi, V. *et al.* Regional differences between grey and white matter in cuprizone induced demyelination. *Brain Res.* **1283**, 127–138 (2009).
2. Gudi, V. *et al.* Spatial and temporal profiles of growth factor expression during CNS demyelination reveal the dynamics of repair priming. *PLoS One* **6**, e22623 (2011).
3. Bai, C. B. *et al.* A mouse model for testing remyelinating therapies. *Exp. Neurol.* **283**, 330–340 (2016).
4. Chang, A. *et al.* Cortical remyelination: A new target for repair therapies in multiple sclerosis. *Ann. Neurol.* **72**, 918–926 (2012).
5. Albert, M., Antel, J., Brück, W. & Stadelmann, C. Extensive cortical remyelination in patients with chronic multiple sclerosis. *Brain Pathol.* **17**, 129–138 (2007).
6. Strijbis, E. M. M., Kooi, E.-J., van der Valk, P. & Geurts, J. J. G. Cortical remyelination is heterogeneous in multiple sclerosis. *J. Neuropathol. Exp. Neurol.* **76**, 390–401 (2017).
7. Kihara, Y. Systematic understanding of bioactive lipids in neuro-immune interactions: Lessons from an animal model of multiple sclerosis. in *Advances in Experimental Medicine and Biology* **1161**, 133–148 (Springer New York LLC, 2019).
8. Falcão, A. M. *et al.* Disease-specific oligodendrocyte lineage cells arise in multiple sclerosis. *Nat. Med.* **24**, 1837–1844 (2018).
9. Jäkel, S. *et al.* Altered human oligodendrocyte heterogeneity in multiple sclerosis. *Nature* **566**, 543–547 (2019).
10. Schober, P., Bossers, S. M. & Schwarte, L. A. Statistical significance versus clinical importance of observed effect sizes: What do p values and confidence intervals really represent? *Anesth. Analg.* **126**, 1068–1072 (2018).