

## University of Groningen

### Genotype correlates with the natural history of severe bile salt export pump deficiency

van Wessel, Daan B E; Thompson, Richard J; Gonzales, Emmanuel; Jankowska, Irena; Sokal, Etienne; Grammatikopoulos, Tassos; Kadaristiana, Agustina; Jacquemin, Emmanuel; Spraul, Anne; Lipiński, Patryk

*Published in:*  
Journal of Hepatology

*DOI:*  
[10.1016/j.jhep.2020.02.007](https://doi.org/10.1016/j.jhep.2020.02.007)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Final author's version (accepted by publisher, after peer review)

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

van Wessel, D. B. E., Thompson, R. J., Gonzales, E., Jankowska, I., Sokal, E., Grammatikopoulos, T., Kadaristiana, A., Jacquemin, E., Spraul, A., Lipiński, P., Czubkowski, P., Rock, N., Shagrani, M., Broering, D., Algoufi, T., Mazhar, N., Nicastro, E., Kelly, D., Nebbia, G., ... Verkade, H. J. (2020). Genotype correlates with the natural history of severe bile salt export pump deficiency. *Journal of Hepatology*, 73(1), 84-93. <https://doi.org/10.1016/j.jhep.2020.02.007>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## Genotype Correlates to the Natural History of Severe Bile Salt Export Pump Deficiency - Results from the NAPPED Consortium

Daan B.E. van Wessel<sup>1</sup>, Richard J. Thompson<sup>2</sup>, Emmanuel Gonzales<sup>3,4</sup>, Irena Jankowska<sup>4,5</sup>, Etienne Sokal<sup>4,6</sup>, Tassos Grammatikopoulos<sup>2</sup>, Agustina Kadaristiana<sup>2</sup>, Emmanuel Jacquemin<sup>3</sup>, Anne Spraul<sup>3b</sup>, Patryk Lipiński<sup>4,5</sup>, Piotr Czubkowski<sup>4,5</sup>, Nathalie Rock<sup>6</sup>, Mohammad Shagrani<sup>7,7b</sup>, Dieter Broering<sup>7</sup>, Talal Algoufi<sup>7</sup>, Nejat Mazhar<sup>7</sup>, Emanuele Nicastro<sup>8</sup>, Deirdre Kelly<sup>4,9</sup>, Gabriella Nebbia<sup>10</sup>, Henrik Arnell<sup>4,11</sup>, Björn Fischler<sup>4,11</sup>, Jan B.F. Hulscher<sup>4,12</sup>, Daniele Serranti<sup>13</sup>, Cigdem Arikan<sup>14</sup>, Esra Polat<sup>15</sup>, Dominique Debray<sup>16</sup>, Florence Lacaille<sup>16</sup>, Cristina Goncalves<sup>4,17</sup>, Loreto Hierro<sup>4,18</sup>, Gema Muñoz Bartolo<sup>4,18</sup>, Yael Mozer-Glassberg<sup>19</sup>, Amer Azaz<sup>20</sup>, Jernej Breclj<sup>21</sup>, Antal Dezsöfi<sup>22</sup>, Pier Luigi Calvo<sup>23</sup>, Enke Grabhorn<sup>24</sup>, Ekkehard Sturm<sup>4,25</sup>, Wendy J. van der Woerd<sup>26</sup>, Binita M. Kamath<sup>27</sup>, Jian-She Wang<sup>28</sup>, Liting Li<sup>28</sup>, Özlem Durmaz<sup>29</sup>, Zerrin Onal<sup>29</sup>, Ton MG Bunt<sup>1</sup>, Bettina Hansen<sup>30,30b</sup>, Henkjan J. Verkade<sup>1,4</sup>

<sup>1</sup>Pediatric Gastroenterology and Hepatology, University Medical Centre Groningen, University of Groningen, The Netherlands, <sup>2</sup>King's College London, United Kingdom, <sup>3</sup>Service d'Hépatologie et de Transplantation Hépatique Pédiatriques, Bicêtre Hôpital, AP-HP, Université Paris-Sud, Paris Saclay, Inserm UMR-S 1174, France, <sup>3b</sup>Service de Biochimie, Bicêtre Hôpital, AP-HP, Université Paris-Sud, Paris Saclay, Inserm UMR-S 1174, France, <sup>4</sup>European Reference Network on Hepatological Diseases (ERN RARE-LIVER), <sup>5</sup>Gastroenterology, Hepatology, Nutritional Disorders and Paediatrics, The Children's Memorial Health Institute, Warsaw, Poland, <sup>6</sup>Université Catholique de Louvain, Cliniques St Luc, Brussels, Belgium, <sup>7</sup>Liver & SB Transplant & Hepatobiliary-Pancreatic Surgery, King Faisal Specialist Hospital & Research Centre, Riyadh, Saudi Arabia, <sup>7b</sup>Alfaisal University, College of Medicine, Riyadh, Saudi Arabia, <sup>8</sup>Pediatric Hepatology, Gastroenterology and Transplantation, Ospedale Papa Giovanni XXIII, Bergamo, Italy, <sup>9</sup>Liver Unit, Birmingham Women's and Children's Hospital, Birmingham, United Kingdom, <sup>10</sup>Servizio Di Epatologia e Nutrizione Pediatrica, Fondazione Irccs Ca' Granda Ospedale Maggiore Policlinico, Milano, Italy, <sup>11</sup>Pediatric Digestive Diseases, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Karolinska Institutet, Stockholm, Sweden <sup>12</sup>Paediatric Surgery, University Medical Centre Groningen, Groningen, The Netherlands, <sup>13</sup>Paediatric and Liver Unit, Meyer Children's University Hospital of Florence, <sup>14</sup>Koc University School of Medicine, Paediatric GI and Hepatology Liver Transplantation Centre, Kuttam System in Liver Medicine, Istanbul Turkey, <sup>15</sup>Hospital Umraniye Training and Research Hospital Istanbul, <sup>16</sup>Unité d'hépatologie Pédiatrique et Transplantation, Hôpital Necker, Paris, France <sup>17</sup>Coimbra University Hospital Center, Coimbra, Portugal, <sup>18</sup>Pediatric Liver Service, La Paz University Hospital, Madrid, Spain, <sup>19</sup>Institute of Gastroenterology, Nutrition and Liver Diseases, Schneider Children's Medical Centre of Israel, <sup>20</sup>Sheikh Khalifa Medical City, Abu Dhabi, United Arab Emirates, <sup>21</sup>Department of Gastroenterology, Hepatology and Nutrition, University Children's Hospital Ljubljana, and Department of Paediatrics, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia, <sup>22</sup>1st Department of Paediatrics, Semmelweis University, Budapest, Hungary, <sup>23</sup>Pediatric Gastroenterology Unit, Regina Margherita Children's Hospital, Azienda Ospedaliera Città Della Salute e Della Scienza University Hospital, Torino, Italy, <sup>24</sup>Klinik Für Kinder- Und Jugendmedizin, Universitätsklinikum Hamburg Eppendorf, Hamburg, Germany, <sup>25</sup>University Children's Hospital Tübingen, Tübingen, Germany, <sup>26</sup>Wilhelmina Children's Hospital, University Medical Centre Utrecht, Paediatric Gastroenterology, Hepatology and Nutrition, Utrecht, The Netherlands, <sup>27</sup>The Hospital for Sick Children and the University of Toronto, Toronto, Canada <sup>28</sup>Children's Hospital of Fudan University, Shanghai, China, <sup>29</sup>Istanbul University, Istanbul Faculty of Medicine, Istanbul, Turkey, <sup>30</sup>Toronto Centre for Liver Disease, University Health Network, Canada <sup>30b</sup>IHPME, University of Toronto, Canada

**Abbreviations:** ABCB11 (ATP-binding cassette, sub-family B member 11), ATP8B1 (ATPase class I type 8B member 1), ALT (alanine-aminotransferase), AST (aspartate-aminotransferase), ATP (Adenosine triphosphate), BSEP (bile salt export pump), DFS (diversion-free survival), GGT (gamma-glutamyltranspeptidase), HCC (hepatocellular carcinoma), IE (ileal exclusion), LTx (liver transplantation), NAPPED (NAtural course and Prognosis of PFIC and Effect of biliary Diversion), NLS (native liver survival), PEBD (partial external biliary diversion), PFIC (Progressive Familial Intrahepatic Cholestasis), PLT (platelet count), REDCap (Research Electronic Data Capture), sBA (serum bile acids), SBD (surgical biliary diversion), TJP2 (tight junction protein 2), TSB (total serum bilirubin), UDCA (ursodeoxycholic acid).

**Corresponding author:** Henkjan J. Verkade MD, PhD, Paediatric Gastroenterology & Hepatology, Department of Paediatrics, Beatrix Children's Hospital, University Medical Centre Groningen, University of Groningen, PO Box 30.001, 9700 RB Groningen, The Netherlands, e-mail: h.j.verkade@umcg.nl

**Keywords:** severe BSEP deficiency; PFIC2; ABCB11; natural history; surgical biliary diversion.

**Electronic word count:** 6000

**Number of figures and tables:** 7

**Conflicts of interest:** Daan B.E. van Wessel: [nothing to disclose], Richard J. Thompson [consultancy for Shire, Albireo, Mirum, Horizon Pharmaceuticals, Sana Biotechnology, GenerationBio, Retrophin and Qing Bile Therapeutics], Emmanuel M. Gonzales [Consultant for CTRS and Mirum Pharmaceuticals], Irena Jankowska [nothing to disclose], Tassos Grammatikopoulos [nothing to disclose], Agustina Kadaristiana [nothing to disclose], Patryk Lipiński [nothing to disclose], Piotr Czubkowski [nothing to disclose], Nathalie Rock [nothing to disclose] Emmanuel Jacquemin [nothing to disclose], Anne Spraul [nothing to disclose], Etienne M. Sokal [founder and CSMO of Promethera Biosciences], Mohammad Shagrani [nothing to disclose], Dieter Broering [nothing to disclose], Talal Algoufi [nothing to disclose], Nejat Mazhar [nothing to disclose], Emanuele Nicastro [nothing to disclose] Deirdre Kelly [Consultant for Albireo], Gabriela Nebbia [nothing to disclose], Henrik Arnell [consultant for Albireo and Mirum Pharmaceuticals], Björn Fischler [has attended one advisory board meeting with Albireo in 2016], Jan Hulscher [nothing to disclose], Daniele Serranti [nothing to disclose], Cigdem Arikan [nothing to disclose], Esra Polat [nothing to disclose], Dominique Debray [consultant for Alexion pharmaceuticals], Florence Lacaille [nothing to disclose], Cristina Goncalves [nothing to disclose], Loreto Hierro [nothing to disclose], Gema Muñoz Bartolo [nothing to disclose], Yael Mozer-Glassberg [nothing to disclose], Amer Azaz [nothing to disclose], Jernej Brecelj [nothing to disclose], Antal Dezsófi [nothing to disclose], Pier Luigi Calvo [nothing to disclose], Enke Grabhorn [nothing to disclose], Ekkehard Sturm [nothing to disclose] Wendy van der Woerd [nothing to disclose], Binita Kamath [consultant for Mirum Pharmaceuticals, Shire and DCI], Jian-She Wang [nothing to disclose], Liting Li [nothing to disclose], Özlem Durmaz [nothing to disclose], Zerrin Onal [nothing to disclose], Ton Bunt [nothing to disclose], Bettina Hansen [consultant for Mirum Pharmaceuticals, Albireo AB, Chemomab, Calliditas, Intercept, Cyma Bay, unrestricted grants from Cyma bay, Intercept, Mirum and Albireo], Henkjan J. Verkade [Consultant for Danone/Nutricia Research, Ausnutria BV, Albireo AB, GMP+Orphan, Mirum Pharmaceuticals, Intercept and Vivet]

**Financial support statement:** 1. MD/PhD scholarship from the University of Groningen, Groningen, The Netherlands 2. Unrestrictive research grant from Albireo.

**Authors' contributions:** Daan van Wessel: study concept and design, acquisition of data, analysis and interpretation of data, statistical analysis, drafting of the manuscript, obtained funding; Richard Thompson: study concept and design, acquisition of data, critical revision of the manuscript for important intellectual content; Emmanuel Gonzales: study concept and design, acquisition of data, critical revision of the manuscript for important intellectual content; Irena Jankowska: study concept and design, acquisition of data, critical revision of the manuscript for important intellectual content; Etienne Sokal: study concept and design, acquisition of data, critical revision of the manuscript for important intellectual content; Tassos Grammatikopoulos: acquisition of data, critical revision of the manuscript for important intellectual content; Agustina Kadaristiana: acquisition of data, critical revision of the manuscript for important intellectual content; Emmanuel Jacquemin: critical revision of the manuscript for important intellectual content; Anne Spraul: acquisition of data, critical revision of the manuscript for important intellectual content; Patryk Lipiński: acquisition of data, critical revision of the manuscript for important intellectual content; Piotr Czubkowski: acquisition of data, critical revision of the manuscript for important intellectual content; Nathalie Rock: acquisition of data, critical revision of the manuscript for important intellectual content; Mohammad Shagrani: acquisition of data, critical revision of the manuscript for important intellectual content; Dieter Broering: acquisition of data, critical revision of the manuscript for important intellectual content; Talal Algoufi: acquisition of data, critical revision of the manuscript for important intellectual content; Nejat Mazhar: acquisition of data, critical

revision of the manuscript for important intellectual content; Emanuele Nicastro: acquisition of data, critical revision of the manuscript for important intellectual content; Deirdre Kelly: acquisition of data, critical revision of the manuscript for important intellectual content; Gabriela Nebbia: acquisition of data, critical revision of the manuscript for important intellectual content; Henrik Arnell: acquisition of data, critical revision of the manuscript for important intellectual content; Björn Fischler: acquisition of data, critical revision of the manuscript for important intellectual content; Jan Hulscher: acquisition of data, critical revision of the manuscript for important intellectual content; Daniele Serranti: acquisition of data, critical revision of the manuscript for important intellectual content; Cigdem Arikan: acquisition of data, critical revision of the manuscript for important intellectual content; Esra Polat: acquisition of data, critical revision of the manuscript for important intellectual content; Dominique Debray: acquisition of data, critical revision of the manuscript for important intellectual content; Florence Lacaille: acquisition of data, critical revision of the manuscript for important intellectual content; Cristina Goncalves: acquisition of data, critical revision of the manuscript for important intellectual content; Loreto Hierro: acquisition of data, critical revision of the manuscript for important intellectual content; Gema Muñoz Bartolo: acquisition of data, critical revision of the manuscript for important intellectual content; Yael Mozer-Glassberg: acquisition of data, critical revision of the manuscript for important intellectual content; Amer Azaz: acquisition of data, critical revision of the manuscript for important intellectual content; Jernej Breclj: acquisition of data, critical revision of the manuscript for important intellectual content; Antal Dezsőfi: acquisition of data, critical revision of the manuscript for important intellectual content; Pier Luigi Calvo: acquisition of data, critical revision of the manuscript for important intellectual content; Enke Grabhorn: acquisition of data, critical revision of the manuscript for important intellectual content; Ekkehard Sturm: acquisition of data, critical revision of the manuscript for important intellectual content; Wendy van der Woerd: acquisition of data, critical revision of the manuscript for important intellectual content; Binita Kamath: acquisition of data, critical revision of the manuscript for important intellectual content; Jian-She Wang: acquisition of data, critical revision of the manuscript for important intellectual content; Liting Li: acquisition of data, critical revision of the manuscript for important intellectual content; Özlem Durmaz: acquisition of data, critical revision of the manuscript for important intellectual content; Zerrin Onal: acquisition of data, critical revision of the manuscript for important intellectual content; Ton Bunt: critical revision of the manuscript for important intellectual content; Bettina Hansen: study concept and design, analysis and interpretation of data, statistical analysis, drafting of the manuscript, obtained funding; Henkjan Verkade: acquisition of data, study concept and design, analysis and interpretation of data, drafting of the manuscript, obtained funding.

## **ABSTRACT**

**Electronic word count: 256**

### **BACKGROUND & AIMS**

Mutations in *ABCB11* can cause deficiency of the bile salt export pump (BSEP), leading to cholestasis and end-stage liver disease. The rarity of the disease has prevented determination of associations between genotype and either natural history or the effect of surgical biliary diversion (SBD) on long-term outcome (liver transplantation (LTx), hepatocellular carcinoma (HCC), death). We aimed to determine these associations by assembling the largest genetically defined cohort of severe BSEP deficiency patients to date.

### **METHODS**

This multicentre, retrospective cohort study of patients with homozygous or compound heterozygous pathological *ABCB11* mutations included 264 patients. Patients were categorized according to genotypic severity (BSEP1, BSEP2, BSEP3). The predicted residual BSEP transport function decreased with each category.

### **RESULTS**

Genotype severity was strongly associated with native liver survival (NLS, BSEP1 median 20.4y; BSEP2, 7.0y; BSEP3, 3.5y;  $P<.001$ ). At age 15y, the proportion of patients with HCC was 4% in BSEP1, 7% in BSEP2 and 34% in BSEP3 ( $P=0.001$ ). SBD was associated with significantly increased NLS (HR:0.50; 95%CI 0.27-0.94,  $P=.03$ ) in BSEP1 and BSEP2. A serum bile acid (sBA) concentration below 102  $\mu\text{mol/L}$  and a decrease in sBA by at least 75%, each shortly after SBD, reliably predicted NLS of at least 15 years after SBD (each  $P<.001$ ).

### **CONCLUSIONS**

The genotype of severe BSEP deficiency strongly predicts the long-term native liver survival, the risk to develop hepatocellular carcinoma, and the chance that surgical biliary diversion increases native liver survival. Serum bile acid parameters shortly after surgical biliary diversion allow the reliable identification of patients with long-term native liver survival.

## Introduction

Deficiency of the bile salt export pump (BSEP) can result from mutations in the *ABCB11* gene and may lead to intrahepatic cholestasis. The most severe form of BSEP deficiency has been labelled progressive familial intrahepatic cholestasis (PFIC) type 2[1,2]. This autosomal disorder, which is (mostly) recessive in nature, belongs to the group of low gamma-glutamyl transpeptidase (GGT) cholestatic diseases[3]. BSEP is a canalicular ATP binding cassette transporter which transports conjugated bile acids into bile[4,5]. Patients with severe BSEP deficiency (i.e. PFIC due to mutations in *ABCB11*) typically present in early childhood with jaundice, pruritus, elevated serum bile acids, malabsorption and failure to thrive. Some patients respond to medical therapy such as ursodeoxycholic acid (UDCA), however, most patients progress into end-stage liver disease[6,7]. It has been observed that patients may benefit from surgical biliary diversion (SBD) procedures, such as partial external biliary diversion (PEBD) or ileal exclusion (IE)[8-17]. SBD aims to decrease the size of the bile acid pool by interrupting the enterohepatic circulation. Unfortunately, not all patients benefit from SBD and, at some point, many require a liver transplantation (LTx) for refractory pruritus or end-stage liver disease. Severe BSEP deficiency has also been associated with the development of hepatocellular carcinoma (HCC) at an early age, which by itself may necessitate LTx[8,18-20].

Severe BSEP deficiency is a rare disease (incidence estimated between 1:50.000 and 1:100.000 births[18,21]), yet relatively high incidences are reported in Saudi-Arabia (approximately 1:7200 [22]). Due to its rarity, the natural course of severe BSEP deficiency and the genotype-phenotype relationships have remained poorly

characterized, precluding clinicians from providing optimal care and counselling for patients suffering this disease.

We aimed to obtain more detailed insights in the natural history of the disease and to determine associations between genotype and phenotype. We set up a global consortium; *NAPPED* (NAtural course and Prognosis of PFIC and Effect of biliary Diversion). *NAPPED* aims to characterize the natural history of severe BSEP deficiency, to assess the effect of genetic, clinical and therapeutic parameters (including SBD) on major surgical and clinical events, such as native liver survival (NLS), LTx and mortality.



## Patients and Methods

### *Data collection, patient selection and genetic categorization*

Since its start in 2017, NAPPED set out to collect retrospective individual data of patients with a clinical phenotype of progressive low GGT cholestasis. The consortium currently comprises 48 tertiary referral centres from all over the globe. Data were collected conforming to the 1975 Declaration of Helsinki. Data collection used a pre-specified case-record form and was performed using REDCap[23]. Demographic, clinical, and outcome data were collected by investigators within each centre, who identified all consecutive patients who had ever been under paediatric care (age 0-18 years) since 1977. Data reported in the present manuscript were exported from REDCap on March 1, 2019. Patients with compound heterozygous or homozygous pathological *ABCB11* mutations were selected. Where available, functional studies were used in the determination of the pathogenicity of the observed mutations[24-28]. ACMG criteria were applied. Variants of unknown significance were evaluated with the Combined Annotation Dependent Depletion score; a score >25 was sufficient for inclusion in the study. Patients were excluded if genetic reports were unavailable, if they had *ABCB11* mutations of no or unknown pathogenicity, or mutations in *ATP8B1* or *TJP2* (Figure 1). Included patients were further categorized based on their predicted mildest mutation, as estimated from functional *in vitro* and/or genetic *in silico* data, if available[2,18,20,24,26,29]. Patients were categorized as BSEP1 (at least one p.D482G (c.1445A>G) or p.E297G (c.890A>G) mutation; two common European mutations associated with residual BSEP functionality[26], BSEP2 (at least one missense mutation, not p.D482G or p.E297G) or BSEP3 (mutations leading to a predicted, non-functional protein). Biochemistry at presentation in the tertiary centre, as well as most recent before SBD

(pre-SBD) and at least two months after SBD (post-SBD, maximally one year after SBD) were collected. Parameters were converted to standardized units. Pruritus was scored as 'absent', 'mild to moderate' or 'severe', at the discretion of the centre, what, for statistical purposes, was dichotomized into 'absent' or 'present'. Effect of SBD on pruritus was noted as 'no improvement in pruritus', 'transient (partial or complete) relief of pruritus' and 'sustained (partial or complete) relief of pruritus'. Outcome parameters were diversion-free survival (DFS, years between birth and SBD, last visit, LTx or death) and native liver survival (NLS, years between birth and either LTx or death, whichever occurred first, or last visit). Follow-up ended at last known visit, LTx or death.

#### *Statistical evaluation*

Continuous variables are expressed as medians [interquartile range] (IQR). Data were analysed using appropriate methods, including Mann-Whitney and Kruskal-Wallis tests. Categorical variables were analysed using Chi-square, McNemar or Mantel's trend-test. We included sex, birth year, age at presentation in the tertiary centre, use of medical therapy prior to presentation (ursodeoxycholic acid (UDCA), rifampicin, phenobarbital, cholestyramine, antihistamines), sBA, total serum bilirubin (TSB), alanine aminotransferase (ALT), aspartate aminotransferase (AST), GGT and platelet count (PLT). Unadjusted differences in DFS, NLS and hepatocellular carcinoma (HCC) between subgroups were assessed using Kaplan-Meier estimates and compared using the log-rank test. Time-dependent Cox-regression was applied for studying the association between SBD and NLS in patients with BSEP1 or BSEP2 genotypes (hazard ratios [HR] and 95% confidence intervals [CI]). The genotype (BSEP1 or BSEP2), sex, birth year and SBD as a time-dependent factor for NLS

were included in the multivariate model. Patients with a BSEP3 genotype were excluded due to low numbers. The model was extended with additional factors one by one to assess the association with the endpoint. Sensitivity analyses (including geographic region and caseload) were performed to control for heterogeneity between sites. A clock-reset approach was used to visualize the association of the time-dependent risk of SBD with NLS: all patients start without SBD. Then, patients that underwent SBD during follow-up are censored at the age of SBD and restart with a new risk in the SBD curve. Receiver Operating Characteristic (ROC) curve analysis was used to determine the cut-off for post-SBD sBAs in relation to predicting NLS after SBD. Imputation of missing data was not attempted.

A two-sided P-value <0.05 was considered statistically significant. All analyses were performed using IBM SPSS Statistics 23.0 (Armonk, NY). Figures were constructed using Prism 7.02, GraphPad Software, La Jolla, CA, USA.

## Results

### *Baseline data*

At time of data export, NAPPED had 590 patients in its database. This study included 264 patients with compound heterozygous or homozygous disease associated mutations in *ABCB11* with a phenotype of low-GGT cholestasis. Of these patients, 84 had been described in previously published literature (Table S1). Three hundred twenty-six patients were excluded since mutations did not meet inclusion criteria (n=222) or genetic analysis results lacked (n=104) (Figure 1). Eleven patients previously presented with a BRIC2 phenotype (i.e. episodic cholestasis and/or pruritus and transient hepatocellular damage). These patients later presented with severe BSEP deficiency phenotypes (i.e. continuous cholestasis and/or pruritus and continuous hepatocellular damage) and had pathological mutations. Within BSEP1, two homozygous subgroups were identified: 17 patients with p.E297G mutations and 11 patients with p.D482G mutations. The final 264 included patients were followed at European (number of patients, n=202), Middle Eastern (n=25) and Asian (n=37) centres. Table S1 provides genetic profiles and corresponding genotype severity allocations. Table S2 provides the predicted effects of all mutations observed on a single allele (n=194).

The median birth year was 2004 [1995–2012]. Half of the patients were male (125/252, 47%). Age at presentation in the tertiary referral centre was 0.7 [0.2-1.9] years. Prior to presentation, 46% patients used or had ever used UDCA (Table 1). Follow-up ended at LTx, death or last visit. The median follow-up was 4.1 [1.5-12.3] years. During follow-up, 61 patients had undergone SBD and 120 patients had undergone LTx. In total, sixteen patients (6%, BSEP1 n=3/72 (4%), BSEP2 n=8/136

(6%), BSEP3 n=5/56 (9%) died prior to LTx (age 1.6 [1.1-3.5] years). Deaths were all related to liver disease. At 18 years of age, 32% of patients were alive with native liver. During adulthood (age  $\geq 18$  years), five patients underwent LTx (aged 19.6-27.5 years).

#### *Associations between genetic severity category and baseline characteristics*

Table 1 depicts patient characteristics and biochemistry at presentation in the referral centre for BSEP1 (n=72), BSEP2 (n=136) and BSEP3 (n=56). The birth year and year of first visit were earlier in BSEP1 than in BSEP2 or BSEP3 (BSEP1 vs. BSEP2, BSEP1 vs. BSEP3,  $P < .001$ ). Biochemistry at presentation did not differ significantly. ALT and AST, however, tended to be higher in BSEP3. Table S3 shows baseline data of BSEP1 patients with homozygous p.E297G or p.D482G mutations. p.D482G patients presented later in hospital (2.9y vs. 0.2y,  $P < .001$ ), with lower ALT levels (66 vs. 231 U/L,  $P = .01$ ), compared to p.E297G. Patients with homozygous p.E297G or p.D482G mutations presented with lower sBA and ALT levels compared to patients with compound heterozygous mutations, although statistical significance was not reached in either of these variables. Patients with compound heterozygous p.E297G mutations presented at younger age than patients with homozygous p.E297G mutations (0.2 [0.2-0.6]y vs. 0.6 [0.3-3.8]y, respectively,  $P = .02$ , Table S4, S5). Patients with compound heterozygous p.D482G/p.E297G mutations (n=3) presented at a median age of 1.2 years.

*Genetic severity category is associated with diversion-free survival and with survival with native liver*

BSEP1 patients had better long-term outcomes than BSEP2 or BSEP3 patients: a median NLS of 20.4 years, versus 7.0 and 3.5 years, respectively (BSEP1 vs. BSEP2  $P=0.009$ ; BSEP1 vs. BSEP3  $P<0.001$ ; BSEP2 vs. BSEP3  $P=0.02$ ; Figure 2). SBD was more often performed in BSEP1, as opposed to BSEP2 and BSEP3 ( $P<0.001$ , % of patients with SBD at 15y: 74%, 38% and 28% respectively; BSEP1 vs. BSEP2  $P<0.001$ , BSEP1 vs. BSEP3  $P=0.004$ , BSEP2 vs. BSEP3  $P=0.90$ , Figure 2). Patients with homozygous p.D482G mutations had comparable NLS to that of the homozygous p.E297G patients (NLS at 15y: 73% vs. 69% respectively;  $P=.41$ , Figure S1), despite undergoing SBD less often (% with SBD at 15y: 26% vs. 90%;  $P=0.006$ , Figure S1). Compound heterozygous p.D482G patients underwent SBD more often compared to homozygous p.D482G patients (% with SBD at 15y: 91% vs. 26% respectively;  $P=.01$ , Figure S2). Each of the three patients with compound heterozygous p.D482G/p.E297G mutations underwent SBD (at 1.4, 8.9 and 9.4 years of age). None underwent LTx at a median age at last follow-up of 17.8y.

#### *Association between the severity of genotype and occurrence of hepatocellular carcinoma*

Severe BSEP deficiency is associated with a significant risk of HCC[19], which was confirmed in our cohort. The incidence of HCC increased with the genotype severity: at 15 years of age, the observed incidence increased from 4% in BSEP1, to 7% and even to 34% in BSEP2 and BSEP3, respectively (derived from the survival curve, Figure 3,  $P=.001$ ). HCC occurred in 2/61 (3%) patients that underwent SBD and in 13/180 (7%) that did not ( $P=.32$ ). In patients with homozygous p.E297G and

p.D482G mutations, HCC was observed in 6% (one patient) and 0%, respectively ( $P=0.41$ ). HCC was not observed in the three patients with compound heterozygous p.D482G/p.E297G mutations

#### *Surgical biliary diversion: baseline data*

Median age at time of SBD was 2.3 [1.2-4.7] years ( $n=61$ ). Of these patients, 47 underwent PEBD, 13 underwent IE and one underwent gallbladder-colic diversion. The observed proportion of patients with an SBD at 5, 10 and 18 years of age was 29%, 37% and 47%, respectively (derived from the diversion-free survival curve). One patient received an SBD during adulthood (age 25.6 years). Follow-up after SBD was 8.4 [1.6-12.0] years. The diversion was surgically closed in 6 patients (BSEP1  $n=2$ , BSEP2  $n=3$ , BSEP3  $n=1$ ) at 2.0 [0.1-4.0] years after SBD. LTx followed closure in 5/6 patients, 6.2 [0.8-10.2] years after initial SBD. LTx was performed in 18 (30%) of the 61 patients at 2.4 [1.3-10.0] years after SBD.

Prior to SBD, pruritus was present in 36 (97%) of the 37 patients for whom paired data was available pre- and post-SBD. After SBD, 17 patients (46%) experienced pruritus ( $P<0.001$ ). The improvement of pruritus post-SBD was semi-quantified: in 12/41 patients (29%), no improvement of pruritus was reported, whereas 7/41 (17%) had transient partial or complete relief of pruritus and 22/41 patients (54%) had sustained partial or complete relief of pruritus. BSEP1 patients achieved sustained partial or complete relief of pruritus (18/27, 66%) more often compared to BSEP2 (4/11, 36%) and BSEP3 (0/3, 0%) ( $P=0.002$ , Table S6).

SBD was associated with a decrease in sBAs (363 [254-452] to 48 [4-258]  $\mu\text{mol/L}$ ,

median 90% decrease,  $P<.001$ ), TSB (59 [21-129] to 15 [9-51]  $\mu\text{mol/L}$ ; median 56% decrease,  $P<.001$ ), ALT (117 [64-247] to 63 [22-111] U/L; median 54% decrease,  $P<.001$ ) and AST (165 [91-358] to 83 [92-176] U/L, median 45% decrease,  $P=.002$ ) (Figure 4). Post-SBD sBAs were significantly lower in BSEP1, compared to BSEP2 (27 [3-210] vs. 249 [43-332]  $\mu\text{mol/L}$ ,  $P=0.02$ ).

#### *Relationship between surgical biliary diversion and native liver survival*

Time-dependent Cox regression analysis (corrected for genotype severity, sex and birth year) showed that SBD was associated with significantly higher NLS (HR 0.50; 95%CI 0.27-0.94,  $P=.03$ , Figure 5) in BSEP1 and BSEP2. Time-dependent Cox-regression for BSEP1 and BSEP2 separately yielded HRs of 0.42 (95%CI 0.16-1.07) and 0.64 (95%CI 0.22-1.82), respectively (difference between HRs:  $P=0.57$ ). To assess the model's robustness and the impact of centre heterogeneity on outcome, we performed sensitivity analyses, stratifying for geographical region and caseload of the centres. Neither region, nor the caseload significantly impacted outcomes (HRs 0.32-0.62). Including BSEP3 patients in the original model yielded comparable results (n=3 added, HR 0.51; 95% CI 0.29-0.91,  $P=0.02$ ).

#### *Follow-up after diversion*

NLS after SBD significantly decreased with genotype severity ( $P=.002-0.03$ , Figure S3). Since sBAs likely play an important role in hepatocellular damage, we performed ROC analyses on post-surgical sBA levels in relation to NLS. A post-SBD sBA level  $<102 \mu\text{mol/L}$  was associated with prolonged NLS after SBD (Figure 6;  $P<.001$ , AUC sBAs: 0.778; cut-off  $102 \mu\text{mol/L}$ : sensitivity 80%, specificity 75%). Additionally, a decrease of at least 75% in sBAs was associated with improved NLS after SBD



(Figure 6;  $P < .001$ ; AUC %change sBAs: 0.774; cut-off 75%: sensitivity 73%, specificity 78%).

Figure 5, as stated before, shows NLS up to and after SBD. The NLS after SBD (dotted line) declines rapidly until 2.5 years. Patients with NLS  $\geq 2.5$  years after SBD harboured more often BSEP1 mutations compared with patients with NLS  $< 2.5$  years after SBD (proportion BSEP1: 71% vs. 31% respectively,  $P = .001$ , Table S7), providing further support for SBD being able to improve NLS in BSEP1 (and BSEP2) patients. Pre-SBD ALT levels were significantly higher in patients with short NLS (160 vs. 101 IU/L,  $P = .005$ ). Post-SBD sBAs in patients with prolonged NLS were significantly lower than in patients with shorter NLS (sBA 29 [3-214] vs. 259 [39-297] respectively ( $P = .03$ ), as were TSB levels (9 [8-19]  $\mu\text{mol/L}$  vs. 41 [12-73]  $\mu\text{mol/L}$  respectively,  $P = .02$ ).

## Discussion

Our aim was to characterize the natural history in severe BSEP deficiency, to determine genotype/phenotype associations and to assess the effect of genetic, clinical and therapeutic parameters on clinical outcome. We succeeded to assemble the largest genetically defined cohort of severe BSEP deficiency patients to date. Our data indicate that the majority of severe BSEP deficiency patients undergo an LTx before reaching adulthood. SBD, however, can postpone or obviate the need for transplantation in selected patients. The probability of undergoing SBD, the NLS, the incidence of HCC and follow-up after SBD were all associated with predicted genotype severity.

To study genotype-phenotype relations we categorized patients based on their predicted mildest mutation. We categorized patients harbouring at least one copy of p.D482G or p.E297G as the mildest genotype (i.e. BSEP1); these mutations are associated with less severe disease[8,20,26,29]. Patients within BSEP3 (i.e. the most severe category) harboured mutations known or predicted to lead to a non-functional protein or to absent BSEP expression. BSEP2 patients harboured at least one missense mutation (yet not p.D482G or p.E297G). While these groups showed distinct courses of disease, we suggest that BSEP2 needs further sub-characterization. A large number of different mutations in this category with 136 patients exists. Studying patients with mutations leading to residual BSEP transport activity[30] other than p.D482G and p.E297G could be one of these initiatives. As stated in the introduction, severe BSEP deficiency is an autosomal recessive disorder, belonging to low-GGT cholestatic diseases (including familial intrahepatic cholestasis protein type 1 and the tight junction protein 2 deficiencies[3]). The

recessive nature of *ABCB11* mutations can be somewhat questioned, based on heterozygous *ABCB11* mutations in patients with, for example, intrahepatic cholestasis of pregnancy[31]. This may suggest that the actual phenotype can be determined by the residual transport capacity in relation to environmental challenges.

Our data indicate that patients generally presented before their first birthday, which is consistent with previous reports[20,32]. However, it should be realized that initial presentation may occur later, even at adult age, e.g. in patients with BRIC phenotypes[29,33,34]. Interestingly, the age and biochemistry at presentation did not differ significantly between the genetic severity groups. Yet, ALT and AST levels tended to be higher in BSEP3 patients. These observations may indicate that biochemical parameters are not sensitive enough to discriminate patients according to genotype. The median year of birth was highest in BSEP3. It might be that these patients were rarely reported before, either due to early mortality or early LTx without full diagnostic work up.

HCC is rare in young children, yet severe BSEP deficiency is associated with relatively high incidences of HCC[19,20]. HCC was indeed encountered in a significant proportion of our patients; in up to 34% in patients with a BSEP3 genotype. Although already high, the here reported incidences may actually be an underestimation of the incidence during childhood or lifetime, since the present values were based on a median follow-up of 4.1 years, and data regarding the diagnosis of HCC at explant were not consistently collected. Our study, in addition to earlier reports, underlines the need to screen for HCC in all severe BSEP deficiency

patients, especially in the BSEP3 category [18].

By adulthood, only a third of severe BSEP deficiency patients were alive with their native liver, among these patients over half had undergone SBD. The relatively low NLS highlights the severity of the disease, illustrated by smaller scale studies describing that up to 50% of patients undergo LTx [20,32]. While patients with a BSEP1, BSEP2 and BSEP3 genotype are at baseline apparently clinically and biochemically indistinguishable, their natural history differed significantly. Research so far has not been able to adequately characterize large groups of patients based on genotype at all, or solely by distinguishing between patients with either one or two copies of p.D482G/p.E297G on the one hand, and patients that did not have one of these mutations on the other hand[18,20,32]. The present study was, besides categorizing patient in three genetic severity groups, able to analyse patients with homozygous p.D482G or p.E297G mutations. Our data indicate that these patients (i.e. harbouring the mildest, BSEP1 genotype) have a milder phenotype than patients not carrying either p.D482G or p.E297G. Apart from confirming earlier findings[8,20,26,29], we add that of these two, p.D482G might be the milder. Homozygous p.D482G patients present in hospital 2.8 years later, with lower ALT levels. To our knowledge, this study is the first to describe the natural history of these mutations in homozygous cohorts. Moreover, patients with homozygous mutations had a slightly favourable prognosis compared to patients with compound heterozygous p.D482G or p.E297G mutations, in regard to baseline biochemistry, proportion undergoing SBD, and NLS. A note of caution is due here since statistical significance was not reached in all variables and a centre bias cannot be ruled out.

Future studies scrutinizing the natural history in these patient groups are therefore recommended.

One of the mechanisms of action of surgical biliary diversion (SBD) involves re-targeting of BSEP to the canalicular membrane, thereby improving bile acid excretion[35]. We show that SBD is associated with increased NLS in BSEP1 and BSEP2 patients. Our data do not support to perform SBD in BSEP3 because of apparent lack of beneficial effect on long-term outcome. In BSEP1 and BSEP2, however, the interruption of the enterohepatic circulation (EHC) seemed to postpone or even remove the need for LTx. Also, BSEP1 patients were most likely to achieve long-term NLS after SBD as opposed to BSEP2 or BSEP3. Similarly, a recent, in-depth study by Bull et al. indicated that patients harbouring at least one p.D482G or p.E297G copy were less likely to undergo LTx after SBD[8].

Apart from postponing or eliminating the need for LTx, SBD aims to relieve patients of pruritus. Pruritus is regarded one of the most burdensome symptoms of severe BSEP deficiency, negatively affecting health related quality of life within the PFIC spectrum[36,37]. The efficacy of current medical antipruritogens is limited. Yet, SBD has been reported to decrease pruritus in some patients [8-17]. In agreement with earlier observations, our data show that SBD was associated with a sustained or complete improvement in pruritus, especially in BSEP1, even though a standardized itch score was absent in this retrospective cohort. None of the analysed BSEP3 patients seemed to benefit from SBD. Our present data therefore also do not support surgical interruption of the EHC as a viable and successful short-term treatment strategy for pruritus in patients with this genotype.

In our study, post-surgical sBA levels were highly prognostic of NLS. This information provides valuable information for expectations towards the need for alternative treatments after SBD (e.g. LTx) and for counselling patients and their families. Furthermore, we propose that sBA parameters could function as markers for long-term outcome after interruption of the EHC. It is interesting to explore whether also biliary bile acid concentrations could provide (more) prognostic information for long-term outcome after SBD[20,38].

Whilst SBD is helpful in a fraction of severe BSEP deficiency patients, it remains an invasive procedure with unwanted cosmetic consequences. Alternative treatments for severe BSEP deficiency are currently being developed and studied, such as medical interruption of the EHC by means of apical sodium-dependent bile acid transporter inhibitors[39,40], and chaperone drugs[30,41]. The present data do provide further support for a personalized strategy to change the natural history of the disease for the better by interrupting the EHC and/or by increasing residual function of BSEP. Especially BSEP1 or BSEP2 patients might benefit from these medical strategies, based on the current observations of SBD with respect to long-term outcome.

We are aware of the limitations of our retrospective study. Inevitably, we encountered missing data. Although this might have influenced our outcomes to some extent, we believe that our numbers are sufficient to nevertheless draw adequate and relevant conclusions. In this study, the age at first presentation was defined as the first visit in the tertiary centre. Although the first presentation was at young age, it is likely that the first symptomatic presentation of disease had been at even younger age. The

present study included eleven patients who initially presented with a BRIC-phenotype. This did not impact our main conclusions; excluding these patients from major analyses yielded comparable results. Our database does currently not include data regarding liver histology. Therefore, we could not assess if histological features at either presentation or at the time of SBD were associated with long-term outcome. The global nature of NAPPED might have resulted in a degree of selection-bias. Therefore, we performed sensitivity analyses regarding the effect of the centre (caseload, geographical region) on outcome. Our main conclusions were not affected by these potential confounders. Although LTx remains presently the most definitive treatment for severe BSEP deficiency, the disease might reoccur after transplant[42,43]. Our study design, with follow-up ending at LTx, precluded analysis of this phenomenon. Despite these limitations NAPPED offers a solution to obtain relevant clinical and follow-up data in a rare disease over an extended period of time.

In conclusion, our study shows that only a third of severe BSEP deficiency patients reach adulthood with their native liver. Moreover, the genotype severity of severe BSEP deficiency strongly predicts long-term NLS. Our results indicate that SBD is associated with significantly prolonged NLS in BSEP1 or BSEP2 patients. Based on the data provided, we propose that serum bile acids after interruption of the enterohepatic circulation can be used as a marker for long-term outcome. The present results increase the understanding and characterization of severe BSEP deficiency, allowing for improved personalized clinical care for these patients throughout childhood and into adulthood, for better targeting of novel therapeutic strategies and, finally, for an improved means to assess their effect via candidate surrogate markers. Our data indicate that that the care for patients with severe BSEP

deficiency can and should, at least partially, be based on the severity of their genotype.

We thank the patients and their families for their important data. Our gratitude goes out to all centres and investigators that are currently or in the near future will be participating in NAPPED. Additionally, we want to thank all our colleagues who provided care for the included patients and/or have supported the present research initiative. Finally, this work was supported by a European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) Networking Grant.



- [1] Elferink RO, Groen AK. Genetic defects in hepatobiliary transport. *Biochim Biophys Acta* 2002;1586:129-145.
- [2] Strautnieks SS, Bull LN, Knisely AS, Kocoshis SA, Dahl N, Arnell H et al. A gene encoding a liver-specific ABC transporter is mutated in progressive familial intrahepatic cholestasis. *Nat Genet* 1998;20:233-238.
- [3] Hadzic N, Verkade HJ. The Changing Spectrum of Neonatal Hepatitis. *JPediatr Gastroenterol Nutr* 2016;63:316-319.
- [4] Noe J, Stieger B, Meier PJ. Functional expression of the canalicular bile salt export pump of human liver. *Gastroenterology* 2002;123:1659-1666.
- [5] Thompson R, Strautnieks S. BSEP: function and role in progressive familial intrahepatic cholestasis. *Semin Liver Dis* 2001;21:545-550.
- [6] Shneider BL. Progressive intrahepatic cholestasis: mechanisms, diagnosis and therapy. *Pediatr Transplant* 2004;8:609-612.
- [7] Whittington PF, Freese DK, Alonso EM, Schwarzenberg SJ, Sharp HL. Clinical and biochemical findings in progressive familial intrahepatic cholestasis. *JPediatr Gastroenterol Nutr* 1994;18:134-141.
- [8] Bull LN, Pawlikowska L, Strautnieks S, Jankowska I, Czubkowski P, Dodge JL et al. Outcomes of surgical management of familial intrahepatic cholestasis 1 and bile salt export protein deficiencies. *Hepatol Commun* 2018;2:515-528.
- [9] Ellinger P, Stindt J, Droge C, Sattler K, Stross C, Kluge S et al. Partial external biliary diversion in bile salt export pump deficiency: Association between outcome and mutation. *World J Gastroenterol* 2017;23:5295-5303.
- [10] Lemoine C, Bhardwaj T, Bass LM, Superina RA. Outcomes following partial external biliary diversion in patients with progressive familial intrahepatic cholestasis. *JPediatr Surg* 2017;52:268-272.
- [11] Ng VL, Ryckman FC, Porta G, Miura IK, de Carvalho E, Servidoni MF et al. Long-term outcome after partial external biliary diversion for intractable pruritus in patients with intrahepatic cholestasis. *JPediatr Gastroenterol Nutr* 2000;30:152-156.
- [12] Yang H, Porte RJ, Verkade HJ, De Langen ZJ, Hulscher JB. Partial external biliary diversion in children with progressive familial intrahepatic cholestasis and Alagille disease. *JPediatr Gastroenterol Nutr* 2009;49:216-221.
- [13] Arnell H, Bergdahl S, Papadogiannakis N, Nemeth A, Fischler B. Preoperative observations and short-term outcome after partial external biliary diversion in 13 patients with progressive familial intrahepatic cholestasis. *JPediatr Surg* 2008;43:1312-1320.

- [14] Kurbegov AC, Setchell KD, Haas JE, Mierau GW, Narkewicz M, Bancroft JD et al. Biliary diversion for progressive familial intrahepatic cholestasis: improved liver morphology and bile acid profile. *Gastroenterology* 2003;125:1227-1234.
- [15] Melter M, Rodeck B, Kardorff R, Hoyer PF, Petersen C, Ballauff A et al. Progressive familial intrahepatic cholestasis: partial biliary diversion normalizes serum lipids and improves growth in noncirrhotic patients. *Am J Gastroenterol* 2000;95:3522-3528.
- [16] Emond JC, Whittington PF. Selective surgical management of progressive familial intrahepatic cholestasis (Byler's disease). *J Pediatr Surg* 1995;30:1635-1641.
- [17] Wang KS, Tiao G, Bass LM, Hertel PM, Mogul D, Kerkar N et al. Analysis of surgical interruption of the enterohepatic circulation as a treatment for pediatric cholestasis. *Hepatology* 2017;65:1645-1654.
- [18] Strautnieks SS, Byrne JA, Pawlikowska L, Cebecauerova D, Rayner A, Dutton L et al. Severe bile salt export pump deficiency: 82 different ABCB11 mutations in 109 families. *Gastroenterology* 2008;134:1203-1214.
- [19] Knisely AS, Strautnieks SS, Meier Y, Stieger B, Byrne JA, Portmann BC et al. Hepatocellular carcinoma in ten children under five years of age with bile salt export pump deficiency. *Hepatology* 2006;44:478-486.
- [20] Davit-Spraul A, Fabre M, Branchereau S, Baussan C, Gonzales E, Stieger B et al. ATP8B1 and ABCB11 analysis in 62 children with normal gamma-glutamyl transferase progressive familial intrahepatic cholestasis (PFIC): phenotypic differences between PFIC1 and PFIC2 and natural history. *Hepatology* 2010;51:1645-1655.
- [21] Davit-Spraul A, Gonzales E, Baussan C, Jacquemin E. Progressive familial intrahepatic cholestasis. *Orphanet JRare Dis* 2009;4:1-1.
- [22] Shagrani M, Burkholder J, Broering D, Abouelhoda M, Faquih T, El-Kalioby M et al. Genetic profiling of children with advanced cholestatic liver disease. *Clin Genet* 2017;92:52-61.
- [23] Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *JBiomed Inform* 2009;42:377-381.
- [24] Byrne JA, Strautnieks SS, Ihrke G, Pagani F, Knisely AS, Linton KJ et al. Missense mutations and single nucleotide polymorphisms in ABCB11 impair bile salt export pump processing and function or disrupt pre-messenger RNA splicing. *Hepatology* 2009;49:553-567.
- [25] Noe J, Kullak-Ublick GA, Jochum W, Stieger B, Kerb R, Haberl M et al. Impaired expression and function of the bile salt export pump due to three novel ABCB11 mutations in intrahepatic cholestasis. *JHepatol* 2005;43:536-543.

- [26] Hayashi H, Takada T, Suzuki H, Akita H, Sugiyama Y. Two common PFIC2 mutations are associated with the impaired membrane trafficking of BSEP/ABCB11. *Hepatology* 2005;41:916-924.
- [27] Plass JR, Mol O, Heegsma J, Geuken M, de Bruin J, Elling G et al. A progressive familial intrahepatic cholestasis type 2 mutation causes an unstable, temperature-sensitive bile salt export pump. *JHepatol* 2004;40:24-30.
- [28] Wang L, Soroka CJ, Boyer JL. The role of bile salt export pump mutations in progressive familial intrahepatic cholestasis type II. *JClin Invest* 2002;110:965-972.
- [29] Droge C, Bonus M, Baumann U, Klindt C, Lainka E, Kathemann S et al. Sequencing of FIC1, BSEP and MDR3 in a large cohort of patients with cholestasis revealed a high number of different genetic variants. *JHepatol* 2017;67:1253-1264.
- [30] Gonzales E, Grosse B, Schuller B, Davit-Spraul A, Conti F, Guettier C et al. Targeted pharmacotherapy in progressive familial intrahepatic cholestasis type 2: Evidence for improvement of cholestasis with 4-phenylbutyrate. *Hepatology* 2015;62:558-566.
- [31] Dixon PH, Sambrotta M, Chambers J, Taylor-Harris P, Syngelaki A, Nicolaidis K et al. An expanded role for heterozygous mutations of ABCB4, ABCB11, ATP8B1, ABCC2 and TJP2 in intrahepatic cholestasis of pregnancy. *Sci Rep* 2017;7:11823-x.
- [32] Pawlikowska L, Strautnieks S, Jankowska I, Czubkowski P, Emerick K, Antoniou A et al. Differences in presentation and progression between severe FIC1 and BSEP deficiencies. *JHepatol* 2010;53:170-178.
- [33] van Ooteghem NA, Klomp LW, van Berge-Henegouwen GP, Houwen RH. Benign recurrent intrahepatic cholestasis progressing to progressive familial intrahepatic cholestasis: low GGT cholestasis is a clinical continuum. *JHepatol* 2002;36:439-443.
- [34] van Mil SW, van der Woerd, WL, van der Brugge G, Sturm E, Jansen PL, Bull LN et al. Benign recurrent intrahepatic cholestasis type 2 is caused by mutations in ABCB11. *Gastroenterology* 2004;127:379-384.
- [35] Varma S, Revencu N, Stephenne X, Scheers I, Smets F, Beleza-Meireles A et al. Retargeting of bile salt export pump and favorable outcome in children with progressive familial intrahepatic cholestasis type 2. *Hepatology* 2015;62:198-206.
- [36] Kamath BM, Chen Z, Romero R, Fredericks EM, Alonso EM, Arnon R et al. Quality of Life and Its Determinants in a Multicenter Cohort of Children with Alagille Syndrome. *JPediatr* 2015;167:390-6.e3.
- [37] Baker A, Kerkar N, Todorova L, Kamath BM, Houwen RHJ. Systematic review of progressive familial intrahepatic cholestasis. *Clin Res Hepatol Gastroenterol* 2019;43:20-36.

[38] Emerick KM, Elias MS, Melin-Aldana H, Strautnieks S, Thompson RJ, Bull LN et al. Bile composition in Alagille Syndrome and PFIC patients having Partial External Biliary Diversion. *BMC Gastroenterol* 2008;8:47-47.

[39] Baghdasaryan A, Fuchs CD, Osterreicher CH, Lemberger UJ, Halilbasic E, Pahlman I et al. Inhibition of intestinal bile acid absorption improves cholestatic liver and bile duct injury in a mouse model of sclerosing cholangitis. *JHepatol* 2016;64:674-681.

[40] Miethke AG, Zhang W, Simmons J, Taylor AE, Shi T, Shanmukhappa SK et al. Pharmacological inhibition of apical sodium-dependent bile acid transporter changes bile composition and blocks progression of sclerosing cholangitis in multidrug resistance 2 knockout mice. *Hepatology* 2016;63:512-523.

[41] Gonzales E, Jacquemin E. Mutation specific drug therapy for progressive familial or benign recurrent intrahepatic cholestasis: a new tool in a near future?. *JHepatol* 2010;53:385-387.

[42] Stindt J, Kluge S, Droge C, Keitel V, Stross C, Baumann U et al. Bile salt export pump-reactive antibodies form a polyclonal, multi-inhibitory response in antibody-induced bile salt export pump deficiency. *Hepatology* 2016;63:524-537.

[43] Maggiore G, Gonzales E, Sciveres M, Redon MJ, Grosse B, Stieger B et al. Relapsing features of bile salt export pump deficiency after liver transplantation in two patients with progressive familial intrahepatic cholestasis type 2. *JHepatol* 2010;53:981-986.

**Fig. 1. Flowchart of patient inclusion from NAatural course and Prognosis of PFIC and Effect of biliary Diversion (NAPPED) database.** \*:heterozygous mutations, *ABCB11* mutations of no (known) clinical consequence or mutations in *ATP8B1/TJP2*.

**Table 1. Baseline characteristics in all patients, and BSEP1, BSEP2 and BSEP3 genotypes separately.** \*Mantel-Haenszel or Kruskal-Wallis test, as appropriate, to test differences between BSEP1, BSEP2 and BSEP3. Genotypic categorization clarified in Methods. *BSEP* (bile salt export pump); *GGT* (gamma-glutamyl transpeptidase); *UDCA* (ursodeoxycholic acid).

**Fig. 2. Observed native liver survival and diversion-free survival per genotypic severity category.** (A) Proportion of patients alive with native liver over time with a BSEP1 (solid line), BSEP2 (dashed line) or BSEP3 (dotted line) genotype. Log-rank test. (B) Proportion of patients with a surgical biliary diversion over time with BSEP1 (solid line), BSEP2 (dashed line) or BSEP3 (dotted line) genotypes. Genotypic categorization clarified in Methods. Log-rank test. *BSEP* (bile salt export pump).

**Fig. 3. Observed proportion of patients with a hepatocellular carcinoma per genotypic severity category.** Genotypic categorization clarified in Methods. Log-rank test. *BSEP* (bile salt export pump).

**Fig. 4. Paired pre- and post-surgical biochemical parameters in all patients undergoing surgical biliary diversion.** Kruskal-Wallis test. *ALT* (alanine aminotransferase); *AST* (aspartate aminotransferase); *SBD* (surgical biliary diversion).

**Fig. 5. Observed native liver survival in BSEP1 or BSEP2 patients undergoing surgical biliary diversion (SBD) or not.** The clock-reset approach allows visualization of native liver survival up to SBD (solid line, all patients) and after SBD (dotted line, only patients that underwent SBD). The estimated hazard ratio (HR) is achieved by Cox-regression with SBD as a time-dependent risk-factor, adjusted for genotype, sex and birth year. Patients in analysis: n=173.

**Fig. 6. Observed native liver survival after surgical biliary diversion, stratified for post-surgical serum bile acids cut-offs** (A) In patients with a post-surgical serum bile acid concentration < or  $\geq$  102 $\mu$ mol/L. (B) In patients with a relative decrease in serum bile acids of < or  $\geq$  75%. Log-rank test. *SBA* (serum bile acids)

## **Lay summary**

By using the largest genetically defined cohort of severe BSEP deficiency (i.e. PFIC due to *ABCB11* mutations) patients known to date, this study increases the understanding and characterization of severe BSEP deficiency. The present data allow for improved personalized clinical care throughout childhood and into adulthood, for better targeting of novel therapeutic strategies and, for an improved means to assess their effect via candidate surrogate markers.