

## University of Groningen

### Letter to the Editor on "Guyuron B, Son JH. String fat/dermis graft for correction of wrinkles and scars. *Plast. Reconstruct. Surg.* 2019; 144

Stenekes, Martin; Van Der Lei, Berend

*Published in:*  
Plastic and Reconstructive Surgery

*DOI:*  
[10.1097/PRS.0000000000006580](https://doi.org/10.1097/PRS.0000000000006580)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Stenekes, M., & Van Der Lei, B. (2020). Letter to the Editor on "Guyuron B, Son JH. String fat/dermis graft for correction of wrinkles and scars. *Plast. Reconstruct. Surg.* 2019; 144: 93-96". *Plastic and Reconstructive Surgery*, 145(3), 653e-654e. <https://doi.org/10.1097/PRS.0000000000006580>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

Department of Surgery  
Albany Medical Center  
Albany, N.Y.

Correspondence to Dr. Cotofana  
Mayo Clinic College of Medicine and Science  
Mayo Clinic  
Stabile Building 9-38  
200 First Street  
Rochester, Minn. 55905  
cotofana.sebastian@mayo.edu  
Facebook: professorsebastiancotofana  
Instagram: @professorsebastiancotofana

### DISCLOSURE

*The authors have no commercial associations or financial disclosures that might pose or create a conflict of interest with the methods applied or the results presented in this communication. No funding was received for this communication.*

### REFERENCES

1. Lee SH, Ha TJ, Koh KS, Song WC. External and internal diameters of the facial artery relevant to intravascular filler injection. *Plast Reconstr Surg*. 2019;143:1031–1037.
2. Phumyoo T, Tansatit T, Rachkeaw N. The soft tissue landmarks to avoid injury to the facial artery during filler and neurotoxin injection at the nasolabial region. *J Craniofac Surg*. 2014;25:1885–1889.
3. Camuzard O, Foissac R, Georgiou C, et al. Facial artery perforator flap for reconstruction of perinasal defects: An anatomical study and clinical application. *J Craniomaxillofac Surg*. 2015;43:2057–2065.
4. Coronel-Banda ME, Serra-Renom JM, Lorente M, Larrea-Terán WP. Mucosal perforators from the facial artery. *Br J Oral Maxillofac Surg*. 2014;52:535–538.
5. Qassemayr Q, Havet E, Sinna R. Vascular basis of the facial artery perforator flap: Analysis of 101 perforator territories. *Plast Reconstr Surg*. 2012;129:421–429.
6. Pinar YA, Bilge O, Govsa F. Anatomic study of the blood supply of perioral region. *Clin Anat*. 2005;18:330–339.
7. Zhao YP, Arijji Y, Gotoh M, et al. Color Doppler sonography of the facial artery in the anterior face. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2002;93:195–201.
8. Ng ZY, Fogg QA, Shoaib T. Where to find facial artery perforators: A reference point. *J Plast Reconstr Aesthet Surg*. 2010;63:2046–2051.
9. Risselada M, Mathews KG, Griffith E. The effect of specimen preparation on post-excision and post-fixation dimensions, translation, and distortion of canine cadaver skin-muscle-fascia specimens. *Vet Surg*. 2016;45:563–570.
10. Dauendorffer JN, Bastuji-Garin S, Guéro S, Brousse N, Freitag S. Shrinkage of skin excision specimens: Formalin fixation is not the culprit. *Br J Dermatol*. 2009;160:810–814.
11. Vent J, Zimmermann Ch, Drebber U, et al. Influence of formalin fixation on tissue dimensions in palatal tonsils. *Pathol Res Pract*. 2014;210:59–61.
12. Hayashi S, Naito M, Kawata S, et al. History and future of human cadaver preservation for surgical training: From formalin to saturated salt solution method. *Anat Sci Int*. 2016;91:1–7.
13. Hołda MK, Klimek-Piotrowska W, Koziej M, Piątek K, Hołda J. Influence of different fixation protocols on the preservation and dimensions of cardiac tissue. *J Anat*. 2016;229:334–340.

## String Fat/Dermis Graft for Correction of Wrinkles and Scars

Sir:

With great interest we have read the article of Bahman Guyuron and Ji H. Son regarding the use of string fat/dermis graft for correction of wrinkles and scars.<sup>1</sup> We have also experienced that treating wrinkles and folds with such a graft will lead to a significant improvement. The video of Guyuron and Son<sup>1</sup> clearly demonstrates how the graft is inserted, an original technique described by Carraway and Mellow.<sup>2</sup>

Two aspects with regard to this way of treating skin impressions are worth mentioning to our readers. First, we would like to mention another source of autograft material: when performing a face lift and removing a little of the superficial musculoaponeurotic system (SMAS) after SMAS-ectomy or SMAS shortening, this resected SMAS material can be used for the same purpose with the same great results. In addition, we have used this SMAS material to improve the nasolabial fold.<sup>3</sup>

Second, it may not always be easy to introduce the graft in the tunnel as demonstrated, especially underneath a long wrinkle or fold. We suggest an alternative solution to insert the graft. First, a small incision is made on one end of the wrinkle or fold; this can be done either with a large bore needle or, as in case of the nasolabial fold, with a surgical blade no. 11 in the transverse fold of ala nasi of the nose. Through this wound, a subdermal tunnel is dissected either with the large bore needle or, as in case of the nasolabial fold, with scissors. After this dissection, a long 20-gauge needle is inserted through this tunnel, exiting through the skin on the other side of the fold. A larger bore, 18-gauge needle is then slid in reverse over the first (20-gauge) needle until it exits in the initial incision, after which the first needle is removed. A suture is attached to one end of the strip of SMAS. The long end of the suture is then inserted into the needle and advanced until it appears at the other side of the needle. Now the needle can be carefully removed, leaving the advanced suture out through the skin. Then, by pulling gently on the suture, the SMAS strip can be introduced in the tunnel and nicely advanced underneath the entire length of the wrinkle or fold.

DOI: 10.1097/PRS.0000000000006580

**Martin W. Stenekes, M.D., Ph.D.**  
University Medical Center Groningen  
University of Groningen  
Groningen, The Netherlands

**Berend van der Lei, M.D., Ph.D.**  
Department of Plastic Surgery  
University Medical Center Groningen  
University of Groningen  
Groningen, The Netherlands  
Bergman Clinics  
Heerenveen, Zwolle, and Hilversum, The Netherlands

Correspondence to Dr. van der Lei  
Department of Plastic Surgery  
University Medical Centre of Groningen  
Hanzeplein 1, P.O. Box 30.001  
9700 RB Groningen, The Netherlands  
[b.van.der.lei@umcg.nl](mailto:b.van.der.lei@umcg.nl)

### DISCLOSURE

*The authors have no financial interest to declare in relation to the content of this communication.*

### REFERENCES

1. Guyuron B, Son JH. String fat/dermis graft for correction of wrinkles and scars. *Plast Reconstr Surg*. 2019;144:93–96.
2. Carraway JH, Mellow CG. Syringe aspiration and fat concentration: A simple technique for autologous fat injection. *Ann Plast Surg*. 1990;24:293–296; discussion 297.
3. Stenekes MW, Van Der Lei B. Nasolabial fold augmentation with SMAS graft. *J Plast Reconstr Aesthet Surg*. 2012;65:1618–1621.

## Secondary Abdominoplasty: Management of the Umbilicus after Prior Stalk Transection

*Sir:*

Every time, we are going to face an increased demand from patients requesting secondary abdominoplasties who had previously undergone a mini-abdominoplasty with supraumbilical detachment and navel detachment.<sup>1</sup> The reasons are that the abdominal area is increasingly more exposed, there are more frequent changes in partners, and there is a greater aesthetic demand from society, thereby causing a greater demand for this type of treatment.

Personally, I have had nine cases of mini-abdominoplasty treated using the supraumbilical diastasis with umbilical transection and three cases of previous treatment of umbilical hernia. In those 12 cases, a secondary abdominoplasty was performed (until 2015) without any suffering or vascular damage to the umbilicus. The probable explanation is that the umbilical pedicle behaved like a randomized flap 21 days after primary surgery, and in the worst-case scenario, partial or total necrosis with closure by secondary-intention healing would be acceptable from an aesthetic point of view.

In my opinion, these cases would be an accurate indication for a transverse plication, no undermining, full liposuction, neoumbilicoplasty, and low transverse abdominal scar, or TULUA abdominoplasty, as published by Dr. Villegas in 2014. I incorporated the TULUA procedure into my routine for secondary abdominoplasty in 2016, and I have operated on 36 patients using this technique. The TULUA technique allows for vigorous and wide liposuction of the supraumbilical area since it will not be detached, a trans-infra-umbilical wide plication, and amputation of the patient's original navel. At the end of the operation, the new location of the

neo-umbilicus is chosen and made through the use of a skin graft.

Our approach is to use epidural anesthesia performed by the anesthesiologist and tumescent infiltration with lidocaine for its bacteriostatic power. Of the 36 patients I have operated on, 14 had previous unsuccessful tummy tucks and 22 patients had extensive lipoaspiration of the anterior abdomen, with adhesions or little elasticity of the supraumbilical flap with which to perform the infraumbilical portion. The TULUA technique allowed us to incorporate a new niche of patients who were dismissed because their cases were complicated.<sup>2</sup> TULUA offers a high degree of safety in its execution. It also allows us to combine other procedures in the same patient without extending the surgical time exceedingly.

With vigorous and extensive lipoaspiration of the supraumbilical flap, it is possible to perform a high-definition liposuction in that area and combine it with the infraumbilical transverse plication of the TULUA, allowing us to offer a very safe high-definition lipoabdominoplasty to our patients.

In the last 2 years, this technique has been disseminated worldwide, beginning in South America. Several colleagues are practicing it and have it within their therapeutic arsenal, allowing us to solve complicated cases of our own or from colleagues who are not specialists in plastic surgery.

DOI: [10.1097/PRS.00000000000006581](https://doi.org/10.1097/PRS.00000000000006581)

**Javier Vera Cucchiario, M.D.**

Javier Vera Cucchiario Aesthetic and Laser Clinica  
Avenue Entre Rios 220  
Salta, Argentina  
[jvc@centrocirurgiaestetica.com](mailto:jvc@centrocirurgiaestetica.com)  
Instagram: @javiervacucchiario  
Facebook: @DrJavierVCucchiario

### DISCLOSURE

*The author has no conflict of interests to report.*

### REFERENCES

1. Dean RA, Dean JA, Matarasso A. Secondary abdominoplasty: Management of the umbilicus after prior stalk transection. *Plast Reconstr Surg*. 2019;143:729e–733e.
2. Villegas F. A novel approach to abdominoplasty: TULUA. *Aesthetic Plast Surg*. 2014; 38:511–520.

## Reply: Secondary Abdominoplasty: Management of the Umbilicus after Prior Stalk Transection

*Sir:*

Managing an umbilicus when the stalk has been previously transected and now requires circumscription (i.e., for a full abdominoplasty) can theoretically lead to ischemia.<sup>1,2</sup> Our report<sup>3</sup> about delaying the umbilicus represents a potential solution and preserves the patient's intrinsic umbilical appearance. Neoumbilization, as Dr. Javier Vera Cucchiario and others have