

University of Groningen

## Fibroblast growth factor 23 mediates the association between iron deficiency and mortality in worsening heart failure

van der Wal, Haye H.; Beverborg, Niels Grote; ter Maaten, Jozine M.; Vinke, Joanna S. J.; de Borst, Martin H.; van Veldhuisen, Dirk J.; Voors, Adriaan A.; van der Meer, Peter

*Published in:*  
European Journal of Heart Failure

*DOI:*  
[10.1002/ejhf.1801](https://doi.org/10.1002/ejhf.1801)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

van der Wal, H. H., Beverborg, N. G., ter Maaten, J. M., Vinke, J. S. J., de Borst, M. H., van Veldhuisen, D. J., Voors, A. A., & van der Meer, P. (2020). Fibroblast growth factor 23 mediates the association between iron deficiency and mortality in worsening heart failure. *European Journal of Heart Failure*, 22(5), 903-906. <https://doi.org/10.1002/ejhf.1801>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

doi:10.1002/ejhf.1801

## Fibroblast growth factor 23 mediates the association between iron deficiency and mortality in worsening heart failure

Iron deficiency (ID) is prevalent in heart failure (HF) and associated with a poor prognosis.<sup>1,2</sup> The pathophysiological mechanisms of ID are not fully understood. While a strong link between ID and anaemia exists, ID is associated with increased mortality in non-anaemic patients as well.<sup>3</sup> Additionally, intravenous iron administration is also beneficial in non-anaemic patients. This implies other, non-haematopoietic effects of ID on outcome. ID has been linked to increased levels of fibroblast growth factor 23 (FGF23), which is a phosphaturic osteocyte-derived hormone. FGF23 inhibits renal phosphate reabsorption and regulates 1,25(OH)<sub>2</sub> vitamin D. The association between iron status and FGF23 originates from studies in FGF23-related autosomal dominant hypophosphataemic rickets, which has an iron-dependent onset.<sup>4</sup> Previously, we have shown that in HF patients, FGF23 is independently associated with congestion, unsuccessful angiotensin-converting enzyme inhibitor and angiotensin receptor blocker up-titration, and poor prognosis.<sup>5</sup> Moreover, FGF23 has been linked to incident HF and mortality in community-based studies and development of left ventricular hypertrophy and mortality in chronic kidney disease patients.<sup>6,7</sup> Recent preclinical data suggest an association between FGF23 and cardiac renin-angiotensin-aldosterone system activation, thereby leading to cardiac fibrosis and hypertrophy.<sup>8</sup> Finally, correction of ID seems related to significant reductions in FGF23 levels in HF patients, a finding which further links iron status and FGF23 together.<sup>9</sup> The association between iron status, FGF23 and outcome in HF is currently unclear. This study therefore focused on the interplay between iron and FGF23, and whether FGF23 mediates the association between iron status and outcome in HF.

This study is a post-hoc analysis of the 'systems BIOlogy Study to Tailored Treatment in Chronic Heart Failure' (BIOSTAT-CHF), which has been described previously.<sup>5</sup> In short, BIOSTAT-CHF prospectively enrolled 2516 patients with worsening signs and/or symptoms of HF and a left ventricular ejection fraction  $\leq 40\%$  or brain natriuretic peptide levels  $>400$  pg/mL or N-terminal prohormone of brain natriuretic peptide  $>2000$  pg/mL. BIOSTAT-CHF was approved by local medical ethics committees at each participating centre. We measured the following biomarkers in 2279 stored samples, drawn at the time of presentation at the emergency department or hospital admission: iron, transferrin saturation (TSAT), ferritin, hepcidin, soluble transferrin receptor (sTfR) and FGF23 [using a c-terminal ELISA (Immutopics, Inc., San Clemente, CA, USA), measuring both intact and c-terminal FGF23 cumulatively].<sup>5</sup> Univariable and multivariable linear regression analyses were performed using log-transformed FGF23 levels as dependent variables and log-transformed TSAT, sTfR, ferritin and hepcidin as independent variables. The multivariable models were adjusted for predictors that have previously been associated with FGF23.<sup>5</sup> Univariable and FGF23-adjusted restricted cubic splines based on Cox proportional hazard regression were constructed to assess the association between iron parameters and all-cause mortality. Mediation analyses were performed according to the methods described by Baron and Kenny.<sup>7</sup>

Baseline characteristics of all patients are depicted in Table 1. Mean ( $\pm$  standard deviation) age of the patients was  $69 \pm 12$  years, 26.1% were female and median (interquartile range) left ventricular ejection fraction was 30% (25–36%). Patients with higher FGF23 levels were more frequently female, had lower ferritin, TSAT, and hepcidin levels and higher levels of inflammatory markers ( $P$  for trend  $<0.001$ ). FGF23 levels were strongly correlated to TSAT (Spearman's  $\rho = -0.42$ ), sTfR ( $\rho = 0.43$ ), ferritin ( $\rho = -0.31$ ) and hepcidin ( $\rho = -0.37$ ; all  $P < 0.0001$ ). Individual levels of TSAT, sTfR, ferritin and hepcidin were the strongest predictors of FGF23 levels compared to previously established determinants of FGF23 levels (all  $P < 0.001$ ).<sup>5</sup> During a median follow-up of 21 months (interquartile range 16–27 months), 596 patients (26%)

died. Continuous iron parameter levels were strongly associated with prognosis in univariable analyses (all  $P < 0.001$ ) (Figure 1). When adjusting for FGF23, all iron parameters lost their predictive value. There was a significant interaction between TSAT and FGF23 on outcome ( $P = 0.012$ ). Moreover, we identified a highly significant interaction between a history of renal disease and FGF23 in the association between iron parameters and all-cause mortality ( $P < 0.01$ ). Finally, we evaluated whether the association between iron parameters and all-cause mortality was mediated by FGF23. FGF23 levels significantly mediated the association between TSAT, ferritin, sTfR, and hepcidin and all-cause mortality [ $P$  for indirect effect (FGF23-mediated)  $<0.0001$ ]. The direct effect (non-FGF23-mediated) was not significant for all iron parameters in these models (Table 1). As a sensitivity analysis, we also evaluated whether inflammation alters the association between iron status and outcome. Adjustment for C-reactive protein and interleukin-6 did not affect the prognostic consequences of iron parameters.

In this study, we found that in a large, multinational cohort of worsening HF patients, iron parameters are independently related to FGF23 levels. Second, the prognostic value of iron parameters is significantly mediated by FGF23. Taken these findings together, our data provide insight into the pathophysiology of ID in HF patients, in which FGF23 may play a prominent role.

Low iron has previously been identified as one of the drivers of FGF23 production *in vivo*.<sup>4,10</sup> By stabilizing hypoxia-inducible factor 1- $\alpha$ , ID up-regulates the expression of the proprotein convertase furin, which in turn increases FGF23 production and cleavage into intact and c-terminal FGF23.<sup>4,11</sup> Observational studies in chronic kidney disease and general population cohorts identified an association between ID and outcome, which was mediated by FGF23. For example, an observational study in renal transplant recipients yielded comparable results to our study, showing that the association between ID and outcome is considerably mediated by FGF23 levels and not by inflammation.<sup>7</sup> Furthermore, FGF23 was strongly linked to

**Table 1** Baseline characteristics, stratified by fibroblast growth factor 23 levels in tertiles, and multivariable mediation analysis of iron status parameters through fibroblast growth factor 23


	Total cohort (n = 2297)	FGF23 ≤ 140 RU/mL (n = 766)	FGF23 141– 386 RU/mL (n = 766)	FGF23 ≥ 387 RU/mL (n = 765)	P-value for trend	Mediation analysis <sup>a</sup>				
						Direct effect coefficient (95% CI)	P-value	Indirect effect coefficient (95% CI)	P-value	
<b>Demographics</b>										
Age (years)	69 ± 12	66 ± 11	70 ± 12	70 ± 12	<0.001					
Women	600 (26.1)	162 (21.1)	201 (26.2)	237 (31.0)	<0.001					
BMI (kg/m <sup>2</sup> )	27.9 ± 5.4	28.0 ± 4.9	27.7 ± 5.7	28.0 ± 5.7	0.381					
LVEF (%)	30 (25–36)	30 (25–35)	30 (25–37)	30 (23–37)	0.247					
HF aetiology										
Ischaemic	1045 (46.3)	345 (45.6)	339 (45.4)	361 (47.9)	0.369					
Cardiomyopathy	577 (25.6)	245 (32.4)	176 (23.6)	156 (20.7)	<0.001					
Hypertension	232 (10.3)	77 (10.2)	89 (11.9)	66 (8.8)	0.13					
Valvular disease	169 (7.5)	23 (3.0)	63 (8.5)	83 (11.0)	<0.001					
Other	130 (5.8)	29 (3.8)	55 (7.4)	46 (6.1)	0.012					
Unknown	113 (5.0)	41 (5.4)	27 (3.6)	45 (6.0)	0.094					
Estimated protein intake (g/day)	55 ± 11	58 ± 12	55 ± 11	52 ± 9	<0.001					
KCCQ score	50 ± 22	59 ± 21	49 ± 21	40 ± 21	<0.001					
<b>Laboratory</b>										
Iron (mg/dL)	45 (28–73)	62 (39–84)	45 (28–73)	34 (22–56)	<0.001					
Transferrin (mg/dL)	200 (160–250)	200 (170–240)	200 (160–250)	210 (160–260)	0.006					
Iron deficiency <sup>b</sup>	1413 (61.5)	323 (42.2)	474 (61.9)	616 (80.5)	<0.001					
Hepcidin (nmol/L)	6.3 (2.2–16.4)	9.0 (5.0–20.7)	7.5 (3.1–18.6)	2.3 (0.8–8.4)	<0.001	0.004 (–0.01 to 0.01)	0.36	–0.02 (–0.02 to –0.01)	<0.001	
sTfR (mg/L)	1.5 (1.2–2.1)	1.3 (1.0–1.6)	1.5 (1.2–2.0)	2.0 (1.4–2.8)	<0.001	–0.02 (–0.05 to 0.004)	0.08	0.04 (0.03–0.05)	<0.001	
Transferrin saturation (%)	17 (11–25)	22 (16–29)	17 (12–24)	12 (8–18)	<0.001	0.01 (–0.02 to 0.03)	0.61	–0.03 (–0.04 to –0.02)	<0.001	
Ferritin (µg/L)	102 (50–193)	143 (83–240)	103 (58–190)	63 (31–139)	<0.001	0.01 (–0.01 to 0.02)	0.44	–0.02 (–0.02 to –0.01)	<0.001	
IL-6 (pg/mL)	5.1 (2.8–10.2)	3.0 (1.8–5.6)	5.1 (3.0–8.9)	8.6 (5.0–16.1)	<0.001					
CRP (mg/L)	13 (6–26)	8 (4–19)	14 (6–26)	18 (9–32)	<0.001					
Haemoglobin (g/dL)	13.2 ± 1.9	13.8 ± 1.6	13.3 ± 1.9	12.6 ± 1.9	<0.001					
Anaemia	753 (36.0)	136 (20.5)	252 (35.6)	365 (50.3)	<0.001					
NT-proBNP (ng/L)	2680 (1173–5696)	1302 (511–2811)	2913 (1429–5357)	4688 (2408–9852)	<0.001					
<b>Medication</b>										
Beta-blockers	1912 (83.2)	666 (86.9)	637 (83.2)	609 (79.6)	<0.001					
Beta-blockers (on target dose)	125 (5.4)	31 (4.0)	41 (5.4)	53 (6.9)	0.013					
ACEi/ARBs	1662 (72.4)	601 (78.5)	565 (73.8)	496 (64.8)	<0.001					
ACEi/ARBs (on target dose)	300 (13.1)	133 (17.4)	99 (12.9)	68 (8.9)	<0.001					
Loop diuretics	2287 (99.6)	760 (99.2)	763 (99.6)	764 (99.9)	0.053					
Aldosterone antagonists	1235 (53.8)	421 (55.0)	403 (52.6)	411 (53.7)	0.628					
Proton pump inhibitors	806 (35.1)	213 (27.8)	269 (35.1)	324 (42.4)	<0.001					
P2Y <sub>12</sub> inhibitors	356 (15.5)	137 (17.9)	118 (15.4)	101 (13.2)	0.011					
Oral anticoagulants	881 (38.4)	225 (29.4)	333 (43.5)	323 (42.2)	<0.001					

Values are given as mean ± standard deviation, n (%), or median (interquartile range).

ACEi, angiotensin-converting enzyme inhibitor; ARB, angiotensin receptor blocker; BMI, body mass index; CI, confidence interval; CRP, C-reactive protein; FGF23, fibroblast growth factor 23; IL-6, interleukin-6; KCCQ, Kansas City Cardiomyopathy Questionnaire; LVEF, left ventricular ejection fraction; NT-proBNP, N-terminal pro B-type natriuretic peptide; sTfR, soluble transferrin receptor.

<sup>a</sup>Mediation analysis was performed for transferrin saturation, sTfR, ferritin and hepcidin. The direct effect is the effect of the respective iron status parameter on all-cause mortality after correcting for FGF23; the indirect effect is the FGF23-mediated effect of the respective iron status parameter on all-cause mortality. The total effect of iron status on mortality is a composite of the indirect (FGF23-related) and direct (non-FGF23-related) pathways. CIs are bootstrapped 2000 times. The mediation model is corrected for age, haemoglobin, NT-proBNP, serum urea, and the use of beta-blockers at baseline.

<sup>b</sup>Iron deficiency was defined as transferrin saturation <20%.



iron parameters in these patients, similar to our findings.

Given the strong link between iron status and FGF23 levels, one may expect changes in FGF23 levels after iron administration. Conflicting data have been published on the effect of intravenous iron administration on FGF23 levels. In a small cohort of iron-deficient patients with HF and reduced

ejection fraction, c-terminal FGF23 levels decreased after ferric carboxymaltose (FCM) administration during 28 days of follow-up, with transient hypophosphataemia and short-term increase in intact FGF23 levels.<sup>9</sup> Contrarily, another study comparing FCM and iron sucrose in haemodialysis patients found a short-term drop in intact FGF23 and a rise in c-terminal FGF23 in patients

receiving FCM, whereas iron sucrose did not affect these parameters.<sup>12</sup> Currently, no large studies on the effect of different iron agents on FGF23 levels in HF have been published. When intravenous iron therapy substantially and persistently lowers FGF23 levels, this might provide an additional explanation for the mode of action of this treatment modality, besides its beneficial haematological effects.

Our study has several strengths and limitations. First, we used a large and well-described cohort of worsening HF patients, in which we measured FGF23 and multiple iron parameters. We acknowledge the observational nature of our study, making it challenging to directly study pathophysiological mechanisms. Second, using the FGF23 assay we used for our study, we could not distinguish c-terminal FGF23 from intact FGF23. Finally, most patients in BIOSTAT-CHF have a reduced ejection fraction. Our data should therefore be cautiously interpreted in HF patients with a preserved ejection fraction.

### Funding

The BIOSTAT-CHF study was supported by the European Commission (FP7-242209-BIOSTAT-CHF).

**Conflict of interest:** The University Medical Center Groningen, which employs several authors, has received research grants and/or fees from AstraZeneca, Abbott, Bristol-Myers Squibb, Novartis, Roche Diagnostics, Trevena, and Thermofisher GmbH. N.G.B. received personal fees from Vifor Pharma. D.J.v.V. received board membership fees or travel expenses from Novartis, Johnson & Johnson, and CorviaMedical. A.A.V. received consultancy fees and/or research grants from Amgen, Applied Therapeutics, AstraZeneca, Bayer, Boehringer Ingelheim, Cytokinetics, GSK, Merck, Myokardia, Novartis, Roche Diagnostics, Servier, Vifor. P.v.d.M. received consultancy fees and/or grants from Novartis, Servier, Vifor Pharma, AstraZeneca, Pfizer and Ionis. The other authors have nothing to disclose.

**Haye H. van der Wal<sup>1</sup>,  
Niels Grote Beverborg<sup>1,2</sup>,  
Jozine M. ter Maaten<sup>1</sup>,  
Joanna S.J. Vinke<sup>3</sup>, Martin H. de Borst<sup>3</sup>,  
Dirk J. van Veldhuisen<sup>1</sup>,  
Adriaan A. Voors<sup>1</sup>, and  
Peter van der Meer<sup>1\*</sup>**

<sup>1</sup>Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; <sup>2</sup>Department of Cell and Molecular Biology, Karolinska Institutet, Stockholm, Sweden; and <sup>3</sup>Department of Internal Medicine, Division of Nephrology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

\*Email: p.van.der.meer@umcg.nl

### References

- van der Wal HH, Grote Beverborg N, Dickstein K, Anker SD, Lang CC, Ng LL, van Veldhuisen DJ, Voors AA, van der Meer P. Iron deficiency in worsening heart failure is associated with reduced estimated protein intake, fluid retention, inflammation, and antiplatelet use. *Eur Heart J* 2019;**40**:3616–3625.
- McDonagh T, Damy T, Doehner W, Lam CSP, Sindone A, van der Meer P, Cohen-Solal A, Kindermann I, Manito N, Pfister O, Pohjantähti-Maaroos H, Taylor J, Comin-Colet J. Screening, diagnosis and treatment of iron deficiency in chronic heart failure: putting the 2016 European Society of Cardiology heart failure guidelines into clinical practice. *Eur J Heart Fail* 2018;**20**:1664–1672.
- Klip IT, Comin-Colet J, Voors AA, Ponikowski P, Enjuanes C, Banasiak W, Lok DJ, Rosentryt P, Torrens A, Polonski L, van Veldhuisen DJ, van der Meer P, Jankowska EA. Iron deficiency in chronic heart failure: an international pooled analysis. *Am Heart J* 2013;**165**:575–582.e3.
- Wolf M, White KE. Coupling fibroblast growth factor 23 production and cleavage: iron deficiency, rickets, and kidney disease. *Curr Opin Nephrol Hypertens* 2014;**23**:411–419.
- Ter Maaten JM, Voors AA, Damman K, van der Meer P, Anker SD, Cleland JG, Dickstein K, Filippatos G, van der Harst P, Hillege HL, Lang CC, Metra M, Navis G, Ng L, Ouwerkerk W, Ponikowski P, Samani NJ, van Veldhuisen DJ, Zannad F, Zwinderman AH, de Borst MH. Fibroblast growth factor 23 is related to profiles indicating volume overload, poor therapy optimization and prognosis in patients with new-onset and worsening heart failure. *Int J Cardiol* 2018;**253**:84–90.
- Almahmoud MF, Soliman EZ, Bertoni AG, Kestenbaum B, Katz R, Lima JA, Ouyang P, Miller PE, Michos ED, Herrington DM. Fibroblast growth factor-23 and heart failure with reduced versus preserved ejection fraction: MESA. *J Am Heart Assoc* 2018;**7**:e008334.
- Eisenga MF, van Londen M, Leaf DE, Nolte IM, Navis G, Bakker SJ, de Borst MH, Gaillard CA. C-terminal fibroblast growth factor 23, iron deficiency, and mortality in renal transplant recipients. *J Am Soc Nephrol* 2017;**28**:3639–3646.
- Bockmann I, Lischka J, Richter B, Deppe J, Rahn A, Fischer DC, Heineke J, Haffner D, Leifheit-Nestler M. FGF23-mediated activation of local RAAS promotes cardiac hypertrophy and fibrosis. *Int J Mol Sci* 2019;**20**:4634.
- Stohr R, Sandstede L, Heine GH, Marx N, Brandenburg V. High-dose ferric carboxymaltose in patients with HFrEF induces significant hypophosphatemia. *J Am Coll Cardiol* 2018;**71**:2270–2271.
- David V, Martin A, Isakova T, Spaulding C, Qi L, Ramirez V, Zumbrennen-Bullough KB, Sun CC, Lin HY, Babitt JL, Wolf M. Inflammation and functional iron deficiency regulate fibroblast growth factor 23 production. *Kidney Int* 2016;**89**:135–146.
- McMahon S, Grondin F, McDonald PP, Richard DE, Dubois CM. Hypoxia-enhanced expression of the proprotein convertase furin is mediated by hypoxia-inducible factor-1: impact on the bioactivation of proproteins. *J Biol Chem* 2005;**280**:6561–6569.
- Roberts MA, Huang L, Lee D, MacGinley R, Troster SM, Kent AB, Bansal SS, Macdougall IC, McMahon LP. Effects of intravenous iron on fibroblast growth factor 23 (FGF23) in haemodialysis patients: a randomized controlled trial. *BMC Nephrol* 2016;**17**:177.