

University of Groningen

## Alternative Sugar Sources for Biobased Chemicals

Abdilla - Santes, Ria

DOI:  
[10.33612/diss.127600956](https://doi.org/10.33612/diss.127600956)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2020

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Abdilla - Santes, R. (2020). *Alternative Sugar Sources for Biobased Chemicals*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.127600956>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Stellingen  
Behorende bij het proefschrift  
Alternative Sugar Sources for Biobased Chemicals  
Ria Mayasari Abdilla-Santes

1. Sugar is the new oil.
2. The exact sugar content in the sugar fraction of pyrolytic liquid is difficult to determine as direct analytical techniques are absent.
3. The kinetic model for 5-HMF synthesis from glucose as developed by Chang *et al.* is unrealistic as it does not consider the consecutive reaction of 5-HMF to humins (Chang, C. Ma, X. and Cen, P., (2006). *Chin. J. Chem. Eng.*, 14, 708-712).
4. The proposed reaction network for humins as proposed by Kruse *et al.* is not correct as it does not involve the direct conversion of sugars to humins (Jung, D., Zimmermann, M., Kruse, A., (2018). *ACS Sustainable Chem. Eng.*, 6, 13877–13887).
5. Amberlyst 16 is preferred above sulfuric acid for the hydrolysis of levoglucosan to glucose (This thesis, Chapter 2 and 3).
6. The procedure used by Dogu *et al.* to calculate effective diffusion coefficients is scientifically not justified (Oktar, N., Mürtezaoğlu, K., Doğu, G., & Doğu, T. (1999). *Can. J. Chem. Eng.*, 77(2), 406-412; Doğu, T. and Doğu, G., (1991). *Chem. Eng. Comm.*, 103, 1-9).
7. Impurities in thick juice surprisingly have a positive impact on HMF selectivity and yield (this thesis, Chapter 5 and 6).
8. The experimental observation that HMF is formed from sucrose at basic conditions (this thesis, Chapter 6) is surprising when considering that HMF is not observed when using fructose as the feed (Kuster, B. F., & Temmink, H. M. (1977). *Carbohydrate research*, 54(2), 185-191; Körner, P., Jung, D., & Kruse, A. (2019). *ChemistryOpen*, 8(8), 1121-1132).
9. Caramel – a seemingly simple food ingredient, is in fact is a complex mixture of several thousand compounds formed by a large number of unselective reactions.
10. Performing experiments in the laboratory is like cooking and baking. All require precision, good temperature distribution and control, patience and experience to achieve best results.