

University of Groningen

Screening for cardiovascular disease risk using traditional risk factor assessment or coronary artery calcium scoring

van der Aalst, Carlijn M; Denissen, Sabine J A M; Vonder, Marleen; Gratama, Jan Willem C; Adriaansen, Henk J; Kuijpers, Dirkjan; Vliegenthart, Rozemarijn; van Lennep, Jeanine E Roeters; van der Harst, Pim; Braam, Richard L

Published in:

European Heart Journal - Cardiovascular Imaging

DOI:

[10.1093/ehjci/jeaa168](https://doi.org/10.1093/ehjci/jeaa168)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Aalst, C. M., Denissen, S. J. A. M., Vonder, M., Gratama, J. W. C., Adriaansen, H. J., Kuijpers, D., Vliegenthart, R., van Lennep, J. E. R., van der Harst, P., Braam, R. L., van Dijkman, P. R. M., van Bruggen, R., Oudkerk, M., & de Koning, H. J. (2020). Screening for cardiovascular disease risk using traditional risk factor assessment or coronary artery calcium scoring: the ROBINSCA trial. *European Heart Journal - Cardiovascular Imaging*, 21(11), 1216-1224. <https://doi.org/10.1093/ehjci/jeaa168>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Screening for cardiovascular disease risk using traditional risk factor assessment or coronary artery calcium scoring: the ROBINSICA trial

Carlijn M. van der Aalst^{1*}, Sabine J.A.M. Denissen^{1*}, Marleen Vonder²,
Jan Willem C. Gratama³, Henk J. Adriaansen⁴, Dirkjan Kuijpers^{5,6},
Rozemarijn Vliegthart², Jeanine E. Roeters van Lennep⁷,
Pim van der Harst^{2,8,9}, Richard L. Braam¹⁰, Paul R.M. van Dijkman^{11,12},
Rykel van Bruggen¹³, Matthijs Oudkerk^{14,15}, and Harry J. de Koning¹

¹Department of Public Health, Erasmus Medical Centre, PO Box 2040, 3000 CA Rotterdam, The Netherlands; ²Centre for Medical Imaging North-East Netherlands (CMI-NEN), University Medical Centre Groningen, Hanzeplein 1, EB45, Groningen 9713 GZ, The Netherlands; ³Department of Radiology and Nuclear Medicine, Gelre Hospitals, Albert Schweitzerlaan 31, 7334 DZ Apeldoorn, The Netherlands; ⁴Department of Clinical Chemistry and Hematology Laboratory, Gelre Hospitals, Albert Schweitzerlaan 31, 7334 DZ Apeldoorn, The Netherlands; ⁵Department of Radiology, University Medical Centre Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands; ⁶Department of Radiology, Haaglanden Medical Centre Bronovo, Bronovolaan 5, 2597 AX Den Haag, The Netherlands; ⁷Department of Internal Medicine, Erasmus Medical Centre, Doctor Molewaterplein 40, 3015 GD Rotterdam, The Netherlands; ⁸Department of Cardiology, University Medical Centre Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands; ⁹Department of Cardiology, University Medical Centre Utrecht, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands; ¹⁰Department of Cardiology, Gelre Hospitals, Albert Schweitzerlaan 31, 7334 DZ Apeldoorn, The Netherlands; ¹¹Department of Cardiology, Leids University Medical Centre, Albinusdreef 2, 2333 ZA Leiden, The Netherlands; ¹²Department of Cardiology, Haaglanden Medical Centre Bronovo, Bronovolaan 5, 2597 AX Den Haag, The Netherlands; ¹³General practice, Arnhemseweg 2 A, 7331 BK Apeldoorn, The Netherlands; ¹⁴University of Groningen, University Medical Centre Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands; and ¹⁵Institute for Diagnostic Accuracy—iDNA, Prof. E.D. Wiersmastraat 5, 9713 GH Groningen, The Netherlands

Received 6 March 2020; editorial decision 14 May 2020; accepted 14 May 2020

Aims

Screening for a high cardiovascular disease (CVD) risk followed by preventive treatment can potentially reduce coronary heart disease-related morbidity and mortality. ROBINSICA (Risk Or Benefit IN Screening for Cardiovascular disease) is a population-based randomized controlled screening trial that investigates the effectiveness of CVD screening in asymptomatic participants using the Systematic COronary Risk Evaluation (SCORE) model or coronary artery calcium (CAC) scoring. This study describes the distributions in risk and treatment in the ROBINSICA trial.

Methods and results

Individuals at expected elevated CVD risk were randomized into screening arm A ($n = 14\,478$; SCORE, 10-year fatal and non-fatal risk); or screening arm B ($n = 14\,450$; CAC scoring). Preventive treatment was largely advised according to current Dutch guidelines. Risk and treatment differences between the screening arms were analysed. A total of 12 185 participants (84.2%) in arm A and 12 950 (89.6%) in arm B were screened. In total, 48.7% were women, and median age was 62 (interquartile range 10) years. SCORE screening identified 45.1% at low risk (SCORE < 10%), 26.5% at intermediate risk (SCORE 10–20%), and 28.4% at high risk (SCORE ≥ 20%). According to CAC screening, 76.0% were at low risk (Agatston < 100), 15.1% at high risk (Agatston 100–399), and 8.9% at very high risk (Agatston ≥ 400). CAC scoring significantly reduced the number of individuals indicated for preventive treatment compared to SCORE (relative reduction women: 37.2%; men: 28.8%).

* Corresponding author. Tel: +31 10 7038673; Fax: +31 10 7038475. E-mail: s.denissen@erasmusmc.nl

† The first two authors contributed equally to this work.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2020. For permissions, please email: journals.permissions@oup.com.

Conclusion

We showed that compared to risk stratification based on SCORE, CAC scoring classified significantly fewer men and women at increased risk, and less preventive treatment was indicated.

Trial registration number

NTR6471.

Keywords

Cardiovascular disease • Computed tomography imaging • Coronary artery calcification • Population-based screening • Risk prediction • ROBINSICA trial

Introduction

Cardiovascular disease (CVD) is the leading cause of death worldwide and it is responsible for 45% of all annual deaths in Europe.^{1,2} Although various preventive measures in terms of lifestyle and timely drug treatment are known to reduce CVD burden, their application is suboptimal and unhealthy lifestyles remain frequent. Population-based screening for cardiovascular risk aims to identify individuals at increased risk in order to stop or delay disease progression by preventive treatment. This might be an appropriate strategy to reduce CVD-related events.^{3–5} However, there is no evidence from randomized controlled trials (RCTs) on the effectiveness of screening and a reliable screening modality yet.

One potentially suitable risk assessment tool is the Dutch Systematic Coronary Risk Evaluation (SCORE) risk model, which predicts 10-year risk for developing fatal and non-fatal CVD.^{4,6,7} Although this model is easy to use and is integrated into current guidelines, it has limited accuracy in predicting the correct risk status. The indication for preventive treatment is often uncertain in intermediate-risk individuals, limiting the ability to prevent coronary heart disease (CHD) in this group.^{4,8} Another potential screening modality is quantification of coronary artery calcification (CAC), expressed as CAC score, using computed tomography (CT) scanning.^{9,10} Evidence shows that CAC scoring is a strong independent predictor of CHD events and improves classification of intermediate-risk individuals, causing a large shift in the distribution of CVD risk.^{11,12} Currently, European and American guidelines recommend considering additional CAC scoring to guide preventive therapy decisions in intermediate-risk adults.^{4,13}

In CVD screening, the expected difference in CVD risk distribution between the SCORE model and CAC scoring might cause an effective shift towards more correctly classified individuals and more accurate risk reduction. In addition, a reduction in preventive overtreatment with cardiovascular medication is expected when CAC scoring is used as the screening modality. This will not only be beneficial for participants as it reduces potential side effects, but it will also save costs.¹⁴ However, the effect of the shift in risk distribution in the setting of CVD screening in an elevated risk population is unknown.

The Risk Or Benefit IN Screening for CArdiovascular diseases (ROBINSICA) trial is a population-based randomized controlled screening trial to investigate whether screening for a high risk of CVD in asymptomatic individuals followed by early treatment will reduce CHD-related morbidity and mortality compared to

no screening.¹⁵ The SCORE model and CAC scoring are used as potential screening modalities. The aim of the present study is to present the CVD risk distributions in both screening arms and to investigate the shift in risk distribution and the potential reduction in preventive (over)treatment due to the use of different risk assessment tools.

Methods**Study population**

The design, objectives and recruitment of the ROBINSICA trial have been described previously.¹⁵ In summary, 394 058 individuals, women aged 55–74 years and men aged 45–74 years from three regions in the Netherlands, were selected from the national population registry, and received an invitation to participate, including an information brochure, a baseline questionnaire, a waist circumference measuring tape and a written informed consent form. Asymptomatic individuals were subsequently selected based on at least one of the following inclusion criteria: (i) a high self-measured waist circumference (≥ 88 cm for women and ≥ 102 cm for men); (ii) a high body mass index (BMI; ≥ 30 kg/m²); (iii) a family history of myocardial infarction or sudden death before the age of 65 years in first- or second-degree relatives; and/or (iv) current smoking. Exclusion criteria were: (i) previously diagnosed CVD; (ii) previous CVD surgery; (iii) prescription of a combination of cholesterol- and blood pressure-lowering medication; (iv) CAC score measurement in the past year; and/or (v) incomplete informed consent. In total, 43 447 eligible individuals were randomized (1:1:1) to either the control arm where usual care was continued, or to one of the two intervention arms where screening was offered. All participants received generic healthy lifestyle recommendations of the Dutch Heart Foundation (Figure 1). The current study focuses only on the screening arms.

Screening

Screening was performed from 2015 to 2018. In intervention arm A, the 10-year risk for fatal and non-fatal CVD was estimated using the adapted version of the SCORE model as described in the Dutch guideline for Cardiovascular Risk Management (CVRM, edition 2011) by the College of General Practitioners.⁷ Participants were invited for blood pressure and cholesterol measurement. The algorithm stratifies participants into low (SCORE <10%), intermediate (SCORE 10–20%), or high (SCORE $\geq 20\%$) risk according to the guideline.¹⁵ In intervention arm B, participants underwent CT scanning using a second-generation dual-source CT system. The CAC imaging protocol has been described elsewhere.¹⁶ In short, images were analysed with semiautomatic identification of calcifications. A calcification was defined as an area with a density of ≥ 130 Hounsfield units and ≥ 2 adjacent voxels. Individual calcifications per




Figure 1 Flowchart of the recruitment, inclusion, and randomization process in the ROBINSICA trial.

Table 1 Baseline characteristics of the study population split for women (a) and men (b)

	Intervention arm A SCORE, n/N (%)	Intervention arm B CAC score, n/N (%)	P-value
a) Women	N = 6009	N = 6223	
Median age (IQR)	64.0 (8)	64.0 (8)	0.786
Educational level ^a			0.278
Low	2692/5987 (45.0)	2699/6200 (43.5)	
Medium	1454/5987 (24.3)	1552/6200 (25.0)	
High	1841/5987 (30.7)	1949/6200 (31.4)	
Current smoker	827/6009 (13.8)	850/6223 (13.7)	0.868
BMI, median (IQR)	25.5 (5.1)	25.5 (5.0)	0.826
Waist circumference, ^b median (IQR)	97.0 (13.5)	96.5 (13.5)	0.277
Family history of CHD ^c	2451/5437 (45.1)	2518/5614 (44.9)	0.810
Diabetes mellitus	152/6009 (2.5)	178/6223 (2.9)	0.259
Hypertension in past year	948/5864 (16.2)	1005/6080 (16.5)	0.592
Hypercholesterolaemia in past year	938/5802 (16.2)	974/5994 (16.2)	0.903
Baseline medical treatment			
Antihypertensive	1306/5989 (21.8)	1370/6203 (22.1)	0.709
Lipid-lowering	449/5987 (7.5)	490/6189 (7.9)	0.388
b) Men	N = 6176	N = 6727	
Age, median (IQR)	59.0 (13)	59.0 (13)	0.095
Educational level ^a			0.976
Low	1900/6165 (30.8)	2057/6705 (30.7)	
Medium	1840/6165 (29.8)	2012/6705 (30.0)	
High	2425/6165 (39.3)	2636/6705 (39.3)	
Current smoker	1525/6176 (24.7)	1736/6727 (25.8)	0.146
BMI, median (IQR)	26.9 (4.3)	26.9 (4.4)	0.758
Waist circumference, ^b median (IQR)	104.5 (12.0)	104.5 (12.0)	0.647
Family history of CHD ^c	2637/5718 (46.1)	2812/6262 (44.9)	0.183
Diabetes mellitus	200/6176 (3.2)	258/6727 (3.8)	0.067
Hypertension in past year	964/6020 (16.0)	1119/6561 (17.1)	0.116
Hypercholesterolaemia in past year	917/5997 (15.3)	1001/6541 (15.3)	0.985
Baseline medical treatment			
Antihypertensive	1017/6157 (16.5)	1187/6710 (17.7)	0.078
Lipid-lowering	493/6152 (8.0)	585/6709 (8.7)	0.149

A P-value of <0.005 was considered statistically significant after application of the Bonferroni correction.

CAC, coronary artery calcium; CHD, coronary heart disease; IQR, interquartile range; SCORE, systematic coronary risk evaluation.

^aEducational levels: low: primary, lower secondary general, or lower vocational education; medium: intermediate vocational or higher secondary education; and high: higher vocational education or university.

^bWaist circumference in centimetres.

^cFamily history of myocardial infarction or sudden death before the age of 65 years in first and second degree relatives.

women according to the SCORE model, compared to 1043 out of 6223 (16.8%) women according to CAC scoring ($P < 0.001$; absolute reduction of 9.9%). The relative reduction in the number of women indicated for preventive drug treatment was estimated to be 37.2% when using CAC scoring compared to SCORE calculation (based on RR 0.628, 95% CI 0.586–0.673). Among men, 2666 out of 6176 (43.2%) were advised to start preventive drug treatment based on SCORE calculation, whereas 2068 out of 6727 (30.7%) received preventive drug treatment advice based on CAC score ($P < 0.001$; absolute reduction of 12.4%). Risk estimation using CAC scoring caused a relative reduction in the number of preventive drug treatment indications of 28.8% in men (based on RR 0.712, 95% CI 0.680–0.746).

Discussion

In this population-based screening RCT for the early detection and treatment of an increased risk for CVD, 25 135 asymptomatic participants were screened by means of either applying the SCORE model or CAC scoring. As expected, the CVD risk distributions differed significantly between the two screening modalities. Risk assessment through CAC scoring identified more low-risk individuals compared to the SCORE model. Follow-up analyses should establish whether the indicated high-risk individuals were treated correctly.

The associations between traditional risk factors and a higher SCORE are a natural result of the SCORE model being based on these risk factors. However, similar associations were not observed

Table 2 Distributions of cardiovascular risks by baseline characteristics in intervention arm A (Dutch SCORE calculation) and intervention arm B (CAC quantification) in women

	Low risk SCORE <10%, ^a n/N (%)	Intermediate risk SCORE 10–20%, ^a n/N (%)	High risk SCORE ≥20%, ^a n/N (%)	P-value	Low risk CAC score 0–99, n/N (%)	High risk CAC score 100–399, n/N (%)	Very high risk CAC score ≥400, n/N (%)	P-value
Age, median (IQR)	3234/6009 (53.8) 60.0 (5)	1479/6009 (24.6) 67.0 (4)	1296/6009 (21.6) 71.0 (4)	<0.001	5180/6223 (83.2) 63.0 (9)	754/6223 (12.1) 66.5 (8)	289/6223 (4.6) 68.0 (8)	<0.001
Current smoker	371/3234 (11.5) 25.3 (5.0)	255/1479 (17.2) 25.7 (5.3)	201/1296 (15.5) 26.0 (5.1)	<0.001	654/5180 (12.6) 25.5 (5.0)	142/754 (18.8) 25.5 (5.3)	54/289 (18.7) 25.6 (5.7)	<0.001
BMI, median (IQR)	96.0 (13.5)	97.0 (13.5)	98.5 (14.5)	<0.001	96.0 (13.5)	97.0 (13.0)	97.5 (14.9)	0.653
Waist circumference, ^b median (IQR)	1352/2968 (45.6) 21/3234 (0.6)	598/1325 (45.1) 37/1479 (2.5)	501/1144 (43.8) 94/1296 (7.3)	0.596	2008/4684 (42.9) 114/5180 (2.2)	359/666 (53.9) 49/754 (6.5)	151/264 (57.2) 15/289 (5.2)	<0.001
Family history of CHD ^c	443/3173 (14.0) 475/3147 (15.1)	236/1445 (16.3) 267/1413 (18.9)	269/1246 (21.6) 196/1242 (15.8)	<0.001	786/5065 (15.5) 774/5004 (15.5)	151/737 (20.5) 141/717 (19.7)	68/278 (24.5) 59/273 (21.6)	<0.001
Diabetes mellitus				0.005				0.001
Hypertension in past year				<0.001				<0.001
Hypercholesterolaemia in past year				<0.001				<0.001
Baseline medical treatment				<0.001				<0.001
Antihypertensive	573/3227 (17.8)	360/1474 (24.4)	373/1288 (29.0)	<0.001	1077/5166 (20.8)	195/750 (26.0)	98/287 (34.1)	<0.001
Lipid-lowering	186/3226 (5.8)	132/1472 (9.0)	131/1289 (10.2)	<0.001	340/5153 (6.6)	107/750 (14.3)	43/286 (15.0)	<0.001

A P-value of <0.005 was considered statistically significant after application of the Bonferroni correction.

The risk distribution categories in intervention arm A are as in the Dutch Cardiovascular Risk Management protocol. The risk distribution categories in intervention arm B are as cut-offs from literature.

BMI, body mass index; CAC, coronary artery calcium; CHD, coronary heart disease; IQR, interquartile range; SCORE, systematic coronary risk evaluation.

^aSCORE calculation according to the 2011 edition of the Cardiovascular Risk Management protocol of the Dutch College of General Practitioners.⁷

^bWaist circumference in centimetres.

^cFamily history of myocardial infarction or sudden death before the age of 65 years in first and second degree relatives.

Table 3 Distributions of cardiovascular risks by baseline characteristics in intervention arm A (Dutch SCORE calculation) and intervention arm B (CAC quantification) in men

	Low risk SCORE <10%, ^a n/N (%)	Intermediate risk SCORE 10–20%, ^a n/N (%)	High risk SCORE ≥20%, ^a n/N (%)	P-value	Low risk CAC score 0–99, n/N (%)	High risk CAC score 100–399, n/N (%)	Very high risk CAC score ≥400, n/N (%)	P-value
Age, median (IQR)	2262/6176 (36.6) 52.0 (6)	1751/6176 (28.4) 60.0 (7)	2163/6176 (35.0) 68.0 (7)	<0.001	4659/6727 (69.3) 56.0 (12)	1200/6727 (17.8) 63.0 (11)	868/6727 (12.9) 66.0 (9)	<0.001
Current smoker	432/2262 (19.1) 26.6 (4.3)	441/1751 (25.2) 27.1 (4.3)	652/2163 (30.1) 26.9 (4.2)	<0.001	1198/4659 (25.7) 26.9 (4.3)	318/1200 (26.5) 27.0 (4.6)	220/868 (25.3) 26.9 (4.4)	0.811
BMI, median (IQR)	104.0 (12.0) 104.6/2141 (48.9)	104.5 (11.5) 755/1640 (46.0)	105.0 (11.0) 836/1937 (43.2)	<0.001	104.0 (12.0) 1880/4341 (43.3)	104.5 (11.0) 512/1119 (45.8)	104.5 (12.0) 420/802 (52.4)	0.605
Family history of CHD ^c	19/2262 (0.8) 264/2202 (12.0)	24/1751 (1.4) 324/1706 (19.0)	157/2163 (7.3) 376/2112 (17.8)	<0.001	138/4659 (3.0) 694/4550 (15.3)	52/1200 (4.4) 237/1169 (20.3)	67/868 (7.7) 188/842 (22.3)	<0.001
Diabetes mellitus	298/2208 (13.5)	289/1694 (17.1)	330/2095 (15.8)	0.007	648/4546 (14.3)	209/1156 (18.1)	144/839 (17.2)	0.002
Hypertension in past year	198/2258 (8.8)	305/1746 (17.5)	514/2153 (23.9)	<0.001	647/4648 (13.9)	288/1196 (24.1)	252/866 (29.1)	<0.001
Hypercholesterolaemia in past year	124/2258 (5.5)	132/1741 (7.6)	237/2153 (11.0)	<0.001	323/4649 (6.9)	135/1194 (11.3)	127/866 (14.7)	<0.001
Baseline medical treatment								
Antihypertensive								
Lipid-lowering								

A P-value of <0.005 was considered statistically significant after application of the Bonferroni correction.


The risk distribution categories in intervention arm A are as in the Dutch Cardiovascular Risk Management protocol. The risk distribution categories in intervention arm B are as cut-offs from literature.

BMI, body mass index; CAC, coronary artery calcium; CHD, coronary heart disease; IQR, interquartile range; SCORE, systematic coronary risk evaluation.

^aSCORE calculation according to the 2011 edition of the Cardiovascular Risk Management protocol of the Dutch College of General Practitioners.⁷

^bWaist circumference in centimetres.

^cFamily history of myocardial infarction or sudden death before the age of 65 years in first and second degree relatives.



in intervention arm B: higher CAC score categories were not associated with increasing waist circumference in women and men, nor with current smoking in men, nor with increasing BMI in women. In men, BMI was not associated with a higher SCORE, nor a higher CAC score. Regarding BMI, previous studies indeed reported that BMI does not predict CAC, which is largely related to the inability of BMI to differentiate between fat and muscle and the assumption that CAC scores can be underestimated in women with large chest size and large patients.^{19,20} In contrast, the lack of an association between waist circumference and CAC contradicts earlier findings indicating that waist circumference is associated with CAC beyond traditional risk factors.²¹ As there is no unambiguous evidence on this subject yet, future research should focus more on this potential association. Furthermore, the proportion of male current smokers did not increase with higher CAC score categories. This is in line with previous research that concluded that the effect of current smoking on CAC might decrease with age.²² Discrepancies in presence of CAC and absence of traditional risk factors, and vice versa, might influence the decision on whether to start preventive drug treatment or not. In particular, current preventive treatment in people with zero CAC may be considered as overtreatment, since this score represents a minimal risk.²³ Current preventive treatment decisions are largely based on traditional risk prediction models, whereas CAC scoring is thought to be better at correctly identifying individuals who would benefit the most from preventive treatment.²⁴

The SCORE model has several limitations, including the limited adaptation for different ethnic groups and age ranges, and the lack of incorporating risk modifiers that potentially reclassify CVD risk, such as socio-economic status, CVD family history and obesity, and therefore lacks discriminative power.^{4,8} CAC scoring has superior discrimination and risk reclassification as compared with other risk indicators.²⁵ Previous studies showed that asymptomatic intermediate-risk individuals were more often downgraded to a lower-risk category after adding CAC scoring to risk prediction, which is in line with our results.^{11,12} In addition, the review of

Greenland *et al.*,⁹ which summarized the results of population-based cohorts, convincingly showed the value of CAC scoring as a single predictive cardiovascular risk marker beyond traditional risk factors. Furthermore, recent literature described that shared decision making guided by CAC scoring in intermediate-risk individuals can be a cost-effective strategy to avoid years of preventive medication.^{9,14} Future analyses on CVD-related events in the ROBINSICA trial might add important evidence on the extent to which preventive treatment decisions should be based on CAC screening.

The observed reduction in the number of individuals indicated to consult their GP for preventive treatment after screening by CAC scoring compared to screening using the SCORE model will potentially influence prevention strategies. However, future analyses on CVD-related events are needed to determine whether the indicated high-risk individuals were treated correctly. Within the screening setting of the current study, the results might imply a reduction in burden for both screening participants and GPs. The improved estimate of a CAC-based CVD risk status might reduce unnecessary stress that participants may experience upon receiving an unfavourable test result, while it might increase adherence to preventive treatment.²⁶ For GPs, risk management in intervention arm B participants is less time consuming since the treatment indication in intervention arm A is not solely based on the SCORE model, but also on additional risk-increasing factors that are not known in the ROBINSICA trial. Furthermore, a potential reduction in unnecessary treatment will reduce costs. However, as CT scanning is more expensive compared to using the SCORE model, the effectiveness of CT screening should first be confirmed.^{14,27}

The strength of this study is its large study population that was randomly selected from the national population registry. The aimed sample size was reached and therefore there should be sufficient power to show a reduction in CHD events of at least 15%.¹⁵ Furthermore, screening results were consistently obtained by adequately trained research personnel. A main limitation was that the presented data analysis is cross-sectional. Therefore, conclusions on the reduction of preventive overtreatment cannot be drawn yet. Future analyses on this subject are required. Another limitation was that recall bias might have caused some inaccuracies in the population characteristics data obtained from the self-reported baseline questionnaire. However, multiple questions were used per health topic to increase the reliability of the answers. Therefore, self-reported questionnaires are the preferred and most cost-effective method for obtaining data in large study populations. Another point is that the described treatment indications in intervention arm A are not completely comparable with preventive treatment based on the SCORE model in current practice. To maintain feasibility, not all risk-increasing factors that co-determine the treatment indication were incorporated in the screening as they are not part of the SCORE calculation itself. Lastly, the final decision regarding preventive treatment was made in consultation with the GP as GPs have access to participants' medical background. The role of GPs in the risk management of increased-risk individuals is important in the feasibility of a potential CVD screening programme.

Within this first population-based RCT on screening for a high risk of CHD, CAC scoring classified significantly fewer individuals at intermediate and high risk in both women and men compared to applying the SCORE model. Subsequently, the potential expected reduction in

preventive overtreatment favours the use of CAC scoring in screening. However, future analyses are required to confirm the effectiveness of CVD screening for reduction of CHD and to incorporate costs of CT scanning and preventive treatment. Should screening for a high risk of CVD be net-effective, large health gains will be achieved.

Supplementary data

Supplementary data are available at *European Heart Journal - Cardiovascular Imaging* online.

Acknowledgements

We would like to thank the Ministry of Health, Welfare and Sports for the ethical approval to perform the trial. Naturally, we thank all participants for their participation. Furthermore, we thank M. Quak for the extensive research assistance, RADventure for developing the data management system and IVA group for handling all questionnaires and letters. Lastly, we thank all employees of the participating centres for facilitating screening and for screening the participants.

Funding

This work was supported by an advanced grant of the European Research Council [294604].

Conflict of interest: H.J.d.K., M.O., and C.M.v.d.A. report grants from the European Research Council. R.V. reports grants from Siemens Healthineers outside the submitted work. No other author reports any conflict of interest.

References

- World Health Organization. Cardiovascular diseases (CVDs): World Health Organization; 2017. [http://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](http://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)) (17 May 2017, date last accessed: 4 June 2020).
- Wilkins E, Wilson L, Wickramasinghe K, Bhatnagar P, Leal J, Luengo-Fernandez R et al. European Cardiovascular Disease Statistics 2017. Brussels: European Heart Network; 2017.
- Kotseva K, De Bacquer D, De Backer G, Ryden L, Jennings C, Gyberg V et al. Lifestyle and risk factor management in people at high risk of cardiovascular disease. A report from the European Society of Cardiology European Action on Secondary and Primary Prevention by Intervention to Reduce Events (EUROASPIRE) IV cross-sectional survey in 14 European regions. *Eur J Prev Cardiol* 2016;**23**:2007–18.
- Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: the Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2016;**37**:2315–81.
- Sillescu H, Falk E. Why not screen for subclinical atherosclerosis? *Lancet* 2011; **378**:645–6.
- Conroy RM, Pyorala K, Fitzgerald AP, Sans S, Menotti A, De Backer G et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;**24**:987–1003.
- Dutch College of General Practitioners [Nederlands Huisartsen Genootschap (NHG)]. Multidisciplinary guideline cardiovascular risk management, revision 2011 [Multidisciplinaire richtlijn Cardiovasculair risicomanagement, herziening 2011]. 2011.
- Yeboah J, McClelland RL, Polonsky TS, Burke GL, Sibley CT, O'Leary D et al. Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA* 2012;**308**:788–95.
- Greenland P, Blaha MJ, Budoff MJ, Erbel R, Watson KE. Coronary calcium score and cardiovascular risk. *J Am Coll Cardiol* 2018;**72**:434–47.
- Budoff MJ, Young R, Burke G, Jeffrey Carr J, Detrano RC, Folsom AR et al. Ten-year association of coronary artery calcium with atherosclerotic cardiovascular disease (ASCVD) events: the multi-ethnic study of atherosclerosis (MESA). *Eur Heart J* 2018;**39**:2401–8.
- Erbel R, Mohlenkamp S, Moebus S, Schmermund A, Lehmann N, Stang A et al. Coronary risk stratification, discrimination, and reclassification improvement based on quantification of subclinical coronary atherosclerosis: the Heinz Nixdorf Recall study. *J Am Coll Cardiol* 2010;**56**:1397–406.
- Polonsky TS, McClelland RL, Jorgensen NW, Bild DE, Burke GL, Guerci AD et al. Coronary artery calcium score and risk classification for coronary heart disease prediction. *JAMA* 2010;**303**:1610–6.
- Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019;**74**:e177–232.
- Hong JC, Blankstein R, Shaw LJ, Padula WV, Arrieta A, Fialkow JA et al. Implications of coronary artery calcium testing for treatment decisions among statin candidates according to the ACC/AHA cholesterol management guidelines: a cost-effectiveness analysis. *JACC Cardiovasc Imaging* 2017;**10**:938–52.
- Van der Aalst CM, Vonder M, Gratama JM, Adriaansen HJ, Kuijpers D, Denissen S et al. Risk or Benefit in Screening for Cardiovascular Disease (ROBINSICA): the rationale and study design of a population-based randomized-controlled screening trial for cardiovascular disease. *J Clin Trials* 2019;**9**:1:1000361
- Vonder M, van der Aalst CM, Vliegenthart R, van Ooijen PMA, Kuijpers D, Gratama JW et al. Coronary artery calcium imaging in the ROBINSICA trial: rationale, design, and technical background. *Acad Radiol* 2018;**25**:118–28.
- Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M Jr, Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;**15**:827–32.
- Budoff MJ, Nasir K, McClelland RL, Detrano R, Wong N, Blumenthal RS et al. Coronary calcium predicts events better with absolute calcium scores than age-sex-race/ethnicity percentiles: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol* 2009;**53**:345–52.
- Roy SK, Zeb I, Kadakia J, Li D, Budoff MJ. Body surface area is a predictor of coronary artery calcium, whereas body mass index is not. *Coron Artery Dis* 2012;**23**:113–7.
- Willeminck MJ, Abramovic B, den Harder AM, van der Werf NR, de Jong PA, Budde RP et al. Coronary calcium scores are systematically underestimated at a large chest size: a multivendor phantom study. *J Cardiovasc Comput Tomogr* 2015; **9**:415–21.
- Bose S, Krishnamoorthy P, Varanasi A, Nair J, Schutta M, Braunstein S et al. Measurement of waist circumference predicts coronary atherosclerosis beyond plasma adipokines. *Obesity (Silver Spring)* 2013;**21**:E118–23.
- Lehmann N, Mohlenkamp S, Mahabadi AA, Schmermund A, Roggenbuck U, Seibel R et al. Effect of smoking and other traditional risk factors on the onset of coronary artery calcification: results of the Heinz Nixdorf recall study. *Atherosclerosis* 2014;**232**:339–45.
- Valenti V, Ó Hartaigh B, Heo R, Cho I, Schulman-Marcus J, Gransar H et al. A 15-year warranty period for asymptomatic individuals without coronary artery calcium: a prospective follow-up of 9,715 individuals. *JACC Cardiovasc Imaging* 2015;**8**:900–9.
- Elias-Smale SE, Proenca RV, Koller MT, Kavousi M, van Rooij FJ, Hunink MG et al. Coronary calcium score improves classification of coronary heart disease risk in the elderly: the Rotterdam study. *J Am Coll Cardiol* 2010;**56**:1407–14.
- Yeboah J, Young R, McClelland RL, Delaney JC, Polonsky TS, Dawood FZ et al. Utility of nontraditional risk markers in atherosclerotic cardiovascular disease risk assessment. *J Am Coll Cardiol* 2016;**67**:139–47.
- Johnson JE, Gulanick M, Penckofer S, Kouba J. Does knowledge of coronary artery calcium affect cardiovascular risk perception, likelihood of taking action, and health-promoting behavior change? *J Cardiovasc Nurs* 2015;**30**:15–25.
- Mamudu HM, Paul TK, Veeranki SP, Budoff M. The effects of coronary artery calcium screening on behavioral modification, risk perception, and medication adherence among asymptomatic adults: a systematic review. *Atherosclerosis* 2014; **236**:338–50.