

University of Groningen

Patient-reported outcomes after cardiac surgery

Zwiers-Blokzijl, Fredrike

DOI:
[10.33612/diss.131754816](https://doi.org/10.33612/diss.131754816)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Zwiers-Blokzijl, F. (2020). *Patient-reported outcomes after cardiac surgery: Things that really matter*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.131754816>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER

6

BARRIERS THAT OBSTRUCT RETURN TO WORK AFTER CORONARY BYPASS SURGERY – A QUALITATIVE STUDY

**Fredrike Blokzijl, Marisa Onrust, Willem Dieperink,
Frederik Keus, Iwan C.C. van der Horst, Wolter Paans,
Massimo A. Mariani, Michiel F. Reneman.**

Submitted

ABSTRACT

Purpose: Coronary artery bypass grafting is the most frequently performed cardiac surgical procedure. Despite its benefits on survival and quality of life, it is associated with a considerable financial burden on society including sick leave. Our study aimed to explore the barriers that obstruct return to work after coronary artery bypass grafting.

Methods: We performed a qualitative study with in-depth interviewing of patients six months after their surgery. We included ten working patients and interviewed them and their spouses at home. The interviews were transcribed and two investigators independently searched the transcriptions for barriers that had obstructed return to work.

Results: Based on the interviews we were able to distinguish four main groups of barriers: 'personal', 'healthcare', 'work' and 'law & regulation.' The personal barriers were subgrouped in affective, physical, cognitive, social and individually determined factors.

Conclusions: In a qualitative study we showed that personal barriers as well as barriers regarding healthcare, work and law & regulation, were perceived by patients as important factors obstructing return to work after coronary artery bypass grafting. To overcome the identified barriers, the process of return to work could preferably be initiated during the hospital phase, started during cardiac rehabilitation, and coordinated by a case-managing professional.

INTRODUCTION

In the Netherlands, over 15,000 cardiac surgeries are performed every year (1). The main reasons to offer cardiac surgery, including coronary artery bypass grafting (CABG) are to improve survival and quality of life (2). A relevant number of patients are at working age when undergoing coronary bypass and return to work (RTW) is an important goal during recovery (3,4). Because ischemic heart disease, including coronary artery disease, can be life-threatening, anxiety and uncertainty have been suggested as barriers for RTW, not only for the patient but also for the occupational physician (3,4). Guidelines of the Dutch society of occupational physicians describe how to guide employees with ischemic heart disease during the process of resuming their work (4). These guidelines are based on internationally conducted studies and recommend to resume work during cardiac rehabilitation (CR) to overcome possible barriers of fear and anxiety (4). The guidelines suggest partial or fully RTW within approximately six weeks for patients with coronary artery disease, including patients after CABG, provided that postoperative recovery was uncomplicated (4).

Only a few studies on return to work after CABG have been conducted. RTW is observed between 8 and 13 weeks after surgery in two studies with CABG patients not participating in cardiac rehabilitation-programs (5,6) while another study reported 64% of patients fully resuming work within six months after cardiac rehabilitation (7). There seems to be a notable discrepancy between guidelines recommendations on RTW after CABG and literature reports, although evidence is scarce. Younger age (6,8,9), higher job satisfaction (5), positive occupational expectations (7,10) and absence of diabetes and myocardial damage (6,8,11) have been suggested to facilitate quicker RTW but may not be considered comprehensive, because they are all personal factors. In RTW models, such as the case-management ecological model, work-related and other factors influencing the disability process are also present (12). While this model was developed for the case management of disability due to low back pain and has been applied in other medical conditions, it has not been used in patients after CABG.

In-depth interviewing of patients after CABG and their spouses may enrich our knowledge on the process of RTW after CABG from the patient perspective including the facilitators and barriers influencing this process. The aim of our study was to identify barriers that obstruct return to work of patients after CABG.

METHODS

Study design

The study is reported according to the Standards for Reporting Qualitative Research (SRQR) (13). A qualitative study design was applied to evaluate barriers that obstruct return to work in patients after CABG. Through in-depth interviewing of patients and their partners we explored personal experiences (14). An essential characteristic of the data collection by in-depth interviewing is that critical issues identified in one interview are used to refine questions and topics in the next

interview. This design of inductive inference provides the opportunity to elaborate each issue with each subsequent interview (14).

Participants

Participants were selected based on distinct inclusion criteria: all patients underwent elective, isolated CABG, were living with a spouse and had paid work before surgery. To ensure that other severe comorbidities or a complicated recovery did not affect the process of RTW, we excluded patients with the following preoperative comorbidities: ejection fraction <30%, stroke, psychiatric illnesses, chronic obstructive pulmonary disease (COPD) GOLD III-IV or renal disease (a reduced renal function prior to surgery with an estimated Glomerular Filtration Rate (eGFR) <60 ml/min/1.73 m² (15). We also excluded patients who participated in the Heart-ROCQ study, a randomized trial on an extended CR-program before and after cardiac surgery (16).

Sample and setting

All interviews were held at the participants' homes between six and seven months after their CABG surgery in line with an expected full recovery. Based on current literature, we defined that data saturation would be reached when in three subsequent interviews, no new additional themes emerged (17).

Data collection

The interviews were semi-structured, using an interview guide which was divided into different topics based on recent literature (13,17)(Appendix 1). The interviewer can use an interview guide as a list of topics and open-ended questions; the guide was not used as a questionnaire but rather as a memory aide during the interview (14). To start, some baseline questions were asked to identify narratives of people's lives (i.e. type of work, education level, working hours). All interviews were conducted by one of the researchers (FB) who works at the university hospital as a nurse practitioner and PhD-student at the department of cardiothoracic surgery. All interviews were audiotaped with permission from the participants.

Data analysis

After verbatim transcriptions, two researchers (FB and MO) independently performed data analyses following the thematic content approach, which focuses on identifying, analyzing and interpreting patterns of meaning within qualitative data (18). All transcriptions were examined to create one large preliminary list of barriers obstructing return to work as derived from the in-depth interviews (19). Both researchers independently created items and after inductive and axial coding using consensus techniques, more collectively shared underlying concepts were analysed leading to a codebook utilized by the qualitative analysis software package ATLAS.ti. (version 8, ATLAS.ti Scientific Software Development, Berlin) (20). Finally, higher-level grouping was guided by the case-management ecological model (12). The frequency of the individual barriers was assessed using

quantitative techniques by determining how frequent each item was mentioned. The frequency of all mentioned barriers was only used in a semi-quantitative approach to guide emphasis in the description of the results section.

Ethical considerations

The institution's ethical committee approved the protocol (reference number 2019/075) and waived the need for formal evaluation according to the Dutch Law on Scientific Medical Research with Humans. All participants were approached by telephone for their participation in this study and formal written informed consent was obtained prior to the interview. To ensure anonymity, all data that may plausibly identify any of the participants were eliminated from the transcripts.

RESULTS

All interviews were held between March and June 2019. Data saturation was reached after ten interviews. Baseline characteristics of the participants are listed in Table 1. Five participants had fully returned to work; two after two months and three within 4-6 months. The other five participants were still in the process of returning to work guided by an occupational physician. Two participants were self-employed. One participant reported that during hospital admittance, a nurse practitioner in the hospital had discussed the expectations regarding RTW with the participant. All patients had participated in a multidisciplinary CR-program after CABG, either a phase I program (inpatient CR-program) (n=2) or a phase II program (an early post-discharge outpatient CR-program (n=8). Nine participants reported that the process of RTW had started after having finished

Table 1. Characteristics of participants

Participant	Age (years)	Gender	Education-level	Occupation (collar)	Employment	Contract hours
1	57	male	high	white	employed	32
2	49	male	high	white	employed	40
3	62	male	low	blue	employed	40
4	61	male	medium	white	employed	40
5	56	male	high	white	employed	36
6	53	male	high	white	self-employed	> 40
7	56	male	medium	pink (service industry)	employed	40
8	63	male	medium	white	self-employed	> 40
9	59	female	high	white	employed	40
10	54	male	low	blue	employed	40

the rehabilitation program, and seven participants reported their impression that there was no communication between the physicians involved.

After verbatim transcription and analyses of the data, four main groups of barriers were defined: 'personal', 'healthcare' 'work' and 'law and regulation'. The personal barriers were divided into five subgroups: 'affective', 'physical', 'individually determined', 'social' and 'cognitive'. All barriers and associated items are listed in table 2.

Within the personal barrier, the subgroup concerning affective factors was mentioned most frequent, i.e. loss of self-confidence, anxiety, bad feeling due to physical complaints & limitations, and being unable to handle a lot of fuss.

Table 2. Barriers that obstruct return to work in patients after CABG

Personal	132 (55%)	Healthcare	63 (26%)
<i>Affective</i>	57	No advice concerning RTW	24
Loss of self-confidence	12	No guidance/follow-up after cardiac rehabilitation	15
Anxiety	10	Discouraging RTW	14
Feeling bad due to physical limitations	9	Employment consultant available but not involved	3
Not being able to handle a lot of fuss	9	Physicians conflicting opinions concerning RTW	3
Feeling of being out of balance	5	Pressure by occupational physician to RTW	3
Fear of becoming physically active again	4	Leaflet concerning RTW not applicable	1
Post-traumatic stress	3		
Awareness of suddenly being a heart patient	3	Work	38 (16%)
Grief/very emotional	2	Factors causing work stress	18
		Communication with supervisor	13
<i>Physical</i>	43	High costs to hire a replacement	3
Fatigue	17	Interaction with colleagues	3
Physical complaints due to other health issues	16	Sole employee in the own company	1
Loss of condition	8		
Chest pain	2	Law & regulation	6 (3%)
		Limited insurance when ill (self-employed)	3
<i>Individually determined</i>	13	Temporary contract: job loss due to illness	2
Feeling obliged to resume work	5	Uncertainty about income	1
No attention for others (focusing on yourself)	3		
Change in personality due to illness	2		
Feeling superfluous at work during reintegration	2		
Not being able to let go of work during reintegration	1		
<i>Social</i>	12		
Family experiencing stress	7		
Social environment advising to take it easy	5		
<i>Cognitive</i>	7		
Memory loss	4		
Concentration disorder	3		

Participants for example mentioned: "... the fear that you visit the cardiologist in the morning and are put in a wheelchair immediately and admitted because you are a ticking time bomb although you felt great, that realization of being a heart patient all of a sudden was very emotional ..." [P (participant) 1]. "...The fact that you simply could not have made it and luckily that is not the case but well, the thought was there because it is your heart..." [R (relative) 5]. One of the partners described the uncertainty as being out of balance: "...because being healthy is a balance and if you are physically out of balance, then this also impacts mentally and right now, the balance is lost in everything..." [R.5]. Among the physical barriers fatigue was often mentioned. Physical complaints due to other health issues and loss of condition were relevant as well. One participant mentioned: "...then I arrive at work and I am very busy, thousand things to do and when I go home I am exhausted while there is still plenty to do at home but I am not capable of doing anything after work..." [P.4].

Among the barriers concerning healthcare, patients mentioned the lack of advice concerning RTW, professionals discouraging RTW, and no follow-up after cardiac rehabilitation, as the most critical barriers in the process of RTW. Participants for example mentioned: "... when I asked my cardiologist about RTW he said to me that work did not interest him, my health was his priority, not work..." [P.3]. "... the occupational physician advised me to take it easy, but I did not know exactly what to do with this advice..." [P.4]. "... during cardiac rehabilitation there was an informative meeting in the hospital for cardiac patients, and we saw a short movie of a man who wanted to RTW after surgery, but that he did not succeed at all. That was not an uplifting video because my wife and I thought I would go back to work pretty soon..." [P.2]. Another participant: "... because actually after the rehabilitation then suddenly there is nothing left. Yes, then there is your job - you have to pick up all vocational and social activities again but there is no guidance or whatsoever..." [P.2].

In the work-related barrier, factors causing work stress (i.e. excessive workload, busy workplace, understaffed) and the communication with the supervisor (i.e. difficult relationship) were mentioned most frequently.

DISCUSSION

This qualitative research showed that personal as well as healthcare-, work- and law & regulation-related factors, are all barriers in returning to work after CABG. Personal barriers concerning affective or physical complaints were most frequently mentioned during the interviews. Remarkably, many participants were dealing with feelings of anxiety, loss of self-confidence and fatigue during RTW, although their CABG was uncomplicated from a medical perspective.

Other recent studies also revealed that affective factors play an essential role in RTW after a cardiac event and suggest to identify this as soon as possible

to allow for targeted psychosocial care (7,10,11). We did not identify other studies confirming our findings on physical barriers such as fatigue that impedes resumption of work after CABG. Previous studies though, focused on patients with coronary artery disease, including also patients after less invasive procedures such as percutaneous coronary interventions or patients treated by medication only. These patients deviate severely from patients after coronary bypass as CABG is a major invasive procedure with impact on the entire body both physically (i.e. a sternal wound, postoperative anemia) as well as mentally, likely leading to prolonged recovery. The invasiveness of the treatment likely explains the long-term complaints of fatigue and feelings of anxiety. Because CR-participants after CABG deviate from other CR-participants in general, further research may possibly lead to adapted guidelines for patients after cardiac surgery. The previously mentioned guidelines suggest RTW within six weeks but based on recent studies (5–7) and based on the findings in our study, this may not be feasible for patients after coronary bypass.

6

86

Another finding of our study are the healthcare-related barriers on RTW after CABG. Remarkably, there is either complete absence or contradictory advice and guidance from the healthcare professionals. Many participants mentioned that surgeons, cardiologists, general practitioners, or occupational physicians did either not advise them concerning RTW, or alternatively, the given advice was discouraging or unclear (e.g. the advice to ‘take it easy’). This pattern of contradictory, or lack of advice was previously recognized for patients after cholecystectomy (21). No or unclear advice may negatively interact with the mentioned feelings of anxiety and uncertainty among patients and their spouses. Also, participants indicated that there appeared to be no communication between the physicians involved. All participants joined a CR-program following international guidelines for patients with CAD, including patients after coronary bypass (22–24). An essential goal of CR is to optimize participation in society regarding different aspects of daily life, such as domestic, occupational, and recreational activities (23,25). While participants were positive about their CR-program, participants also mentioned that the lack of follow-up after CR induced feelings of uncertainty about their activities at home and in resuming work. These feelings of uncertainty became even stronger because participants were advised to start the process of RTW only *after* completing CR, which is in contrast with guideline recommendations. Returning to work during CR requires coordination between the occupational physician and the cardiac rehabilitation team. Our findings are in line with the suggestions for improvement mentioned in the Dutch rehabilitation guidelines (3). A case-managing professional as a central contact is insufficiently imbedded in current practice resulting in inadequately aligned treatment- and RTW-plans (3). Surprisingly, the only participants returning to work during CR and within two months after surgery were the participants who were self-employed.

After the interviews many participants reported that they felt the importance to improve the process of RTW after CABG, which was their incentive to participate

in this study. We fully agree with the participants as there is a lot to improve. RTW itself can improve quality of life and economic security from both the individual (26) and the societal perspective; a quicker RTW can lead to reductions in sick-leave and substantial savings in indirect costs. Two recent reviews on RTW for workers with musculoskeletal pain-related conditions and on workers with coronary heart disease revealed that there is evidence suggesting that long term absence is significantly reduced by multi-domain interventions (i.e., health-focused, service coordination and work modification interventions) (27,28). This model of care may also be of added value in RTW after CABG. Feelings of anxiety and uncertainty can possibly be reduced or overcome by an earlier start of the process of RTW causing positive feedback (self-efficacy) or by proper counseling by healthcare professionals.

This study has several limitations. We focused on individual aspects using a qualitative design and results should, therefore, be considered in an explorative perspective. Further quantitative studies are necessary on RTW after coronary bypass to demonstrate the quantitative importance of the identified barriers. We included patients with paid work and without severe comorbidities limiting generalisability. Also, we excluded patients with complicated recovery. Although we did not use member checking or participant feedback to see whether the participants recognized the results, it is reassuring that most factors reflect the items from the case-management ecological model and another Dutch study on barriers influencing RTW after surgery (21,29). The results may be different in countries without a social security system, so that factors influencing RTW differ as well in these countries. This study focused on perspectives of patients and their spouses. Future studies should also focus on perspectives of supervisors and coworkers, as well as healthcare professionals.

CONCLUSION

Several barriers play a role in the process of return to work after coronary bypass. Affective and physical barriers and the absence or contradictory guidance from the healthcare professionals involved were mentioned most often. As guidelines suggest, the process of RTW could preferably be initiated during the hospital phase, started during CR, and coordinated by a case-managing professional to overcome the barriers of RTW after CABG.

Acknowledgements

We thank Karin Havinga and Peter Zwiers for their help with processing the data.

Conflicts of interest

Dr. Mariani has received grants from AtriCure, Edwards Lifesciences, Abbott and Getinge, and has provided training for Livanova. All other authors declare that they have no conflict of interest.

REFERENCES

1. Vaartjes I, Dis I Van, Visseren FLJ, Bots ML. Heart- and Vascular Disease in the Netherlands 2015. Available at: <http://www.hartstichting.nl>
2. Windecker S, Kolh P, Alfonso F, Collet JP, Cremer J, Falk V, et al. 2014 ESC/EACTS Guidelines on myocardial revascularization: The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J*. 2014;35:2541–619.
3. Rehabilitation committee Dutch Association for Cardiology & Dutch Heart Foundation. Multidisciplinary Guideline for Cardiac Rehabilitation 2011. Available at: <https://www.nvvc.nl>
4. Dutch Association for Occupational and Occupational Medicine. Guideline for Ischemic heart diseases for occupational physicians. Quality Agency Dutch Association for Occupational and Occupational Medicine 2006. Available at: <https://www.nvab-online.nl>
5. Fiabane E, Argentero P, Calsamiglia G, Candura SM, Giorgi I, Scafa F, et al. Does job satisfaction predict early return to work after coronary angioplasty or cardiac surgery? *Int Arch Occup Environ Health*. 2013;86:561–9.
6. Maznyczka AM, Howard JP, Banning AS, Gershlick AH. A propensity matched comparison of return to work and quality of life after stenting or coronary artery bypass surgery. *Open Hear*. 2016;3:e000322.
7. Salzwedel A, Reibis R, Heidler MD, Wegscheider K, Völler H. Determinants of Return to Work After Multicomponent Cardiac Rehabilitation. *Arch Phys Med Rehabil*. 2019;4–7.
8. Hällberg V, Palomäki A, Kataja M, Tarkka M. Return to work after coronary artery bypass surgery. A 10-year follow-up study. *Scand Cardiovasc J*. 2009;43:277–84.
9. Salzwedel A, Reibis R, Wegscheider K, Eichler S, Buhler H, Kaminski S, et al. Cardiopulmonary exercise testing is predictive of return to work in cardiac patients after multicomponent rehabilitation. *Clin Res Cardiol*. 2016;105:257–67.
10. Salzwedel A, Reibis R, Hadzic M, Buhler H, Völler H. Patients' expectations of returning to work, co-morbid disorders and work capacity at discharge from cardiac rehabilitation. *Vasc Health Risk Manag*. 2019;15:301–8.
11. Reibis R, Salzwedel A, Abreu A, Corra U, Davos C, Doehner W, et al. The importance of return to work: How to achieve optimal reintegration in ACS patients. *Eur J Prev Cardiol*. 2019;26:1358–1369.
12. Costa-Black, Katia M; Feuerstein, M; Loisel. Handbook of work disability: Prevention and management. *Handbook of Work Disability: Prevention and Management*. New York:Springer;2013.
13. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting qualitative research: A synthesis of recommendations. *Acad Med*. 2014;89:1245–51.
14. Hennink M, Hutter I, Bailey A. *Qualitative Research Methods*. 2015th ed. London:Sage Publications; 2011.
15. Stevens PE. Evaluation and Management of Chronic Kidney Disease: Synopsis of the Kidney Disease: Improving Global Outcomes 2012 Clinical Practice Guideline. *Ann Intern Med*. 2013 Jun 4;158:825.
16. Hartog J, Blokzijl F, Dijkstra S, DeJongste MJL, Reneman MF, Dieperink W, et al. Heart Rehabilitation in patients awaiting Open heart surgery targeting to prevent

- Complications and to improve Quality of life (Heart-ROCQ): study protocol for a prospective, randomised, open, blinded endpoint (PROBE) trial. *BMJ Open*. 2019;9:e031738.
17. Saunders B, Sim J, Kingstone T, Baker S, Waterfield J, Bartlam B, et al. Saturation in qualitative research: exploring its conceptualization and operationalization. *Qual Quant*. 2018;52:1893–907.
 18. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006;3(2):77–101.
 19. Boeije H. *Analysis in Qualitative Research*. 2nd ed. London:Sage; 2010.
 20. ATLAS.ti 8 Windows User Manual. ATLAS.ti Scientific Software Development GmbH, Berlin 2019.
 21. Keus F, de Vries J, Gooszen HG, van Laarhoven CJHM. Assessing factors influencing return back to work after cholecystectomy: A qualitative research. *BMC Gastroenterol*. 2010;10:12.
 22. Woodruffe S, Neubeck L, Clark RA, Gray K, Ferry C, Finan J, et al. Australian Cardiovascular Health and Rehabilitation Association (ACRA) Core Components of Cardiovascular Disease Secondary Prevention and Cardiac Rehabilitation 2014. *Vol. 24, Heart Lung and Circulation*. 2015;24:430-41.
 23. Piepoli MF, Corrà U, Benzer W, Bjarnason-Wehrens B, Dendale P, Gaita D, et al. Secondary prevention through cardiac rehabilitation: from knowledge to implementation. A position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil*. 2010;17:1–17.
 24. Piepoli MF, Corrà U, Abreu A, Cupples M, Davos C, Doherty P, et al. Challenges in secondary prevention of cardiovascular diseases: A review of the current practice. *International Journal of Cardiology*. 2015;180:114-9.
 25. Balady GJ, Williams MA, Ades PA, Bittner V, Comoss P, Foody JAM, et al. Core Components of Cardiac Rehabilitation/Secondary Prevention Programs: 2007 Update. *J Cardiopulm Rehabil Prev*. 2007;27:121–9.
 26. Bhattacharyya MR, Perkins-Porras L, Whitehead DL, Steptoe A. Psychological and clinical predictors of return to work after acute coronary syndrome. *Eur Heart J*. 2007;28(2):160–5.
 27. Cullen KL, Irvin E, Collie A, Clay F, Gensby U, Jennings PA, et al. Effectiveness of Workplace Interventions in Return-to-Work for Musculoskeletal, Pain-Related and Mental Health Conditions: An Update of the Evidence and Messages for Practitioners. *J Occup Rehabil*. 2018;28:1–15.
 28. Hegewald J, Wegewitz UE, Euler U, Van Dijk JL, Adams J, Fishta A, et al. Interventions to support return to work for people with coronary heart disease. *Cochrane Database Syst Rev*. 2019; DOI: 10.1002/14651858.CD010748.pub2.
 29. Loisel P, Buchbinder R, Hazard R, Keller R, Scheel I, Van Tulder M, et al. Prevention of work disability due to musculoskeletal disorders: The challenge of implementing evidence. *J Occup Rehabil*. 2005;15:507–24.

6
