

University of Groningen

Reply to "Comment on 'Charge transport in disordered semiconducting polymers driven by nuclear tunneling'"

van der Kaap, N. J.; Katsouras, I.; Asadi, K.; Blom, P. W. M.; Koster, L. J. A.; de Leeuw, D. M.

Published in:
Physical Review B

DOI:
[10.1103/PhysRevB.102.066202](https://doi.org/10.1103/PhysRevB.102.066202)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van der Kaap, N. J., Katsouras, I., Asadi, K., Blom, P. W. M., Koster, L. J. A., & de Leeuw, D. M. (2020). Reply to "Comment on 'Charge transport in disordered semiconducting polymers driven by nuclear tunneling'". *Physical Review B*, 102(6), Article 066202. <https://doi.org/10.1103/PhysRevB.102.066202>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Reply to “Comment on ‘Charge transport in disordered semiconducting polymers driven by nuclear tunneling’”N. J. van der Kaap,¹ I. Katsouras,^{2,3} K. Asadi,² P. W. M. Blom,² L. J. A. Koster,¹ and D. M. de Leeuw²¹*Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands*²*Max Planck Institute for Polymer Research, Ackermannweg 10, 55128 Mainz, Germany*³*Holst Centre, High Tech Campus 31, 5656 AE Eindhoven, The Netherlands*

(Received 21 April 2020; accepted 16 July 2020; published 27 August 2020)

The authors reply to the Comment by Nenashev *et al.*DOI: [10.1103/PhysRevB.102.066202](https://doi.org/10.1103/PhysRevB.102.066202)

We welcome the comment by Nenashev *et al.* [1]. Indeed, as long as the characteristic frequency of the bath is large enough, nuclear tunneling causes an increasing mobility with electric field, in agreement with experiments, and in contrast to the Miller-Abrahams and Marcus mechanisms. Nenashev *et al.* are, however, correct in pointing out that in the limit of very large electric fields, the exponential

term will cause a decrease in the mobility. This decrease in mobility is neither seen in the experimental data in our paper, nor in the data in the paper by Asadi *et al.* [2], where the nuclear tunneling hop rate was also applied. This suggests that the energy difference in the hopping process is still smaller than the characteristic frequency of the bath.

[1] A. V. Nenashev, F. Gebhard, and S. D. Baranovskii, *Phys. Rev. B* **102**, 066201 (2020).

[2] K. Asadi, A. J. Kronemeijer, T. Cramer, L. J. A. Koster, P. W. M. Blom, and D. M. de Leeuw, *Nat. Commun.* **4**, 1710 (2013).