

University of Groningen

Parkinson's disease

Koerts, J.

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2009

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koerts, J. (2009). *Parkinson's disease: neuroimaging and clinical studies on cognition and depression*. [Thesis fully internal (DIV), University of Groningen, Faculty of Medical Sciences]. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

- DATATOP: a multicenter controlled clinical trial in early Parkinson's disease. Parkinson Study Group (1989). *Arch.Neurol.*, 46, 1052-1060.
- Aarsland, D., Alves, G., & Larsen, J. P. (2005). Disorders of motivation, sexual conduct, and sleep in Parkinson's disease. *Adv.Neurol.*, 96, 56-64.
- Abe, K., Takanashi, M., & Yanagihara, T. (2000). Fatigue in patients with Parkinson's disease. *Behav.Neurol.*, 12, 103-106.
- Adinoff, B., Devous, M. D., Sr., Best, S. E., Chandler, P., Alexander, D., Payne, K. et al. (2003). Gender differences in limbic responsiveness, by SPECT, following a pharmacologic challenge in healthy subjects. *Neuroimage.*, 18, 697-706.
- Alexander, G. E., DeLong, M. R., & Strick, P. L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu.Rev.Neurosci.*, 9, 357-381.
- Altgassen, M., Zolig, J., Kopp, U., Mackinlay, R., & Kliegel, M. (2007). Patients with Parkinson's disease can successfully remember to execute delayed intentions. *J.Int.Neuropsychol.Soc.*, 13, 888-892.
- Alves, G., Wentzel-Larsen, T., & Larsen, J. P. (2004). Is fatigue an independent and persistent symptom in patients with Parkinson disease? *Neurology*, 63, 1908-1911.
- American Psychiatric Association (2000). *Diagnostic and statistical manual of mental disorders, fourth edition, text-revised (DSM-IV-TR)*. Washington: American Psychiatric Association.
- Anderson, J. R. (1993). *Rules of the mind*. Hillsdale, New Jersey.
- Baudic, S., Tzortzis, C., Barba, G. D., & Traykov, L. (2004). Executive deficits in elderly patients with major unipolar depression. *J.Geriatr.Psychiatry Neurol.*, 17, 195-201.
- Beauchamp, M. H., Dagher, A., Aston, J. A., & Doyon, J. (2003). Dynamic functional changes associated with cognitive skill learning of an adapted version of the Tower of London task. *Neuroimage.*, 20, 1649-1660.
- Bergman, H. & Deuschl, G. (2002). Pathophysiology of Parkinson's disease: from clinical neurology to basic neuroscience and back. *Mov Disord.*, 17 Suppl 3, S28-S40.
- Bialystok, E., Craik, F. I., & Stefurak, T. (2008). Planning and task management in Parkinson's disease: differential emphasis in dual-task performance. *J.Int.Neuropsychol.Soc.*, 14, 257-265.

References

- Bloem, B. R., Grimbergen, Y. A., van Dijk, J. G., & Munneke, M. (2006). The "posture second" strategy: a review of wrong priorities in Parkinson's disease. *J.Neurol.Sci.*, *248*, 196-204.
- Bohnen, N. I., Kaufer, D. I., Hendrickson, R., Constantine, G. M., Mathis, C. A., & Moore, R. Y. (2007). Cortical cholinergic denervation is associated with depressive symptoms in Parkinson's disease and parkinsonian dementia. *J.Neurol.Neurosurg.Psychiatry*, *78*, 641-643.
- Boller, F., Marcie, P., Starkstein, S., & Traykov, L. (1998). Memory and depression in Parkinson's disease. *Eur.J.Neurol.*, *5*, 291-295.
- Brett, M., Anton, J. L., Valbregue, R., & Poline, J. B. (2002). Region of interest analysis using an SPM toolbox. *Neuroimage*, *6*.
- Broussolle, E., Dentresangle, C., Landais, P., Garcia-Larrea, L., Pollak, P., Croisile, B. et al. (1999). The relation of putamen and caudate nucleus 18F-Dopa uptake to motor and cognitive performances in Parkinson's disease. *J.Neurol.Sci.*, *166*, 141-151.
- Brouwer, W. H. & Schmidt, I. W. (2002). Executieve functies: de touwtjes in handen. *Leidraad psychogeriatric*, *C3*, 24-59.
- Brown, R. G. & Marsden, C. D. (1991). Dual task performance and processing resources in normal subjects and patients with Parkinson's disease. *Brain*, *114 (Pt 1A)*, 215-231.
- Brown, R. G. & Pluck, G. (2000). Negative symptoms: the 'pathology' of motivation and goal-directed behaviour. *Trends Neurosci.*, *23*, 412-417.
- Bruck, A., Aalto, S., Nurmi, E., Bergman, J., & Rinne, J. O. (2005). Cortical 6-[18F]fluoro-L-dopa uptake and frontal cognitive functions in early Parkinson's disease. *Neurobiol.Aging*, *26*, 891-898.
- Bruck, A., Portin, R., Lindell, A., Laihin, A., Bergman, J., Haaparanta, M. et al. (2001). Positron emission tomography shows that impaired frontal lobe functioning in Parkinson's disease is related to dopaminergic hypofunction in the caudate nucleus. *Neurosci.Lett.*, *311*, 81-84.
- Burgess, P. W., Alderman, N., Evans, J., Emslie, H., & Wilson, B. A. (1998). The ecological validity of tests of executive function. *J.Int.Neuropsychol.Soc.*, *4*, 547-558.
- Cloninger, C. R. (1987). A systematic method for clinical description and classification of personality variants. A proposal. *Arch.Gen.Psychiatry*, *44*, 573-588.

- Cloninger, C. R., Svrakic, D. M., & Przybeck, T. R. (1993). A psychobiological model of temperament and character. *Arch.Gen.Psychiatry*, *50*, 975-990.
- Coleman, P. G. (1999). Identity management in later life. In R.T.Woods (Ed.), *Psychological problems of aging* (pp. 49-72). West Sussex: John Wiley & Sons Ltd.
- Colman, K. S. F., Koerts, J., Van Beilen, M., Leenders, K. L., & Bastiaanse, Y. R. M. (2008). The impact of executive functions on verb production in patients with Parkinson's disease. *Cortex*, *accepted for publication*.
- Cools, R. (2006). Dopaminergic modulation of cognitive function-implications for L-DOPA treatment in Parkinson's disease. *Neurosci.Biobehav.Rev.*, *30*, 1-23.
- Cools, R. (2008). Role of dopamine in the motivational and cognitive control of behavior. *Neuroscientist.*, *14*, 381-395.
- Cools, R., Barker, R. A., Sahakian, B. J., & Robbins, T. W. (2001). Mechanisms of cognitive set flexibility in Parkinson's disease. *Brain*, *124*, 2503-2512.
- Cools, R., Barker, R. A., Sahakian, B. J., & Robbins, T. W. (2003). L-Dopa medication remediates cognitive inflexibility, but increases impulsivity in patients with Parkinson's disease. *Neuropsychologia*, *41*, 1431-1441.
- Costa, A., Peppe, A., Carlesimo, G. A., Pasqualetti, P., & Caltagirone, C. (2006). Major and minor depression in Parkinson's disease: a neuropsychological investigation. *Eur.J.Neurol.*, *13*, 972-980.
- Cubo, E., Bernard, B., Leurgans, S., & Raman, R. (2000). Cognitive and motor function in patients with Parkinson's disease with and without depression. *Clin.Neuropharmacol.*, *23*, 331-334.
- Cummings, J. L. & Masterman, D. L. (1999). Depression in patients with Parkinson's disease. *Int.J.Geriatr.Psychiatry*, *14*, 711-718.
- Cunnington, R., Egan, G. F., O'Sullivan, J. D., Hughes, A. J., Bradshaw, J. L., & Colebatch, J. G. (2001). Motor imagery in Parkinson's disease: a PET study. *Mov Disord.*, *16*, 849-857.
- Dagher, A., Owen, A. M., Boecker, H., & Brooks, D. J. (2001). The role of the striatum and hippocampus in planning: a PET activation study in Parkinson's disease. *Brain*, *124*, 1020-1032.
- Dauer, W. & Przedborski, S. (2003). Parkinson's disease: mechanisms and models. *Neuron*, *39*, 889-909.

- de la Fuente-Fernandez, R., Kishore, A., Calne, D. B., Ruth, T. J., & Stoessl, A. J. (2000). Nigrostriatal dopamine system and motor lateralization. *Behav.Brain Res.*, *112*, 63-68.
- de la Fuente-Fernandez, R., Ruth, T. J., Sossi, V., Schulzer, M., Calne, D. B., & Stoessl, A. J. (2001). Expectation and dopamine release: mechanism of the placebo effect in Parkinson's disease. *Science*, *293*, 1164-1166.
- de la Fuente-Fernandez, R., Schulzer, M., & Stoessl, A. J. (2004). Placebo mechanisms and reward circuitry: clues from Parkinson's disease. *Biol.Psychiatry*, *56*, 67-71.
- de Lau, L. M., Giesbergen, P. C., de Rijk, M. C., Hofman, A., Koudstaal, P. J., & Breteler, M. M. (2004). Incidence of parkinsonism and Parkinson disease in a general population: the Rotterdam Study. *Neurology*, *63*, 1240-1244.
- de Rijk, M. C., Tzourio, C., Breteler, M. M., Dartigues, J. F., Amaducci, L., Lopez-Pousa, S. et al. (1997). Prevalence of parkinsonism and Parkinson's disease in Europe: the EUROPARKINSON Collaborative Study. European Community Concerted Action on the Epidemiology of Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, *62*, 10-15.
- Debaere, F., Wenderoth, N., Sunaert, S., Van Hecke P., & Swinnen, S. P. (2004). Changes in brain activation during the acquisition of a new bimanual coordination task. *Neuropsychologia*, *42*, 855-867.
- Dhawan, V., Ma, Y., Pillai, V., Spetsieris, P., Chaly, T., Belakhlef, A. et al. (2002). Comparative analysis of striatal FDOPA uptake in Parkinson's disease: ratio method versus graphical approach. *J.Nucl.Med.*, *43*, 1324-1330.
- Draaisma, D. (2006). *Ontregelde geesten*. Groningen: Historische Uitgeverij.
- Dubois, B. & Pillon, B. (1997). Cognitive deficits in Parkinson's disease. *J.Neurol.*, *244*, 2-8.
- Dujardin, K., Blairy, S., Defebvre, L., Duhem, S., Noel, Y., Hess, U. et al. (2004a). Deficits in decoding emotional facial expressions in Parkinson's disease. *Neuropsychologia*, *42*, 239-250.
- Dujardin, K., Blairy, S., Defebvre, L., Krystkowiak, P., Hess, U., Blond, S. et al. (2004b). Subthalamic nucleus stimulation induces deficits in decoding emotional facial expressions in Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, *75*, 202-208.
- Edwards, E., Kitt, C., Oliver, E., Finkelstein, J., Wagster, M., & McDonald, W. M. (2002). Depression and Parkinson's disease: a new look at an old problem. *Depress.Anxiety.*, *16*, 39-48.

- Ehmann, T. S., Beninger, R. J., Gawel, M. J., & Riopelle, R. J. (1990). Depressive symptoms in Parkinson's disease: a comparison with disabled control subjects. *J.Geriatri.Psychiatry Neurol.*, *3*, 3-9.
- Ehrt, U., Bronnick, K., Leentjens, A. F., Larsen, J. P., & Aarsland, D. (2006). Depressive symptom profile in Parkinson's disease: a comparison with depression in elderly patients without Parkinson's disease. *Int.J.Geriatri.Psychiatry*, *21*, 252-258.
- Esselink, R. A., de Bie, R. M., de Haan, R. J., & et al. (2004). Unilateral pallidotomy versus bilateral subthalamic nucleus stimulation in PD: a randomized trial. *Neurology*, *62*, 201-207.
- Fahn, S., Elton, R. L., & UPDRS program members (1987). Unified Parkinsons Disease Rating Scale. In S.Fahn, C. D. Marsden, M. Goldstein, & D. B. Calne (Eds.), *Recent developments in Parkinsons disease* (vol 2 ed., pp. 153-163). Florham Park, NJ: Macmillan Healthcare Information.
- Fischer, C., Schweizer, T. A., Atkins, J. H., Bozanovic, R., Norris, M., Herrmann, N. et al. (2008). Neurocognitive profiles in older adults with and without major depression. *Int.J.Geriatri.Psychiatry*.
- Flowers, K. A. & Robertson, C. (1985). The effect of Parkinson's disease on the ability to maintain a mental set. *Journal of Neurology, Neurosurgery & Psychiatry*, *48*, 517-529.
- Foltynie, T., Brayne, C. E., Robbins, T. W., & Barker, R. A. (2004). The cognitive ability of an incident cohort of Parkinson's patients in the UK. The CamPaIGN study. *Brain*, *127*, 550-560.
- Forrester, J. W. (1968). *Principles of systems*. Cambridge, Massachusetts: Wright-Allen Press, Inc.
- Frank, M. J. (2005). Dynamic dopamine modulation in the basal ganglia: a neurocomputational account of cognitive deficits in medicated and nonmedicated Parkinsonism. *J.Cogn Neurosci.*, *17*, 51-72.
- Frank, M. J., Samanta, J., Moustafa, A. A., & Sherman, S. J. (2007). Hold your horses: impulsivity, deep brain stimulation, and medication in parkinsonism. *Science*, *318*, 1309-1312.
- Frank, M. J., Seeberger, L. C., & O'Reilly, R. C. (2004). By carrot or by stick: cognitive reinforcement learning in parkinsonism. *Science*, *306*, 1940-1943.
- Friedman, J. H., Brown, R. G., Comella, C., Garber, C. E., Krupp, L. B., Lou, J. S. et al. (2007). Fatigue in Parkinson's disease: a review. *Mov Disord.*, *22*, 297-308.

- Friston, K. J. (1995). Commentary and Opinion 2. Statistical Parametric Mapping - Ontology and Current Issues. *Journal of Cerebral Blood Flow and Metabolism*, *15*, 361-370.
- Garber, C. E. & Friedman, J. H. (2003). Effects of fatigue on physical activity and function in patients with Parkinson's disease. *Neurology*, *60*, 1119-1124.
- Georgiou, N., Bradshaw, J. L., Iansek, R., Phillips, J. G., Mattingley, J. B., & Bradshaw, J. A. (1994). Reduction in external cues and movement sequencing in Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, *57*, 368-370.
- Gjerstad, M. D., Wentzel-Larsen, T., Aarsland, D., & Larsen, J. P. (2007). Insomnia in Parkinson's disease: frequency and progression over time. *J.Neurol.Neurosurg.Psychiatry*, *78*, 476-479.
- Goerendt, I. K., Lawrence, A. D., & Brooks, D. J. (2004). Reward processing in health and Parkinson's disease: neural organization and reorganization. *Cereb.Cortex*, *14*, 73-80.
- Green, J., McDonald, W. M., Vitek, J. L., Evatt, M., Freeman, A., Haber, M. et al. (2002). Cognitive impairments in advanced PD without dementia. *Neurology*, *59*, 1320-1324.
- Groenewegen, H. J. & Dongen, Y. C. (2007). Role of the basal ganglia. In E.Ch.Wolters, T. van Laar, & H. W. Berendse (Eds.), *Parkinsonism and Related Disorders* (Amsterdam: VU University Press.
- Gupta, A. & Bhatia, S. (2000). Psychological functioning in patients with Parkinson's disease. *Parkinsonism.Relat Disord.*, *6*, 185-190.
- Haruno, M. & Kawato, M. (2006). Different neural correlates of reward expectation and reward expectation error in the putamen and caudate nucleus during stimulus-action-reward association learning. *J.Neurophysiol.*, *95*, 948-959.
- Helmich, R. C., de Lange, F. P., Bloem, B. R., & Toni, I. (2007). Cerebral compensation during motor imagery in Parkinson's disease. *Neuropsychologia*, *45*, 2201-2215.
- Hershey, T., Revilla, F. J., Wernle, A., Gibson, P. S., Dowling, J. L., & Perlmutter, J. S. (2004). Stimulation of STN impairs aspects of cognitive control in PD. *Neurology*, *62*, 1110-1114.
- Herzog, J., Volkmann, J., Krack, P., Kopper, F., Potter, M., Lorenz, D. et al. (2003). Two-year follow-up of subthalamic deep brain stimulation in Parkinson's disease. *Mov Disord.*, *18*, 1332-1337.

- Hilker, R., Portman, A. T., Voges, J., Staal, M. J., Burghaus, L., van Laar, T. et al. (2005). Disease progression continues in patients with advanced Parkinson's disease and effective subthalamic nucleus stimulation. *J.Neurol.Neurosurg.Psychiatry*, *76*, 1217-1221.
- Hoefl, F., Watson, C. L., Kesler, S. R., Bettinger, K. E., & Reiss, A. L. (2008). Gender differences in the mesocorticolimbic system during computer game-play. *J.Psychiatr.Res.*, *42*, 253-258.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, *6*, 65-70.
- Huber, S. J., Freidenberg, D. L., Paulson, G. W., Shuttleworth, E. C., & Christy, J. A. (1990). The pattern of depressive symptoms varies with progression of Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, *53*, 275-278.
- Hughes, T. A., Ross, H. F., Mindham, R. H., & Spokes, E. G. (2004). Mortality in Parkinson's disease and its association with dementia and depression. *Acta Neurol.Scand.*, *110*, 118-123.
- Jacobs, H., Heberlein, I., Vieregge, A., & Vieregge, P. (2001). Personality traits in young patients with Parkinson's disease. *Acta Neurol.Scand.*, *103*, 82-87.
- Jahanshahi, M., Jenkins, I. H., Brown, R. G., Marsden, C. D., Passingham, R. E., & Brooks, D. J. (1995). Self-initiated versus externally triggered movements. I. An investigation using measurement of regional cerebral blood flow with PET and movement-related potentials in normal and Parkinson's disease subjects. *Brain*, *118* (Pt 4), 913-933.
- Janvin, C., Aarsland, D., Larsen, J. P., & Hugdahl, K. (2003). Neuropsychological profile of patients with Parkinson's disease without dementia. *Dement.Geriatr.Cogn Disord.*, *15*, 126-131.
- Kaasinen, V., Nurmi, E., Bergman, J., Eskola, O., Solin, O., Sonninen, P. et al. (2001). Personality traits and brain dopaminergic function in Parkinson's disease. *Proc.Natl.Acad.Sci.U.S.A*, *98*, 13272-13277.
- Kamali, M., Oquendo, M. A., & Mann, J. J. (2001). Understanding the neurobiology of suicidal behavior. *Depress.Anxiety.*, *14*, 164-176.
- Karlsen, K., Larsen, J. P., Tandberg, E., & Jorgensen, K. (1999). Fatigue in patients with Parkinson's disease. *Mov Disord.*, *14*, 237-241.
- Kerenyi, L., Ricaurte, G. A., Schretlen, D. J., McCann, U., Varga, J., Mathews, W. B. et al. (2003). Positron emission tomography of striatal serotonin transporters in Parkinson disease. *Arch.Neurol.*, *60*, 1223-1229.

References

- Kirsch-Darrow, L., Fernandez, H. H., Marsiske, M., Okun, M. S., & Bowers, D. (2006). Dissociating apathy and depression in Parkinson disease. *Neurology*, *67*, 33-38.
- Kliegel, M., Phillips, L. H., Lemke, U., & Kopp, U. A. (2005). Planning and realisation of complex intentions in patients with Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, *76*, 1501-1505.
- Knutson, B., Westdorp, A., Kaiser, E., & Hommer, D. (2000). FMRI visualization of brain activity during a monetary incentive delay task. *Neuroimage.*, *12*, 20-27.
- Koerts, J., Leenders, K. L., Koning, M., Bouma, A., & van, B. M. (2008). The assessment of depression in Parkinson's disease. *Eur.J.Neurol.*, *15*, 487-492.
- Koerts, J., Leenders, K. L., Koning, M., Portman, A. T., & van, B. M. (2007). Striatal dopaminergic activity (FDOPA-PET) associated with cognitive items of a depression scale (MADRS) in Parkinson's disease. *Eur.J.Neurosci.*, *25*, 3132-3136.
- Krupp, L. B. & Pollina, D. A. (1996). Mechanisms and management of fatigue in progressive neurological disorders. *Curr.Opin.Neurol.*, *9*, 456-460.
- Kunig, G., Leenders, K. L., Martin-Solch, C., Missimer, J., Magyar, S., & Schultz, W. (2000). Reduced reward processing in the brains of Parkinsonian patients. *Neuroreport*, *11*, 3681-3687.
- Kuzis, G., Sabe, L., Tiberti, C., Leiguarda, R., & Starkstein, S. E. (1997). Cognitive functions in major depression and Parkinson disease. *Arch.Neurol.*, *54*, 982-986.
- Lauterbach, E. C. (2004). The neuropsychiatry of Parkinson's disease and related disorders. *Psychiatr.Clin.North Am.*, *27*, 801-825.
- Leentjens, A. F. (2004). Depression in Parkinson's disease: conceptual issues and clinical challenges. *J.Geriatr.Psychiatry Neurol.*, *17*, 120-126.
- Leentjens, A. F., Marinus, J., Van Hilten, J. J., Lousberg, R., & Verhey, F. R. (2003a). The contribution of somatic symptoms to the diagnosis of depressive disorder in Parkinson's disease: a discriminant analytic approach. *J.Neuropsychiatry Clin.Neurosci.*, *15*, 74-77.
- Leentjens, A. F., Van den, A. M., Metsemakers, J. F., Lousberg, R., & Verhey, F. R. (2003b). Higher incidence of depression preceding the onset of Parkinson's disease: a register study. *Mov Disord.*, *18*, 414-418.
- Leentjens, A. F., Verhey, F. R., Lousberg, R., Spitsbergen, H., & Wilmlink, F. W. (2000). The validity of the Hamilton and Montgomery-Asberg depression rating scales as

- screening and diagnostic tools for depression in Parkinson's disease. *Int.J.Geriatr.Psychiatry*, 15, 644-649.
- Lees, A. J. (1992). When did Ray Kennedy's Parkinson's disease begin? *Mov Disord.*, 7, 110-116.
- Lees, A. J., Blackburn, N. A., & Campbell, V. L. (1988). The nighttime problems of Parkinson's disease. *Clin.Neuropharmacol.*, 11, 512-519.
- Levy, R. & Dubois, B. (2006). Apathy and the functional anatomy of the prefrontal cortex-basal ganglia circuits. *Cereb.Cortex*, 16, 916-928.
- Lewis, S. J., Dove, A., Robbins, T. W., Barker, R. A., & Owen, A. M. (2003). Cognitive impairments in early Parkinson's disease are accompanied by reductions in activity in frontostriatal neural circuitry. *J.Neurosci.*, 23, 6351-6356.
- Lozza, C., Baron, J. C., Eidelberg, D., Mentis, M. J., Carbon, M., & Marie, R. M. (2004). Executive processes in Parkinson's disease: FDG-PET and network analysis. *Hum.Brain Mapp.*, 22, 236-245.
- Macht, M., Schwarz, R., & Ellgring, H. (2005). Patterns of psychological problems in Parkinson's disease. *Acta Neurol.Scand.*, 111, 95-101.
- Manchester, D., Priestley, N., & Jackson, H. (2004). The assessment of executive functions: coming out of the office. *Brain Inj.*, 18, 1067-1081.
- Martin-Soelch, C., Leenders, K. L., Chevalley, A. F., Missimer, J., Kunig, G., Magyar, S. et al. (2001). Reward mechanisms in the brain and their role in dependence: evidence from neurophysiological and neuroimaging studies. *Brain Res.Brain Res.Rev.*, 36, 139-149.
- Martinot, M., Bragulat, V., Artiges, E., Dolle, F., Hinnen, F., Jouvent, R. et al. (2001). Decreased presynaptic dopamine function in the left caudate of depressed patients with affective flattening and psychomotor retardation. *Am.J.Psychiatry*, 158, 314-316.
- Mayberg, H. S., Starkstein, S. E., Sadzot, B., Preziosi, T., Andrezejewski, P. L., Dannals, R. F. et al. (1990). Selective hypometabolism in the inferior frontal lobe in depressed patients with Parkinson's disease. *Ann.Neurol.*, 28, 57-64.
- McClure, S. M., York, M. K., & Montague, P. R. (2004). The neural substrates of reward processing in humans: the modern role of FMRI. *Neuroscientist.*, 10, 260-268.
- Menza, M. A., Golbe, L. I., Cody, R. A., & Forman, N. E. (1993a). Dopamine-related personality traits in Parkinson's disease. *Neurology*, 43, 505-508.

References

- Menza, M. A., Mark, M. H., Burn, D. J., & Brooks, D. J. (1995). Personality correlates of [18F]dopa striatal uptake: results of positron-emission tomography in Parkinson's disease. *J.Neuropsychiatry Clin.Neurosci.*, *7*, 176-179.
- Menza, M. A., Robertson-Hoffman, D. E., & Bonapace, A. S. (1993b). Parkinson's disease and anxiety: comorbidity with depression. *Biol.Psychiatry*, *34*, 465-470.
- Meyer, J. H., Kruger, S., Wilson, A. A., Christensen, B. K., Goulding, V. S., Schaffer, A. et al. (2001). Lower dopamine transporter binding potential in striatum during depression. *Neuroreport*, *12*, 4121-4125.
- Monchi, O., Petrides, M., Doyon, J., Postuma, R. B., Worsley, K., & Dagher, A. (2004). Neural bases of set-shifting deficits in Parkinson's disease. *J.Neurosci.*, *24*, 702-710.
- Monchi, O., Petrides, M., Petre, V., Worsley, K., & Dagher, A. (2001). Wisconsin Card Sorting revisited: distinct neural circuits participating in different stages of the task identified by event-related functional magnetic resonance imaging. *J.Neurosci.*, *21*, 7733-7741.
- Monchi, O., Petrides, M., Strafella, A. P., Worsley, K. J., & Doyon, J. (2006). Functional role of the basal ganglia in the planning and execution of actions. *Ann.Neurol.*, *59*, 257-264.
- Montgomery, S. A. & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *Br.J.Psychiatry*, *134*, 382-389.
- Muller, U., Wachter, T., Barthel, H., Reuter, M., & von Cramon, D. Y. (2000). Striatal [123I]beta-CIT SPECT and prefrontal cognitive functions in Parkinson's disease. *J.Neural Transm.*, *107*, 303-319.
- Murray, J. B. (1996). Depression in Parkinson's disease. *J.Psychol.*, *130*, 659-667.
- Muslimovic, D., Post, B., Speelman, J. D., & Schmand, B. (2005). Cognitive profile of patients with newly diagnosed Parkinson disease. *Neurology*, *65*, 1239-1245.
- Muslimovic, D., Post, B., Speelman, J. D., & Schmand, B. (2007). Motor procedural learning in Parkinson's disease. *Brain*, *130*, 2887-2897.
- Myers, C. E., Shohamy, D., Gluck, M. A., Grossman, S., Kluger, A., Ferris, S. et al. (2003). Dissociating hippocampal versus basal ganglia contributions to learning and transfer. *J.Cogn Neurosci.*, *15*, 185-193.
- Nelson, H. E. (1991). *The revised National Adult Reading Test - Test manual*. Windsor, UK.

- Norman, D. A. & Shallice, T. (1986). Attention to action: willed and automatic control of behaviour. In G.E.Schwarz & D. Shapiro (Eds.), *Consciousness and self-regulation* (New York: Plenum Press.
- Norman, S., Troster, A. I., Fields, J. A., & Brooks, R. (2002). Effects of depression and Parkinson's disease on cognitive functioning. *J.Neuropsychiatry Clin.Neurosci.*, *14*, 31-36.
- Papapetropoulos, S., Ellul, J., Argyriou, A. A., Chroni, E., & Lekka, N. P. (2006). The effect of depression on motor function and disease severity of Parkinson's disease. *Clin.Neurol.Neurosurg.*, *108*, 465-469.
- Poldrack, R. A., Prabhakaran, V., Seger, C. A., & Gabrieli, J. D. (1999). Striatal activation during acquisition of a cognitive skill. *Neuropsychology.*, *13*, 564-574.
- Pomati, S., Farina, E., Magni, E., Laiacona, M., & Mariani, C. (1996). Normative data for two neuropsychological tests sensitive to frontal dysfunction. *Ital.J.Neurol.Sci.*, *17*, 201-209.
- Portman, A. T., van Laar, T., Staal, M. J., Rutgers, A. W., Journee, H. L., & Leenders, K. L. (2006). Chronic stimulation of the subthalamic nucleus increases daily on-time without dyskinesia in advanced Parkinson's disease. *Parkinsonism.Relat Disord.*, *12*, 143-148.
- Post, B., Merkus, M. P., de Haan, R. J., & Speelman, J. D. (2007). Prognostic factors for the progression of Parkinson's disease: a systematic review. *Mov Disord.*, *22*, 1839-1851.
- Rascol, O., Brooks, D. J., Melamed, E., Oertel, W., Poewe, W., Stocchi, F. et al. (2005). Rasagiline as an adjunct to levodopa in patients with Parkinson's disease and motor fluctuations (LARGO, Lasting effect in Adjunct therapy with Rasagiline Given Once daily, study): a randomised, double-blind, parallel-group trial. *Lancet*, *365*, 947-954.
- Reitan, R. M. (1958). Validity of the Trail Making Test as an indicator of organic braindamage. *Perceptual and Motor Skills*, *8*, 271-276.
- Remy, P., Doder, M., Lees, A., Turjanski, N., & Brooks, D. (2005). Depression in Parkinson's disease: loss of dopamine and noradrenaline innervation in the limbic system. *Brain*, *128*, 1314-1322.
- Richard, I. H. (2005). Anxiety disorders in Parkinson's disease. *Adv.Neurol.*, *96*, 42-55.
- Ridderinkhof, K. R., Ullsperger, M., Crone, E. A., & Nieuwenhuis, S. (2004). The role of the medial frontal cortex in cognitive control. *Science*, *306*, 443-447.

References

- Rojo, A., Aguilar, M., Garolera, M. T., Cubo, E., Navas, I., & Quintana, S. (2003). Depression in Parkinson's disease: clinical correlates and outcome. *Parkinsonism.Relat Disord.*, *10*, 23-28.
- Rubinstein, T. C., Giladi, N., & Hausdorff, J. M. (2002). The power of cueing to circumvent dopamine deficits: a review of physical therapy treatment of gait disturbances in Parkinson's disease. *Mov Disord.*, *17*, 1148-1160.
- Saan, R. J. & Deelman, B. G. (1986). *De 15-Woorden Test A (manual)*. Groningen: Afdeling neuropsychologie UMCG (internal publication).
- Saling, L. L. & Phillips, J. G. (2007). Automatic behaviour: efficient not mindless. *Brain Res.Bull.*, *73*, 1-20.
- Schapira, A. H., Mann, V. M., Cooper, J. M., Krige, D., Jenner, P. J., & Marsden, C. D. (1992). Mitochondrial function in Parkinson's disease. The Royal Kings and Queens Parkinson's Disease Research Group. *Ann.Neurol.*, *32 Suppl*, S116-S124.
- Schmand, B. (2005). *Normen 15-Woordentest*. Nederlands Instituut van Psychologen.
- Schmand, B., Groenink, S. C., & van den, D. M. (2008). [Letterfluency: psychometric properties and Dutch normative data]. *Tijdschr.Gerontol.Geriatr.*, *39*, 64-76.
- Schmand, B., Houx, P., & De Koning, I. (2003). *Normen voor Stroop kleur-woord tests, Trail Making test, en Story Recall van de Rivermead Behavioural Memory Test*. Nederlands Instituut van Psychologen.
- Schneider, J. S. (2007). Behavioral persistence deficit in Parkinson's disease patients. *Eur.J.Neurol.*, *14*, 300-304.
- Schott, B. H., Niehaus, L., Wittmann, B. C., Schutze, H., Seidenbecher, C. I., Heinze, H. J. et al. (2007). Ageing and early-stage Parkinson's disease affect separable neural mechanisms of mesolimbic reward processing. *Brain*, *130*, 2412-2424.
- Schrag, A., Jahanshahi, M., & Quinn, N. (2000). What contributes to quality of life in patients with Parkinson's disease? *J.Neurol.Neurosurg.Psychiatry*, *69*, 308-312.
- Schroeder, U., Kuehler, A., Haslinger, B., Erhard, P., Fogel, W., Tronnier, V. M. et al. (2002). Subthalamic nucleus stimulation affects striato-anterior cingulate cortex circuit in a response conflict task: a PET study. *Brain*, *125*, 1995-2004.
- Schultz, W. (1997). Dopamine neurons and their role in reward mechanisms. *Curr.Opin.Neurobiol.*, *7*, 191-197.
- Schultz, W. (2001). Reward signaling by dopamine neurons. *Neuroscientist.*, *7*, 293-302.
- Shallice, T. (1988). *From neuropsychology to mental structure*. Cambridge: Cambridge University Press.

- Shallice, T. & Burgess, P. (1996). The domain of supervisory processes and temporal organization of behaviour. *Philos.Trans.R.Soc.Lond B Biol.Sci.*, 351, 1405-1411.
- Shallice, T. & Burgess, P. W. (1991). Deficits in strategy application following frontal lobe damage in man. *Brain*, 114 (Pt 2), 727-741.
- Shohamy, D., Myers, C. E., Gekhman, K. D., Sage, J., & Gluck, M. A. (2006). L-dopa impairs learning, but spares generalization, in Parkinson's disease. *Neuropsychologia*, 44, 774-784.
- Shohamy, D., Myers, C. E., Grossman, S., Sage, J., Gluck, M. A., & Poldrack, R. A. (2004). Cortico-striatal contributions to feedback-based learning: converging data from neuroimaging and neuropsychology. *Brain*, 127, 851-859.
- Silberman, C. D., Laks, J., Capita, C. F., Rodrigues, C. S., Moreira, I., Vasconcellos, L. F. et al. (2007). Frontal functions in depressed and nondepressed Parkinson's disease patients: impact of severity stages. *Psychiatry Res.*, 149, 285-289.
- Smeding, H. M., Speelman, J. D., Koning-Haanstra, M., Schuurman, P. R., Nijssen, P., van, L. T. et al. (2006). Neuropsychological effects of bilateral STN stimulation in Parkinson disease: a controlled study. *Neurology*, 66, 1830-1836.
- Starkstein, S. E., Bolduc, P. L., Mayberg, H. S., Preziosi, T. J., & Robinson, R. G. (1990a). Cognitive impairments and depression in Parkinson's disease: a follow up study. *J.Neurol.Neurosurg.Psychiatry*, 53, 597-602.
- Starkstein, S. E., Petracca, G., Chemerinski, E., & Merello, M. (2001). Prevalence and correlates of parkinsonism in patients with primary depression. *Neurology*, 57, 553-555.
- Starkstein, S. E., Petracca, G., Chemerinski, E., Teson, A., Sabe, L., Merello, M. et al. (1998). Depression in classic versus akinetic-rigid Parkinson's disease. *Mov Disord.*, 13, 29-33.
- Starkstein, S. E., Preziosi, T. J., Forrester, A. W., & Robinson, R. G. (1990b). Specificity of affective and autonomic symptoms of depression in Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, 53, 869-873.
- Stevens, J. P. (2002). *Applied Multivariate Statistics for the Social Sciences*. (4th ed.) New Jersey: Mahwah.
- Stinissen, J., Willems, P. J., Coetsier, P., & Hulsman, W. L. L. (1970). *Handleiding bij de Nederlandstalige bewerking van de Wechsler Adult Intelligence Scale (WAIS; manual)*. Lisse: Swets & Zeitlinger.

References

- Stocchi, F., Barbato, L., Nordera, G., Berardelli, A., & Ruggieri, S. (1998). Sleep disorders in Parkinson's disease. *J.Neurol.*, *245 Suppl 1*, S15-S18.
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, *18*, 643-662.
- Swainson, R., Rogers, R. D., Sahakian, B. J., Summers, B. A., Polkey, C. E., & Robbins, T. W. (2000). Probabilistic learning and reversal deficits in patients with Parkinson's disease or frontal or temporal lobe lesions: possible adverse effects of dopaminergic medication. *Neuropsychologia*, *38*, 596-612.
- Talairach, J. & Tournoux, P. (1988). *Co-planar stereotactic atlas of the human brain*. (1 ed.) New York: Thieme Medical Publishers.
- Tandberg, E., Larsen, J. P., Aarsland, D., Laake, K., & Cummings, J. L. (1997). Risk factors for depression in Parkinson disease. *Arch.Neurol.*, *54*, 625-630.
- Taylor, A. E., Saint-Cyr, J. A., & Lang, A. E. (1990). Subcognitive processing in the frontocaudate "complex loop": the role of the striatum. *Alzheimer Dis.Assoc.Disord.*, *4*, 150-160.
- Thut, G., Schultz, W., Roelcke, U., Nienhusmeier, M., Missimer, J., Maguire, R. P. et al. (1997). Activation of the human brain by monetary reward. *Neuroreport*, *8*, 1225-1228.
- Tomer, R. & Aharon-Peretz, J. (2004). Novelty seeking and harm avoidance in Parkinson's disease: effects of asymmetric dopamine deficiency. *J.Neurol.Neurosurg.Psychiatry*, *75*, 972-975.
- Tremblay, L. K., Naranjo, C. A., Graham, S. J., Herrmann, N., Mayberg, H. S., Hevenor, S. et al. (2005). Functional neuroanatomical substrates of altered reward processing in major depressive disorder revealed by a dopaminergic probe. *Arch.Gen.Psychiatry*, *62*, 1228-1236.
- Troster, A. I., Stalp, L. D., Paolo, A. M., Fields, J. A., & Koller, W. C. (1995). Neuropsychological impairment in Parkinson's disease with and without depression. *Arch.Neurol.*, *52*, 1164-1169.
- Uekermann, J., Daum, I., Peters, S., Wiebel, B., Przuntek, H., & Muller, T. (2003). Depressed mood and executive dysfunction in early Parkinson's disease. *Acta Neurol.Scand.*, *107*, 341-348.
- Van Beilen, M., Portman, A. T., Kiers, H. A., Maguire, R. P., Kaasinen, V., Koning, M. et al. (2008). Striatal FDOPA uptake and cognition in advanced non-demented Parkinson's disease: A clinical and FDOPA-PET study. *Parkinsonism.Relat Disord.*

- Van Beilen, M., van Zomeren, E. H., van den Bosch, R. J., Withaar, F. K., & Bouma, A. (2005). Measuring the executive functions in schizophrenia: the voluntary allocation of effort. *J.Psychiatr.Res.*, *39*, 585-593.
- Van der Elst, W., Van Boxtel, M. P., Van Breukelen, G. J., & Jolles, J. (2006). Normative data for the Animal, Profession and Letter M Naming verbal fluency tests for Dutch speaking participants and the effects of age, education, and sex. *J.Int.Neuropsychol.Soc.*, *12*, 80-89.
- Van der Graaf, F. H., de Jong, B. M., Maguire, R. P., Meiners, L. C., & Leenders, K. L. (2004). Cerebral activation related to skills practice in a double serial reaction time task: striatal involvement in random-order sequence learning. *Brain Res.Cogn Brain Res.*, *20*, 120-131.
- Van der Graaf, F. H., Maguire, R. P., Leenders, K. L., & de Jong, B. M. (2006). Cerebral activation related to implicit sequence learning in a Double Serial Reaction Time task. *Brain Res.*, *1081*, 179-190.
- Van Hilten, J. J., Weggeman, M., van der Velde, E. A., Kerkhof, G. A., van Dijk, J. G., & Roos, R. A. (1993). Sleep, excessive daytime sleepiness and fatigue in Parkinson's disease. *J.Neural Transm.Park Dis.Dement.Sect.*, *5*, 235-244.
- Verbrugge, L. M. & Jette, A. M. (1994). The disablement process. *Soc.Sci.Med.*, *38*, 1-14.
- Verleden, S., Vingerhoets, G., & Santens, P. (2007). Heterogeneity of cognitive dysfunction in Parkinson's disease: a cohort study. *Eur.Neurol.*, *58*, 34-40.
- Visser, M., Leentjens, A. F., Marinus, J., Stiggelbout, A. M., & Van Hilten, J. J. (2006). Reliability and validity of the Beck depression inventory in patients with Parkinson's disease. *Mov Disord.*, *21*, 668-672.
- Weintraub, D., Moberg, P. J., Culbertson, W. C., Duda, J. E., Katz, I. R., & Stern, M. B. (2005a). Dimensions of executive function in Parkinson's disease. *Dement.Geriatr.Cogn Disord.*, *20*, 140-144.
- Weintraub, D., Newberg, A. B., Cary, M. S., Siderowf, A. D., Moberg, P. J., Kleiner-Fisman, G. et al. (2005b). Striatal dopamine transporter imaging correlates with anxiety and depression symptoms in Parkinson's disease. *J.Nucl.Med.*, *46*, 227-232.
- Wichowicz, H. M., Slawek, J., Derejko, M., & Cubala, W. J. (2006). Factors associated with depression in Parkinson's disease: a cross-sectional study in a Polish population. *Eur.Psychiatry*.

References

- Williams-Gray, C. H., Foltynie, T., Brayne, C. E., Robbins, T. W., & Barker, R. A. (2007). Evolution of cognitive dysfunction in an incident Parkinson's disease cohort. *Brain, 130*, 1787-1798.
- Wilson, B. A., Alderman, N., Burgess, P., Emslie, H., & Evans, J. J. (1996). *Behavioural Assessment of the Dysexecutive syndrome*. Bury St. Edmunds, Suffolk: Thames Valley Test Company.
- Wu, T. & Hallett, M. (2005). A functional MRI study of automatic movements in patients with Parkinson's disease. *Brain, 128*, 2250-2259.
- Wu, T. & Hallett, M. (2007). Neural correlates of dual task performance in patients with Parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*.
- Zald, D. H., Boileau, I., El Dearedy, W., Gunn, R., McGlone, F., Dichter, G. S. et al. (2004). Dopamine transmission in the human striatum during monetary reward tasks. *J.Neurosci., 24*, 4105-4112.