

University of Groningen

Chronic kidney disease

Thio, C. H. L.

DOI:
[10.33612/diss.133648108](https://doi.org/10.33612/diss.133648108)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Thio, C. H. L. (2020). *Chronic kidney disease: Insights from social and genetic epidemiology*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.133648108>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Educational level and risk of chronic kidney disease: Longitudinal data from the PREVEND study

Chris HL Thio, Priya Vart, Lyanne M Kieneker, Harold Snieder, Ron T Gansevoort, Ute Bültmann

Nephrol Dial Transplant 2018 (in press)

CHAPTER

A large, light gray, stylized number '2' is positioned on the right side of the page, partially overlapping the 'CHAPTER' text. The number is composed of a thick, rounded top curve and a straight vertical stem that ends in a horizontal base.

ABSTRACT

Introduction. The longitudinal association between low education and chronic kidney disease (CKD) and its underlying mechanisms are poorly characterized. We therefore examined the association of low education with incident CKD and change in kidney function, and explored potential mediators of this association.

Methods. We analyzed data on 6078 participants from the community-based PREVEND Study. Educational level was categorized into low, medium, and high (<secondary, secondary/equivalent, >secondary schooling). Kidney function was assessed by estimating glomerular filtration rate (eGFR) by serum creatinine and cystatin C at five examinations during ~11 years of follow-up. Incident CKD was defined as new-onset eGFR<60mL/min/1.73m² and/or urinary albumin≥30mg/24h in those free of CKD at baseline. We estimated main effects with Cox regression and linear mixed models. In exploratory causal mediation analyses, we examined mediation by several potential risk factors.

Results. Incident CKD was observed in 861 (17%) participants. Lower education was associated with higher rates of incident CKD (low vs high education; HR[95%CI]=1.25 [1.05 to 1.48], $p_{\text{trend}}=0.009$) and accelerated eGFR decline (B[95%CI]=-0.15 [-0.21 to -0.09] mL/min/1.73m² per year, $p_{\text{trend}}<0.001$). The association between education and incident CKD was mediated by smoking, potassium excretion, BMI, WHR, and hypertension. Analysis on annual eGFR change in addition suggested mediation by magnesium excretion, protein intake, and diabetes.

Conclusions. In the general population, we observed an inverse association of educational level with CKD. Diabetes, and the modifiable risk factors smoking, poor diet, BMI, WHR, and hypertension are suggested to underlie this association. These findings provide support for targeted preventive policies to reduce socioeconomic disparities in kidney disease.

Keywords: chronic kidney disease, educational level, socioeconomic status, health disparities

ABBREVIATIONS

BMI = body-mass index

CKD = chronic kidney disease

eGFR = estimated glomerular filtration rate

PREVEND = Study Prevention of renal and vascular end-stage disease study

SES = socioeconomic status

UAE = urinary albumin excretion

WHR = waist-to-hip ratio

Supplementary material available at www.doi.org/10.1093/ndt/gfy361

BACKGROUND

Chronic kidney disease (CKD) is a heterogeneous group of disorders characterized by sustained diminished kidney function and/or kidney damage. CKD affects ~10-15% of the global population, and its incidence is increasing¹⁻⁴. CKD can progress to end-stage renal disease (ESRD), and is associated with an increased incidence of cardiovascular disease and all-cause mortality^{5,6}. As such, CKD poses a major burden on patients and global health resources.

CKD is unequally distributed across socioeconomic groups: higher prevalence and incidence rates of CKD and ESRD have consistently been observed among those with lower socioeconomic status (SES). Socioeconomic gradients have also been observed for eGFR and urinary albumin. However, large heterogeneity exists between studies of the SES-CKD association^{7,8}. One possible explanation for this heterogeneity is that factors underlying the SES-CKD association vary between populations due to differences in e.g. ethnicity, lifestyle, prevalence of comorbid conditions, or healthcare^{9,10}. Currently, the available literature is limited: 1) most observations were made in US-based cross-sectional data^{7,8} and 2) European studies established cross-sectional associations of SES measures with CKD¹¹⁻¹³; however no European study explicitly examined the association of SES with CKD, or mediators of this association, in a longitudinal setting. Hence, it is uncertain to what extent SES conveys risk of CKD in the European general population, and which factors underlie this association. Characterization of underlying mechanisms may help identify targets for disease prevention and management, thus help alleviate the burden of CKD and its consequences among disadvantaged populations. Our aim was therefore to examine the strength of the association of SES with the longitudinal outcomes, CKD incidence and annual change in eGFR, in a sample of the Dutch general population. Furthermore, we explored health-related behaviors and comorbid conditions that potentially mediate this association.

MATERIALS AND METHODS

Study design and population

We used data from the Prevention of REnal and Vascular ENdstage Disease (PREVEND) cohort study. PREVEND was initiated to investigate the natural course of increased urinary albumin levels and its association to renal and vascular outcomes. Details of this study have been described elsewhere¹⁴. Briefly, 8592 individuals, sampled from the general population of Groningen,

the Netherlands, underwent extensive examination between 1997-1998. Four follow-up examinations were completed in 2003, 2006, 2008, and 2012. All subjects gave written informed consent. PREVEND was approved by the medical ethics committee of the University Medical Center Groningen and conducted in accordance with the Helsinki Declaration guidelines. In the present study, we excluded participants with incomplete data on educational level, kidney outcomes, or important covariates.

Measures

We defined CKD according to Kidney Disease: Improving Global Outcomes guidelines (eGFR < 60 mL/min/1.73 m² or UAE ≥ 30 mg/24h)¹⁵. Incident cases were those participants free of CKD at baseline who developed CKD during follow-up. We calculated eGFR from serum creatinine and serum cystatin C, using the corresponding CKD-EPI equation¹⁶.

Collection procedures of blood and two consecutive 24h-urine specimens at each examination has been described previously¹⁷. Measurement of serum creatinine was performed by an enzymatic method on a Roche Modular analyzer using reagents and calibrators from Roche (Roche Diagnostics, Mannheim, Germany), with intra- and interassay coefficients of variation of 0.9% and 2.9%, respectively. Serum cystatin C concentration was measured by a Gentian cystatin C Immunoassay (Gentian AS Moss, Norway) on a Modular analyzer (Roche Diagnostics). Cystatin C was calibrated directly using the standard supplied by the manufacturer (traceable to the International Federation of Clinical Chemistry Working Group for Standardization of Serum Cystatin C)¹⁸. Intra- and interassay coefficients of variation were < 4.1% and < 3.3%, respectively. Urinary albumin concentration (UAC) was measured by nephelometry with a lower threshold of detection of 2.3 mg/L, and intra- and interassay coefficient of variation of 2.2% and 2.6%, respectively (Dade Behring Diagnostic, Marburg, Germany). UAC was multiplied by urine volume to obtain a value of UAE in mg/24h. The two 24h-urinary albumin values of each subject per examination were averaged.

SES was measured by educational level, categorized into low (no, primary, basic vocational, and secondary education), medium (senior secondary vocational and general senior secondary education), and high (higher professional and higher academic education) according to the International Standard Classification of

Education¹⁹. Furthermore, we examined associations of income as alternative measure of SES. For this, we categorized income into low, medium, and high according to tertiles of the ratio between reported income and the 1998 poverty line (1658 guilders per month).

Age, sex, and baseline eGFR were included as potential confounders. Included as potential mediators were: current smoking (self-reported yes/no), alcohol consumption (labelled as none, occasional {<10g/wk}, light (10-69.9g/wk), moderate (70-210g/wk), heavier (>210g/wk)}, 24h urinary excretions of sodium (Na⁺), potassium (K⁺), and magnesium (Mg²⁺) (as surrogates for dietary intake of sodium, potassium, and magnesium), 24h protein intake (estimated from 24h urea excretion by the Maroni formula^{20,21}), body-mass index (BMI, weight/height²), waist-to-hip ratio (WHR, waist/hip circumference), diabetes (fasting glucose>7.0mmol/L, non-fasting glucose>11.0mmol/L, anti-diabetic treatment, or self-reported), hypertension (systolic blood pressure>140mmHg, diastolic blood pressure>90mmHg, blood pressure lowering treatment, or self-reported), hypercholesterolemia (total cholesterol≥6.21mmol/L, lipid lowering treatment, or self-reported). Covariates were collected at baseline by questionnaires, anthropometry, urine collections, or pharmacy records. Urinary concentrations of Na⁺, K⁺, and Mg²⁺ were determined as previously described^{17,22}.


Statistical analyses

Statistical analyses were performed using R v3.4.1²³ and SPSS v23 software (IBM corp, Amonk, NY, USA) during years 2017 and 2018. Two-sided significance level was set at =0.05 unless otherwise stated. Baseline characteristics were examined for the total population and compared across categories of education using one-way ANOVA, Jonckheere-Terpstra, or χ^2 -tests for linear trend. We used the *survival* R-package²⁴ for Cox proportional hazards modelling of time to CKD. Time of CKD was estimated using a midpoint imputation method. Crude effects were examined in an unadjusted model. Next, we adjusted for age, sex, and baseline eGFR. In a final model, we introduced potential mediators. We calculated p for linear trend by analyzing education as a continuous rather than an ordinal variable. Using the *lme4* R-package²⁵, we estimated eGFR change by modelling eGFR as a function of time in a random intercept, random coefficient linear mixed model. To examine the crude effect of SES on annual eGFR change, an interaction term between time and SES was introduced. Next, we adjusted for age and sex, as well as their

interaction with time. Finally, we introduced all potential mediators, and the interaction of each with time.

Next, we performed exploratory mediation analyses. **Figure 1** shows a graph of hypothesized pathways tested in the present study. Main effects of potential mediators on kidney outcomes were examined with Cox proportional hazards and linear mixed models adjusting for age, sex, and baseline eGFR. Next, we used the *mediation* R-package²⁶ to estimate mediation within the counterfactual framework described by Imai et al²⁷. Here, we simplified our statistical models by using one contrast for education (low vs high education). Furthermore, we used individual eGFR slopes (extracted from a linear mixed-effects model) as outcome variable in mediation analysis of eGFR change. Finally, we used parametric survival models implemented in the *survival* R-package. Due to these alternative methods, effects may deviate slightly from those of our main effects analyses. Each potential mediator was analyzed separately, adjusting for age, sex, and baseline eGFR. Any significant SES x mediator interaction was controlled for in the mediation

Figure 1. Graph of tested pathways through which low education could potentially lead to chronic kidney disease. BMI, body-mass index; WHR, waist-to-hip ratio. Poorer diet: high in sodium, low in potassium, low in magnesium, high in protein. Black arrows indicate a posited causal pathway; grey arrows indicate potential confounding pathways



model. Non-parametric bootstrap CIs and p-values were estimated from 1000 simulations. One-sided hypotheses were tested to assess potential mediators (i.e. current smoking²⁸, higher alcohol consumption, higher Na⁺ excretion²⁹, lower K⁺ excretion¹⁷, lower Mg²⁺ excretion³⁰, higher estimated protein intake³¹, higher BMI³², higher WHR³³, diabetes, hypertension, and hypercholesterolemia).

In secondary analyses, we examined associations of education with incident CKD_{eGFR} (eGFR<60 mL/min/1.73m²), incident CKD_{UAE} (UAE≥30mg/24h) and annual change in UAE (natural-log transformed to approximate normality, *ln*UAE).

RESULTS

Baseline characteristics by educational level for 6078 participants with complete baseline data are presented in **Table 1**. Traditional risk factors (i.e. diabetes, hypertension, high cholesterol, smoking, higher BMI) were more prevalent in participants with low education. At baseline, low education participants were more likely to have CKD, lower eGFR, and higher UAE compared to high education participants. A higher attrition rate was observed for participants with low education: follow-up time was shorter for these participants. Low education was univariably associated with lower dietary quality as indicated by higher Na⁺ excretion, lower K⁺ excretion, lower Mg²⁺ excretion, and higher protein intake. Low education participants reported less alcohol consumption.

After excluding N=883 participants with baseline CKD, N=5195 remained for time-to-CKD analysis. Among these, 861 (17%) experienced new-onset CKD, with a significant socioeconomic gradient (low; med; high education: 22%; 14%; 12%, ²[df] =62.8[1], $p_{\text{trend}} < 0.001$). In the crude model, we observed an inverse association of education with CKD, again with a significant gradient (low vs high education: HR [95%CI] =1.97 [1.67 to 2.32], $p_{\text{trend}} < 0.001$; **Table 2**). After adjusting for age, sex, and baseline eGFR, the association was attenuated, but significance remained (low vs high education: HR [95%CI] =1.25 [1.05 to 1.48], $p_{\text{trend}} = 0.009$). After introducing all potential mediators to the model, the education-CKD association was no longer significant, suggesting mediation within our hypothesized framework (**Figure 1**).

Average estimated annual eGFR change for the total N=6078 population was -0.93 (95%CI: -0.95 to -0.91) mL/min/1.73m² per year. Low education was associated with accelerated eGFR change, with a significant gradient (low vs high education: B

Table 1. Baseline characteristics by categories of educational level.

	Total	Educational level			P _{trend}
		Low (< secondary)	Medium (secondary or equivalent)	High (> secondary)	
N	6078	2637	1565	1876	
Males	3071 (51%)	1223 (46%)	855 (55%)	993 (53%)	0.005
Age, years	48 [39-59]	54 [45-63]	44 [36-54]	43 [37-51]	<0.001
BMI, kg/m²	26 (4.1)	27 (4.3)	26 (4.0)	25 (3.3)	<0.001
WHR	0.88 (0.09)	0.90 (0.09)	0.87 (0.09)	0.86 (0.09)	<0.001
Current smoking	1956 (32%)	951 (36%)	529 (34%)	476 (25%)	<0.001
Alcohol					
None	1438 (24%)	881 (33%)	346 (22%)	211 (11%)	<0.001
Occasional (<10 g/wk)	956 (16%)	436 (17%)	263 (17%)	257 (14%)	
Light (10-69.9 g/wk)	2120 (35%)	782 (30%)	573 (37%)	765 (41%)	
Moderate (70-210 g/wk)	1252 (21%)	404 (15%)	303 (19%)	545 (29%)	
Heavier (>210 g/wk)	312 (5%)	134 (5%)	80 (5%)	98 (5%)	
Na⁺ excretion (mmol/24h)	143 (51)	143 (52)	145 (51)	140 (48)	0.021
K⁺ excretion (mmol/24h)	72 (21)	69 (20)	73 (22)	76 (21)	<0.001
Mg²⁺ excretion (mmol/24h)	3.9 (1.5)	3.8 (1.5)	4.0 (1.6)	4.1 (1.5)	<0.001
Estimated protein intake (g/kg/24h)	1.16 (0.26)	1.18 (0.28)	1.15 (0.26)	1.16 (0.24)	0.005
Diabetes	202 (3%)	130 (5%)	43 (3%)	29 (2%)	<0.001
Hypertension	1912 (31%)	1112 (42%)	418 (27%)	382 (20%)	<0.001
High cholesterol	1879 (31%)	1062 (40%)	423 (27%)	4394 (21%)	<0.001
Creatinine (μmol/L)	72 (16)	72 (17)	72 (15)	73 (14)	0.015
Cystatin C (mg/L)	0.89 (0.17)	0.91 (0.19)	0.88 (0.16)	0.86 (0.14)	<0.001
eGFR, ml/min/1.73m²	95 (17)	91 (17)	98 (16)	99 (15)	<0.001
UAE, mg/24h	9.1 [6.3-16]	10 [6.3-20]	8.9 [6.2-15]	8.4 [6.2-13]	<0.001
CKD at baseline	883 (15%)	510 (19%)	199 (13%)	174 (9%)	<0.001
CKD_{eGFR} at baseline	167 (3%)	109 (4%)	37 (2%)	21 (1%)	<0.001
CKD_{UAE} at baseline	805 (13%)	457 (17%)	181 (12%)	167 (9%)	<0.001
Follow-up time, yrs	11.2 [8.5-12.1]	11.1 [7.0-11.8]	11.3 [9.3-12.2]	11.4 [10.6-12.4]	<0.001

Baseline characteristics by categories of educational level. Data is presented as mean (standard deviation), median (interquartile range), and number (%) where appropriate. P-values reflect significance of a linear trend across categories of educational level, using one-way ANOVA, χ^2 , or Jonckheere-Terpstra tests where appropriate.

Abbreviations: BMI, body-mass index; WHR, waist-to-hip ratio; eGFR, estimated glomerular filtration rate; UAE, urinary albumin excretion; CKD, chronic kidney disease

[95%CI] = -0.15 [-0.21 to -0.09], $p_{\text{trend}} < 0.001$, adjusted for age and sex; **Table 2**). Addition of potential mediators to the model attenuated the association, although significance remained (low vs high education: B [95%CI] = -0.11 [-0.16 to -0.04], $p_{\text{trend}} < 0.001$).

All potential mediators were associated with either CKD or annual eGFR change (Supplementary Table S4). We tested interactions of education with each potential mediator separately (age, sex, and baseline eGFR adjusted); none were significant

Table 2. Association of education with incident CKD (Panel A) and annual change in eGFR (Panel B).				
A) Incident CKD				
	Educational level			
	Low (<secondary)	Medium (secondary/equivalent)	High (>secondary)	
N=5195	N=2127	N=1366	N=1702	P_{trend}
Events N=861	460 (22%)	193 (14%)	208 (12%)	<0.001
	HR (95%CI)			
Model 1	1.97 (1.67 to 2.32)	1.17 (0.96 to 1.42)	(ref.)	<0.001
Model 2 ^a	1.25 (1.05 to 1.48)	1.07 (0.88 to 1.30)	(ref.)	0.009
Model 3	1.02 (0.85 to 1.22)	0.97 (0.80 to 1.19)	(ref.)	0.789
B) Annual eGFR change				
	Educational level			
	Low (<secondary)	Medium (secondary/equivalent)	High (>secondary)	
N=6078	N=2637	N=1565	N=1876	P_{trend}
	B (95%CI)			
Model 1 ^b	-0.30 (-0.36 to -0.24)	-0.10 (-0.17 to -0.04)	(ref.)	<0.001
Model 2 ^b	-0.15 (-0.21 to -0.09)	-0.08 (-0.14 to -0.02)	(ref.)	<0.001
Model 3 ^b	-0.11 (-0.16 to -0.04)	-0.06 (-0.12 to 0.00)	(ref.)	<0.001
Data are presented as hazard ratio (95%CI) or unstandardized regression coefficient (95%CI, in mL/min/1.73m ² per year). Abbreviations: CKD, chronic kidney disease; eGFR, estimated glomerular filtration rate. Model 1: Crude Educational level (high educational level is reference category) Model 2: Model 1 + age, sex, ^a (and in addition baseline eGFR), ^b (and in addition their interaction with time) Model 3: Model 2 + potential mediators (body-mass index, waist-to-hip ratio, smoking, alcohol use, Na ⁺ excretion, K ⁺ excretion, Mg ²⁺ excretion, estimated protein intake, diabetes, hypertension, high cholesterol) ^b (and in addition their interaction with time)				

($p > 0.05$). The association of low education and CKD was mediated by higher likelihood of smoking (proportion mediated [95%CI] = 0.14 [0.02 to 0.51], $p = 0.009$; **Table 3**), lower 24h K⁺-excretion (0.12 [0.02 to 0.45], $p = 0.008$), higher BMI (0.29 [0.13 to 0.96], $p = 0.004$), higher WHR (0.31 [0.14 to 1.16], $p = 0.002$), and higher prevalence of hypertension (0.14 [0.05 to 0.47], $p = 0.006$) in this subpopulation. We observed no significant mediation by 24h Na⁺ excretion, Mg²⁺ excretion, protein intake, diabetes, or hypercholesterolemia.

There were no education x mediator interactions with eGFR change as outcome except for smoking (low education [vs high education] x smoking: B [95%CI] = -0.07 [-0.10 to -0.05], $p = 0.01$). The association of low education and accelerated eGFR decline was mediated by lower 24h K⁺ excretion (**Table 3**, proportion mediated [95%CI] = 0.08 [0.03 to 0.16], one-sided $p < 0.001$), higher BMI (0.22 [0.12 to 0.42], $p < 0.001$), higher WHR (0.09 [0.01 to 0.18], $p = 0.008$), and higher prevalence of

hypertension (0.13 [0.08 to 0.24], $p < 0.001$). Additionally, lower Mg^{2+} excretion (0.03 [-0.003 to 0.07], $p = 0.030$), higher protein intake (0.01 [-0.001 to 0.04], $p = 0.032$) and higher prevalence of diabetes (0.04 [0.01 to 0.10], $p = 0.009$) mediated the association of low education with accelerated eGFR decline. A protective effect of smoking on eGFR change was observed; higher prevalence of smoking in those with low education appeared to offset risk of accelerated eGFR decline (proportion mediated [95%CI] = -0.12 [-0.22 to -0.06], $p = 1.000$). Higher alcohol consumption was not a mediating risk factor, rather, alcohol seemed protective of CKD and accelerated eGFR decline (Supplementary Table S4). Estimates of average causal mediation effects and direct effects are listed in Supplementary Tables S5-6.

No significant associations between education with CKD_{eGFR} or CKD_{UAE} were found, although directions of effect for these outcomes were consistent with our main analysis (Supplementary Table S1-S2). Average estimated increase in UAE for

Mediators	Incident CKD		Annual change in eGFR	
	Proportion mediated (95%CI)	p	Proportion mediated (95%CI)	p
Health-related behaviors				
<i>Smoking</i>	0.14 (0.02 to 0.51)	0.009	-0.12 (-0.23 to -0.05) ^o	1.000
<i>Alcohol</i>	0.24 (0.05 to 0.99)	0.989	0.26 (0.16 to 0.49)	1.000
<i>24h Na⁺ excretion</i>	-0.01 (-0.09 to 0.09)	0.431	0.01 (-0.02 to 0.06)	0.216
<i>24h K⁺ excretion</i>	0.12 (0.02 to 0.45)	0.008	0.08 (0.03 to 0.16)	<0.001
<i>24h Mg²⁺ excretion</i>	0.04 (-0.03 to 0.18)	0.098	0.03 (-0.003 to 0.07)	0.030
<i>Estimated 24h protein intake</i>	0.001 (-0.002 to 0.04)	0.447	0.01 (-0.001 to 0.04)	0.032
Comorbid conditions				
<i>BMI</i>	0.29 (0.13 to 0.96)	0.004	0.22 (0.12 to 0.42)	<0.001
<i>WHR</i>	0.31 (0.14 to 1.16)	0.002	0.09 (0.01 to 0.18)	0.008
<i>Diabetes</i>	0.08 (-0.005 to 0.06)	0.064	0.04 (0.01 to 0.10)	0.009
<i>Hypertension</i>	0.14 (0.05 to 0.47)	0.006	0.13 (0.08 to 0.24)	<0.001
<i>Hypercholesterolemia</i>	0.02 (-0.05 to 0.13)	0.223	-0.04 (-0.10 to 0.01)	0.962
Results from causal mediation analysis. N=6078. Effects are reported as proportion mediated of the association between education (low vs high) and kidney outcomes. Non-parametric bootstrap confidence intervals and one-sided p-values are estimated from 1000 simulations. Estimates are conditioned on age, sex, and baseline eGFR.				
One-sided hypotheses were that low education leads to steeper eGFR decline through: current smoking, higher alcohol consumption, higher Na ⁺ excretion, lower K ⁺ excretion, lower Mg ²⁺ excretion, higher protein intake, higher BMI, higher WHR, diabetes, hypertension, and hypercholesterolemia.				
^o In addition adjusted for the interaction term educational level x smoking.				

the total population was 1.1% (95%CI: 0.9% to 1.3%) per year. Low education was associated with accelerated increase in UAE (low vs high education: 0.7% [0.2% to 0.11%] accelerated increase in UAE per year, $p_{\text{trend}}=0.003$), but no longer significantly after adjusting for age and sex (Supplementary Table S3). There were no significant associations of household income, as alternative measure of socioeconomic status, with kidney outcomes after confounder adjustment (data not shown).

DISCUSSION

In a middle-aged community-based cohort, we examined the associations of SES, as indicated by educational level, with the longitudinal kidney outcomes, incident CKD and eGFR decline. Low education was associated with higher incidence rates of CKD, independent of age, sex, and baseline eGFR, but not of potential mediators. Furthermore, low education was associated with accelerated eGFR decline, independent of age, sex, and potential mediators. Exploratory longitudinal mediation analysis suggested that the association between education and CKD can partly be explained by diabetes and the modifiable risk factors, BMI, WHR, smoking, potassium, and hypertension. No significant associations of household income with kidney outcomes were observed.

With this longitudinal study, we corroborate previous cross-sectional observations that in the Netherlands, education, not income, is associated with kidney outcomes¹². Recent longitudinal data from the US-based Atherosclerosis Risk in Communities study show effects of education on CKD incidence and eGFR decline comparable to the present data³⁴. However, in contrast to the Netherlands, income is associated to CKD in the US^{12,34}. Possible explanations for this discrepancy are: 1) in the US, healthcare access is income-dependent³⁵, and 2) there is larger income inequality compared to the Netherlands³⁶.

Our results are generally consistent with a previous mediation analysis on the SES-CKD association. This study assessed SES by household income, and was performed in a cross-sectional sample of the general US population³⁷. Similar to that study, we observed mediation by smoking, (abdominal) obesity, diabetes, and hypertension. However, we could not corroborate a mediation effect of hypercholesterolemia. Vart et al³⁷ used questionnaires on availability of fruits and vegetables at home to assess dietary quality but did not observe mediation. In contrast, we used urinary measures to objectively assess dietary intake of various nutrients. We found strong

mediation effects of lower potassium intake on both incident CKD and accelerated eGFR decline, as well as suggestions for effects of lower magnesium intake and higher protein intake.

Of all nutrients examined in the present study, lower potassium intake was the strongest mediator. A large body of epidemiological data shows that low SES is associated with poor diet, especially with a lower consumption of micronutrients such as potassium³⁸. Low potassium intake was previously observed to associate with an increased risk of incident hypertension³⁹, but also with incident CKD independent of hypertension¹⁷. A proposed mechanism involves induction of tubulointerstitial injury by ammoniogenesis caused by potassium deficiency^{40,41}. Furthermore, potassium itself might be renoprotective by upregulating renal kinins⁴². On the other hand, potassium intake might reflect dietary quality more generally. The main dietary sources of potassium are fruits/vegetables, legumes, whole grains, and dairy products⁴³. These potassium-rich foods contain fibers, polyphenols, antioxidants, and vitamins, which have health benefits⁴⁴ that may be renoprotective.

Interestingly, no mediation through sodium intake was observed. High sodium intake reflects poor diet due to its high content in processed foods^{45,46}, and is associated with the major renal risk factor, hypertension⁴⁷; we therefore expected sodium to mediate the relation between education and CKD. However, we did not observe a strong educational gradient in sodium at baseline (**Table 1**). Furthermore, sodium intake was not found to be associated with CKD in PREVEND¹⁷, which likely explains the observed lack of mediation in the present study.

Three counterintuitive findings need to be addressed. Firstly, despite its association with an elevated risk of CKD (concordant with literature²⁸), smoking was associated with decelerated eGFR decline. We therefore further examined the main effect of smoking on eGFR decline in fully adjusted models: compared to non-smokers, smokers had lower baseline eGFR, and despite decelerated decline, eGFR on average remained lower in these participants (data not shown). Therefore, this finding is likely the result of a floor effect. Secondly, alcohol consumption was inversely associated with risk of CKD and eGFR decline. Moreover, lower alcohol consumption among low education participants partly explained the elevated risk of CKD. This may be due to residual confounding, a sick quitter/sick non-starter effect⁴⁸, or a cohort-specific effect; for a detailed discussion we refer to a study

by Koning et al. that previously observed this association in PREVEND⁴⁹. Thirdly, mediation effects of diabetes were significant on eGFR decline, but only borderline significant ($p=0.064$) on incident CKD. This is likely the result of reduced statistical power of a dichotomous outcome compared to a continuous outcome, and low prevalence of diabetes (3% at baseline) in the PREVEND sample.

The mechanisms underlying the education-CKD association are incompletely understood. We therefore tested several biologically plausible mediating pathways. However, some are overlapping (e.g. BMI, WHR), or on the same causal pathway (e.g. low potassium intake leading to CKD possibly through hypertension). Hence, we examined each mediator separately, correcting only for age, sex, and baseline eGFR to prevent overadjustment. Due to sparse adjustment and the observational nature of PREVEND, we cannot exclude residual confounding. However, results were broadly concordant with the literature, i.e. effects were generally in the hypothesized direction. Therefore, any confounding has likely only biased magnitude, not direction, of mediation effects. Future work may involve further characterization of the education-CKD association by estimating effects of multiple mediators relative to one another using multivariable techniques (e.g. structural equation modelling or the counterfactual approach described by Lange et al^{50,51}).

To the best of our knowledge, the present study is the first in Europe examining the longitudinal association between education and CKD in the general population, and the first exploring its underlying mechanisms in a longitudinal setting. Strengths of this study are its considerable size ($N=6078$) and follow-up time (~11 years). GFR was estimated from serial measurements of serum creatinine and cystatin C, currently considered the best proxy of kidney function in population-based studies. Furthermore, data on urinary albumin was available for all included participants. Finally, dietary variables were objectively measured from 24h urinary collections. Several limitations should be addressed. Firstly, PREVEND consists of >95% whites; we therefore could not address the influence of ethnicity in the education-CKD association. Secondly, we observed a higher attrition rate of participants with low education, which may have resulted in a bias towards the null. Thirdly, we lacked baseline information on several potential mediators (e.g. physical activity/sedentary time, healthcare access, health literacy, psychological factors). Finally, only individual-level socioeconomic data were available; we therefore could not examine effects of area-level SES.

In an effort to characterize socioeconomic disparities in CKD, we explored a number

of plausible mediating pathways (i.e. health behaviors and clinical risk factors) that link education to CKD. Future research may focus on e.g. 1) confirming the pathways suggested in the present study; 2) exploring other potential mediating factors such as health care access, health literacy, and psychological factors; 3) establishing the interrelationship between these factors. Understanding how and why socioeconomically disadvantaged groups (e.g. those with a lower educational level) show higher vulnerability to CKD may prove helpful in designing interventions to reduce socioeconomic disparities in CKD. Given the challenges of intervening on education itself, managing and/or modifying downstream effects of low education may be a more promising approach.

To conclude: in the Dutch general population, low SES, as indicated by educational level, is associated with elevated risk of CKD. This association is suggested to be driven by higher rates of diabetes and the modifiable risk factors, (abdominal) obesity, smoking, low potassium intake, and hypertension, in those with lower education. The data presented are a first step towards potential targeted public health interventions to reduce socioeconomic health disparities.

ACKNOWLEDGEMENTS

The PREVEND Study in general was funded by the Dutch Kidney Foundation (grant E.033). The funding source had no role in study design; in collection, analysis, or interpretation of the data; in writing of the report; or in the decision to submit for publication. Results have been previously presented as an abstract at the ERA-EDTA conference in Copenhagen, Denmark, May 2018. CT and PV conceived and designed the study. RG contributed to data acquisition. All authors contributed to either analysis or interpretation of the data. CT and LK drafted the manuscript. PV, LK, HS, RG, UB revised the article. HS, RG, UB supervised the work. All authors approved of the final version of the manuscript.

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

1. Grams ME, Chow EKH, Segev DL, Coresh J. Lifetime incidence of CKD stages 3-5 in the united states. *Am J Kidney Dis.* 2013;62(2):245-252.
2. Zhang QL, Rothenbacher D. Prevalence of chronic kidney disease in population-based studies: Systematic review. *BMC Public Health.* 2008;8:117-2458-8-117.
3. Jha V, Garcia-Garcia G, Iseki K, et al. Chronic kidney disease: Global dimension and perspectives. *The Lancet.* 2013;382(9888):260-272.
4. El Nahas AM, Bello AK. Chronic kidney disease: The global challenge. *The Lancet.* 2005;365(9456):331-340.
5. Gansevoort RT, Correa-Rotter R, Hemmelgarn BR, et al. Chronic kidney disease and cardiovascular risk: Epidemiology, mechanisms, and prevention. *The Lancet.* 2013;382(9889):339-352.
6. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med.* 2004;351(13):1296-1305.
7. Vart P, Gansevoort RT, Joosten MM, Bultmann U, Reijneveld SA. Socioeconomic disparities in chronic kidney disease: A systematic review and meta-analysis. *Am J Prev Med.* 2015;48(5):580-592.
8. Zeng X, Liu J, Tao S, Hong HG, Li Y, Fu P. Associations between socioeconomic status and chronic kidney disease: A meta-analysis. *J Epidemiol Community Health.* 2018;72(4):270-279.
9. Bruck K, Stel VS, Gambaro G, et al. CKD prevalence varies across the European general population. *J Am Soc Nephrol.* 2016;27(7):2135-2147.
10. Patzer RE, McClellan WM. Influence of race, ethnicity and socioeconomic status on kidney disease. *Nature Reviews Nephrology.* 2012;8(9):533.
11. Bello AK, Peters J, Rigby J, Rahman AA, El Nahas M. Socioeconomic status and chronic kidney disease at presentation to a renal service in the United Kingdom. *Clin J Am Soc Nephrol.* 2008;3(5):1316-1323.
12. Vart P, Gansevoort RT, Coresh J, Reijneveld SA, Bultmann U. Socioeconomic measures and CKD in the United States and the Netherlands. *Clin J Am Soc Nephrol.* 2013;8(10):1685-1693.
13. Chudek J, Wieczorowska-Tobis K, Zejda J, et al. The prevalence of chronic kidney disease and its relation to socioeconomic conditions in an elderly Polish population: Results from the national population-based study PolSenior. *Nephrol Dial Transplant.* 2013;29(5):1073-1082.
14. Pinto-Sietsma SJ, Janssen WM, Hillege HL, Navis G, De Zeeuw D, De Jong PE. Urinary albumin excretion is associated with renal functional abnormalities in a nondiabetic population. *J Am Soc Nephrol.* 2000;11(10):1882-1888.
15. Levey AS, de Jong PE, Coresh J, et al. The definition, classification, and prognosis of chronic kidney disease: A KDIGO controversies conference report. *Kidney Int.* 2011;80(1):17-28.
16. Inker LA, Schmid CH, Tighiouart H, et al. Estimating glomerular filtration rate from serum creatinine and cystatin C. *N Engl J Med.* 2012;367(1):20-29.
17. Kieneker LM, Bakker SJ, de Boer RA, Navis GJ, Gansevoort RT, Joosten MM. Low potassium excretion but not high sodium excretion is associated with increased risk of developing chronic kidney disease. *Kidney Int.* 2016;90(4):888-896.
18. Grubb A, Bliurup-Jensen S, Lindstrom V, et al. First certified reference material for cystatin C in human serum ERM-DA471/IFCC. *Clin Chem Lab Med.* 2010;48(11):1619-1621.
19. UNESCO Institute for Statistics. International standard classification of education. . 2011. Maroni BJ, Steinman TI, Mitch WE. A method for estimating nitrogen intake of patients with chronic renal failure. *Kidney Int.* 1985;27(1):58-65.
20. Halbesma N, Bakker SJ, Jansen DF, et al. High protein intake associates with cardiovascular events but not with loss of renal function. *J Am Soc Nephrol.* 2009;20(8):1797-1804.
21. Joosten MM, Gansevoort RT, Mukamal KJ, et al. Urinary magnesium excretion and risk of hypertension: The prevention of renal and vascular end-stage disease study. *Hypertension.* 2013;61(6):1161-1167.

22. R Core Team. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. www.R-project.org/. 2014.
23. Therneau T. A package for survival analysis in S. R package version 2.38. Retrieved from www.CRAN.R-project.org/package=survival. 2015.
24. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*. 2014.
25. Tingley D, Yamamoto T, Hirose K, Keele L, Imai K. Mediation: R package for causal mediation analysis. 2014.
26. Imai K, Keele L, Tingley D. A general approach to causal mediation analysis. *Psychol Methods*. 2010;15(4):309.
27. Xia J, Wang L, Ma Z, et al. Cigarette smoking and chronic kidney disease in the general population: A systematic review and meta-analysis of prospective cohort studies. *Nephrology Dialysis Transplantation*. 2017;32(3):475-487.
28. Smyth A, O'donnell MJ, Yusuf S, et al. Sodium intake and renal outcomes: A systematic review. *Am J Hypertension*. 2014;27(10):1277-1284.
29. Rebholz CM, Tin A, Liu Y, et al. Dietary magnesium and kidney function decline: The healthy aging in neighborhoods of diversity across the life span study. *Am J Nephrol*. 2016;44(5):381-387.
30. Malhotra R, Lipworth L, Cavanaugh KL, et al. Protein intake and long-term change in glomerular filtration rate in the jackson heart study. *J Renal Nutrition*. 2018.
31. Herrington WG, Smith M, Bankhead C, et al. Body-mass index and risk of advanced chronic kidney disease: Prospective analyses from a primary care cohort of 1.4 million adults in England. *PloS one*. 2017;12(3):e0173515.
32. Elsayed EF, Sarnak MJ, Tighiouart H, et al. Waist-to-hip ratio, body mass index, and subsequent kidney disease and death. *Am J Kidney Dis*. 2008;52(1):29-38.
33. Vart P, Grams ME, Ballew SH, Woodward M, Coresh J, Matsushita K. Socioeconomic status and risk of kidney dysfunction: The Atherosclerosis Risk in Communities study. *Nephrol Dial Transplant*. 2018.
34. Agrawal V, Jaar BG, Frisby XY, et al. Access to health care among adults evaluated for CKD: Findings from the kidney early evaluation program (KEEP). *Am J Kidney Dis*. 2012;59(3):S5-S15.
35. Zheng H. Do people die from income inequality of a decade ago? *Soc Sci Med*. 2012;75(1):36-45.
36. Vart P, Gansevoort RT, Crews DC, Reijneveld SA, Bultmann U. Mediators of the association between low socioeconomic status and chronic kidney disease in the United States. *Am J Epidemiol*. 2015;181(6):385-396.
37. Darmon N, Drewnowski A. Does social class predict diet quality?. *Am J Clin Nutr*. 2008;87(5):1107-1117.
38. Kieneker LM, Gansevoort RT, Mukamal KJ, et al. Urinary potassium excretion and risk of developing hypertension: The prevention of renal and vascular end-stage disease study. *Hypertension*. 2014;64(4):769-776.
39. Nath KA, Hostetter MK, Hostetter TH. Increased ammoniogenesis as a determinant of progressive renal injury. *Am J Kidney Dis*. 1991;17(6):654-657.
40. Tolins JP, Hostetter MK, Hostetter TH. Hypokalemic nephropathy in the rat. role of ammonia in chronic tubular injury. *J Clin Invest*. 1987;79(5):1447-1458.
41. Ardiles L, Cardenas A, Burgos ME, et al. Antihypertensive and renoprotective effect of the kinin pathway activated by potassium in a model of salt sensitivity following overload proteinuria. *Am J Physiol-Renal Physiol*. 2013;304(12):F1399-F1410.
42. Hendriksen M, van Rossum C. Kalium inname: Risico van hyperkaliëmie?: Overzicht van beschikbare gegevens in Nederland. 2015.
43. Slavin JL, Lloyd B. Health benefits of fruits and vegetables. *Adv Nutr*. 2012;3(4):506-516.

44. Anderson CA, Appel LJ, Okuda N, et al. Dietary sources of sodium in China, Japan, the United Kingdom, and the United States, women and men aged 40 to 59 years: The INTERMAP study. *J Am Diet Assoc.* 2010;110(5):736-745.
45. Van Rossum C, Fransen H, Verkaik-Kloosterman J, Buurma-Rethans E, Ocké M. Dutch national food consumption survey 2007-2010: Diet of children and adults aged 7 to 69 years. 2011.
46. Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower sodium intake on health: Systematic review and meta-analyses. *BMJ.* 2013;346:f1326.
47. Marmot M, Brunner E. Alcohol and cardiovascular disease: The status of the U shaped curve. *BMJ.* 1991;303(6802):565-568.
48. Koning SH, Gansevoort RT, Mukamal KJ, et al. Alcohol consumption is inversely associated with the risk of developing chronic kidney disease. *Kidney Int.* 2015.
49. Lange T, Hansen JV. Direct and indirect effects in a survival context. *Epidemiology.* 2011;22(4):575-581.
50. Lange T, Rasmussen M, Thygesen LC. Assessing natural direct and indirect effects through multiple pathways. *Am J Epidemiol.* 2013;179(4):513-518.

