

University of Groningen

DNAJ proteins: more than just “co-chaperones”

Kakkar, Vaishali

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kakkar, V. (2014). *DNAJ proteins: more than just “co-chaperones”: Implicaties voor eiwit-aggregatie ziektes.* [S.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the “Taverne” license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

DNAJ proteins: more than just “co-chaperones”

Implications for protein aggregation diseases

Vaishali Kakkar
2014

The research described in this thesis was conducted at the Department of Cell Biology (Division: Radiation & Stress Cell Biology), University Medical Center Groningen, University of Groningen, The Netherlands.

The research was supported by grants from Senter Novem (IOP-IGE07004), a discovery initiative grants sponsored by the High Q foundation (now CHDI) and a grant by the Nederlandse Hersenstichting (grant nr. 15F07(2).58).

Printing of this thesis was financially supported by University of Groningen and Graduate School for Drug Exploration (GUIDE) at University Medical Center Groningen.

Cover Idea/Concept: Vaishali Kakkar

Cover Design: Randy Wind; contact: wind.randy@gmail.com

Thesis Layout: Midas Mentink; contact: www.midasmentink.nl

Thesis printed by Ipskamp Drukkers, Enschede



ISBN (print): 978-90-367-7310-2

ISBN (electronic): 978-90-367-7311-9

Copyright © 2014 by Vaishali Kakkar. All rights reserved. No part of this thesis may be reproduced, stored in a retrieval system, or transmitted in any form or by any means without permission of the author.



university of
 groningen

DNAJ proteins: more than just “co-chaperones”

Implications for protein aggregation diseases

PhD thesis

to obtain the degree of PhD at the
 University of Groningen
 on the authority of the
 Rector Magnificus Prof. E. Sterken
 and in accordance with
 the decision by the College of Deans.

This thesis will be defended in public on

Monday 13 October 2014 at 11.00 hours

by

Vaishali Kakkar

born on 22 May 1985
 in Hisar, India

Supervisor

Prof. dr. H.H. Kampinga

Assessment committee

Prof. dr. C. Emanuelsson

Prof. dr. G.J. van Ommen

Prof. dr. ir. E.A.A. Nollen

Paranymphs

Melanie Meister-Broekema

Peter W. K. Nagle

Martti Maimets

TABLE OF CONTENTS

	PROLOGUE	9
Chapter 1	Aim and Outline of the thesis	11
Chapter 2	Introduction- Barcoding heat shock proteins to human diseases: looking beyond the heat shock response	15
Chapter 3	DNAJB6, a peptide chaperone, is the most potent suppressor of aggregation of polyglutamine-containing proteins within the DNAJ family	49
Chapter 4	DNAJB6 is unable to prevent seeding/initiation of aggregation mediated by extracellular polyglutamine peptide fragments	81
Chapter 5	DNAJB6 ameliorates Huntington's disease phenotype in R6/2 mouse model	99
Chapter 6	DNAJ proteins and protein aggregation diseases – a review	121
Chapter 7	DNAJ proteins act as rate limiting step in C209G parkin aggregation	143
Chapter 8	General Discussion and future perspectives	165
	APPENDIX:	179
	Nederlandse samenvatting	180
	Acknowledgements	184
	List of Publications	188
	EPILOGUE	189

PROLOGUE

More than 50 years ago, Ferruccio Ritossa, while studying nucleic acid synthesis in *Drosophila melanogaster* salivary glands, observed a different “puffing” pattern in the chromosomes of the flies when the incubators were accidentally shifted to elevated temperature. Following up this serendipity, he showed that cells can activate a specific transcriptional program when exposed to elevated temperatures, later referred to as “heat shock response”. A decade later, Alfred Tissieres discovered the Heat Shock Proteins (HSPs) that were the main products induced by this transcriptional program. The understanding that heat unfolds proteins and they next can form toxic aggregates which could be counteracted by this transient induction of HSPs (via their chaperone actions) provides the basis for the concept of cellular protein homeostasis. All cells are continuously subjected to changes in intra and extra-cellular conditions that impede on protein folding. Further, given the macromolecular crowding within the cells it runs the risk to cause protein aggregation. Protein aggregates hallmark almost all the major neurodegenerative diseases known today, suggesting that an imbalance in the maintenance of protein homeostasis underlies these diseases. This thesis aim at providing an insight into the role of one subset of heat shock protein families, i.e. the class of DNAJ (Hsp40) proteins and the role they play both as cause for and potential therapeutic target in age-related neurodegenerative diseases, with a particular focus on Huntington’s Disease.

