

University of Groningen

Hydrogen sulfide

Snijder, Pauline

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Snijder, P. (2014). *Hydrogen sulfide: Protective properties in models of disease*. [Thesis fully internal (DIV), University of Groningen]. [S.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER

INTRODUCTION

1

Historically, hydrogen sulfide (H_2S) has been perceived as a highly toxic gas with the distinctive smell of rotten eggs (Table 1). Recently, this pungent gas was discovered to be synthesized enzymatically in mammalian and human tissues,^{1,2} which positively transformed the apparent image of H_2S into a promising therapeutic compound. In a relatively small period of time, H_2S gained considerable interest and the amount of research on this subject has immensely expanded. Over these past years, many (patho-) physiological functions of H_2S have been discovered.² Like nitric oxide (NO) and carbon monoxide (CO), H_2S fulfills all of the criteria to be considered a gasotransmitter. Its functions range from modulation of oxidative stress, vasodilatation as an endothelial derived hyperpolarizing factor to signaling through posttranslational protein modification.³⁻⁶ Beside this wide range of uncovered functions, there are still many effects of H_2S that have not been elucidated yet. However, it is clear that H_2S is an important mediator in cellular physiology.

ENDOGENOUS PRODUCTION AND FUNCTION

H_2S is generated in mammalian cells via both enzymatic and nonenzymatic pathways, although the nonenzymatic pathway only accounts for a small portion of H_2S production. Among the enzymes involved in the production of H_2S , cystathionine β -synthase (CBS) and cystathionine γ -lyase (CSE) have been investigated most extensively. Both enzymes produce H_2S by using pyridoxal 5'-phosphate (vitamin B6) as a cofactor. CBS converts homocysteine into cystathionine, which is then converted to L-cysteine by CSE. Both enzymes transform L-cysteine to H_2S (Figure 1).^{8,9} H_2S is also synthesized from D-cysteine by tandem enzymes cysteine aminotransferase (CAT) and 3-mercaptopyruvate sulfurtransferase (3-MST) (Figure 1).¹⁰ H_2S is then serially oxidized to persulfide, thiosulfate, sulfite and sulfate (Figure 1). CBS and CSE are

Table 1 – Overview of the known toxic effects of H_2S

| Dose (parts per million) | Effect |
|--------------------------|--|
| 0.02 ppm - 0.03 ppm | Detectable odor |
| 5 ppm - 10 ppm | Offensive, unpleasant odor |
| 20 ppm | Maximum allowable concentration for daily 8 hour exposure |
| 50 ppm | Irritation of the ocular conjunctiva |
| 100 ppm | Loss of smell / olfactory nerve paralysis |
| 100 ppm - 200 ppm | Upper respiratory tract irritation |
| 300 ppm - 500 ppm | Pulmonary edema |
| 500 ppm | Headache, dizziness, unconsciousness after 30-60 minutes of exposure |
| 500 ppm - 700 ppm | Unconsciousness, respiratory paralysis eventually leading to death |
| 1000 ppm | Rapid collapse, death within minutes |

Adapted from Beauchamp et al.⁷

present in a variety of tissues such as blood vessels, heart, liver, kidneys and the brain, whereas MPST expression has been mainly demonstrated in the vascular endothelium. It has been widely suggested that the expression of CSE, CBS, 3-MST (and CAT) in rodents and humans shows a marked degree of tissue specificity. However, as more researchers investigate the emerging role of H₂S in their particular disease model, this simple though convenient distinction is no longer as clear as once thought.

The majority of evidence for a physiological role of H₂S has been obtained from studies in vascular tissue. These studies convincingly show that H₂S is a vasodilatory intermediate. Genetic knockout of CSE in mice leads to an age-dependent increase in blood pressure. This hypertensive phenotype is likely due to the lack of endogenous H₂S, since injection of H₂S rescues these mice from an increasing blood pressure.⁵ While homozygous CBS knockout mice are lethal during the embryonic phase, heterozygous variants similarly develop an elevated blood pressure.¹¹ Mechanisms thought to be responsible for the vasodilatory effects of H₂S include the activation of ATP-sensitive potassium channels (K_{ATP} channels) in vascular smooth muscle cells through direct cysteine-S-sulfhydration.⁴ Furthermore, it is thought that the vasodilatory effects of H₂S are partially NO dependent. Inhibition of NO using L-NG-nitroargininemethyl ester (L-NAME) decreased the potency of H₂S in aortic rings,¹² these findings suggest crosstalk between gasotransmitters is involved in the regulation of blood pressure. Last, it has been shown that H₂S is able to prevent renovascular hypertension by affecting the renin-angiotensin-aldosterone system through direct inhibition of renin and angiotensin-converting enzyme activity.¹³

Another important feature of H₂S is its ability to reduce and modulate oxidative stress by directly scavenging reactive oxygen species (ROS), increasing the formation of the intracellular antioxidant glutathione and activating the anti-oxidative transcription factor Nuclear factor (erythroid-derived 2)-like 2 (Nrf-2).¹⁴⁻¹⁶ An additional strategy in reducing the amount of oxidative stress is through modulation of mitochondrial ROS production⁶. It has become apparent that H₂S is able to protect the integrity of mitochondria and supports mitochondrial function, thereby contributing to the preservation of cellular energetics.^{17,18} H₂S can be used as an inorganic energy substrate for mitochondria.¹⁹ Under stress, such as increased intracellular calcium concentrations, H₂S producing enzymes can translocate to the mitochondria.^{18,20,21} This translocation might increase mitochondrial H₂S levels to be used as an electron donor in the electron transport chain.

Furthermore, H₂S seems to regulate inflammatory responses. In literature, both pro- and anti-inflammatory effects of H₂S have been reported. These discrepancies may reflect the varying effects of dose–response relationships. At low, physiological concentrations, H₂S is predominantly anti-inflammatory, whereas at high concentrations H₂S promotes inflammation, a pattern that is also observed for NO and CO.^{1,22} The mechanisms by which H₂S exerts anti-inflammatory effects are the inhibition of pro-inflammatory transcription factors like nuclear factor kappa B and the inhibition of pro-inflammatory enzyme activity iNOS, cyclooxygenase-2 and tumor necrosis factor- α converting enzyme.²³

Also, H_2S is essential for vascular endothelial growth factor (VEGF) mediated angiogenesis. VEGF-stimulated in vitro sprouting is reduced in blood vessels after CSE silencing, whereas it is enhanced in vessels overexpressing CSE. The finding that genetic knockout of CSE in mice causes impaired wound healing, further indicates that endogenous H_2S production by CSE is a dynamic regulator of angiogenesis.^{24,25} Exogenous treatment with H_2S improves wound healing in rats, and attenuates heart failure by inducing angiogenesis via an increase in VEGF and eNOS/NO.^{24,26}

The capacity of H_2S to exert intracellular signals by sulfhydrating target proteins, a form of posttranslational modification, is particularly striking.²⁷ During this process a SH-group is added to a reactive cysteine residue and thereby the activity of the target protein can be altered. Sulfhydration by H_2S seems to be more widespread than nitrosylation, and the prevalence and ratio of sulfhydration is perhaps comparable to that of phosphorylation. This is a field still in its infancy, but it seems that it can become a meaningful aspect of intracellular signaling pathways. Table 2 gives an overview of the proposed mechanisms of action of H_2S .

EXOGENOUS ADMINISTRATION

The application of H_2S in biomedical experiments is challenging with regard to its chemical properties. The vast majority of studies that have examined the potential role of H_2S in health and disease have invariably utilized commercially available sulfide salts such as sodium sulfide (Na_2S) and sodium hydrosulfide ($NaSH$). Over the course of the years, it has become clear that these sodium salts have disadvantages, such as the rapid peak in H_2S levels and short half-life when dissolved or injected. This makes it hard to monitor actual H_2S levels and achieve controlled, stable and therapeutic levels of this gas in vitro and in vivo. Interestingly, H_2S donors with a regulated release have been developed recently.²⁸ With these slow-release H_2S donor molecules, such as GYY4137, it is possible to expose animals, tissue and cells to H_2S generated in a more physiological manner.²⁹

Another interesting approach is the use of thiosulfate (TS), an intermediate of sulfur metabolism from cysteine and a metabolite of H_2S that can also produce H_2S through the action of thiosulfate reductase.^{30,31} Since TS seems to have similar protective properties as other H_2S donors, this provides us with exciting possibilities for the translation into clinical use. While short-term treatment with STS is well tolerated in humans for the treatment of calciphylaxis,³² the long-term side effects should be further explored.

Gaseous administration of H_2S is less frequently used than treatment with sulfide salts or other H_2S donor compounds, but has a very interesting accessory feature. In high, sub-toxic concentrations (80-100 ppm) gaseous H_2S can induce a reversible hibernation-like state in mice. During treatment, oxygen consumption and carbon dioxide production can be reduced by 90-95% within minutes. Also, the core body temperature decreases to around 2°C above ambient temperature. After cessation of H_2S exposure, this massive reduction in oxidative metabolism is reversible without




Figure 1 – Overview of endogenous H_2S production. Abbreviations: CSE – cystathionine γ -lyase; CBS – cystathionine – β -synthase (CBS); CAT – cysteine aminotransferase; DAO – D-amino acid oxidase; 3-MST – 3-mercaptopyruvate sulfurtransferase; TSR – thiosulfate reductase

apparent toxic effects.^{33,34} This brilliant discovery led to the finding that H_2S -induced hypometabolism is protective during lethal hypoxia possibly by buffering oxygen consumption, altering the redox environment within cells, and preventing a lethal imbalance between energy supply and demand.³⁵ The hypothesized mechanism behind H_2S -induced hypometabolism is the reversible inhibition of cytochrome c oxidase (COX), the terminal enzyme in the mitochondrial electron transport chain.³⁶ This capacity of H_2S makes it an extremely interesting protective agent in ischemia-reperfusion related damage where oxygen demand exceeds oxygen availability.

ISCHEMIA-REPERFUSION INJURY

Ischemia-reperfusion injury (IRI) is one of the major damaging processes responsible for a worsened outcome after organ transplantation. IRI is implicated in other processes than organ transplantation, such as myocardial infarction or surgical procedures like cardiopulmonary bypass. One of the strategies to reduce IRI-related damage is cooling, thereby reducing metabolism and enzymatic activity. However, cooling does

Table 2 – Proposed mechanisms for protective effects of H₂S

| |
|---|
| Increase in GSH levels |
| Reduction of mitochondrial superoxide production |
| Activation of ROS-scavengers (e.g. MnSOD, CuZnSOD) |
| Preservation of mitochondrial integrity |
| Acting as an electron donor in the electron transport chain |
| Modulation of mitochondrial activity / Hypometabolism |
| Induction of hypothermia |
| Modulation of vascular proliferation |
| Modulation of leukocyte adhesion / rolling |
| Modulation of cytokine production |
| Activation of KATP-channels |
| Activation of calcium activated K ⁺ channels |
| Modulation of protein activity through sulphydration |
| Modulation of platelet function |
| Modulation of hypoxia inducible factor (HIF) |
| Modulation of Nrf2 signaling |
| Increase in eNOS / iNOS / HO-1 |
| Increase in HSP90 |
| Modulation of COX-2 activity |
| Modulation of GSK3beta activity |
| Modulation of apoptotic signaling |
| Modulation of autophagy |

have detrimental effects as well. Using H₂S to reduce oxidative metabolism might be a promising strategy to circumvent these detrimental effects. In addition, H₂S is also considered protective during other processes critically involved in IRI, such as oxidation, inflammation and apoptosis. Changes in renal CSE and CBS expression and/or activity during ischemia and in the early phase of reperfusion suggest a role for H₂S production in this process.^{6,37} Furthermore, CBS and CSE both are involved in the response to oxidative stress. CSE deficient mice are more susceptible to ischemic damage,⁶ and overexpression of CSE in cardiac tissue protects from myocardial infarction.³⁸ The protective effect of H₂S administration against IRI has been shown in various organs, including heart,³⁹ kidney,³³ liver⁴⁰ and brain.⁴¹ In these experimental studies NaSH is often used as a H₂S-donating compound. So far, the exogenous administration, endogenous manipulation and use of genetically modified animals have been successful in demonstrating H₂S-mediated cytoprotection in models of IRI.

HYPERTENSION

H₂S is an important mediator in the vascular system, contributing to vascular regulation and protection against vascular injury that leads to vascular dysfunction. The vasodilatory effects of H₂S were first reported when administration of H₂S donors to anaesthetized rats was found to induce a transient hypotensive effect.⁴² Endogenous H₂S shortage is suggested to be involved in the pathogenesis of hypertension and a significant down regulation of the H₂S/CSE system has been reported in several hypertensive models. Spontaneously hypertensive rats (SHR) have been widely used as a model for human essential hypertension because of the similarity of the main cardiovascular characteristics. The CSE – L-cysteine pathway is downregulated in SHR and treating them with a H₂S donor is protective, reducing blood pressure and vascular remodeling.⁴³ The most compelling evidence for the importance of H₂S in blood pressure regulation is that mice deficient in CSE develop endothelial dysfunction and hypertension within 8 weeks of birth and that H₂S replacement decreases systolic blood pressure in both CSE^{-/-} and CSE^{+/-} mice.⁵ H₂S is also reported to have preventive and therapeutic effects on renovascular hypertension by inhibiting plasma renin activity.¹³ Because H₂S is a major endothelium derived hyperpolarizing factor and a primary determinant of vasorelaxation in numerous vascular beds, drugs that alter CSE activity or H₂S-mediated channel sulfhydration may be effective therapeutic agents for treating hypertension.

AGING

Aging has been a topic of great interest to the scientific community for a long time. This biological process is characterized by time-dependent progressive decline of physiological functions accompanied by increased incidence of age-associated diseases. One of the proposed theories to explain the process of aging is that aging and age-related diseases are caused by the deleterious consequences of free radicals exposure on cell constituents and connective tissues.⁴⁴ Substantial evidence shows that H₂S is involved in aging by inhibiting free-radical reactions, activating SIRT1, and probably interacting with the age-related gene *Klotho*. One study reported that the plasma H₂S level in humans over 50 to 80 years of age declines with age.⁴⁵ In *Drosophila*, CBS is crucial for the increased lifespan linked to dietary restriction. When exposed to H₂S, nematodes are apparently healthy and do not exhibit phenotypes consistent with metabolic inhibition. Instead, animals exposed to H₂S are long-lived, and this phenotype requires Sir2 activity, which may translate environmental changes into physiological alterations to improve survival.⁴⁶ Moreover, H₂S has been shown to have therapeutic potential in age-associated diseases, such as hypertension and several neurodegenerative diseases. Decreased levels of H₂S in brain tissue are associated with neurodegenerative age-related diseases like Parkinson's and administration of H₂S has been shown protective in experimental models for this disease.^{47,48} The exact mechanisms of action of the role of H₂S in the pathology of

aging have not yet been sufficiently characterized and await elucidation by further studies. A better understanding of the roles of H₂S in aging can provide insights into potential therapeutic interventions against aging and reduce age-associated diseases.

AIM OF THE THESIS

The aim of this thesis was to investigate the therapeutic potential of H₂S and its producing enzymes in the setting of hypertensive renal and heart disease, ischemia and neurodegeneration. Our work underlines the broad implications of H₂S by showing protective effects in several distinct disease models in various organisms.

Since H₂S has been established as the third gasotransmitter alongside NO and CO, many overlapping actions as well as differences have been reported. Crosstalk between the three gasotransmitters has been shown, although the evidence is still scarce. Recently, it has become clear that the vasorelaxant effects of H₂S are partly NO-dependent. In **Chapter 2** an outline of the mechanisms of gas-mediated cytoprotection and complex interactions between gasotransmitters in renal IRI and transplantation is presented. Protective effects of sulfide salts (NaSH and Na₂S) in IRI have been investigated in several organs. Gaseous administration of H₂S has been used only a few times in renal and hepatic IRI. In **Chapter 3** we investigated whether gaseous administration of H₂S is protective in a model for transient myocardial infarction in mice and whether non-hypometabolic concentrations of H₂S have similar protective properties as hypometabolic concentrations. Although the exogenous protective effects of H₂S have been investigated in several IRI models, the role of endogenous H₂S-producing enzymes in IRI has been less researched. **Chapter 4** therefore investigates the antioxidant role of endogenous H₂S production by CSE in ischemic settings in vitro, in vivo and in human renal transplant tissue, using models of CSE deficiency and overexpression. The convincing protective effects of H₂S in IRI and the fact that there are many overlapping underlying damaging processes between IRI and hypertension, led us to investigate its alleged beneficial effects in hypertensive disease. The translation of H₂S to the clinical setting unfortunately is still out of our reach. For that purpose we applied sodium thiosulfate (STS), a novel H₂S-related endogenous substance with clinical potential. STS is an intermediate of sulfur metabolism from cysteine and a metabolite of H₂S that can also release sulfide. In **Chapter 5** and **6** we studied the protective effects of STS in angiotensin II-induced hypertensive renal and cardiac disease. The known role of oxidative stress in aging related neurodegenerative disease instigated a project that led to the creation of CSE overexpressing *Drosophila*. In **Chapter 7** we studied the protective potential of CSE overexpression and STS in a model of spinocerebellar ataxia type 3 in *Drosophila*. In addition, we investigated CSE expression in pontine tissue of SCA3 patients. In **Chapter 8**, all results are summarized and discussed, followed by a view on the future possibilities of H₂S-related research and therapeutic applications.

REFERENCES

- 1 Szabó C: Hydrogen sulphide and its therapeutic potential. *Nat Rev Drug Discov* 6: 917–935, 2007
- 2 Wang R: Physiological implications of hydrogen sulfide: a whiff exploration that blossomed. *Physiol. Rev.* 92: 791–896, 2012
- 3 Mustafa AK, Gadalla MM, Sen N, Kim S, Mu W, Gazi SK, Barrow RK, Yang G, Wang R, Snyder SH: H₂S signals through protein S-sulfhydration. *Sci Signal* 2: ra72, 2009
- 4 Mustafa AK, Sikka G, Gazi SK, Steppan J, Jung SM, Bhunia AK, Barodka VM, Gazi FK, Barrow RK, Wang R, Amzel LM, Berkowitz DE, Snyder SH: Hydrogen sulfide as endothelium-derived hyperpolarizing factor sulfhydrates potassium channels. *Circ. Res.* 109: 1259–1268, 2011
- 5 Yang G, Wu L, Jiang B, Yang W, Qi J, Cao K, Meng Q, Mustafa AK, Mu W, Zhang S, Snyder SH, Wang R: H₂S as a physiologic vasorelaxant: hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 322: 587–590, 2008
- 6 Bos EM, Wang R, Snijder PM, Boersema M, Damman J, Fu M, Moser J, Hillebrands J-L, Ploeg RJ, Yang G, Leuvenink HGD, van Goor H: Cystathionine γ -lyase protects against renal ischemia/reperfusion by modulating oxidative stress. *J. Am. Soc. Nephrol.* 24: 759–770, 2013
- 7 Beauchamp RO, Bus JS, Popp JA, Boreiko CJ, Andjelkovich DA: A critical review of the literature on hydrogen sulfide toxicity. *Crit. Rev. Toxicol.* 13: 25–97, 1984
- 8 Singh S, Padovani D, Leslie RA, Chiku T, Banerjee R: Relative contributions of cystathionine beta-synthase and gamma-cystathionase to H₂S biogenesis via alternative trans-sulfuration reactions. *J. Biol. Chem.* 284: 22457–22466, 2009
- 9 Abe K, Kimura H: The possible role of hydrogen sulfide as an endogenous neuromodulator. *J. Neurosci.* 16: 1066–1071, 1996
- 10 Shibuya N, Kimura H, Koike S, Tanaka M, Ishigami-Yuasa M, Kimura Y, Ogasawara Y, Fukui K, Nagahara N: A novel pathway for the production of hydrogen sulfide from D-cysteine in mammalian cells. *Nat Commun* 4: 1366, 2013
- 11 Sen U, Munjal C, Qipshidze N, Abe O, Gargoum R, Tyagi SC: Hydrogen sulfide regulates homocysteine-mediated glomerulosclerosis. *Am. J. Nephrol.* 31: 442–455, 2010
- 12 Zhao W, Wang R: H₂S-induced vasorelaxation and underlying cellular and molecular mechanisms. *Am. J. Physiol. Heart Circ. Physiol.* 283: H474–80, 2002
- 13 Lu M, Liu Y-H, Goh HS, Wang JJX, Yong Q-C, Wang R, Bian J-S: Hydrogen sulfide inhibits plasma renin activity. *J. Am. Soc. Nephrol.* 21: 993–1002, 2010
- 14 Kimura Y, Goto Y-I, Kimura H: Hydrogen sulfide increases glutathione production and suppresses oxidative stress in mitochondria. *Antioxid. Redox Signal.* 12: 1–13, 2010
- 15 Kimura Y, Kimura H: Hydrogen sulfide protects neurons from oxidative stress. *FASEB J.* 18: 1165–1167, 2004
- 16 Calvert JW, Jha S, Gundewar S, Elrod JW, Ramachandran A, Pattillo CB, Keivil CG, Lefer DJ: Hydrogen sulfide mediates cardioprotection through Nrf2 signaling. *Circ. Res.* 105: 365–374, 2009
- 17 Bouillaud F, Blachier F: Mitochondria and sulfide: a very old story of poisoning, feeding, and signaling? *Antioxid. Redox Signal.* 15: 379–391, 2011
- 18 Módos K, Coletta C, Erdélyi K, Papapetropoulos A, Szabó C: Intramitochondrial hydrogen sulfide production by 3-mercaptopyruvate sulfurtransferase maintains mitochondrial electron flow and supports cellular bioenergetics. *FASEB J.* 27: 601–611, 2013
- 19 Gubern M, Andriamihaja M, Nübel T, Blachier F, Bouillaud F: Sulfide, the first inorganic substrate for human cells. *FASEB J.* 21: 1699–1706, 2007
- 20 Fu M, Zhang W, Wu L, Yang G, Li H, Wang R: Hydrogen sulfide (H₂S) metabolism in mitochondria and its regulatory role in energy production. *Proc. Natl. Acad. Sci. U.S.A.* 109: 2943–2948, 2012
- 21 Teng H, Wu B, Zhao K, Yang G, Wu L, Wang R: Oxygen-sensitive mitochondrial accumulation of cystathionine β -synthase mediated by Lon protease. *Proc. Natl. Acad. Sci. U.S.A.* 110: 12679–12684, 2013
- 22 Yang G: Hydrogen sulfide in cell survival: a double-edged sword. *Expert Rev Clin Pharmacol* 4: 33–47, 2011
- 23 Li L, Fox B, Keeble J, Salto-Tellez M, Winyard PG, Wood ME, Moore PK, Whiteman M: The complex effects of the slow-releasing hydrogen sulfide donor GYY4137 in a model of acute joint inflammation and in human cartilage cells. *J. Cell. Mol. Med.* 17: 365–376, 2013
- 24 Papapetropoulos A, Pyriochou A, Altaany Z, Yang G, Marazioti A, Zhou Z, Jeschke MG, Branski LK, Herndon DN, Wang R, Szabó C: Hydrogen sulfide is an endogenous stimulator of angiogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 106: 21972–21977, 2009

- 25 Coletta C, Papapetropoulos A, Erdélyi K, Olah G, Módis K, Panopoulos P, Asimakopoulou A, Gerö D, Sharina I, Martin E, Szabó C: Hydrogen sulfide and nitric oxide are mutually dependent in the regulation of angiogenesis and endothelium-dependent vasorelaxation. *Proc. Natl. Acad. Sci. U.S.A.* 109: 9161–9166, 2012
- 26 Polhemus DJ, Kondo K, Bhushan S, Bir SC, Kevil CG, Murohara T, Lefer DJ, Calvert JW: Hydrogen sulfide attenuates cardiac dysfunction after heart failure via induction of angiogenesis. *Circ Heart Fail* 6: 1077–1086, 2013
- 27 Paul BD, Snyder SH: H₂S signalling through protein sulfhydration and beyond. *Nat. Rev. Mol. Cell Biol.* 13: 499–507, 2012
- 28 Zhao Y, Bhushan S, Yang C, Otsuka H, Stein JD, Pacheco A, Peng B, Devarie-Baez NO, Aguilar HC, Lefer DJ, Xian M: Controllable hydrogen sulfide donors and their activity against myocardial ischemia-reperfusion injury. *ACS Chem. Biol.* 8: 1283–1290, 2013
- 29 Li L, Whiteman M, Guan YY, Neo KL, Cheng Y, Lee SW, Zhao Y, Baskar R, Tan C-H, Moore PK: Characterization of a novel, water-soluble hydrogen sulfide-releasing molecule (GYY4137): new insights into the biology of hydrogen sulfide. *Circulation* 117: 2351–2360, 2008
- 30 Kolluru GK, Shen X, Bir SC, Kevil CG: Hydrogen sulfide chemical biology: pathophysiological roles and detection. *Nitric Oxide* 35: 5–20, 2013
- 31 Olson KR, Deleon ER, Gao Y, Hurley K, Sadauskas V, Batz C, Stoy GF: Thiosulfate: a readily accessible source of hydrogen sulfide in oxygen sensing. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 305: R592–603, 2013
- 32 Singh RP, Derendorf H, Ross EA: Simulation-based sodium thiosulfate dosing strategies for the treatment of calciphylaxis. *Clin J Am Soc Nephrol* 6: 1155–1159, 2011
- 33 Bos EM, Leuvenink HGD, Snijder PM, Kloosterhuis NJ, Hillebrands J-L, Leemans JC, Florquin S, van Goor H: Hydrogen sulfide-induced hypometabolism prevents renal ischemia/reperfusion injury. *J. Am. Soc. Nephrol.* 20: 1901–1905, 2009
- 34 Blackstone E, Morrison M, Roth MB: H₂S induces a suspended animation-like state in mice. *Science* 308: 518–518, 2005
- 35 Blackstone E, Roth MB: Suspended animation-like state protects mice from lethal hypoxia. *Shock* 27: 370–372, 2007
- 36 Cooper CE, Brown GC: The inhibition of mitochondrial cytochrome oxidase by the gases carbon monoxide, nitric oxide, hydrogen cyanide and hydrogen sulfide: chemical mechanism and physiological significance. *J. Bioenerg. Biomembr.* 40: 533–539, 2008
- 37 Xu Z, Prathapasinghe G, Wu N, Hwang S-Y, Siow YL, O K: Ischemia-reperfusion reduces cystathionine-beta-synthase-mediated hydrogen sulfide generation in the kidney. *Am. J. Physiol. Renal Physiol.* 297: F27–35, 2009
- 38 Elrod JW, Calvert JW, Morrison J, Doeller JE, Kraus DW, Tao L, Jiao X, Scalia R, Kiss L, Szabó C, Kimura H, Chow C-W, Lefer DJ: Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc. Natl. Acad. Sci. U.S.A.* 104: 15560–15565, 2007
- 39 Snijder PM, de Boer RA, Bos EM, van den Born JC, Ruifrok W-PT, Vreeswijk-Baudoin I, van Dijk MCRF, Hillebrands J-L, Leuvenink HGD, van Goor H: Gaseous hydrogen sulfide protects against myocardial ischemia-reperfusion injury in mice partially independent from hypometabolism. *PLoS ONE* 8: e63291, 2013
- 40 Bos EM, Snijder PM, Jekel H, Weij M, Leemans JC, van Dijk MCRF, Hillebrands J-L, Lisman T, van Goor H, Leuvenink HGD: Beneficial effects of gaseous hydrogen sulfide in hepatic ischemia/reperfusion injury. *Transpl. Int.* 25: 897–908, 2012
- 41 Qu K, Chen CPLH, Halliwell B, Moore PK, Wong PT-H: Hydrogen sulfide is a mediator of cerebral ischemic damage. *Stroke* 37: 889–893, 2006
- 42 Zhao W, Zhang J, Lu Y, Wang R: The vasorelaxant effect of H(2)S as a novel endogenous gaseous K(ATP) channel opener. *EMBO J.* 20: 6008–6016, 2001
- 43 Yan H, Du J, Tang C: The possible role of hydrogen sulfide on the pathogenesis of spontaneous hypertension in rats. *Biochem. Biophys. Res. Commun.* 313: 22–27, 2004
- 44 HARMAN D: Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 11: 298–300, 1956
- 45 Chen Y-H, Yao W-Z, Geng B, Ding Y-L, Lu M, Zhao M-W, Tang C-S: Endogenous hydrogen sulfide in patients with COPD. *Chest* 128: 3205–3211, 2005
- 46 Kabil H, Kabil O, Banerjee R, Harshman LG, Pletcher SD: Increased transsulfuration mediates longevity and dietary restriction in *Drosophila*. *Proc. Natl. Acad. Sci. U.S.A.* 108: 16831–16836, 2011
- 47 Kida K, Yamada M, Tokuda K, Marutani E, Kakinohana M, Kaneki M, Ichinose F: Inhaled hydrogen sulfide prevents neurodegeneration and movement disorder in a mouse model of

Parkinson's disease. *Antioxid. Redox Signal.* 15: 343–352, 2011

- 48 Paul BD, Sbodio JI, Xu R, Vandiver MS, Cha JY, Snowman AM, Snyder SH: Cystathionine γ -lyase deficiency mediates neurodegeneration in Huntington's disease. *Nature* 2014

