

University of Groningen

## Computational inference of regulator activity in a single input motif from gene expression data

Khanin, Raya; Vinciotti, Veronica; Mersinias, Vassilis; Smith, Colin; Wit, Ernst

*Published in:*  
Bmc Bioinformatics

*DOI:*  
[10.1186/1471-2105-6-S3-S7](https://doi.org/10.1186/1471-2105-6-S3-S7)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2005

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Khanin, R., Vinciotti, V., Mersinias, V., Smith, C., & Wit, E. (2005). Computational inference of regulator activity in a single input motif from gene expression data. *Bmc Bioinformatics*, 6 Suppl 3(57).  
<https://doi.org/10.1186/1471-2105-6-S3-S7>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

Oral presentation

Open Access

## Computational inference of regulator activity in a single input motif from gene expression data

Raya Khanin\*<sup>1</sup>, Veronica Vinciotti<sup>2</sup>, Vassilis Mersinias<sup>3</sup>, Colin Smith<sup>3</sup> and Ernst Wit<sup>1</sup>

Address: <sup>1</sup>Department of Statistics, 15 University Gardens, University of Glasgow, Glasgow G12 8QQ, UK., <sup>2</sup>School of Information Systems, Computing and Mathematics, Brunel University, Uxbridge, UB8 3PH, UK. and <sup>3</sup>School of Biomedical and Molecular Sciences, University of Surrey, Guildford, GU2 7XH, UK.

Email: Raya Khanin\* - raya@stats.gla.ac.uk

\* Corresponding author

from BioSysBio: Bioinformatics and Systems Biology Conference  
Edinburgh, UK, 14–15 July 2005

Published: 21 September 2005

*BMC Bioinformatics* 2005, **6**(Suppl 3):S7

This work deals with the computational modelling of gene regulation by transcription factor (regulator) in a single input motif (one regulator-many targets). The basic underlying problem is that the expression of a gene encoding the regulator provides only limited information about the activity of the regulator, which can be modified post-transcriptionally, for example, by phosphorylation. In the absence of reliable technology, it is of great importance to understand whether regulator activity can be inferred from gene expression data. We here develop a computational approach to reconstruct the regulator activity from gene expression data of the target genes in the SIM. We put forward a statistical framework that embeds the quantitative Michaelis-Menten model of the kinetics of gene regulation. The model uses the gene expression data on its original (rather than log) scale, employing the assumption of a log-normal noise distribution. For each target gene the maximum likelihood kinetic parameters of its regulation can be inferred independently given the profile of their common regulator. For a set of regulated target genes in the SIM, the regulator profile is reconstructed by maximizing the likelihood of the whole SIM.

The model is applied to two time series of the *Streptomyces coelicolor* bacterium: in one of them the transcriptional activator, *cdaR*, that is partly responsible for the production of one type of antibiotic, is knocked out. Our approach can be extended to include more complex regulatory relationships: multiple regulatory factors, competition, cooperativity.