

University of Groningen

Strategies to improve the outcome of biliary atresia

Vries, Willemien de

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Vries, W. D. (2011). *Strategies to improve the outcome of biliary atresia: lessons from the Dutch national database*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Bibliography

Bibliography

1. A-Kader H, Santangelo JD, Setchell KDR, et al. The effects of ursodeoxycholic acid (UDCA) therapy in biliary atresia (BA): a double blind, randomized, placebo controlled trial. *Ped Res* 1993;33:97A.
2. A-Kader HH, El-Ayyouti M, Hawas S, et al. HLA in Egyptian children with biliary atresia. *J Pediatr* 2002;141:432-434.
3. Alonso EM, Limbers CA, Neighbors K, et al. Cross-sectional analysis of health-related quality of life in pediatric liver transplant recipients. *J Pediatr* 2010;156:270-276.
4. Altman RP, Lily JR, Greenfeld J, et al. A multivariate risk factor analysis of the portoenterostomy (Kasai) procedure for biliary atresia. *Ann Surg* 1997;226:348-355.
5. Andrew M. The relevance of developmental hemostasis to hemorrhagic disorders of newborns. *Semin Perinatol* 1997;21:70-85.
6. Arrington-Sanders R, Yi MS, Tsevat J, et al. Gender differences in health-related quality of life of adolescents with cystic fibrosis. *Health Qual Life Outcomes* 2006;24:5.
7. Asthana S, McClean P, Stringer MD. Does the pediatric end-stage liver disease score or hepatic artery resistance index predict outcome after liver transplantation for biliary atresia? *Pediatr Surg Int* 2006;22:697-700.
8. Azar G, Beneck D, Lane B, et al. Atypical morphological presentation of biliary atresia and value of serial biopsies. *J Ped Gastroenterol Nutr* 2002;34:212-215.
9. Bardin R, Danon D, Tor R, et al. Reference values for gamma-glutamyl-transferase in amniotic fluid in normal pregnancies. *Prenat Diagn* 2009;29:703-706.
10. Barshes NR, Lee TC, Udell IW, et al. The pediatric end-stage liver disease (PELD) model as a predictor of survival benefit and posttransplant survival in pediatric liver transplant recipients. *Liver Transpl* 2006;12:475-480.
11. Beath S, Pearmain G, Kelly D, et al. Liver transplantation in babies and children with extrahepatic biliary atresia. *J Pediatr Surg* 1993;28:1044-1047.
12. Becker NS, Barshes NR, Aloia TA, et al. Analysis of recent pediatric orthotopic liver transplantation outcomes indicates that allograft type is no longer a predictor of survival. *Liver Transpl* 2008;14:1125-1132.
13. Bill AH, Haas JE, Foster GL. Biliary atresia: histopathologic observations and reflections upon its natural history. *J Ped Surg* 1977;12:977-982.
14. Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Chest* 1992;101:1644-1655.
15. Bourdeaux C, Tri TT, Gras J, et al. PELD score and posttransplant outcome in pediatric liver transplantation: a retrospective study of 100 recipients. *Transplantation* 2005;79:1273-1276.
16. Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology* 2005;42:1208-1236.
17. Bu LN, Chen HL, Chang CJ, et al. Prophylactic oral antibiotics in prevention of recurrent cholangitis after the Kasai portoenterostomy. *J Ped Surg* 2003;38:590-593.
18. Cakir M, Arikan C, Akman SA, et al. Infectious complications in pediatric liver transplantation candidates. *Pediatr Transpl* 2010;14:82-86.
19. Caudle SE, Katzenstein JM, Karpen SJ, et al. Language and motor skills are impaired in

- infants with biliary atresia before transplantation. *J Pediatr* 2010;156:940.
20. Chardot C, Carton M, Spire-Bendelac N, et al. Prognosis of biliary atresia in the era of liver transplantation: French national study from 1986-1996. *Hepatology* 1999;30:606-611.
 21. Chardot C, Carton M, Spire-Bendelac N, et al. Is the Kasai operation still indicated in children older than 3 months diagnosed with biliary atresia? *J Pediatr* 2001;138:224-228.
 22. Chardot C, Carton M, Spire-Bendelac N, et al. Epidemiology of biliary atresia in France: a national study 1986-96. *Journal of Hepatology* 1999;31:1006-1013.
 23. Choi SS, Diehl AM. Epithelial-to-mesenchymal transitions in the liver. *Hepatology* 2009;50:2007-2013.
 24. Cohen J. *Statistical power analysis for the behavioural sciences*. New York: Academy Press, 1988.
 25. Cowles R, Lobritto S, Ventura K, et al. Timing of liver transplantation in biliary atresia—results in 71 children managed by a multidisciplinary team. *J Ped Surg* 2008;43:1605-1609.
 26. Davenport M, Caponcelli E, Livesey E, et al. Surgical outcome in biliary atresia: etiology affects the influence of age at surgery. *Ann Surg* 2008;247:694-698.
 27. Davenport M, Gonde C, Redkar R, et al. Immunohistochemistry of the liver and biliary tree in extrahepatic biliary atresia. *J Ped Surg* 2001;36:1017-1025.
 28. Davenport M, Puricelli V, Farrant P, et al. The outcome of the older (>100 days) infant with biliary atresia. *J Ped Surg* 2004;39:575-581.
 29. Davenport M, Savage M, Mowat AP, et al. Biliary atresia splenic malformation syndrome: an etiologic and prognostic subgroup. *Surgery* 1993;113:662-668.
 30. Davenport M, Stringer MD, Tizzard SA, et al. Randomized, double-blind, placebo-controlled trial of corticosteroids after Kasai portoenterostomy for biliary atresia. *Hepatology* 2007;46:1821-1827.
 31. Davenport M, Tizzard SA, Underhill J, et al. The biliary splenic malformation syndrome: a 28-year single center retrospective study. *J Pediatr* 2006;149:393-400.
 32. Davenport M, Ville de Goyet J, Stringer MD, et al. Seamless management of biliary atresia in England and Wales (1999-2002). *Lancet* 2004;363:1354-1357.
 33. de Vries W, de Langen ZJ, Aronson DC, et al. Mortality of biliary atresia in children not undergoing liver transplantation in the Netherlands. *Pediatr Transplant* 2011;15:176-183.
 34. de Vries W, J Homan-van der Veen, Hulscher JB, et al. PD-H-105 Outcome of biliary atresia patients with 20 years native liver survival. *J Pediatr Gastroenterol Nutr* 2010;50:E32.
 35. Dellovade T, Romer JT, Curran T, et al. The Hedgehog pathway and neurological disorders. *Annu Rev Neurosci* 2006;29:563.
 36. DeRusso PA, Ye W, Shepherd R, et al. Growth failure and outcomes in infants with biliary atresia: a report from the biliary atresia research consortium. *Hepatology* 2007;46:1632-1638.
 37. Deurloo JA, Ekkelkamp S, Hartman EE, et al. Quality of life in adult survivors of correction of esophageal atresia. *Arch Surg* 2005;140:976-980.
 38. Diamond IR, Fecteau A, Millis JM, et al. Impact of graft type on outcome in pediatric liver transplantation. *Ann Surg* 2007;246:301-310.
 39. Diaz R, Kim J, Hui J, et al. Evidence for the epithelial to mesenchymal transition in biliary atresia fibrosis. *Human Pathology* 2008;39:102-115.
 40. DiNapoli PP, Murphy D. The marginalization of chronically ill adolescents. *Nurs Clin North*

- Am 2002;37:565-572.
41. Donaldson PT, Clare M, Constantini PK, et al. HLA and cytokine gene polymorphisms in biliary atresia. *Liver* 2002;22:213-219.
 42. Duffy JP, Kao K, KO CY, et al. Long-term patient outcome and quality of life after liver transplantation: analysis of 20-year survivors. *Ann Surg* 2010;252:652-661.
 43. Ecoffey C, Rothman E, Bernard O, et al. Bacterial cholangitis after surgery for biliary atresia. *J Pediatr* 1987;111:824-829.
 44. Fickert P, Fuchsichler A, Moustafa T, et al. Farnesoid X Receptor Critically Determines the Fibrotic Response in Mice but Is Expressed to a Low Extent in Human Hepatic Stellate Cells and Periductal Myofibroblasts. *Am J Pathol* 2009;175:2392-2405.
 45. Fickert P, Wagner M, Marschall H, et al. 24-norUrsodeoxycholic Acid Is Superior to Ursodeoxycholic Acid in the Treatment of Sclerosing Cholangitis in Mdr2 (Abcb4) Knockout Mice. *Gastroenterology* 2006;130:465-481.
 46. Fischler B, Ehrnst A, Forsgren M, et al. The viral association of neonatal cholestasis in Sweden: a possible link between cytomegalovirus infection and extrahepatic biliary atresia. *J Ped Gastroenterol Nutr* 1998;27:57-64.
 47. Fouquet V, Alves A, Branchereau S, et al. Long-term outcome of pediatric liver transplantation for biliary atresia: a 10-year follow-up in a single center. *Liver Transpl* 2005;11:152-160.
 48. Freeman RR, Wiesner RH, Edwards EB, et al. Results of the first year of the new liver allocation plan. *Liver Transpl* 2004;10:7-15.
 49. Garber J. Classification of childhood psychopathology: a developmental perspective. *Child Dev* 1984;55:30-48.
 50. Garcia-Barceló MM, Yeung MY, Miao XP, et al. Genome-wide association study identifies a susceptibility locus for biliary atresia on 10q24.2. *Hum Mol Gen* 2010;19:2917-2925.
 51. Gerver WJM, de Bruin R. Paediatric morphometrics, a reference manual. Utrecht, The Netherlands: Wetenschappelijke Uitgeverij Bunge, 1996.
 52. Glaser SS, Gaudio E, Miller T, et al. Cholangiocyte proliferation and liver fibrosis. *Exp Rev* 2009;11.
 53. Goodman E, Amick BC, Rezendes MO, et al. Influence of gender and social class on adolescents' perceptions of health. *Arch Pediatr Adolesc Med* 1997;151:899-904.
 54. Grootenhuis MA, Stam H, Destrée-Vonk A. Course of life questionnaire for young adults. *Gedrag Gezond* 2003;31:336-350.
 55. Gross CR, Malinchoc M, Kim WR, et al. Quality of life before and after liver transplantation for cholestatic liver disease. *Hepatology* 1999;29:356-364.
 56. Gutteling JJ, Duivenvoorden HJ, Bussbach JJV, et al. Psychological determinants of health-related quality of life in patients with chronic liver disease. *Psychosomatics* 2010;51:157-165.
 57. Hadzic N, Davenport M, Tizzard S, et al. Long-term survival following Kasai portoenterostomy: is chronic liver disease inevitable? *J Pediatr Gastroenterol Nutr* 2003;37:430-433.
 58. Hanada S, Harada M, Koga H, et al. Tumor necrosis factor-alpha and interferon-gamma directly impair epithelial barrier function in cultured mouse cholangiocytes. *Liver Int* 2003;23:3-11.

59. Harada K, Nakanuma Y. Biliary innate immunity: function and modulation. *Mediators Inflamm* 2010;Article ID 373878.
60. Hartley JL, Davenport M, Kelly DA. Biliary atresia. *Lancet* 2009;374:1704-1713.
61. Haupt R, Byrne J, Connelly RR. Smoking habits in survivors of childhood and adolescent cancer. *Med Pediatr Oncol* 1992;20:301-306.
62. Hays DM, Kimura K. Biliary atresia: new concepts of management. *Curr Probl Surg* 1981;18:541-608.
63. Hofmann A, Zakko S, Lira M, et al. Novel biotransformation and physiological properties of norursodeoxycholic acid in humans. *Hepatology* 2005;42:1391-1398.
64. Hollander D, Rim E. Factors affecting the absorption of vitamin K-1 in vitro. *Gut* 1976;17:450-455.
65. Hollen PJ, Hobbie WL. Decision making and risk behaviors of cancer-surviving adolescents and their peers. *Decision making and risk behaviors of cancer-surviving adolescents and their peers* 1996;13:121-133, discussion 135-137.
66. Host A, Halken S. Primary prevention of food allergy in infants who are at risk. *Curr Opin Allergy Clin Immunol* 2005;5:255-259.
67. Houwen RH, Kerremans II, van Steensel-Moll HA, et al. Time-space distribution of extrahepatic biliary atresia in the Netherlands and West Germany. *Z Kinderchir* 1988;43:68-71.
68. Houwen RH, Zwierstra RP, Severijnen RS, et al. Prognosis of extrahepatic biliary atresia. *Arch Dis Child* 1989;64:214-218.
69. Houwen RHJ, Bijleveld CMA, De Vries-Hospers HG. Ciprofloxacin for cholangitis after hepatic portoenterostomy. *Lancet* 1987;1:1367.
70. Howard ER, MacLean G, Nio M, et al. Survival patterns in biliary atresia and comparison of quality of life of long-term survivors in Japan and England. *J Ped Surg* 2001;36:892-897.
71. Hsiao CH, Chang MH, Chen HL, et al. Universal screening for biliary atresia using an infant stool color card in Taiwan. *Hepatology* 2008;47:1233-1240.
72. Humphrey TM, Stringer MD. Biliary atresia: US diagnosis. *Radiology* 2007;244:845-851.
73. Hyams JS, Glaser JH, Leichtner AM, et al. Discordance for biliary atresia in two sets of monozygotic twins. *J Pediatr* 1985;107:420-422.
74. Jain A, Mazariegos G, Kashyap R, et al. Pediatric liver transplantation. A single center experience spanning 20 years. *Transplantation* 2002;73:941-947.
75. Jay CL, Butt Z, Ladner DP, et al. A review of quality of life instruments used in liver transplantation. *J Hepatol* 2009;51:949-959.
76. Jones EA. Fatigue Complicating Chronic Liver Disease. *Metabolic Brain Disease* 2004;19:421-429.
77. Jung Y, Brown KD, Witek RP, et al. Accumulation of hedgehog-responsive progenitors parallels alcoholic liver disease severity in mice and humans. *Gastroenterology* 2008;134:1543.
78. Jung Y, McCall SJ, Li YX, and Diehl AM. Bile ductules and stromal cells express hedgehog ligands and/or hedgehog target genes in primary biliary cirrhosis. *Hepatology* 45, 1091-1096. 2007.
79. Kahn E. Biliary atresia revisited. *Pediatr Dev Pathol* 2004;7:109-124.
80. Karrer FM, Hall RJ, Lilly JR. Biliary atresia and the polysplenia syndrome. *J Ped Surg*

- 1991;26:524-527.
81. Karrer FM, Price MR, Bensard DD, et al. Long-term results with the Kasai operation for biliary atresia. *Arch Surg* 1996;131:493-496.
 82. Kasai M, Suzuki S. A new operation for 'non-correctable' biliary atresia: portoenterostomy. *Shujutsu* 1959;13:733-739.
 83. Keil R, Snajdauf J, Rygl M, et al. Diagnostic efficacy of ERCP in cholestatic infants and neonates - a retrospective study on a large series. *Endoscopy* 2010;42:121-126.
 84. Kelly D, Davenport M. Current management of biliary atresia. *Arch Dis Child* 2007;92:1132-1135.
 85. Kobayashi K, Kubota M, Okuyama N, et al. Mother-to-daughter occurrence of biliary atresia: a case report. *J Ped Surg* 2008;43:1566-1568.
 86. Kobayashi H, Tamatani T, Tamura T, et al. Maternal microchimerism in biliary atresia. *Journal of Pediatric Surgery* 2007;42:987-991.
 87. Kremer A, Beuers U, Oude-Elferink R, et al. Pathogenesis and Treatment of Pruritus in Cholestasis. *Drugs* 2008;68:2163-2182.
 88. Kuroda T, Saeki M, Nakano M, et al. Biliary atresia, the next generation: a review of liver function, social activity, and sexual development in the late postoperative period. *J Ped Surg* 2002;37:1709-1712.
 89. Landing BH. Considerations of the pathogenesis of neonatal hepatitis, biliary atresia and choledochal cyst--the concept of infantile obstructive cholangiopathy. *Prog Pediatr Surg* 1974;6:113-139.
 90. Larossa-Haro A, Caro-López AM, Coello-Ramírez P, et al. Duodenal tube test in the diagnosis of biliary atresia. *J Pediatr Gastroenterol Nutr* 2001;32:311-315.
 91. Lauth M, Rohnalter V, Bergström A, et al. Antipsychotic drugs regulate hedgehog signaling by modulation of 7-dehydrocholesterol reductase levels. *Mol Pharm* 2010;78:486-496.
 92. Lauth M, Toftgard R. Non-canonical activation of GLI transcription factors. *Cell Cycle* 2007;6:2458-2463.
 93. Lazaridis K, Gores GJ, Lindor KD. Ursodeoxycholic acid 'mechanisms of action and clinical use in hepatobiliary disorders'. *J Hepatol* 2001;35:134-146.
 94. Lee JM, Dedhar S, Kalluri R, et al. The epithelial-mesenchymal transition: new insights in signaling, development, and disease. *J Cell Biol* 2006;172:973-981.
 95. Lee MJ, Kim MJ, Yoon CS, et al. Gadopentetate dimeglumin-enhanced MR cholangiopancreatography in infants with cholestasis. *Pediatr Radiol* 2011;41:488-494.
 96. Lewis M, Miller AM. *Handbook of developmental psychopathology*. New York, USA: Plenum Press, 1990.
 97. Li MK, Crawford JM. The pathology of cholestasis. *Sem Liver Dis* 2004;24:21-42.
 98. Lien TH, Chang MH, Wu JF, et al. Effects of the infant stool color card screening program on 5-year outcome of biliary atresia in Taiwan. *Hepatology* 2011;53:202-208.
 100. Lindor KD, Therneau TM, Jorgensen RA, et al. Effects of ursodeoxycholic acid on survival in patients with primary biliary cirrhosis. *Gastroenterology* 1996;110:1515-1518.
 101. Lindor KD, Kowdley KV, Luketic VAC, et al. High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009;50:808-814.
 102. Lu B, Brindley S, Tucker R, et al. α -Enolase autoantibodies cross-reactive to viral proteins

- in a mouse model of biliary atresia. *Gastroenterology* 2010;139:1753-1761.
103. Lykavieris P, Chardot C, Sokhn M, et al. Outcome in adulthood of biliary atresia: A study of 63 patients who survived for over 20 years with their native liver. *Hepatology* 2005;41:366-371.
 104. Mack CL. The pathogenesis of biliary atresia: evidence for a virus-induced immune disease. *Sem Liver Dis* 2007;27:233-242.
 105. Mack C, Tucker R, Lu B, et al. Cellular and humoral autoimmunity directed at bile duct epithelia in murine biliary atresia. *Hepatology* 2006;44:1231-1239.
 106. Marchesini G, Bianchi G, Amodio P, et al. Factors associated with poor health-related quality of life of patients with cirrhosis. *Gastroenterology* 2001;120:170-178.
 107. Matthews R, EauClaire S, Mugnier M, et al. DNA hypomethylation causes bile duct defects in zebrafish and is a distinguishing feature of infantile biliary atresia. *Hepatology* 2011;53:905-914.
 108. McDiarmid SV, Anand R, Lindblad AS. Development of a pediatric end-stage liver disease score to predict poor outcome in children awaiting liver transplantation. *Transplantation* 2002;74:173-181.
 109. McDiarmid SV, Merion RM, Dykstra DM, et al. Selection of pediatric candidates under the PELD system. *Liver Transpl* 2004;10:s23-s30.
 110. McKiernan PJ, Baker AJ, Lloyd C, et al. British paediatric surveillance unit study of biliary atresia: outcome at 13 years. *J Ped Gastroenterol Nutr* 2009;48:78-81.
 111. McKiernan P, Baker A, Kelly D. The frequency and outcome of biliary atresia in the UK and Ireland. *Lancet* 2000;355:25-29.
 112. McNinch AW, Tripp JH. Haemorrhagic disease of the newborn in the British isles: two year prospective study. *BMJ* 1991;303:1105-1109.
 113. McNinch A, Busfield A, Tripp J. Vitamin K deficiency bleeding in Great Britain and Ireland: British Paediatric Surveillance Unit Surveys, 1993-1994 and 2001-2002. *Arch Dis Child* 2007;92:759-766.
 114. Mehta G, Abraldes JG, Bosch J. Developments and controversies in the management of oesophageal and gastric varices. *Gut* 2010;59:701-705.
 115. Miethke AG, Saxena V, Shivakumar P, et al. Post-natal paucity of regulatory T cells and control of NK cell activation in experimental biliary atresia. *J Hepatology* 2010;52:718-726.
 116. Mihatsch WA, Hogel J, Pohlandt F. Hydrolysed protein accelerates the gastrointestinal transport of formula in preterm infants. *Acta Paediatr* 2001;90:196-198.
 117. Mizuta K, Sanada T, Wakiya T, et al. Living-donor liver transplantation in 126 patients with biliary atresia: single-center experience. *Transplant Proc* 2010;42:4127-4131.
 118. Moss MH. Hypoprothrombinemic bleeding in a young infant. Association with a soy protein formula. *Am J Dis Child* 1969;117:540-542.
 119. Mrus JM, Williams PL, Tsevat J, et al. Gender differences in health-related quality of life in patients with HIV/AIDS. *Qual Life Res* 2005;14:479-491.
 120. Münz C, Lünemann JD, Teague Getts M, et al. Antiviral immune responses: triggers or triggered by immunity? *Nat Rev Immunol* 2009;9:246-258.
 121. Muraji T, Harada T, Miki K, et al. Urinary sulfated bile acid concentrations in infants with biliary atresia and breast-feeding jaundice. *Pediatr Int* 2003;45:283.
 122. Muraji T, Hosaka N, Irie N, et al. Maternal microchimerism in underlying pathogenesis of

- biliary atresia: quantification and phenotypes of maternal cells in the liver. *Pediatrics* 2008;121:517-521.
123. Mushtaq I, Logan S, Morris M, et al. Screening of newborn infants for cholestatic hepatobiliary disease with tandem mass spectrometry. *BMJ* 1999;319:471-477.
124. Narayanaswamy B, Gonde C, Tredger JM, et al. Serial circulating markers of inflammation in biliary atresia: Evolution of the post-operative inflammatory process. *Hepatology* 2007;46:180-187.
125. Nederlandse Vereniging voor Kindergeneeskunde. Richtlijn preventie, diagnostiek en behandeling van hyperbilirubinemie bij de pasgeborene, geboren na een zwangerschapsduur van meer dan 35 weken. CBO richtlijn, 2008.
126. Ng VL, Fecteau A, Shepherd R, et al. Outcomes of 5-year survivors of pediatric liver transplantation: report on 461 children from a north american multicenter registry. *Pediatrics* 2008;122:e1128-e1135.
127. Nietgen GW, Vacanti JP, Perez-Atayde AR. Intrahepatic bile duct loss in biliary atresia despite portoenterostomy: a consequence of ongoing obstruction. *Gastroenterology* 1992;102:2126-2133.
128. Nio M, Ohi R, Miyano T, et al. Five- and 10-year survival rates after surgery for biliary atresia: a report from the Japanese biliary atresia registry. *J Ped Surg* 2003;38:997-1000.
129. Nio M, Ohi R, Hayashi Y, et al. Current status of 21 patients who have survived more than 20 years since undergoing surgery for biliary atresia. *J Ped Surg* 1996;31:381-384.
130. Omenetti A, Porrello A, Jung Y, et al. Hedgehog signalling regulates epithelial-mesenchymal transition during biliary fibrosis in rodents and humans. *J Clin Invest* 2008;118:3331-3342.
131. Omenetti A, Yang L, Li Y, et al. Hedgehog-mediated mesenchymal-epithelial interactions modulate hepatic response to bile duct ligation. *Lab Invest* 2007;87:499-514.
132. Pakkarinen MP, Lampela H, Gylling H, et al. Surrogate markers of cholesterol metabolism in children with native liver after successful portoenterostomy for biliary atresia. *J Ped Surg* 2010;45:1659-1664.
133. Peeters PMJG, Sieders E, vd Heuvel M, et al. Predictive factors for portal fibrosis in pediatric liver transplant recipients. *Transplantation* 2000;70:1581-1587.
134. Pellegrino M, Sacco M, D'Altilia MR. Hemorrhagic disease in an infant fed on a vitamin K-deficient soya formula. *J Pediatr Gastroenterol Nutr* 1996;23:413-414.
135. Petersen C, Harder D, Melter M, et al. Postoperative high-dose steroids do not improve mid-term survival with native liver in biliary atresia. *Am J Gastroenterol* 2008;103:712-719.
136. Petersen C, Meier PN, Schneider A, et al. Endoscopic retrograde cholangiopancreatography prior to explorative laparotomy avoids unnecessary surgery in patients suspected for biliary atresia. *J Hepatol* 2009;51:1055-1060.
137. Petersen C, Biermanns D, Kuske M, et al. New aspects in a murine model for extrahepatic biliary atresia. *J Ped Surg* 1997;32:1190-1195.
138. Riepenhoff-Talty M, Gouvea V, Evans MJ, et al. Detection of group C rotavirus in infants with extrahepatic biliary atresia. *J Infect Dis* 1996;174:8-15.
139. Riepenhoff-Talty M, Schäkel K, Clark HF, et al. Group A rotaviruses produce extrahepatic biliary obstruction in orally inoculated newborn mice. *Pediatr Res* 1993;33:394-399.
140. Roderfeld M, Rath T, Voswinckel R, et al. Bone marrow transplantation demonstrates

- medullar origin of CD34+ fibrocytes and ameliorates hepatic fibrosis in *Abcb4* $-/-$ mice. *Hepatology* 2010;51:267-276.
141. Rudin C, Hann C, Lartera J, et al. Treatment of Medulloblastoma with Hedgehog Pathway Inhibitor GDC-0449. *New England Journal of Medicine* 2009;361:1173-1178.
 142. Rygiel KA, Robertson H, Marshall HL, et al. Epithelial mesenchymal transition contributes to portal tract fibrogenesis during human chronic liver disease. *Lab Invest* 2008;88:112-123.
 143. Scheenstra R, Peeters P, Verkade H, et al. Graft fibrosis after pediatric liver transplantation: Ten years of follow-up. *Hepatology* 2009;49:880-886.
 144. Schippers EI. Beleidsreactie minister van Volksgezondheid, Welzijn en Sport aan de voorzitter van de Tweede Kamer. www.rijksoverheid.nl, 2011.
 145. Scholten D, Österreicher CH, Scholten A, et al. Genetic labeling does not detect epithelial to mesenchymal transition of cholangiocytes in liver fibrosis in mice. *Gastroenterology* 2010;39:987-998.
 146. Schreiber RA, Barker CC, Roberts EA, et al. Biliary atresia in Canada: the effect of centre caseload experience on outcome. *J Pediatr Gastroenterol Nutr* 2010;51:61-65.
 147. Schreiber R, Barker C, Roberts E, et al. Biliary Atresia: The Canadian Experience. *J Pediatr* 2007;151:659-665.
 148. Schurgers LJ, Vermeer C. Determination of phylloquinone and menaquinones in food. Effect of food matrix on circulating vitamin K concentrations. *Haemostasis* 2000;30:298-307.
 149. Schwartz CE, Sprangers MAG. Methodological approaches for assessing response shift in longitudinal health-related quality of life research. *Soc Sci Med* 1999;48:1531-1548.
 150. Sebagh M, Rifai K, Féray C, et al. All liver recipients benefit from the protocol 10-year liver biopsies. *Hepatology* 2003;37:1293-1301.
 151. Serinet MO, Broué P, Jacquemin E, et al. Management of patients with biliary atresia in France: Results of a decentralized policy 1986-2002. *Hepatology* 2006;44:75-84.
 152. Serinet M, Wildhaber B, Broué P, et al. Impact of age at Kasai operation on its results in late childhood and adolescence: a rational basis for biliary atresia screening. *Pediatrics* 2009;123:1280-1286.
 153. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Ped Neurol* 2003;29:410-421.
 154. Shinkai M, Ohhama Y, Take H, et al. Long-term outcome of children with biliary atresia who were not transplanted after the Kasai operation: > 20-year experience at a children's hospital. *J Pediatr Gastroenterol Nutr* 2009;48:443-450.
 155. Shinkai M, Ohhama Y, Take H, et al. Evaluation of the PELD risk score as a severity index of biliary atresia. *J Pediatr Surg* 2003;38:1001-1004.
 156. Shivakumar P, Campbell KM, Sabla GE, et al. Obstruction of extrahepatic bile ducts by lymphocytes is regulated by IFN- γ in experimental biliary atresia. *J Clin Invest* 2004;114:322-329.
 157. Shivakumar P, Sabla GE, Whittington P, et al. Neonatal NK cells target the mouse duct epithelium via Nkg2d and drive tissue-specific injury in experimental biliary atresia. *J Clin Invest* 2009;119:2281-2290.
 158. Shneider BL, Abel RB, Raghumathan T, et al. A prospective multi-centered investigation

- of vitamin supplementation in infants with biliary atresia: interim analysis from the biliary atresia reserach consortium (BARC). *Hepatology* 2009;50:631A-632A.
159. Shneider B, Brown M, Haber B, et al. A multicenter study of the outcome of biliary atresia in the United States, 1997 to 2000. *J Pediatr* 2006;148:467-474.
160. Sicklick JK, Choi SS, Bustamante M, et al. Evidence for epithelial-mesenchymal transitions in adult liver cells. *Am J Gastrointest Liver Physiol* 2006;291:G575-G583.
161. Silveira TR, Salzano FM, Donaldson PT, et al. Association between HLA and extrahepatic biliary atresia. *J Ped Gastroenterol Nutr* 1993;16:114-117.
162. Singh AM, Dalglish DG. The emulsifying properties of hydrolysates of whey proteins. *J Dairy Sci* 1998;81:919-924.
163. Sokol RJ, Mack C, Narkewicz MR, et al. Pathogenesis and outcome of biliary atresia: current concepts. *J Pediatr Gastroenterol Nutr* 2003;37:4-21.
164. Stam H, Grootenhuis MA, Caron HN, et al. Quality of life and current coping in young adult survivors of childhood cancer: positive expectations about the further course of the disease were correlated with a better quality of life. *Psycho-oncology* 2006;15:31-43.
165. Stam H, Grootenhuis MA, Last BF. The course of life of survivors of childhood cancer. *Psycho-oncology* 2005;14:227-238.
166. Stam H, Hartman E, Deurloo J, et al. Young adult patients with a history of pediatric disease: impact on course of life and transition into adulthood. *Journal of Adolescent Health* 2006;39:4-13.
167. Strickland AD, Shannon K. Studies in etiology of extrahepatic biliary atresia: tim-space clustering. *J Pediatr* 1982;100:749-753.
168. Strickland AD, Shannon K, Coln CD. Biliary atresia in two sets of twins. *J Pediatr* 1985;107:18-20.
169. Sumida W, Kaneko K, Ono Y, et al. Different polyunsaturated fatty acid profiles in patients with biliary atresia after successful Kasai operation and liver transplantation. *Pediatr Surg Int* 2009;25:255-259.
170. Suskind DL, Rosenthal P, Heyman MB, et al. Maternal microchimerism in the livers of patients with biliary atresia. *BMC Gastroenterology* 2004;4:14.
171. Sutor AH, von Kries R, Cornelissen EA. Vitamin K deficiency bleeding (VKDB) in infancy. ISTH Pediatric/Perinatal Subcommittee. International Society on Thrombosis and Haemostasis. *Thromb Haemost* 1999;81:456-461.
172. Sutor AH, Dages N, Niederhoff H. Late form of Vitamin K Deficiency Bleeding in Germany. *Klin Padiatr* 1995;207:89,97.
173. Syn W, Jung Y, Omenetti A, et al. Hedgehog-mediated epithelial-to-mesenchymal transition and fibrogenic repair in nonalcoholic fatty liver disease. *Gastroenterology* 2009;137:1478-1488.
174. Tainaka T, Kaneko K, Seo T, et al. Hepatolithiasis after hepatic portoenterostomy for biliary atresia. *J Ped Surg* 2006;41:808-811.
175. Tao ML, Zeltzer LK, Byrne J, et al. Smoking in adult survivors of childhood acute lymphoblastic leukemia. *J Natl Cancer Inst* 1998;90:219-225.
176. Tatekawa Y, Asonuma K, Uemoto S, et al. Liver transplantation for biliary atresia associated with malignant hepatic tumors. *J Pediatr Surg* 2001;36:436-439.

177. Taura K, Miura K, Iwaisako K, et al. Hepatocytes do not undergo epithelial-mesenchymal transition in liver fibrosis in mice. *Hepatology* 2010;51:1027-1036.
178. Taylor RM, Cheeseman P, Davenport M, et al. Humoral immunity in children with biliary atresia splenic malformation syndrome. *Eur J Pediatr* 2003;162:539-540.
179. Tiao MM, Tsai SS, Kuo HW, et al. Epidemiological features of biliary atresia in Taiwan, a national study 1996-2003. *J Gastroenterol Hepatol* 2008;34:62-66.
180. Tome S, Wells JT, Said A, et al. Quality of life after liver transplantation. A systematic review. *J Hepatol* 2008;48:567-577.
181. Tuinman M, Gazendam-Donofrio S, Hoekstra-Weebers J. Screening and referral for psycho social distress in oncologic practice. *Cancer* 2008;113:870-878.
182. Tyler KL, Sokol RJ, Oberhaus SM, et al. Detection of reovirus RNA in hepatobiliary tissues from patients with extrahepatic biliary atresia and choledochal cysts. *Hepatology* 1998;27:1475-1482.
183. Ullrich D, Rating D, Schröter W, et al. Treatment with ursodeoxycholic acid renders children with biliary atresia suitable for liver transplantation. *Lancet* 1987;2:1324.
184. Utterson E, Shepherd R, Sokol R, et al. Biliary atresia: Clinical profiles, risk factors, and outcomes of 755 patients listed for liver transplantation. *J Pediatr* 2005;147:180-185.
185. van der Plas SM, Hansen BE, de Boer JB, et al. The Liver Disease Symptom Index 2.0; validation of a disease-specific questionnaire. *Qual Life Res* 2004;13:1469-1481.
186. van der Steeg AFW, de Vries J, Roukema JA. The value of quality of life and health status measurements in the evaluation of the well-being of breast cancer survivors. *Eur J Surg Oncol* 2008;34:1225-1230.
187. van Hasselt P, de Koning T, Kvist N, et al. Prevention of vitamin K deficiency bleeding in breastfed infants: lessons from the Dutch and Danish biliary atresia registries. *Pediatrics* 2008;121:e857-e863.
188. van Hasselt PM, Kok K, Vorselaars ADM, et al. Vitamin K deficiency bleeding in cholestatic infants with alpha-1-antitrypsin deficiency. *Arch Dis Child - Fet Neonat Ed* 2009;94:F456-F460.
189. van Heurn LW, Saing H, Tam PK. Cholangitis after hepatic portoenterostomy for biliary atresia: a multivariate analysis of risk factors. *J Pediatr* 2003;142:566-571.
190. VanderZee KI, Sanderman R, Heyink JW, et al. Psychometric qualities of the RAND 36-Item health survey 1.0: A multidimensional measure of general health status. *Int J Behav Med* 1996;3:104-122.
191. Varela-Fascinetto G, Castaldo P, Fox IJ, et al. Biliary atresia-Polysplenia syndrome. *Ann Surg* 1998;227:583-589.
192. Vazquez J, López Gutierrez JC, Gámez M, et al. Biliary atresia and the polysplenia syndrome: its impact on final outcome. *J Ped Surg* 1995;30:485-487.
193. Vegting IL, Tabbers MM, Taminiou JAJM, et al. Is endoscopic retrograde cholangiopancreatography valuable and safe in children of all ages? *J Ped Gastroenterol Nutr* 2009;48:66-71.
194. Verrill JR, Schafer J, Vanatta K. Aggression, antisocial behavior, and substance abuse in survivors of pediatric cancer: possible protective effects of cancer and its treatment. *J Pediatr Psychol* 2000;25:493-502.
195. Vo Thi Diem H, Evrard V, Tran Vinh H, et al. Pediatric liver transplantation for biliary atresia:

- results of primary grafts in 328 recipients. *Transplantation* 2003;75:1692-1697.
196. Vongwiwatana A, Tasanarong A, Rayner DC, et al. Epithelial to mesenchymal transition during late deterioration of human kidney transplants: the role of tubular cells in fibrogenesis. *Am J Transplant* 2005;5:1367-1374.
197. Wada H, Muraji T, Yokoi A, et al. Insignificant seasonal and geographical variation in incidence of biliary atresia in Japan: a regional survey of over 20 years. *J Ped Surg* 2007;42:2090-2092.
198. Wayman KI, Cox KL, Esquivel CO. Neurodevelopmental outcome of young children with extrahepatic biliary atresia 1 year after liver transplantation. *J Pediatr* 1997;131:898.
199. Weinstein ND. Testing four competing theories of health-protective behaviour. *Health psychol* 1993;12:324-333.
200. Wessels M, De Graaf B, Cohen-Overbeek T, et al. A new syndrome with noncompaction cardiomyopathy, bradycardia, pulmonary stenosis, atrial septal defect and heterotaxy with suggestive linkage to chromosome 6p. *Hum Genet* 2008;122:595-603.
201. Wiesner RH, McDiarmid SV, Kamath PS, et al. MELD and PELD: application of survival models to liver allocation. *Liver Transpl* 2001;7:567-580.
202. Wijnhoven HAH, Kriegsman DMW, Snoek FJ, et al. Gender differences in health-related quality of life among asthma patients. *J Asthma* 2003;40:189-199.
203. Wildhaber BE, Majno P, Mayr J, et al. Biliary Atresia: Swiss National Study, 1994-2004. *J Ped Gastroenterol Nutr* 2008;46:299-307.
204. Willot S, Uhlen S, Michaud L, et al. Effect of ursodeoxycholic acid on liver function in children after successful surgery for biliary atresia. *Pediatrics* 2008;122:1232-1241.
205. Wong KKY, Chung PHY, Chan IHY, et al. Performing Kasai portoenterostomy beyond 60 days of life is not necessarily associated with a worse outcome. *J Ped Gastroenterol Nutr* 2010;51:631-634.
206. Wood RP, Langnas AN, Stratta RJ, et al. Optimal therapy for patients with biliary atresia: portoenterostomy ("Kasai" procedures) versus primary transplantation. *J Ped Surg* 1990;25:153-162.
207. Yang H, Groen H, Verkade H, et al. Steroids should be administered after Kasai portoenterostomy for biliary atresia: a systematic review and meta-analysis (abstract). *United European Gastroenterology Federation* 2008;Vienna; Oct 18-22:poster 0645.
208. Yang H, Plosch T, Lisman T, et al. Inflammation mediated down-regulation of hepatobiliary transporters contributes to intrahepatic cholestasis and liver damage in murine biliary atresia. *Pediatr Res* 2009;66:380-385.
209. Yauch R, Dijkgraaf G, Alicke B, et al. Smoothed mutation confers resistance to a hedgehog pathway inhibitor in medulloblastoma. *Science* 2009;326:572-574.
210. Yoon PW, Bresee JS, Olney RS, et al. Epidemiology of biliary atresia: a population-based study. *Pediatrics* 1997;99:376-382.
211. Younossi ZM, Boparai N, Price LL, et al. Health-related quality of life in chronic liver disease: the impact of type and severity of disease. *Am J Gastroenterol* 2001;96:2199-2205.
212. Younossi ZM, Kiwi ML, Boparai N, et al. Cholestatic liver diseases and health-related quality of life. *Am J Gastroenterol* 2000;95:497-502.
213. Yssaad-Fesselier R, Lion-François L, Herbillon V, et al. Intellectual and visuo-spatial assessment in long-term pediatric liver transplantation for biliary atresia. *Transplantation*

- 2009;87:1427-1428.
214. Yuasa T, Tsuji H, Kimura S, et al. Human leukocyte antigens in Japanese patients with biliary atresia:retrospective analysis of patients who underwent living donor liver transplantation. *Hum Immunol* 2005;66:295-300.
215. Zeisberg M, Neilson EG. Biomarkers for epithelial-mesenchymal transitions. *J Clin Invest* 2009;119:1429-1437.
216. Zhang SJ, Huang LH, Wen YL, et al. Impact of personality and coping mechanisms on health-related quality of life in liver transplantation recipients. *Hepatobiliary Pancreat Dis Int* 2005;4:356-359.

