

University of Groningen

## Microvillus Inclusion Disease. Lessons about the apical plasma membrane.

Golachowska, Magdalena Renata

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2011

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Golachowska, M. R. (2011). *Microvillus Inclusion Disease. Lessons about the apical plasma membrane.*  
s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## REFERENCES

---

## References

---

- Aikawa Y, Martin TF** (2003) ARF6 regulates a plasma membrane pool of phosphatidylinositol(4,5)bisphosphate required for regulated exocytosis. *J Cell Biol* 162, 647–659.
- Aït Slimane T, Trugnan G, van IJzendoorn SCD, Hoekstra D** (2003). Raft-mediated trafficking of apical resident proteins occurs in both direct and transcytotic pathways in polarized hepatic cells: role of distinct lipid microdomains. *Mol Biol Cell* 14, 611-624.
- Ambu R, Riva A, Lai ML, Loffredo F, Riva FT, Tandler B** (2000). Scanning electron microscopy of the interior of cells in Hürthle cell tumors. *Ultrastruct Pathol* 24(4), 211-219.
- Ameen NA, Salas PJ** (2000). Microvillus inclusion disease: a genetic defect affecting apical membrane protein traffic in intestinal epithelium. *Traffic* 1, 76-83.
- An T, Ghatak N, Kastner R, Kay S, Lee HM** (1983). Hyaline globules and intracellular lumina in a hepatocellular carcinoma. *Am J Clin Pathol* 79(3), 392-6.
- Ang AL, Taguchi T, Francis S, Fölsch H, Murrells LJ, Pypaert M, Warren G, Mellman I** (2004). Recycling endosomes can serve as intermediates during transport from the Golgi to the plasma membrane of MDCK cells. *J Cell Biol* 167, 531–543.
- Apodaca G, Katz LA, Mostov KE** (1994). Receptor-mediated transcytosis of IgA in MDCK cells is via apical recycling endosomes. *J Cell Biol* 125, 67–86.
- Babbey CM, Ahktar N, Wang E, Chen CC, Grant BD, Dunn KW** (2006). Rab10 regulates membrane transport through early endosomes of polarized Madin-Darby canine kidney cells. *Mol Biol Cell* 17, 3156–3175.
- Babia T, Kok JW, van der Haar M, Kalicharan R, Hoekstra D** (1994). Transport of biosynthetic sphingolipids from Golgi to plasma membrane in HT29 cells: involvement of different carrier vesicle populations. *Eur J Cell Biol* 63(2), 172-81.
- Bacallao R, Antony C, Dotti C, Karsenti E, Stelzer EH, Simons K** (1989). The subcellular organization of Madin-Darby canine kidney cells during the formation of a polarized epithelium. *J Cell Biol* 109(6 Pt 1), 2817-32.
- Badano JL, Mitsuma N, Beales PL, Katsanis N** (2006). The ciliopathies: an emerging class of human genetic disorders. *Annu Rev Genomics Hum Genet* 7, 125-148.
- Badano JL, Teslovich TM, Katsanis N** (2005). The centrosome in human genetic disease. *Nat Rev Genet* 6(3), 194-205.
- Balklava Z, Pant S, Fares H, Grant BD** (2007). Genome-wide analysis identifies a general requirement for polarity proteins in endocytic traffic. *Nat Cell Biol* 9, 1066–1073.

- Barroso M, Sztul ES** (1994). Basolateral to apical transcytosis in polarized cells is indirect and involves BFA and trimeric G proteinsensitive passage through the apical endosome. *J Cell Biol* 124, 83–100.
- Bartoloni L, Blouin JL, Maiti AK, Sainsbury A, Rossier C, Gehrig C, She JX, Marron MP, Lander ES, Meeks M, Chung E, Armengot M, Jorissen M, Scott HS, Delozier-Blanchet CD, Gardiner RM, Antonarakis SE** (2001). Axonemal beta heavy chain dynein DNAH9: cDNA sequence, genomic structure, and investigation of its role in primary ciliary dyskinesia. *Genomics* 72(1), 21-33.
- Beaudoin AR, Grondin G** (1991). Shedding of vesicular material from the cell surface of eukaryotic cells: different cellular phenomena. *Biochim Biophys Acta* 1071(3), 203-19.
- Beech PL, Pagh-Roehl K, Noda Y, Hirokawa N, Burnside B, Rosenbaum JL** (1996). Localization of kinesin superfamily proteins to the connecting cilium of fish photoreceptors. *J Cell Sci* 109 (Pt 4), 889-97.
- Benesh AE, Nambiar R, McConnell RE, Mao S, Tabb DL, Tyska MJ** (2010). Differential Localization and Dynamics of Class I Myosins in the Enterocyte Microvillus, *Mol Biol Cell* 21, 970–978.
- Benninga MA, Lilien M, de Koning TJ, Duran M, Versteegh FG, Goldschmeding R, Poll-The BT** (2007). Renal Fanconi syndrome with ultrastructural defects in lysinuric protein intolerance, *J Inherit Metab Dis* 30, 402-403.
- Berryman M, Franck Z, Bretscher A** (1993). Ezrin is concentrated in the apical microvilli of a wide variety of epithelial cells whereas moesin is found primarily in endothelial cells. *J Cell Sci* 105 (Pt 4), 1025-43.
- Biemesderfer D, DeGray B, Aronson PS** (2001). Active (9.6 s) and inactive (21 s) oligomers of NHE-3 in microdomains of the renal brush border. *J Biol Chem* 276, 10161–10167.
- Bijlsma PB, Kiliaan AJ, Scholten G, van der Wal A, Heyman M, Heymans HS, Groot JA, Taminiou JA** (2000). Increased paracellular macromolecular transport and subnormal glucose uptake in duodenal biopsies of patients with microvillus inclusion disease. Comparisons to other chronic diarrhea patients and to nondiarrhea patients. *Ann N Y Acad Sci* 915, 267-269.
- Black BL, Yoneyama Y, Moog F** (1980). Microvillous membrane vesicle accumulation in media during culture of intestine of chick embryo. *Biochim Biophys Acta* 601(2), 343-8.
- Blankenship JT, Fuller MT, Zallen JA** (2007). The *Drosophila* homolog of the Exo84 exocyst subunit promotes apical epithelial identity. *J Cell Sci* 120, 3099–3110.
- Bligh EG, Dyer WJ** (1959). A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37(8), 911-7.
-

## References

---

- Bonilha VL, Rayborn ME, Saotome I, McClatchey AI, Hollyfield JG** (2006). Microvilli defects in retinas of ezrin knockout mice. *Exp Eye Res* 82(4), 720-9.
- Boron WF, McCormick WC, Roos A** (1981). pH regulation in barnacle muscle fibers: dependence on extracellular sodium and bicarbonate. *Am J Physiol* 240(1), C80-9.
- Brailov I, Bancila M, Brisogueil M, Miquel M, Hamon M, Verge D** (2000). Localization of 5-HT<sub>6</sub> receptors at the plasma membrane of neuronal cilia in the rat brain. *Brain Res* 872, 271–275.
- Breitfeld PP, Harris JM, Mostov KE** (1989). Postendocytotic sorting of the ligand for the polymeric immunoglobulin receptor in Madin-Darby canine kidney cells. *J Cell Biol* 109, 475–486.
- Bretscher A, Chambers D, Nguyen R, Reczek D** (2000). ERM-Merlin and EBP50 protein families in plasma membrane organization and function. *Annu Rev Cell Dev Biol* 16, 113-43.
- Bretscher A, Edwards K, Fehon RG** (2002). ERM proteins and merlin: integrators at the cell cortex. *Nat Rev Mol Cell Biol* 3(8), 586-99.
- Brignoni M, Podesta EJ, Mele P, Rodriguez ML, Vega-Salas DE, Salas PJ** (1993). Exocytosis of vacuolar apical compartment (VAC) in Madin-Darby canine kidney epithelial cells: cAMP is involved as second messenger. *Exp Cell Res* 205(1), 171-8.
- Brown DA, Rose JK** (1992). Sorting of GPI-anchored proteins to glycolipid enriched membrane subdomains during transport to the apical cell surface. *Cell* 68, 533-544.
- Brown FD, Rozelle AL, Yin HL, Balla T, Donaldson JG** (2001). Phosphatidylinositol 4,5-bisphosphate and Arf6-regulated membrane traffic. *J Cell Biol* 154(5), 1007-17.
- Brown PS, Wang E, Aroeti B, Chapin SJ, Mostov KE, Dunn KW** (2000). Definition of distinct compartments in polarized Madin-Darby canine kidney (MDCK) cells for membrane volume sorting, polarized sorting and apical recycling. *Traffic* 1, 124–140.
- Bubb MR, Spector I, Beyer BB, Fosen KM** (2000). Effects of jasplakinolide on the kinetics of actin polymerization. An explanation for certain in vivo observations. *J Biol Chem* 275(7), 5163-70.
- Buchman AL, Moukarzel A, Ament ME, Gornbein J, Goodson B, Carlson C, Hawkins RA** (1993). Serious renal impairment is associated with long-term parenteral nutrition. *J Parenter Enteral Nutr* 17, 438-444.
- Burkhardt JK, Echeverri CJ, Nilsson T, Vallee RB** (1997). Overexpression of the dynamitin (p50) subunit of the dynactin complex disrupts dynein-dependent maintenance of membrane organelle distribution. *J Cell Biol* 139(2), 469-84.

- Caballero T, Aneiros J, Lopez-Caballero J, Gomez-Morales M, Nogales F** (1985). Fibrolamellar hepatocellular carcinoma. An immunohistochemical and ultrastructural study. *Histopathology* 9(4), 445-56.
- Campbell K, Knust E, Skaer H** (2009) Crumbs stabilises epithelial polarity during tissue remodelling. *J Cell Sci* 122, 2604–2612.
- Cancino J, Torrealba C, Soza A, Yuseff MI, Gravotta D, Henklein P, Rodriguez-Boulan E, González A** (2007). Antibody to AP1B adaptor blocks biosynthetic and recycling routes of basolateral proteins at recycling endosomes. *Mol Biol Cell* 18, 4872–4884.
- Casanova JE, Wang X, Kumar R, Bhartur SG, Navarre J, Woodrum JE, Altschuler Y, Ray GS, Goldenring JR** (1999). Association of Rab25 and Rab11a with the apical recycling system of polarized Madin-Darby canine kidney cells. *Mol Biol Cell* 10, 47–56.
- Cecchini M, Houdusse A, Karplus M** (2008). Allosteric communication in myosin V: from small conformational changes to large directed movements. *PLoS Comput Biol* 4, e1000129.
- Charras GT, Yarrow JC, Horton MA, Mahadevan L, Mitchison TJ** (2005). Non-equilibration of hydrostatic pressure in blebbing cells. *Nature* 435(7040), 365-9.
- Charras GT** (2008). A short history of blebbing. *Microsc* 231(3), 466-78.
- Chen W, Feng Y, Chen D, Wandinger-Ness A** (1998). Rab11 is required for trans-golgi network-toplasma membrane transport and a preferential target for GDP dissociation inhibitor. *Mol Biol Cell* 9, 3241–3257.
- Cheong KH, Zacchetti D, Schneeberger EE, Simons K** (1999). VIP17/MAL, a lipid raft-associated protein, is involved in apical transport in MDCK cells. *Proc Natl Acad Sci USA* 96, 6241-6248.
- Cole NB, Sciaky N, Marotta A, Song J, Lippincott-Schwartz J** (1996). Golgi dispersal during microtubule disruption: regeneration of Golgi stacks at peripheral endoplasmic reticulum exit sites. *Mol Biol Cell* 7(4), 631-50.
- Coluccio LM** (1997). Myosin I. *Am J Physiol* 273(2 Pt 1), C347-59.
- Corbeil D, Marzesco AM, Fargeas CA, Huttner WB** (2010). Prominin-1: a distinct cholesterol-binding membrane protein and the organisation of the apical plasma membrane of epithelial cells. *Subcell Biochem* 51, 399-423.
- Coureur PD, Sweeney HL, Houdusse A** (2004). Three myosin V structures delineate essential features of chemo-mechanical transduction. *EMBO J* 23, 4527-37.
- Cramm-Behrens CI, Dienst M, Jacob R** (2008). Apical cargo traverses endosomal compartments on the passage to the cell surface. *Traffic* 9, 2206-30.

## References

---

- Cramm-Behrens CI, Dienst M, Jacob R** (2008). Apical cargo traverses endosomal compartments on the passage to the cell surface. *Traffic* 9, 2206–2220.
- Cresawn KO, Potter BA, Oztan A, Guerriero CJ, Ihrke G, Goldenring JR, Apodaca G, Weisz OA** (2007). Differential involvement of endocytic compartments in the biosynthetic traffic of apical proteins. *EMBO J* 26, 3737–3748.
- Cullinane AR, Straatman-Iwanowska A, Zaucker A, Wakabayashi Y, Bruce CK, Luo G, Rahman F, Gürakan F, Utine E, Ozkan TB, Denecke J, Vukovic J, Di Rocco M, Mandel H, Cangul H, Matthews RP, Thomas SG, Rappoport JZ, Arias IM, Wolburg H, Knisely AS, Kelly DA, Müller F, Maher ER, Gissen P** (2010). Mutations in VIPAR cause an arthrogyposis, renal dysfunction and cholestasis syndrome phenotype with defects in epithelial polarization. *Nat Genet* 42, 303–312.
- Cutz E, Rhoads JM, Drumm B, Sherman PM, Durie PR, Forstner GG** (1989). Microvillus inclusion disease: an inherited defect of brush-border assembly and differentiation. *N Engl J Med* 320, 646–51.
- Cutz E, Sherman PM, Davidson GP** (1997). Enteropathies associated with protracted diarrhea of infancy: Clinicopathological features, cellular and molecular mechanisms. *Pediatric Pathology & Laboratory Medicine* 17, 335–368.
- D'Angelo A, Franco B** (2009). The dynamic cilium in human diseases. *PathoGenetics* 2, 3.
- D'Angelo G, Polishchuk E, Di Tullio G, Santoro M, Di Campli A, Godi A, West G, Bielawski J, Chuang CC, van der Spoel AC, Platt FM, Hannun YA, Polishchuk R, Mattjus P, De Matteis MA** (2007). Glycosphingolipid synthesis requires FAPP2 transfer of glucosylceramide. *Nature* 449(7158), 62–7.
- Danielsen EM, Hansen GH** (2003). Lipid rafts in epithelial brush borders: atypical membrane microdomains with specialized functions. *Biochim Biophys Acta* 1617, 1–9.
- Danielsen EM, van Deurs B, Hansen GH** (2003). “Nonclassical” secretion of annexin A2 to the luminal side of the enterocyte brush border. *Membr Biochem* 42, 14670–14676.
- Daugherty BL, Mateescu M, Patel AS, Wade K, Kimura S, Gonzales LW, Guttentag S, Ballard PL, Koval M** (2004). Developmental regulation of claudin localization by fetal alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 287(6), L1266–73.
- Davidson GP, Cutz E, Hamilton JR, Gall DG** (1978). Familial enteropathy: a syndrome of protracted diarrhea from birth, failure to thrive, and hypoplastic villus atrophy. *Gastroenterology* 75, 783–790.
- de Graaf P, Zwart WT, van Dijken RA, Deneka M, Schulz TK, Geijsen N, Coffier PJ, Gadella BM, Verkleij AJ, van der Sluijs P, van Bergen en Henegouwen PM** (2004). Phosphatidylinositol 4-kinase beta is critical for functional association of rab11 with the Golgi complex. *Mol Biol Cell* 15, 2038–2047.

- Deborde S, Perret E, Gravotta D, Deora A, Salvarezza S, Schreiner R, Rodriguez-Boulan E** (2008). Clathrin is a key regulator of basolateral polarity. *Nature* 452, 719–723.
- Deretic D, Papermaster DS** (1993). Rab6 is associated with a compartment that transports rhodopsin from the trans-Golgi to the site of rod outer segment disk formation in frog retinal photoreceptors. *J Cell Sci* 106 ( Pt 3), 803-13.
- DeSchryver-Kecsckemeti K, Eliakim R, Carroll S, Stenson WF, Moxley MA, Alpers DH** (1989). Intestinal surfactant-like material. A novel secretory product of the rat enterocyte. *J Clin Invest* 84(4), 1355-61.
- Desclozeaux M, Venturato J, Wylie FG, Kay JG, Joseph SR, Le HT, Stow JL** (2008). Active Rab11 and functional recycling endosome are required for E-cadherin trafficking and lumen formation during epithelial morphogenesis. *Am J Physiol Cell Physiol* 295, C545–C556.
- Diao A, Rahman D, Pappin DJ, Lucocq J, Lowe M** (2003). The coiled-coil membrane protein golgin-84 is a novel rab effector required for Golgi ribbon formation. *J Cell Biol* 160(2), 201-12.
- Diaz F, Gravotta D, Deora A, Schreiner R, Schoggins J, Falck-Pedersen E, Rodriguez-Boulan E** (2009). Clathrin adaptor AP1B controls adenovirus infectivity of epithelial cells. *Proc Natl Acad Sci USA* 106, 11143–11148.
- Doi Y, Itoh M, Yonemura S, Ishihara S, Takano H, Noda T, Tsukita S** (1999). Normal development of mice and unimpaired cell adhesion/cell motility/actin-based cytoskeleton without compensatory up-regulation of ezrin or radixin in moesin gene knockout. *J Biol Chem* 274(4), 2315-21.
- Donaudy F, Ferrara A, Esposito L, Hertzano R, Ben-David O, Bell RE, Melchionda S, Zelante L, Avraham KB, Gasparini P** (2003). Multiple mutations of MYO1A, a cochlear-expressed gene, in sensorineural hearing loss. *Am J Hum Genet* 72(6), 1571-7.
- Dumont RA, Zhao YD, Holt JR, Bähler M, Gillespie PG** (2002). Myosin-I isozymes in neonatal rodent auditory and vestibular epithelia. *J Assoc Res Otolaryngol* 3(4), 375-89.
- Eliakim R, DeSchryver-Kecsckemeti K, Nogee L, Stenson WF, Alpers DH** (1989). Isolation and characterization of a small intestinal surfactant-like particle containing alkaline phosphatase and other digestive enzymes. *J Biol Chem* 264(34), 20614-9.
- Erickson RP, Larson-Thomé K, Valenzuela RK, Whitaker SE, Shub MD** (2008). Navajo microvillous inclusion disease is due to a mutation in MYO5B. *Am J Med Genet Part A* 146A, 3117-19.
- Fan J, Hu Z, Zeng L, Lu W, Tang X, Zhang J, Li T** (2008). Golgi apparatus and neurodegenerative diseases. *Int J Dev Neurosci* 26(6), 523-34.



## References

---

- Fan S, Hurd TW, Liu CJ, Straight SW, Weimbs T, Hurd EA, Domino SE, Margolis B** (2004). Polarity proteins control ciliogenesis via kinesin motor interactions. *Curr Biol* 14(16), 1451-61.
- Farquhar MG, Palade GE** (1963). Junctional complexes in various epithelia. *J Cell Biol* 17, 375-412.
- Farr GA, Hull M, Mellman I, Caplan MJ** (2009). Membrane proteins follow multiple pathways to the basolateral cell surface in polarized epithelial cells. *J Cell Biol* 186, 269–282.
- Ferrari A, Veligodskiy A, Berge U, Lucas MS, Kroschewski R** (2008). ROCK-mediated contractility, tight junctions and channels contribute to the conversion of a preapical patch into apical surface during isochoric lumen initiation. *J Cell Sci* 121(Pt 21), 3649-63.
- Fields IC, Shteyn E, Pypaert M, Proux-Gillardeaux V, Kang RS, Galli T, Fölsch H** (2007). v-SNARE cellubrevin is required for basolateral sorting of AP-1B-dependent cargo in polarized epithelial cells. *J Cell Biol* 177, 477–488.
- Fields IC, King SM, Shteyn E, Kang RS, Fölsch H** (2010). Phosphatidylinositol 3,4,5-trisphosphate localization in recycling endosomes is necessary for AP-1B-dependent sorting in polarized epithelial cells. *Mol Biol Cell* 21, 95–105.
- Fliegauf M, Benzing T, Omran H** (2007). When cilia go bad: cilia defects and ciliopathies. *Nat Rev Mol Cell Biol* 8, 880-893.
- Fölsch H, Ohno H, Bonifacino JS, Mellman I** (1999). A novel clathrin adaptor complex mediates basolateral targeting in polarized epithelial cells. *Cell* 99, 189–198.
- Forte JG, Black JA, Forte TM, Machen TE, Wolosin JM** (1981). Ultrastructural changes related to functional activity in gastric oxyntic cells. *Am J Physiol* 241(5), G349-58.
- Forte TM, Machen TE, Forte JG** (1977). Ultrastructural changes in oxyntic cells associated with secretory function: a membrane-recycling hypothesis. *Gastroenterology* 73(4 Pt 2), 941-55.
- Fujita Y, Ohama E, Takatama M, Al-Sarraj S, Okamoto K** (2006). Fragmentation of Golgi apparatus of nigral neurons with alpha-synuclein-positive inclusions in patients with Parkinson's disease. *Acta Neuropatho* 112(3), 261-5.
- Fujiwara T, Oda K, Yokota S, Takatsuki A, Ikehara Y** (1988). Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *J Biol Chem* 263(34), 18545-52.
- Futter CE, Gibson A, Allchin EH, Maxwell S, Ruddock LJ, Odorizzi G, Domingo D, Trowbridge IS, Hopkins CR** (1998). In polarized MDCK cells basolateral vesicles arise

- from clathrin-gamma-adaptin-coated domains on endosomal tubules. *J Cell Biol* 141, 611–623.
- Gagescu R, Demaux N, Parton RG, Hunziker W, Huber LA, Gruenberg J** (2000). The recycling endosome of Madin-Darby canine kidney cells is a mildly acidic compartment rich in raft components. *Mol Biol Cell* 11, 2775–2791.
- Gambarara M, Diamanti A, Ferretti F, Papadatou B, Knafelz D, Pirotto A, Castro M** (2003). Intractable diarrhea of infancy with congenital intestinal mucosa abnormalities: outcome of four cases. *Transplant Proc* 35, 3052-3053.
- Gan Y, McGraw TE, Rodriguez-Boulan E** (2002). The epithelial-specific adaptor AP1B mediates post-endocytic recycling to the basolateral membrane. *Nat Cell Biol* 4, 605–609.
- Gassama-Diagne A, Yu W, ter Beest M, Martin-Belmonte F, Kierbel A, Engel J, Mostov K** (2006). Phosphatidylinositol-3,4,5-trisphosphate regulates the formation of the basolateral plasma membrane in epithelial cells. *Nat. Cell. Biol* 8, 963–970.
- Gerdes JM, Davis EE, Katsanis N** (2009). The Vertebrate Primary Cilium in Development, Homeostasis, and Disease. *Cell* 137, 32-45.
- Gherman A, Davis EE, Katsanis N** (2006). The ciliary proteome database: an integrated community resource for the genetic and functional dissection of cilia. *Nat Genet* 38(9), 961-2.
- Gibson A, Futter CE, Maxwell S, Allchin EH, Shipman M, Kraehenbuhl JP, Domingo D, Odorizzi G, Trowbridge IS, Hopkins CR** (1998). Sorting mechanisms regulating membrane protein traffic in the apical transcytotic pathway of polarized MDCK cells. *J Cell Biol* 143, 81–94.
- Giepmans BN, van Ijzendoorn SC** (2009). Epithelial cell-cell junctions and plasma membrane domains. *Biochim Biophys Acta* 1788(4), 820-31.
- Gilbert T, Rodriguez-Boulan E** (1991). Induction of vacuolar apical compartments in the Caco-2 intestinal epithelial cell line. *J Cell Sci* 100 (Pt 3), 451-8.
- Gissen P, Johnson CA, Morgan NV, Stapelbroek JM, Forshew T, Cooper WN, McKiernan PJ, Klomp LW, Morris AA, Wraith JE, McClean P, Lynch SA, Thompson RJ, Lo B, Quarrell OW, Di Rocco M, Trembath RC, Mandel H, Wali S, Karet FE, Knisely AS, Houwen RH, Kelly DA, Maher ER** (2004). Mutations in VPS33B, encoding a regulator of SNARE-dependent membrane fusion, cause arthrogyposis-renal dysfunction-cholestasis (ARC) syndrome. *Nat Genet* 36, 400–404.
- Godi A, Di Campli A, Konstantakopoulos A, Di Tullio G, Alessi DR, Kular GS, Daniele T, Marra P, Lucocq JM, De Matteis MA** (2004). FAPPs control Golgi-to-cell-surface membrane traffic by binding to ARF and PtdIns(4)P. *Nat Cell Biol* 6(5), 393-404.
-

## References

---

- Golachowska MR, Hoekstra D, van IJendoorn SC** (2010). Recycling endosomes in apical plasma membrane domain formation and epithelial cell polarity. *Trends Cell Biol* 20(10), 618-26.
- Goldenring JR, Smith J, Vaughan HD, Cameron P, Hawkins W, Navarre J** (1996). Rab11 is an apically located small GTP-binding protein in epithelial tissues. *Am J Physiol* 270(3 Pt 1), G515-25.
- Gonatas NK, Stieber A, Gonatas JO** (2006). Fragmentation of the Golgi apparatus in neurodegenerative diseases and cell death. *J Neurol Sci* 246(1-2), 21-30.
- Gorelik J, Shevchuk AI, Frolenkov GI, Diakonov IA, Lab MJ, Kros CJ, Richardson GP, Vodyanoy I, Edwards CR, Klenerman D, Korchev YE** (2003). Dynamic assembly of surface structures in living cells. *Proc Natl Acad Sci USA* 100(10), 5819-22.
- Goulet O, Ruemmele F, Lacaille F, Colomb V** (2004). Irreversible intestinal failure. *J Pediatr Gastroenterol Nutr* 38, 250-69.
- Gravotta D, Deora A, Perret E, Oyanadel C, Soza A, Schreiner R, Gonzalez A, Rodriguez-Boulan E** (2007). AP1B sorts basolateral proteins in recycling and biosynthetic routes of MDCK cells. *Proc Natl Acad Sci USA* 104, 1564-1569.
- Grindstaff KK, Bacallao RL, Nelson WJ** (1998). Apiconuclear organization of microtubules does not specify protein delivery from the trans-Golgi network to different membrane domains in polarized epithelial cells. *Mol Biol Cell* 9(3), 685-99.
- Groisman GM, Amar M, Livne E** (2002). CD10: A valuable tool for the light microscopic diagnosis of microvillous inclusion disease (familial microvillous atrophy). *Am J Surg Pathol* 26(7), 902-907.
- Gu F, Gruenberg** (2000). ARF1 regulates pH-dependent COP functions in the early endocytic pathway. *J Biol Chem* 275, 8154-8160.
- Halbhuber KJ, Schulze M, Rhode H, Bublitz R, Feuerstein H, Walter M, Linss W, Meyer HW, Horn A** (1994). Is the brush border membrane of the intestinal mucosa a generator of "chymosomes"? *Cell Mol Biol (Noisy-le-grand)* 40(8), 1077-96.
- Hall J, Hall A, Pursifull N, Barbar E** (2008). Differences in dynamic structure of LC8 monomer, dimer, and dimer-peptide complexes. *Biochemistry* 47(46), 11940-52.
- Handel M, Schulz S, Stanarius A, Schreff M, Erdtmann-Vourliotis M, Schmidt H, Wolf G, Holtt V** (1999). Selective targeting of somatostatin receptor 3 to neuronal cilia. *Neuroscience* 89, 909-926.
- Hannan LA, Lisanti MP, Rodriguez-Boulan E, Edidin M** (1993). Correctly sorted molecules of a GPI-anchored protein are clustered and immobile when they arrive at the apical surface of MDCK cells. *J Cell Biol* 120, 353-358.

- Hanono A, Garbett D, Reczek D, Chambers DN, Bretscher A** (2006). EPI64 regulates microvillar subdomains and structure. *J Cell Biol* 175(5), 803-13.
- Hansen GH, Pedersen J, Niels-Christiansen LL, Immerdal L, Danielsen EM** (2003). Deep-apical tubules: dynamic lipid-raft microdomains in the brush-border region of enterocytes. *Biochem J* 373, 125-132.
- Helander HF, Hirschowitz BI** (1972). Quantitative ultrastructural studies on gastric parietal cells. *Gastroenterology* 63(6), 951-61.
- Heller B, Adu-Gyamfi E, Smith-Kinnaman W, Babbey C, Vora M, Xue Y, Bittman R, Stahelin RV, Wells CD** (2010). Amot recognizes a juxtannuclear endocytic recycling compartment via a novel lipid binding domain. *J Biol Chem* 285, 12308-12320.
- Hehny H, Xu W, Chen JL, Stamnes M** (2010). Cdc42 regulates microtubule-dependent Golgi positioning. *Traffic* 11(8), 1067-78.
- Henkel JR** (1999). Selective perturbation of early endosome and/or trans-Golgi network pH but not lysosome pH by dose-dependent expression of influenza M2 protein. *J Biol Chem* 274, 9854-9860.
- Herrema H, Czajkowska D, Théard D, van der Wouden JM, Kalicharan D, Zolghadr B, Hoekstra D, van Ijzendoorn SC** (2006). Rho kinase, myosin-II, and p42/44 MAPK control extracellular matrix-mediated apical bile canalicular lumen morphogenesis in HepG2 cells. *Mol Biol Cell* 17(7), 3291-303.
- Hirao M, Sato N, Kondo T, Yonemura S, Monden M, Sasaki T, Takai Y, Tsukita S, Tsukita S** (1996). Regulation mechanism of ERM (ezrin/radixin/moesin) protein/plasma membrane association: possible involvement of phosphatidylinositol turnover and Rho-dependent signaling pathway. *J Cell Biol* 135(1), 37-51.
- Hoekstra D, Maier O, van der Wouden JM, Slimane TA, van IJzendoorn SC** (2003). Membrane dynamics and cell polarity: the role of sphingolipids. *J Lipid Res* 44(5), 869-77.
- Hoekstra D, Tyteca D, van IJzendoorn SC** (2004). The subapical compartment: a traffic center in membrane polarity development. *J Cell Sci* 117, 2183-92.
- Holmes KC, Schroder RR, Sweeney HL, Houdusse A** (2004). The structure of the rigor complex and its implications for the power stroke. *Philos Trans R Soc Lond B Biol Sci* 359, 1819-28.
- Holzer P** (2003). Acid-sensitive ion channels in gastrointestinal function. *Curr Opin Pharmacol* 3(6), 618-25.

## References

---

- Hong DH, Pawlyk B, Sokolov M, Strissel KJ, Yang J, Tulloch B, Wright AF, Arshavsky VY, Li T** (2003). RPGR isoforms in photoreceptor connecting cilia and the transitional zone of motile cilia. *Invest Ophthalmol Vis Sci* 44, 2413-2421.
- Huang S, Carlson GM, Cheung WY** (1994). Calmodulin-dependent enzymes undergo a protein-induced conformational change that is associated with their interactions with calmodulin. *J Biol Chem* 269, 7631-7638.
- Huang S, Cheung WY** (1994).  $H^+$  is involved in the activation of calcineurin by calmodulin. *J Biol Chem* 269, 22067-22074.
- Iancu TC, Mahajnah M, Manov I, Shaoul R** (2007). Microvillous inclusion disease: ultrastructural variability. *Ultrastruct Pathol* 31, 173-88.
- In JG, Tuma PL** (2010). MAL2 selectively regulates polymeric IgA receptor delivery from the Golgi to the plasma membrane in WIF-B cells. *Traffic* 11, 1056-1066.
- Ingraffea J, Reczek D, Bretscher A** (2002). Distinct cell type-specific expression of scaffolding proteins EBP50 and E3KARP: EBP50 is generally expressed with ezrin in specific epithelia, whereas E3KARP is not. *Eur J Cell Biol* 81(2), 61-8.
- Isik N, Brzostowski JA, Jin T** (2008). An Elmo-like protein associated with myosin-II restricts spurious F-actin events to coordinate phagocytosis and chemotaxis. *Dev Cell* 15(4), 590-602.
- Isken O, Maquat LE** (2000). The multiple lives of NMD factors: balancing roles in gene and genome regulation. *Nat Rev Genet* 9, 699-712.
- Jacobs LR** (1983). Biochemical and ultrastructural characterization of the molecular topography of the rat intestinal microvillous membrane. Asymmetric distribution of hydrophilic groups and anionic binding sites. *Gastroenterology* 85(1), 46-54.
- Janich P, Corbeil D** (2007). GM1 and GM3 gangliosides highlight distinct lipid microdomains within the apical domain of epithelial cells. *FEBS Lett* 581, 1783-1787.
- Jerdeva GV, Tesar DB, Huey-Tubman KE, Ladinsky MS, Fraser SE, Bjorkman PJ** (2010). Comparison of FcRn- and pIgR-mediated transport in MDCK cells by fluorescence confocal microscopy. *Traffic* 11, 1205-1220.
- Johnson D, Nehrke K** (2010). Mitochondrial fragmentation leads to intracellular acidification in *Caenorhabditis elegans* and mammalian cells. *Mol Biol Cell* 21(13), 2191-201.
- Kagitani K, Yamamoto T, Miki K, Matsumoto S, Shima M, Tajiri H, Harada T, Okada S** (1998). Hypophosphatemic rickets accompanying congenital microvillous atrophy. *J Bone Miner Res* 13, 1946-1952.

- Kawano N, Ohba Y, Nagashima K** (2000). Eosinophilic inclusions in ependymoma represent microlumina: a light and electron microscopic study. *Acta Neuropathol* 99(2), 214-8.
- Keller LC, Marshall WF** (2008). Isolation and proteomic analysis of *Chlamydomonas* centrioles. *Methods Mol Biol* 432, 289-300.
- Kikuchi S, Hata M, Fukumoto K, Yamane Y, Matsui T, Tamura A, Yonemura S, Yamagishi H, Keppler D, Tsukita S, Tsukita S** (2002). Radixin deficiency causes conjugated hyperbilirubinemia with loss of Mrp2 from bile canalicular membranes. *Nat Genet* 31(3), 320-5.
- Kimura N, Imamura O, Ono F, Terao K** (2007). Aging attenuates dynactin-dynein interaction: down-regulation of dynein causes accumulation of endogenous tau and amyloid precursor protein in human neuroblastoma cells. *J Neurosci Res* 85(13), 2909-16.
- Kimura N, Inoue M, Okabayashi S, Ono F, Negishi T** (2009). Dynein dysfunction induces endocytic pathology accompanied by an increase in Rab GTPases: a potential mechanism underlying age-dependent endocytic dysfunction. *J Biol Chem* 284(45), 31291-302.
- Kinjo M, Yohena C, Kunishima N** (2003). Intracytoplasmic lumina in medullary carcinoma of the thyroid gland. Report of a case with cytologic and immunocytochemical features. *Acta Cytol* 47(4), 663-7.
- Kipp H, Arias IM** (2000). Intracellular trafficking and regulation of canalicular ATP-binding cassette transporters. *Semin Liver Dis* 20(3), 339-51.
- Kipp H, Pichetshote N, Arias IM** (2001). Transporters on demand: intrahepatic pools of canalicular ATP binding cassette transporters in rat liver. *J Biol Chem* 276(10), 7218-24.
- Klemm RW, Ejsing CS, Surma MA, Kaiser HJ, Gerl MJ, Sampaio JL, de Robillard Q, Ferguson C, Proszynski TJ, Shevchenko A et al.** (2009). Segregation of sphingolipids and sterols during formation of secretory vesicles at the trans-Golgi network. *J Cell Biol* 185, 601-612.
- Klopfenstein DR, Tomishige M, Stuurman N, Vale RD** (2002). Role of phosphatidylinositol(4,5)bisphosphate organization in membrane transport by the Unc104 kinesin motor. *Cell* 109, 347-358.
- Knott AB, Perkins G, Schwarzenbacher R, Bossy-Wetzel E** (2008). Mitochondrial fragmentation in neurodegeneration. *Nat Rev Neurosci* 9(7), 505-18.
- Koegler E, Bonnon C, Waldmeier L, Mitrovic S, Halbeisen R, Hauri HP** (2010). p28, a novel ERGIC/cis Golgi protein, required for Golgi ribbon formation. *Traffic* 11(1), 70-89.

## References

---

- Kok JW, Babia T, Hoekstra D** (1991). Sorting of sphingolipids in the endocytic pathway of HT29 cells. *J Cell Biol* 114(2), 231-9.
- Lapierre LA, Goldenring JR** (2005). Interactions of myosin vb with rab11 family members and cargoes traversing the plasma membrane recycling system. *Methods Enzymol* 403, 715-23.
- Lapierre LA, Kumar R, Hales CM, Navarre J, Bhartur SG, Burnette JO, Provance DW Jr, Mercer JA, Bähler M, Goldenring JR** (2001). Myosin Vb is associated with plasma membrane recycling systems. *Mol Biol Cell* 12, 1843-57.
- Lauverjat M, Hadj Aissa A, Vanhems P, Boulétreau P, Fouque D, Chambrier C** (2006). Chronic dehydration may impair renal function in patients with chronic intestinal failure on long-term parenteral nutrition. *Clin Nutr* 25, 75-81.
- Lee HJ, Khoshaghideh F, Lee S, Lee SJ** (2006). Impairment of microtubule-dependent trafficking by overexpression of alpha-synuclein. *Eur J Neurosci* 24(11), 3153-62.
- Li BX, Satoh AK, Ready DF** (2007). Myosin V, Rab11, and dRip11 direct apical secretion and cellular morphogenesis in developing *Drosophila* photoreceptors. *J Cell Biol* 177, 659-669.
- Lin F, Hiesberger T, Cordes K, Sinclair AM, Goldstein LS, Somlo S, Igarashi P** (2003). Kidney-specific inactivation of the KIF3A subunit of kinesin-II inhibits renal ciliogenesis and produces polycystic kidney disease. *Proc Natl Acad Sci USA* 100(9), 5286-91.
- Ling K, Bairstow SF, Carbonara C, Turbin DA, Huntsman DG, Anderson RA** (2007). Type I gamma phosphatidylinositol phosphate kinase modulates adherens junction and E-cadherin trafficking via a direct interaction with mu 1B adaptin. *J Cell Biol* 176, 343-353.
- Lippincott-Schwartz J, Yuan LC, Bonifacino JS, Klausner RD** (1989). Rapid redistribution of Golgi proteins into the ER in cells treated with brefeldin A: evidence for membrane cycling from Golgi to ER. *Cell* 56(5), 801-13.
- Lisanti M, Sargiacomo M, Graeve L, Saltiel A, Rodriguez-Boulan E** (1988). Polarized apical distribution of glycosyl phosphatidylinositol anchored proteins in a renal epithelial line. *Proc Natl Acad Sci USA* 85, 9557-9561.
- Lock JG, Stow JL** (2005). Rab11 in recycling endosomes regulates the sorting and basolateral transport of E-cadherin. *Mol Biol Cell* 16, 1744-1755.
- Loomis PA, Zheng L, Sekerková G, Changyaleket B, Mugnaini E, Bartles JR** (2003). Espin cross-links cause the elongation of microvillus-type parallel actin bundles in vivo. *J Cell Biol* 163(5), 1045-55.

- Louvard D, Kedinger M, Hauri HP** (1992). The differentiating intestinal epithelial cell: establishment and maintenance of functions through interactions between cellular structures. *Annu Rev Cell Biol* 8, 157-95.
- Low SH, Miura M, Roche PA, Valdez AC, Mostov KE, Weimbs T** (2000). Intracellular redirection of plasma membrane trafficking after loss of epithelial cell polarity. *Mol Biol Cell* 11(9), 3045-60.
- Maier O, Hoekstra D** (2003). Trans-Golgi network and subapical compartment of HepG2 cells display different properties in sorting and exiting of sphingolipids. *J Biol Chem* 278, 164-173.
- Manderson AP, Kay JG, Hammond LA, Brown DL, Stow JL** (2007). Subcompartments of the macrophage recycling endosome direct the differential secretion of IL-6 and TNF alpha. *J Cell Biol* 178, 57-69.
- Marra P, Salvatore L, Mironov A Jr, Di Campi A, Di Tullio G, Trucco A, Beznoussenko G, Mironov A, De Matteis MA** (2007). The biogenesis of the Golgi ribbon: the roles of membrane input from the ER and of GM130. *Mol Biol Cell* 18(5), 1595-608.
- Marshansky V, Ausiello DA, Brown D** (2002). Physiological importance of endosomal acidification: potential role in proximal tubulopathies. *Curr Opin Nephrol Hypertens* 11, 527-537.
- Martin-Belmonte F, Gassama A, Datta A, Yu W, Rescher U, Gerke V, Mostov K** (2007). PTEN-mediated apical segregation of phosphoinositides controls epithelial morphogenesis through Cdc42. *Cell* 128, 383-397.
- Martin-Belmonte F, Mostov K** (2008). Regulation of cell polarity during epithelial morphogenesis. *Curr Opin Cell Biol* 20(2), 227-34.
- Martin-Belmonte F, Yu W, Rodríguez-Fraticelli AE, Ewald AJ, Werb Z, Alonso MA, Mostov K** (2008). Cell-polarity dynamics controls the mechanism of lumen formation in epithelial morphogenesis. *Curr Biol* 18(7), 507-13.
- Martin-Belmonte M, Mostov K** (2007) Phosphoinositides control epithelial development. *Cell Cycle* 6, 1957-1961.
- Marzesco AM, Janich P, Wilsch-Brauninger M, Dubreuil V, Langenfeld K, Corbeil D, Huttner WB** (2005). Release of extracellular membrane particles carrying the stem cell marker prominin-1 (CD133) from neural progenitors and other epithelial cells. *J Cell Sci* 118, 2849-2858.
- Massol RH, Larsen JE, Kirchhausen T** (2005). Possible role of deep tubular invaginations of the plasma membrane in MHC-I trafficking, *Experimental Cell Research* 306, 142-149.



## References

---

- Matsui T, Maeda M, Doi Y, Yonemura S, Amano M, Kaibuchi K, Tsukita S, Tsukita S** (1998). Rho-kinase phosphorylates COOH-terminal threonines of ezrin/radixin/moesin (ERM) proteins and regulates their head-to-tail association. *J Cell Biol* 140(3), 647-57.
- Matter K, Mellman I.** (1994). Mechanisms of cell polarity: sorting and transport in epithelial cells. *Curr Opin Cell Biol* 6, 545-554.
- Mattila PE, Kinlough CL, Bruns JR, Weisz OA, Hughey RP** (2009). MUC1 traverses apical recycling endosomes along the biosynthetic pathway in polarized MDCK cells. *Biol Chem* 390, 551–556.
- Maudsley S, Zamah AM, Rahman N, Blitzer JT, Luttrell LM, Lefkowitz RJ, Hall RA** (2000). Platelet-derived growth factor receptor association with Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor potentiates receptor activity. *Mol Cell Biol* 20(22), 8352-63.
- McClatchey AI** (2003). Merlin and ERM proteins: unappreciated roles in cancer development? *Nat Rev Cancer* 3(11), 877-83.
- McConnell RE, Tyska MJ** (2007). Myosin-1a powers the sliding of apical membrane along microvillar actin bundles. *J Cell Biol* 177, 671–681.
- McDonough AA, Biemesderfer D** (2003). Does membrane trafficking play a role in regulating the sodium/hydrogen exchanger isoform 3 in the proximal tubule? *Curr Opin Nephrol Hypertens* 12, 533–541.
- Meder D, Shevchenko A, Simons K, Fullekrug J** (2005) Gp135/podocalyxin and NHERF-2 participate in the formation of a preapical domain during polarization of MDCK cells. *J Cell Biol* 168, 303–313.
- Meiss HK, Green RF, Rodriguez-Boulan E** (1982). Lectin-resistant mutants of polarized epithelial cells. *Mol Cell Biol* 2, 1287-1294.
- Mellman I, Warren G,** (2000). The road taken: past and future foundations of membrane traffic, *Cell* 100(1), 99-112.
- Mellman I, Nelson WJ** (2008). Coordinated protein sorting, targeting and distribution in polarized cells. *Nat Rev Mol Cell Biol* 9, 833–845.
- Michail S, Collins JF, Xu H, Kaufman S, Vanderhoof J, Ghishan FK** (1998). Abnormal expression of brush-border membrane transporters in the duodenal mucosa of two patients with microvillus inclusion disease. *J Pediatr Gastroenterol Nutr* 27, 536-42.
- Michaud EJ, Yoder BK** (2006). The primary cilium in cell signaling and cancer. *Cancer Res* 66(13), 6463-7.

- Miller PM, Folkmann AW, Maia AR, Efimova N, Efimov A, Kaverina I** (2009). Golgi-derived CLASP-dependent microtubules control Golgi organization and polarized trafficking in motile cells. *Nat Cell Biol* 11(9), 1069-80.
- Mohan PM, Barve M, Chatterjee A, Hosur RV** (2006). pH driven conformational dynamics and dimer-to-monomer transition in DLC8. *Protein Sci* 15(2), 335-42.
- Mooseker MS, Bonder EM, Grimwade BG, Howe CL, Keller TC 3rd, Wasserman RH, Wharton KA** (1982). Regulation of contractility, cytoskeletal structure, and filament assembly in the brush border of intestinal epithelial cells. *Cold Spring Harb Symp Quant Biol* 46 Pt 2, 855-70.
- Moritz OL, Tam BM, Hurd LL, Peränen J, Deretic D, Papermaster DS** (2001). Mutant rab8 Impairs docking and fusion of rhodopsin-bearing post-Golgi membranes and causes cell death of transgenic Xenopus rods. *Mol Biol Cell* 12(8), 2341-51.
- Mostov K, Su T, ter Beest M** (2003). Polarized epithelial membrane traffic: conservation and plasticity. *Nat Cell Biol* 5, 287-293.
- Müller T, Hess MW, Schiefermeier N, Pfaller K, Ebner HL, Heinz-Erian P, Ponstingl H, Partsch J, Röllinghoff B, Köhler H, Berger T, Lenhartz H, Schlenck B, Houwen RJ, Taylor CJ, Zoller H, Lechner S, Goulet O, Utermann G, Ruemmele FM, Huber LA, Janecke AR** (2008). MYO5B mutations cause microvillus inclusion disease and disrupt epithelial cell polarity. *Nat Genet* 40(10), 1163-5.
- Murthy A, Gonzalez-Agosti C, Cordero E, Pinney D, Candia C, Solomon F, Gusella J, Ramesh V** (1998). NHE-RF, a regulatory cofactor for Na<sup>+</sup>-H<sup>+</sup> exchange, is a common interactor for merlin and ERM (MERM) proteins. *J Biol Chem* 273(3), 1273-6.
- Nachury MV, Loktev AV, Zhang Q, Westlake CJ, Peranen J, Merdes A, Slusarski DC, Scheller RH, Bazan JF, Sheffield VC, Jackson PK** (2007). A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. *Cell* 129, 1201-1213.
- Nakagomi S, Barsoum MJ, Bossy-Wetzel E, Sütterlin C, Malhotra V, Lipton SA** (2008). A Golgi fragmentation pathway in neurodegeneration. *Neurobiol Dis* 29(2), 221-31.
- Nambiar R, McConnell RE, Tyska MJ** (2009). Control of cell membrane tension by myosin-I. *Proc Natl Acad Sci USA* 106(29), 11972-7.
- Nedvetsky PI, Stefan E, Frische S, Santamaria K, Wiesner B, Valenti G, Hammer JA 3rd, Nielsen S, Goldenring JR, Rosenthal W, Klusmann E** (2007). A Role of myosin Vb and Rab11-FIP2 in the aquaporin-2 shuttle. *Traffic* 8, 110-23.
- Nielsen JS, Graves ML, Chelliah S, Vogl AW, Roskelley CD, McNagny KM** (2007). The CD34-related molecule podocalyxin is a potent inducer of microvillus formation. *PLoS One* 2(2), e237.

## References

---

- Niggli V, Rossy J** (2008). Ezrin/radixin/moesin: versatile controllers of signaling molecules and of the cortical cytoskeleton. *Int J Biochem Cell Biol* 40(3), 344-9.
- Nürnberg J, Bacallao RL, Phillips CL** (2002). Inversin forms a complex with catenins and N-cadherin in polarized epithelial cells. *Mol Biol Cell* 13(9), 3096-106.
- Nyarko A, Cochrun L, Norwood S, Pursifull N, Voth A, Barbar E** (2005). Ionization of His 55 at the dimer interface of dynein light-chain LC8 is coupled to dimer dissociation. *Biochemistry* 44(43), 14248-55.
- Odorizzi G, Pearse A, Domingo D, Trowbridge IS, Hopkins CR** (1996). Apical and basolateral endosomes of MDCK cells are interconnected and contain a polarized sorting mechanism. *J Cell Biol* 135, 139-152.
- Ohgaki R, Matsushita M, Kanazawa H, Ogihara S, Hoekstra D, van Ijzendoorn SC** (2010). The Na<sup>+</sup>/H<sup>+</sup> exchanger NHE6 in the endosomal recycling system is involved in the development of apical bile canalicular surface domains in HepG2 cells. *Mol Biol Cell* 21, 1293-1304.
- Orzech E, Cohen S, Weiss A, Aroeti B** (2000). Interactions between the exocytic and endocytic pathways in polarized Madin-Darby canine kidney cells. *J Biol Chem* 275, 15207-15219.
- Ostedgaard LS, Randak C, Rokhlina T, Karp P, Vermeer D, Ashbourne Excoffon KJ, Welsh MJ** (2003). Effects of C-terminal deletions on cystic fibrosis transmembrane conductance regulator function in cystic fibrosis airway epithelia. *Proc Natl Acad Sci USA* 100(4), 1937-42.
- Oztan A, Silvis M, Weisz OA, Bradbury NA, Hsu SC, Goldenring JR, Yeaman C, Apodaca G** (2007). Exocyst requirement for endocytic traffic directed toward the apical and basolateral poles of polarized MDCK cells. *Mol Biol Cell* 18, 3978-3992.
- Paladino S, Sarnataro D, Pillich R, Tivodar S, Nitsch L, Zurzolo C** (2004). Protein oligomerization modulates raft partitioning and apical sorting of GPI-anchored proteins. *J Cell Biol* 167, 699-709.
- Pazour GJ, Witman GB** (2003). The vertebrate primary cilium is a sensory organelle. *Curr Opin Cell Biol* 15, 105-10.
- Peden AA, Schonteich E, Chun J, Junutula JR, Scheller RH, Prekeris R** (2004). The RCP-Rab11 complex regulates endocytic protein sorting. *Mol Biol Cell* 15, 3530-41.
- Phillips AD, Jenkins P, Raafat F, Walker-Smith JA** (1985). Congenital microvillous atrophy: specific diagnostic features. *Arch Dis Child* 60(2), 135-40.
- Phillips AD, Schmitz J** (1992). Familial Microvillous Atrophy - A Clinicopathological Survey of 23 Cases. *J Pediatr Gastroenterol Nutr* 14, 380-396.

- Phillips MJ, Oshio C, Miyairi M, Katz H, Smith CR** (1982). A study of bile canalicular contractions in isolated hepatocytes. *Hepatology* 2(6), 763-8.
- Pirchl M, Marksteiner J, Humpel C** (2006). Effects of acidosis on brain capillary endothelial cells and cholinergic neurons: relevance to vascular dementia and Alzheimer's disease. *Neurol Res* 28(6), 657-64.
- Potter BA, Hughey RP, Weisz OA** (2006). Role of N- and O-glycans in polarized biosynthetic sorting. *Am J Physiol Cell Physiol* 290, C1-C10.
- Praetorius HA, Spring KR** (2003). The renal cell primary cilium functions as a flow sensor. *Curr Opin Nephrol Hypertens* 12(5), 517-20.
- Praetorius HA, Spring KR** (2005). A physiological view of the primary cilium. *Annu Rev Physiol* 67, 515-29.
- Prekeris R** (2003). Rabs, Rips, FIPs, and endocytic membrane traffic. *Scientific World Journal* 3, 870-80.
- Prekeris R, Klumperman J, Scheller RH** (2000). A Rab11/Rip11 protein complex regulates apical membrane trafficking via recycling endosomes. *Mol Cell* 6, 1437-48.
- Preuss D, Mulholland J, Franzusoff A, Segev N, Botstein D** (1992). Characterization of the *Saccharomyces* Golgi complex through the cell cycle by immunoelectron microscopy. *Mol Biol Cell* 3(7), 789-803.
- Puertollano R, Alonso MA** (1999). MAL, an integral element of the apical sorting machinery, is an itinerant protein that cycles between the trans-Golgi network and the plasma membrane. *Mol Biol Cell* 10, 3435-3447.
- Puertollano R, Martin-Belmonte F, Millan J, de Marco MC, Albar JP, Kremer L, Alonso MA** (1999). The MAL proteolipid is necessary for normal apical transport and accurate sorting of the influenza virus hemagglutinin in Madin-Darby canine kidney cells. *J Cell Biol* 145, 141-151.
- Puthenveedu MA, Bachert C, Puri S, Lanni F, Linstedt AD** (2006). GM130 and GRASP65-dependent lateral cisternal fusion allows uniform Golgi-enzyme distribution. *Nat Cell Biol* 8(3), 238-48.
- Quintyne NJ, Gill SR, Eckley DM, Crego CL, Compton DA, Schroer TA** (1999). Dynactin is required for microtubule anchoring at centrosomes. *J Cell Biol* 147(2), 321-34.
- Reczek D, Berryman M, Bretscher A** (1997). Identification of EBP50: A PDZ-containing phosphoprotein that associates with members of the ezrin-radixin-moesin family. *J Cell Biol* 139(1), 169-79.

## References

---

- Reinshagen K, Naim HY, Zimmer KP** (2002). Autophagocytosis of the apical membrane in microvillus inclusion disease. *Gut* 51(4), 514-21.
- Reinshagen K, Naim H, Heusipp G, Zimmer KP** (2006). Pathophysiology in Microvillus inclusion disease. *Z Gastroenterol* 44(8), 667-71.
- Remy L** (1986). The intracellular lumen: origin, role and implications of a cytoplasmic neostructure. *Biol Cell* 56(2), 97-105.
- Riquier AD, Lee DH, McDonough AA** (2009). Renal NHE-3 and NaPi2 partition into distinct membrane domains. *Am J Physiol Cell Physiol* 296(4), C900-10.
- Riquier-Brison AD, Leong PK, Pihakaski-Maunsbach K, McDonough AA** (2010). Angiotensin II stimulates trafficking of NHE-3, NaPi2, and associated proteins into the proximal tubule microvilli. *Am J Physiol Renal Physiol* 298(1), F177-86.
- Rodriguez-Boulant E, Kreitzer G, Müsch A** (2005). Organization of vesicular trafficking in epithelia. *Nat Rev Mol Cell Biol* 6, 233–247.
- Roeth JF, Sawyer JK, Wilner DA, Peifer M** (2009). Rab11 helps maintain apical crumbs and adherens junctions in the Drosophila embryonic ectoderm. *PLoS One* 4, e7634.
- Roland JT, Kenworthy AK, Peranen J, Caplan S, Goldenring JR** (2007). Myosin Vb interacts with Rab8a on a tubular network containing EHD1 and EHD3. *Mol Biol Cell* 18, 2828-37.
- Roland JT, Lapierre LA, Goldenring JR** (2009). Alternative splicing in class V myosins determines association with Rab10. *J. Biol. Chem.* 284, 1213–1223.
- Rosenbaum JL, Witman GB** (2002). Intraflagellar transport. *Nat Rev Mol Cell Biol* 3(11), 813-25.
- Royle SJ, Lagnado L** (2003). Endocytosis at the synaptic terminal. *J Physiol* 553, 345–355.
- Rozelle AL, Machesky LM, Yamamoto M, Driessens MH, Insall RH, Roth MG, Luby-Phelps K, Marriott G, Hall A, Yin HL** (2000) Phosphatidylinositol 4,5-bisphosphate induces actin-based movement of raft-enriched vesicles through WASP-Arp2/3. *Curr Biol* 10, 311–320.
- Rozen S, Skaletsky H** (2000). Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol* 132, 365-86.
- Ruemmele FM, Jan D, Lacaille F, Cézard JP, Canioni D, Phillips AD, Peuchmaur M, Aigrain Y, Brousse N, Schmitz J, Revillon Y, Goulet O** (2004). New perspectives for children with microvillous inclusion disease: early small bowel transplantation. *Transplantation* 77, 1024-1028.

- Ruemmele FM, Müller T, Schiefermeier N, Ebner HL, Lechner S, Pfaller K, Thöni CE, Goulet O, Lacaille F, Schmitz J, Colomb V, Sauvat F, Revillon Y, Canioni D, Brousse N, de Saint-Basile G, Lefebvre J, Heinz-Erian P, Enninger A, Utermann G, Hess MW, Janecke AR, Huber LA** (2010). Loss-of-function of MYO5B is the main cause of microvillus inclusion disease: 15 novel mutations and a CaCo-2 RNAi cell model. *Hum Mutat* 31(5), 544-51.
- Ruemmele FM, Schmitz J, Goulet O** (2006). Microvillous inclusion disease (microvillous atrophy). *Orphanet J Rare Dis* 1, 22.
- Sandvig K, Olsnes S, Petersen OW, Van Deurs B** (1987). Acidification of the cytosol inhibits endocytosis from coated pits. *J Biol Chem* 105, 679-689.
- Saotome I, Curto M, McClatchey AI** (2004). Ezrin is essential for epithelial organization and villus morphogenesis in the developing intestine. *Dev Cell* 6, 855-64.
- Satir P, Christensen ST** (2007). Overview of structure and function of mammalian cilia. *Annu Rev Physiol* 69:377-400.
- Sato T, Mushiake S, Kato Y, Sato K, Sato M, Takeda N, Ozono K, Miki K, Kubo Y, Tsuji A, Harada R, Harada A** (2007). The Rab8 GTPase regulates apical protein localization in intestinal cells. *Nature* 448, 366-9.
- Sato K, Hattori S, Irie S, Kawashima S** (2003). Spike formation by fibroblasts adhering to fibrillar collagen I gel. *Cell Struct Funct* 28(4), 229-41.
- Sato S, Masuda T, Oikawa H, Satodate R, Suzuki K, Sato S, Suzuki A, Monma N** (1997). Bile canaliculi-like lumina in fibrolamellar carcinoma of the liver: a light- and electron-microscopic study and three-dimensional examination of serial sections. *Pathol Int* 47(11), 763-8.
- Satoh AK, O'Tousa JE, Ozaki K, Ready DF** (2005) Rab11 mediates post-Golgi trafficking of rhodopsin to the photosensitive apical membrane of *Drosophila* photoreceptors. *Development* 132, 1487-1497.
- Schlüter MA, Pfarr CS, Pieczynski J, Whiteman EL, Hurd TW, Fan S, Liu CJ, Margolis B** (2009). Trafficking of Crumbs3 during cytokinesis is crucial for lumen formation. *Mol Biol Cell* 20, 4652-4663.
- Scholey JM** (2008). Intraflagellar transport motors in cilia: moving along the cell's antenna. *J Cell Biol* 180(1), 23-9.
- Schuck S, Simons K** (2004). Polarized sorting in epithelial cells: raft clustering and the biogenesis of the apical membrane. *J Cell Sci* 117, 5955-5964.

## References

---

- Shaw RJ, Henry M, Solomon F, Jacks T** (1998). RhoA-dependent phosphorylation and relocalization of ERM proteins into apical membrane/actin protrusions in fibroblasts. *Mol Biol Cell* 9(2), 403-19.
- Sheff DR, Daro EA, Hull M, Mellman I** (1999). The receptor recycling pathway contains two distinct populations of early endosomes with different sorting functions. *J Cell Biol* 145, 123–213.
- Sherman PM, Mitchell DJ, Cutz E** (2004). Neonatal enteropathies: defining the causes of protracted diarrhea of infancy. *J Pediatr Gastroenterol Nutr* 38, 16-26.
- Shivas JM, Morrison HA, Bilder D, Skop AR** (2010). Polarity and endocytosis: reciprocal regulation. *Trends Cell Biol* 20, 445–452.
- Short DB, Trotter KW, Reczek D, Kreda SM, Bretscher A, Boucher RC, Stutts MJ, Milgram SL** (1998). An apical PDZ protein anchors the cystic fibrosis transmembrane conductance regulator to the cytoskeleton. *J Biol Chem* 273(31), 19797-801.
- Silvis MR, Bertrand CA, Ameen N, Golin-Bisello F, Butterworth MB, Frizzell RA, Bradbury NA** (2009). Rab11b regulates the apical recycling of the cystic fibrosis transmembrane conductance regulator in polarized intestinal epithelial cells. *Mol Biol Cell* 20, 2337–2350.
- Skowron JF, Bement WM, Mooseker MS** (1998). Human brush border myosin-I and myosin-Ic expression in human intestine and Caco-2BBE cells. *Cell Motil Cytoskeleton* 41(4), 308-24.
- Skowron JF, Mooseker MS** (1999). Cloning and characterization of mouse brush border myosin-I in adult and embryonic intestine. *J Exp Zool* 283(3), 242-57.
- Slimane TA, Trugnan G, van IJendoorn SC, Hoekstra D** (2003). Raft-mediated trafficking of apical resident proteins occurs in both direct and transcytotic pathways in polarized hepatic cells: role of distinct lipid microdomains. *Mol Biol Cell* 14(2), 611-24.
- Smythe E** (2002). Direct interactions between rab GTPases and cargo. *Mol Cell* 9, 205–206.
- Spektor A, Tsang WY, Khoo D, Dynlacht BD** (2007). Cep97 and CP110 suppress a cilia assembly program. *Cell* 130(4), 678-90.
- Spinak M, Dembitzer HM** (1978). Unusual intracytoplasmic inclusions in a case of idiopathic scleritis. *Ophthalmologica* 177(1), 53-60.
- Stechly L, Morelle W, Dessen AF, André S, Grard G, Trinel D, Dejonghe MJ, Leteurtre E, Drobecq H, Trugnan G, Gabius HJ, Huet G** (2009). Galectin-4-regulated delivery of glycoproteins to the brush border membrane of enterocyte-like cells. *Traffic* 10(4), 438-50.

- Stock Ch, Schwab A** (2009). Protons make tumor cells move like clockwork. *Eur J Physiol* 458, 981-992.
- St-Pierre MV, Dufour JF, Arias IM** (1997). Disruption of actin organization by cytochalasin D does not impair biliary secretion of organic anions in the rat. *Hepatology* 25(4), 970-5.
- Sung CH, Tai AW** (2000). Rhodopsin trafficking and its role in retinal dystrophies. *Int Rev Cytol* 195, 215-67.
- Sütterlin C, Colanzi A** (2010). The Golgi and the centrosome: building a functional partnership. *J Cell Biol* 188(5):621-8.
- Swiatecka-Urban A, Duhaime M, Coutermarsh B, Karlson KH, Collawn J, Milewski M, Cutting GR, Guggino WB, Langford G, Stanton BA** (2002). PDZ domain interaction controls the endocytic recycling of the cystic fibrosis transmembrane conductance regulator. *J Biol Chem* 277(42), 40099-105.
- Swiatecka-Urban A, Talebian L, Kanno E, Moreau-Marquis S, Coutermarsh B, Hansen K, Karlson KH, Barnaby R, Cheney RE, Langford GM, Fukuda M, Stanton BA** (2007). Myosin Vb is required for trafficking of the cystic fibrosis transmembrane conductance regulator in Rab11a-specific apical recycling endosomes in polarized human airway epithelial cells. *J Biol Chem* 282(32), 23725-36.
- Szperl AM, Golachowska MR, Bruinenberg M, Prekeris R, Thunnissen AM, Karrenbeld A, Dijkstra G, Hoekstra D, Mercer D, Ksiazyk J, Wijmenga C, Wapenaar MC, Rings EH, van Ijzendoorn SC** (2011). Functional characterization of mutations in the Myosin Vb gene associated with Microvillus Inclusion Disease. *J Pediatr Gastroenterol Nutr* 52(3), 307-313.
- Tang S, Liao JC, Dunn AR, Altman RB, Spudich JA, Schmidt JP** (2007). Predicting allosteric communication in myosin via a pathway of conserved residues. *J Mol Biol* 373, 1361-73.
- Taulman PD, Haycraft CJ, Balkovetz DF, Yoder BK** (2001). Polaris, a protein involved in left-right axis patterning, localizes to basal bodies and cilia. *Mol Biol Cell* 12(3), 589-99.
- ten Klooster JP, Jansen M, Yuan J, Oorschot V, Begthel H, Di Giacomo V, Colland F, de Koning J, Maurice MM, Hornbeck P, Clevers H** (2009). Mst4 and Ezrin Induce Brush Borders Downstream of the Lkb1/Strad/Mo25 Polarization Complex. *Dev Cell* 16, 551-562.
- Teuling E, van Dis V, Wulf PS, Haasdijk ED, Akhmanova A, Hoogenraad CC, Jaarsma D** (2008). A novel mouse model with impaired dynein/dynactin function develops amyotrophic lateral sclerosis (ALS)-like features in motor neurons and improves lifespan in SOD1-ALS mice. *Hum Mol Genet* 17(18), 2849-62.



## References

---

- Théard D, Steiner M, Kalicharan D, Hoekstra D, van IJzendoorn SC** (2007). Cell polarity development and protein trafficking in hepatocytes lacking E-cadherin/beta-catenin-based adherens junctions. *Mol Biol Cell* 18(6), 2313-21.
- Thompson A, Nessler R, Wisco D, Anderson E, Winckler B, Sheff D** (2007). Recycling endosomes of polarized epithelial cells actively sort apical and basolateral cargos into separate subdomains. *Mol Biol Cell* 18, 2687-2697.
- Tsukita S, Furuse M, Itoh M** (2001). Multifunctional strands in tight junctions. *Nat Rev Mol Cell Biol* 2(4), 285-93.
- Tyska MJ, Mackey AT, Huang J-D, Copeland NG, Jenkins NA, Mooseker MS** (2005). Myosin-1a is critical for normal brush border structure and composition. *Mol Biol Cell* 16, 2443-2457.
- Tyska MJ, Mooseker MS** (2002). Myo1a (brush border myosin I) dynamics in the brush border of LLC-PK1-CL4 cells. *Biophys J* 82, 1869-1883.
- Tyska MJ, Nambiar R** (2010). Myosin-1a: A motor for microvillar membrane movement and mechanics. *Commun Integr Biol* 3(1), 64-6.
- Tzaban S, Massol RH, Yen E, Hamman W, Frank SR, Lapierre LA, Hansen SH, Goldenring JR, Blumberg RS, Lencer WI** (2009). The recycling and transcytotic pathways for IgG transport by FcRn are distinct and display an inherent polarity. *J Cell Biol* 185, 673-684.
- Ukarapol N, Chotinaruemol S, Lertprasertsuk N, Wongsawasdi L** (2001). Microvillus inclusion disease as a cause of severe protracted diarrhea in infants. *J. Med. Assoc. Thai.* 84(9), 1356-60.
- Utech M, Ivanov AI, Samarin SN, Bruewer M, Turner JR, Mrsny RJ, Parkos CA, Nusrat A** (2005). Mechanism of IFN-gamma-induced endocytosis of tight junction proteins: myosin-II-dependent vacuolarization of the apical plasma membrane. *Mol Biol Cell* 16(10), 5040-52.
- van der Wouden JM, Maier O, van IJzendoorn SC, Hoekstra D** (2003). Membrane dynamics and the regulation of epithelial cell polarity. *Int Rev Cytol* 226, 127-64.
- van IJzendoorn SC, Zegers MM, Kok JW, Hoekstra D** (1997). Segregation of glucosylceramide and sphingomyelin occurs in the apical to basolateral transcytotic route in HepG2 cells. *J Cell Biol* 137(2), 347-57.
- van IJzendoorn SC** (2006). Recycling endosomes. *J Cell Sci* 119, 1679-81.
- van IJzendoorn SC, Hoekstra D** (1998). (Glyco)sphingolipids are sorted in sub-apical compartments in HepG2 cells: a role for non-Golgi-related intracellular sites in the polarized distribution of (glyco)sphingolipids. *J Cell Biol* 142, 683-96.

- van IJzendoorn SC, Hoekstra D** (1999). The subapical compartment: a novel sorting centre? *Trends Cell Biol* 9(4), 144-9.
- van IJzendoorn SC, Hoekstra D** (2000). Polarized sphingolipid transport from the subapical compartment changes during cell polarity development. *Mol Biol Cell* 11, 1093-101.
- van IJzendoorn SC, Tuvim MJ, Weimbs T, Dickey BF, Mostov KE** (2002). Direct interaction between Rab3b and the polymeric immunoglobulin receptor controls ligand-stimulated transcytosis in epithelial cells. *Dev Cell* 2, 219–228.
- van IJzendoorn SC, Théard D, Van Der Wouden JM, Visser W, Wojtal KA, Hoekstra D** (2004). Oncostatin M-stimulated apical plasma membrane biogenesis requires p27(Kip1)-regulated cell cycle dynamics. *Mol Biol Cell* 15, 4105–4114.
- van Meer G, Simons K** (1986). The function of tight junctions in maintaining differences in lipid composition between the apical and the basolateral cell surface domains of MDCK cells. *EMBO J* 5, 1455–1464.
- van Meer G, Simons K** (1988). Lipid polarity and sorting in epithelial cells. *J Cell Biochem* 36, 51-58.
- van Niel G, Raposo G, Candalh C, Boussac M, Hershberg R, Cerf-Bensussan N, Heyman M** (2001). Intestinal epithelial cells secrete exosome-like vesicles. *Gastroenterology* 121(2), 337-49.
- Varma D, Dawn A, Ghosh-Roy A, Weil SJ, Ori-McKenney KM, Zhao Y, Keen J, Vallee RB, Williams JC** (2010). Development and application of in vivo molecular traps reveals that dynein light chain occupancy differentially affects dynein-mediated processes. *Proc Natl Acad Sci USA* 107(8), 3493-8.
- Vega-Salas DE, Salas PJ, Rodriguez-Boulan E** (1987). Modulation of the expression of an apical plasma membrane protein of Madin-Darby canine kidney epithelial cells: cell-cell interactions control the appearance of a novel intracellular storage compartment. *J Cell Biol* 104(5), 1249-59.
- Vega-Salas DE, Salas PJ, Rodriguez-Boulan E** (1988). Exocytosis of vacuolar apical compartment (VAC): a cell-cell contact controlled mechanism for the establishment of the apical plasma membrane domain in epithelial cells. *J Cell Biol* 107(5), 1717-28.
- Vega-Salas DE, San Martino JA, Salas PJ, Baldi A** (1993). Vacuolar apical compartment (VAC) in breast carcinoma cell lines (MCF-7 and T47D): failure of the cell-cell regulated exocytosis mechanism of apical membrane. *Differentiation* 54(2), 131-41.
- Vieira OV, Gaus K, Verkade P, Fullekrug J, Vaz WLC, Simons K** (2006). FAPP2, cilium formation, and compartmentalization of the apical membrane in polarized Madin–Darby canine kidney (MDCK) cells. *PNAS* 103 (49), 18556–18561.
-

## References

---

- Waharte F, Brown CM, Coscoy S, Coudrier E, Amblard F** (2005). A Two-Photon FRAP Analysis of the Cytoskeleton Dynamics in the Microvilli of Intestinal Cells, *Biophys J* 88, 1467–1478.
- Wakabayashi Y, Dutt P, Lippincott-Schwartz J, Arias IM** (2005). Rab11a and myosin Vb are required for bile canalicular formation in WIF-B9 cells. *Proc Natl Acad Sci USA* 102 (42), 15087-92.
- Wang W, Soroka CJ, Mennone A, Rahner C, Harry K, Pypaert M, Boyer JL** (2006). Radixin is required to maintain apical canalicular membrane structure and function in rat hepatocytes. *Gastroenterology* 131(3), 878-84.
- Wang E, Brown PS, Aroeti B, Chapin SJ, Mostov KE, Dunn KW** (2000) Apical and basolateral endocytic pathways of MDCK cells meet in acidic common endosomes distinct from a nearly-neutral apical recycling endosome. *Traffic* 1, 480–493.
- Watanabe S, Miyazaki A, Hirose M, Takeuchi M, Ohide H, Kitamura T, Ueno T, Kominami E, Sato N** (1991). Myosin in hepatocytes is essential for bile canalicular contraction. *Liver* 11(3), 185-9.
- Weisz OA, Rodriguez-Boulan E** (2009). Apical trafficking in epithelial cells: signals, clusters and motors, *Journal of Cell Science* 122, 4253-4266.
- Wells CD, Fawcett JP, Traweger A, Yamanaka Y, Goudreau M, Elder K, Kulkarni S, Gish G, Virag C, Lim C, Colwill K, Starostine A, Metalnikov P, Pawson T** (2006). A Rich1/Amot complex regulates the Cdc42 GTPase and apical-polarity proteins in epithelial cells. *Cell* 125, 535–548.
- Westerlund B, Slotte JP** (2009). How the molecular features of glycosphingolipids affect domain formation in fluid membranes. *Biochim Biophys Acta* 1788, 194–201.
- Wilson W, Scott RB, Pinto A, Robertson MA** (2001). Intractable diarrhea in a newborn infant: microvillous inclusion disease. *Can J Gastroenterol* 15(1) 61-4.
- Wojtal KA, Hoekstra D, van IJendoorn SC** (2007). Anchoring of protein kinase A-regulatory subunit IIalpha to subapically positioned centrosomes mediates apical bile canalicular lumen development in response to oncostatin M but not cAMP. *Mol Biol Cell* 18(7), 2745-54.
- Woolner S, Bement WM** (2009). Unconventional myosins acting unconventionally. *Trends Cell Biol* 6, 245-252.
- Xiao F, Weng J, Fan K, Wang W** (2010). Mechanism of Ser88 phosphorylation-induced dimer dissociation in dynein light chain LC8. *J Phys Chem B* 114(47), 15663-72.
- Yadav S, Puri S, Linstedt AD** (2009). A primary role for Golgi positioning in directed secretion, cell polarity, and wound healing. *Mol Biol Cell* 20(6), 1728-36.

- Yang LE, Maunsbach AB, Leong PK, McDonough AA** (2004). Differential traffic of proximal tubule Na<sup>+</sup> transporters during hypertension or PTH: NHE-3 to base of microvilli vs. NaPi2 to endosomes. *Am J Physiol Renal Physiol* 287, F896–F906.
- Yang LE, Maunsbach AB, Leong PK, McDonough AA** (2005). Redistribution of Myosin VI from Top to Base of Proximal Tubule Microvilli during Acute Hypertension. *J Am Soc Nephrol* 16, 2890–2896.
- Yap AS, Briher WM, Gumbiner BM** (1997). Molecular and functional analysis of cadherin-based adherens junctions. *Annu Rev Cell Dev Biol* 13, 119-46.
- Yeaman C, Le Gall AH, Baldwin AN, Monlauzeur L, Le Bivic A, Rodriguez-Boulan E** (1997). The O-glycosylated stalk domain is required for apical sorting of neurotrophin receptors in polarized MDCK cells. *J Cell Biol* 139, 929-940.
- Yoshimura S, Egerer J, Fuchs E, Haas AK, Barr FA** (2007). Functional dissection of Rab GTPases involved in primary cilium formation. *J Cell Biol* 178(3), 363-9.
- Yu J, Ustach C, Kim HR** (2003). Platelet-derived growth factor signaling and human cancer. *J Biochem Mol Biol* 36(1), 49-59.
- Yun CH, Lamprecht G, Forster DV, Sidor A** (1998). NHE-3 kinase A regulatory protein E3KARP binds the epithelial brush border Na<sup>+</sup>/H<sup>+</sup> exchanger NHE-3 and the cytoskeletal protein ezrin. *J Biol Chem* 273(40), 25856-63.
- Zegers MM, Hoekstra D** (1997). Sphingolipid transport to the apical plasma membrane domain in human hepatoma cells is controlled by PKC and PKA activity: a correlation with cell polarity in HepG2 cells. *J Cell Biol* 138(2), 307-21.
- Zegers MM, Hoekstra D** (1998). Mechanisms and functional features of polarized membrane traffic in epithelial and hepatic cells. *Biochem J* 336 (Pt 2), 257-69.
- Zegers MM, Zaal KJ, van IJendoorn SC, Klappe K, Hoekstra D** (1998). Actin filaments and microtubules are involved in different membrane traffic pathways that transport sphingolipids to the apical surface of polarized HepG2 cells. *Mol Biol Cell* 9(7), 1939-49.

