

University of Groningen

SecYEG

Lijcklama a Nijeholt, Jelger

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2012

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Lijcklama a Nijeholt, J. (2012). *SecYEG: Plug-and-Play!* s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

References

1. von Heijne G. 1990 The signal peptide. *J Membr Biol* **115**, 195-201
2. Beck K., Wu L.F., Brunner J. & Muller M. 2000 Discrimination between SRP- and SecA/SecB-dependent substrates involves selective recognition of nascent chains by SRP and trigger factor. *EMBO J* **19**, 134-143
3. Ullers R.S., Houben E.N., Brunner J., Oudega B., Harms N. & Luirink J. 2006 Sequence-specific interactions of nascent Escherichia coli polypeptides with trigger factor and signal recognition particle. *J Biol Chem* **281**, 13999-14005
4. Gilmore R., Blobel G. & Walter P. 1982 Protein translocation across the endoplasmic reticulum. I. Detection in the microsomal membrane of a receptor for the signal recognition particle. *J Cell Biol* **95**, 463-469
5. Walter P. & Blobel G. 1981 Translocation of proteins across the endoplasmic reticulum III. Signal recognition protein (SRP) causes signal sequence-dependent and site-specific arrest of chain elongation that is released by microsomal membranes. *J Cell Biol* **91**, 557-561
6. Randall L.L., Topping T.B., Hardy S.J., Pavlov M.Y., Freistroffer D.V. & Ehrenberg M. 1997 Binding of SecB to ribosome-bound polypeptides has the same characteristics as binding to full-length, denatured proteins. *Proc Natl Acad Sci U S A* **94**, 802-807
7. Brundage L., Hendrick J.P., Schiebel E., Driessen A.J. & Wickner W. 1990 The purified E. coli integral membrane protein SecY/E is sufficient for reconstitution of SecA-dependent precursor protein translocation. *Cell* **62**, 649-657
8. van den Berg B., Clemons W.M.Jr., Collinson I., Modis Y., Hartmann E., Harrison S.C. & Rapoport T.A. 2004 X-ray structure of a protein-conducting channel. *Nature* **427**, 36-44
9. Hanada M., Nishiyama K.I., Mizushima S. & Tokuda H. 1994 Reconstitution of an efficient protein translocation machinery comprising SecA and the three membrane proteins, SecY, SecE, and SecG (p12). *J Biol Chem* **269**, 23625-23631
10. Nishiyama K., Mizushima S. & Tokuda H. 1993 A novel membrane protein involved in protein translocation across the cytoplasmic membrane of Escherichia coli. *EMBO J* **12**, 3409-3415

11. Nishiyama K., Hanada M. & Tokuda H. 1994 Disruption of the gene encoding p12 (SecG) reveals the direct involvement and important function of SecG in the protein translocation of *Escherichia coli* at low temperature. *EMBO J* **13**, 3272-3277
12. Hanada M., Nishiyama K. & Tokuda H. 1996 SecG plays a critical role in protein translocation in the absence of the proton motive force as well as at low temperature. *FEBS Lett* **381**, 25-28
13. Bost S., Silva F., Rudaz C. & Belin D. 2000 Both transmembrane domains of SecG contribute to signal sequence recognition by the *Escherichia coli* protein export machinery. *Mol Microbiol* **38**, 575-587
14. Flower A.M., Hines L.L. & Pfennig P.L. 2000 SecG is an auxiliary component of the protein export apparatus of *Escherichia coli*. *Mol Gen Genet* **263**, 131-136
15. Flower A.M. 2001 SecG function and phospholipid metabolism in *Escherichia coli*. *J Bacteriol* **183**, 2006-2012
16. Homma T., Yoshihisa T. & Ito K. 1997 Subunit interactions in the *Escherichia coli* protein translocase: SecE and SecG associate independently with SecY. *FEBS Lett* **408**, 11-15
17. Mori H., Shimokawa N., Satoh Y. & Ito K. 2004 Mutational analysis of transmembrane regions 3 and 4 of SecY, a central component of protein translocase. *J Bacteriol* **186**, 3960-3969
18. Satoh Y., Matsumoto G., Mori H. & Ito K. 2003 Nearest neighbor analysis of the SecYEG complex. 1. Identification of a SecY-SecG interface. *Biochemistry* **42**, 7434-7441
19. van der Sluis E.O., Nouwen N. & Driessen A.J. 2002 SecY-SecY and SecY-SecG contacts revealed by site-specific crosslinking. *FEBS Lett* **527**, 159-165
20. Duong F. & Wickner W. 1997 Distinct catalytic roles of the SecYE, SecG and SecDFyajC subunits of preprotein translocase holoenzyme. *EMBO J* **16**, 2756-2768
21. Kato Y., Nishiyama K. & Tokuda H. 2003 Depletion of SecDF-YajC causes a decrease in the level of SecG: implication for their functional interaction. *FEBS Lett* **550**, 114-118
22. Matsumoto G., Mori H. & Ito K. 1998 Roles of SecG in ATP- and SecA-dependent protein translocation. *Proc Natl Acad Sci U S A* **95**, 13567-13572

-
23. Nagamori S., Nishiyama K. & Tokuda H. 2002 Membrane topology inversion of SecG detected by labeling with a membrane-impermeable sulfhydryl reagent that causes a close association of SecG with SecA. *J Biochem* **132**, 629-634
 24. Nishiyama K., Suzuki T. & Tokuda H. 1996 Inversion of the membrane topology of SecG coupled with SecA-dependent preprotein translocation. *Cell* **85**, 71-81
 25. Sugai R., Takemae K., Tokuda H. & Nishiyama K. 2007 Topology inversion of SecG is essential for cytosolic SecA-dependent stimulation of protein translocation. *J Biol Chem* **282**, 29540-29548
 26. van der Sluis E.O., van der Vries E., Berrelkamp G., Nouwen N. & Driessen A.J. 2006 Topologically fixed SecG is fully functional. *J Bacteriol* **188**, 1188-1190
 27. Mori H., Sugiyama H., Yamanaka M., Sato K., Tagaya M. & Mizushima S. 1998 Amino-terminal region of SecA is involved in the function of SecG for protein translocation into Escherichia coli membrane vesicles. *J Biochem* **124**, 122-129
 28. Suzuki H., Nishiyama K. & Tokuda H. 1998 Coupled structure changes of SecA and SecG revealed by the synthetic lethality of the secAcsR11 and delta secG::kan double mutant. *Mol Microbiol* **29**, 331-341
 29. Zimmer J., Nam Y. & Rapoport T.A. 2008 Structure of a complex of the ATPase SecA and the protein-translocation channel. *Nature* **455**, 936-943
 30. Das S. & Oliver D.B. 2011 Mapping of the SecA.SecY and SecA.SecG interfaces by site-directed in vivo photocross-linking. *J Biol Chem* **286**, 12371-12380
 31. Mori H. & Ito K. 2006 The long alpha-helix of SecA is important for the ATPase coupling of translocation. *J Biol Chem* **281**, 36249-36256
 32. Tokuda H., Akimaru J., Matsuyama S., Nishiyama K. & Mizushima S. 1991 Purification of SecE and reconstitution of SecE-dependent protein translocation activity. *FEBS Lett* **279**, 233-236
 33. Nishiyama K., Mizushima S. & Tokuda H. 1992 The carboxyl-terminal region of SecE interacts with SecY and is functional in the reconstitution of protein translocation activity in Escherichia coli. *J Biol Chem* **267**, 7170-7176

34. Schatz P.J., Bieker K.L., Ottemann K.M., Silhavy T.J. & Beckwith J. 1991 One of three transmembrane stretches is sufficient for the functioning of the SecE protein, a membrane component of the E. coli secretion machinery. *EMBO J* **10**, 1749-1757
35. Cao T.B. & Saier M.H.Jr. 2003 The general protein secretory pathway: phylogenetic analyses leading to evolutionary conclusions. *Biochim Biophys Acta* **1609**, 115-125
36. Murphy C.K. & Beckwith J. 1994 Residues essential for the function of SecE, a membrane component of the Escherichia coli secretion apparatus, are located in a conserved cytoplasmic region. *Proc Natl Acad Sci U S A* **91**, 2557-2561
37. Pohlschroder M., Murphy C. & Beckwith J. 1996 In vivo analyses of interactions between SecE and SecY, core components of the Escherichia coli protein translocation machinery. *J Biol Chem* **271**, 19908-19914
38. Kihara A., Akiyama Y. & Ito K. 1995 FtsH is required for proteolytic elimination of uncomplexed forms of SecY, an essential protein translocase subunit. *Proc Natl Acad Sci U S A* **92**, 4532-4536
39. Gumbart J. & Schulten K. 2008 The roles of pore ring and plug in the SecY protein-conducting channel. *J Gen Physiol* **132**, 709-719
40. Saparov S.M., Erlandson K., Cannon K., Schaletzky J., Schulman S., Rapoport T.A. & Pohl P. 2007 Determining the conductance of the SecY protein translocation channel for small molecules. *Mol Cell* **26**, 501-509
41. Park E. & Rapoport T.A. 2011 Preserving the membrane barrier for small molecules during bacterial protein translocation. *Nature* **473**, 239-242
42. Plath K., Mothes W., Wilkinson B.M., Stirling C.J. & Rapoport T.A. 1998 Signal sequence recognition in posttranslational protein transport across the yeast ER membrane. *Cell* **94**, 795-807
43. Plath K., Wilkinson B.M., Stirling C.J. & Rapoport T.A. 2004 Interactions between Sec complex and prepro-alpha-factor during posttranslational protein transport into the endoplasmic reticulum. *Mol Biol Cell* **15**, 1-10

-
44. Frauenfeld J., Gumbart J., van der Sluis E.O., Funes S., Gartmann M., Beatrix B., Mielke T., Berninghausen O., Becker T., Schulten K. & Beckmann R. 2011 Cryo-EM structure of the ribosome-SecYE complex in the membrane environment. *Nat Struct Mol Biol* **18**, 614-621
 45. Menetret J.F., Schaletzky J., Clemons W.M.Jr., Osborne A.R., Skanland S.S., Denison C., Gygi S.P., Kirkpatrick D.S., Park E., Ludtke S.J., Rapoport T.A. & Akey C.W. 2007 Ribosome binding of a single copy of the SecY complex: implications for protein translocation. *Mol Cell* **28**, 1083-1092
 46. van der Sluis E.O., Nouwen N., Koch J., de Keyzer J., van der Does C., Tampe R. & Driessen A.J. 2006 Identification of two interaction sites in SecY that are important for the functional interaction with SecA. *J Mol Biol* **361**, 839-849
 47. Tsukazaki T., Mori H., Fukai S., Ishitani R., Mori T., Dohmae N., Perederina A., Sugita Y., Vassylyev D.G., Ito K. & Nureki O. 2008 Conformational transition of Sec machinery inferred from bacterial SecYE structures. *Nature* **455**, 988-991
 48. Mori H. & Ito K. 2006 Different modes of SecY-SecA interactions revealed by site-directed in vivo photo-cross-linking. *Proc Natl Acad Sci U S A* **103**, 16159-16164
 49. Egea P.F. & Stroud R.M. 2010 Lateral opening of a translocon upon entry of protein suggests the mechanism of insertion into membranes. *Proc Natl Acad Sci U S A* **107**, 17182-17187
 50. Gumbart J. & Schulten K. 2007 Structural determinants of lateral gate opening in the protein translocon. *Biochemistry* **46**, 11147-11157
 51. du Plessis D.J., Berrelkamp G., Nouwen N. & Driessen A.J. 2009 The lateral gate of SecYEG opens during protein translocation. *J Biol Chem* **284**, 15805-15814
 52. Bonardi F., London G., Nouwen N., Feringa B.L. & Driessen A.J. 2010 Light-induced control of protein translocation by the SecYEG complex. *Angew Chem Int Ed Engl* **49**, 7234-7238
 53. Junne T., Schwede T., Goder V. & Spiess M. 2006 The plug domain of yeast Sec61p is important for efficient protein translocation, but is not essential for cell viability. *Mol Biol Cell* **17**, 4063-4068
 54. Li W., Schulman S., Boyd D., Erlandson K., Beckwith J. & Rapoport T.A. 2007 The plug domain of the SecY protein stabilizes the closed state of the translocation channel and maintains a membrane seal. *Mol Cell* **26**, 511-521

55. Maillard A.P., Lalani S., Silva F., Belin D. & Duong F. 2007 Deregulation of the SecYEG translocation channel upon removal of the plug domain. *J Biol Chem* **282**, 1281-1287
56. Harris C.R. & Silhavy T.J. 1999 Mapping an interface of SecY (PrlA) and SecE (PrlG) by using synthetic phenotypes and in vivo cross-linking. *J Bacteriol* **181**, 3438-3444
57. Tam P.C., Maillard A.P., Chan K.K. & Duong F. 2005 Investigating the SecY plug movement at the SecYEG translocation channel. *EMBO J* **24**, 3380-3388
58. Lycklama a Nijeholt J.A., Bulacu M., Marrink S.J. & Driessen A.J. 2010 Immobilization of the plug domain inside the SecY channel allows unrestricted protein translocation. *J Biol Chem* **285**, 23747-23754
59. Zhang B. & Miller T.F., III 2010 Hydrophobically stabilized open state for the lateral gate of the Sec translocon. *Proc Natl Acad Sci U S A* **107**, 5399-5404
60. Lycklama a Nijeholt J.A., Wu Z.C. & Driessen A.J. 2011 Conformational dynamics of the plug domain of the SecYEG protein-conducting channel. *J Biol Chem*
61. Bost S. & Belin D. 1995 A new genetic selection identifies essential residues in SecG, a component of the Escherichia coli protein export machinery. *EMBO J* **14**, 4412-4421
62. Michaelis S. & Beckwith J. 1982 Mechanism of incorporation of cell envelope proteins in Escherichia coli. *Annu Rev Microbiol* **36**, 435-465
63. Shiba K., Ito K. & Yura T. 1984 Mutation that suppresses the protein export defect of the secY mutation and causes cold-sensitive growth of Escherichia coli. *J Bacteriol* **160**, 696-701
64. Derman A.I., Puziss J.W., Bassford P.J.Jr. & Beckwith J. 1993 A signal sequence is not required for protein export in prlA mutants of Escherichia coli. *EMBO J* **12**, 879-888
65. Emr S.D., Hanley-Way S. & Silhavy T.J. 1981 Suppressor mutations that restore export of a protein with a defective signal sequence. *Cell* **23**, 79-88
66. Flower A.M., Doebele R.C. & Silhavy T.J. 1994 PrlA and PrlG suppressors reduce the requirement for signal sequence recognition. *J Bacteriol* **176**, 5607-5614
67. Smith M.A., Clemons W.M.Jr., DeMars C.J. & Flower A.M. 2005 Modeling the effects of prl mutations on the Escherichia coli SecY complex. *J Bacteriol* **187**, 6454-6465
68. Flower A.M., Osborne R.S. & Silhavy T.J. 1995 The allele-specific synthetic lethality of prlA-prlG double mutants predicts interactive domains of SecY and SecE. *EMBO J* **14**, 884-893

-
69. Osborne R.S. & Silhavy T.J. 1993 PrlA suppressor mutations cluster in regions corresponding to three distinct topological domains. *EMBO J* **12**, 3391-3398
70. van der Wolk J.P., Fekkes P., Boorsma A., Huie J.L., Silhavy T.J. & Driessen A.J. 1998 PrlA4 prevents the rejection of signal sequence defective preproteins by stabilizing the SecA-SecY interaction during the initiation of translocation. *EMBO J* **17**, 3631-3639
71. Bondar A.N., del Val C., Freitas J.A., Tobias D.J. & White S.H. 2010 Dynamics of SecY translocons with translocation-defective mutations. *Structure* **18**, 847-857
72. Nouwen N., de Kruijff B. & Tommassen J. 1996 prlA suppressors in Escherichia coli relieve the proton electrochemical gradient dependency of translocation of wild-type precursors. *Proc Natl Acad Sci U S A* **93**, 5953-5957
73. Bonardi F., Halza E., Walko M., du Plessis F., Nouwen N., Feringa B.L. & Driessen A.J. 2011 Probing the SecYEG translocation pore size with preproteins conjugated with sizable rigid spherical molecules. *Proc Natl Acad Sci U S A* **108**, 7775-7780
74. Gouridis G., Karamanou S., Gelis I., Kalodimos C.G. & Economou A. 2009 Signal peptides are allosteric activators of the protein translocase. *Nature* **462**, 363-367
75. Hartl F.U., Lecker S., Schiebel E., Hendrick J.P. & Wickner W. 1990 The binding cascade of SecB to SecA to SecY/E mediates preprotein targeting to the E. coli plasma membrane. *Cell* **63**, 269-279
76. Hendrick J.P. & Wickner W. 1991 SecA protein needs both acidic phospholipids and SecY/E protein for functional high-affinity binding to the Escherichia coli plasma membrane. *J Biol Chem* **266**, 24596-24600
77. Lill R., Dowhan W. & Wickner W. 1990 The ATPase activity of SecA is regulated by acidic phospholipids, SecY, and the leader and mature domains of precursor proteins. *Cell* **60**, 271-280
78. Osborne A.R., Clemons W.M.Jr. & Rapoport T.A. 2004 A large conformational change of the translocation ATPase SecA. *Proc Natl Acad Sci U S A* **101**, 10937-10942
79. Robson A., Booth A.E., Gold V.A., Clarke A.R. & Collinson I. 2007 A large conformational change couples the ATP binding site of SecA to the SecY protein channel. *J Mol Biol* **374**, 965-976

80. Zimmer J. & Rapoport T.A. 2009 Conformational flexibility and peptide interaction of the translocation ATPase SecA. *J Mol Biol* **394**, 606-612
81. Gelis I., Bonvin A.M., Keramisanou D., Koukaki M., Gouridis G., Karamanou S., Economou A. & Kalodimos C.G. 2007 Structural basis for signal-sequence recognition by the translocase motor SecA as determined by NMR. *Cell* **131**, 756-769
82. Musial-Siwiek M., Rusch S.L. & Kendall D.A. 2007 Selective photoaffinity labeling identifies the signal peptide binding domain on SecA. *J Mol Biol* **365**, 637-648
83. Papanikou E., Karamanou S., Baud C., Frank M., Sianidis G., Keramisanou D., Kalodimos C.G., Kuhn A. & Economou A. 2005 Identification of the preprotein binding domain of SecA. *J Biol Chem* **280**, 43209-43217
84. Fekkes P., van der Does C. & Driessen A.J. 1997 The molecular chaperone SecB is released from the carboxy-terminus of SecA during initiation of precursor protein translocation. *EMBO J* **16**, 6105-6113
85. Fekkes P., de Wit J.G., Boorsma A., Friesen R.H. & Driessen A.J. 1999 Zinc stabilizes the SecB binding site of SecA. *Biochemistry* **38**, 5111-5116
86. Zhou J. & Xu Z. 2003 Structural determinants of SecB recognition by SecA in bacterial protein translocation. *Nat Struct Biol* **10**, 942-947
87. Breukink E., Nouwen N., van Raalte A., Mizushima S., Tommassen J. & de Kruijff B. 1995 The C terminus of SecA is involved in both lipid binding and SecB binding. *J Biol Chem* **270**, 7902-7907
88. Karamanou S., Vrontou E., Sianidis G., Baud C., Roos T., Kuhn A., Politou A.S. & Economou A. 1999 A molecular switch in SecA protein couples ATP hydrolysis to protein translocation. *Mol Microbiol* **34**, 1133-1145
89. Fak J.J., Itkin A., Ciobanu D.D., Lin E.C., Song X.J., Chou Y.T., Gierasch L.M. & Hunt J.F. 2004 Nucleotide exchange from the high-affinity ATP-binding site in SecA is the rate-limiting step in the ATPase cycle of the soluble enzyme and occurs through a specialized conformational state. *Biochemistry* **43**, 7307-7327

-
90. Sianidis G., Karamanou S., Vrontou E., Boulias K., Repanas K., Kyrpides N., Politou A.S. & Economou A. 2001 Cross-talk between catalytic and regulatory elements in a DEAD motor domain is essential for SecA function. *EMBO J* **20**, 961-970
91. Akita M., Shinkai A., Matsuyama S. & Mizushima S. 1991 SecA, an essential component of the secretory machinery of Escherichia coli, exists as homodimer. *Biochem Biophys Res Commun* **174**, 211-216
92. Kusters I., van den Bogaart G., Kedrov A., Krasnikov V., Fulyani F., Poolman B. & Driessen A.J. 2011 Quaternary structure of SecA in solution and bound to SecYEG probed at the single molecule level. *Structure* **19**, 430-439
93. Bu Z., Wang L. & Kendall D.A. 2003 Nucleotide binding induces changes in the oligomeric state and conformation of Sec A in a lipid environment: a small-angle neutron-scattering study. *J Mol Biol* **332**, 23-30
94. Benach J., Chou Y.T., Fak J.J., Itkin A., Nicolae D.D., Smith P.C., Wittrock G., Floyd D.L., Golsaz C.M., Gierasch L.M. & Hunt J.F. 2003 Phospholipid-induced monomerization and signal-peptide-induced oligomerization of SecA. *J Biol Chem* **278**, 3628-3638
95. Driessen A.J. 1993 SecA, the peripheral subunit of the Escherichia coli precursor protein translocase, is functional as a dimer. *Biochemistry* **32**, 13190-13197
96. Hunt J.F., Weinkauff S., Henry L., Fak J.J., McNicholas P., Oliver D.B. & Deisenhofer J. 2002 Nucleotide control of interdomain interactions in the conformational reaction cycle of SecA. *Science* **297**, 2018-2026
97. Papanikolaou Y., Papadovasilaki M., Ravelli R.B., McCarthy A.A., Cusack S., Economou A. & Petratos K. 2007 Structure of dimeric SecA, the Escherichia coli preprotein translocase motor. *J Mol Biol* **366**, 1545-1557
98. Sharma V., Arockiasamy A., Ronning D.R., Savva C.G., Holzenburg A., Braunstein M., Jacobs W.R.Jr. & Sacchettini J.C. 2003 Crystal structure of Mycobacterium tuberculosis SecA, a preprotein translocating ATPase. *Proc Natl Acad Sci U S A* **100**, 2243-2248
99. Vassylyev D.G., Mori H., Vassylyeva M.N., Tsukazaki T., Kimura Y., Tahirov T.H. & Ito K. 2006 Crystal structure of the translocation ATPase SecA from Thermus thermophilus reveals a parallel, head-to-head dimer. *J Mol Biol* **364**, 248-258

100. Zimmer J., Li W. & Rapoport T.A. 2006 A novel dimer interface and conformational changes revealed by an X-ray structure of *B. subtilis* SecA. *J Mol Biol* **364**, 259-265
101. Woodbury R.L., Hardy S.J. & Randall L.L. 2002 Complex behavior in solution of homodimeric SecA. *Protein Sci* **11**, 875-882
102. de Keyzer J., van der Sluis E.O., Spelbrink R.E., Nijstad N., de K.B., Nouwen N., van der D.C. & Driessen A.J. 2005 Covalently dimerized SecA is functional in protein translocation. *J Biol Chem* **280**, 35255-35260
103. Ding H., Hunt J.F., Mukerji I. & Oliver D. 2003 *Bacillus subtilis* SecA ATPase exists as an antiparallel dimer in solution. *Biochemistry* **42**, 8729-8738
104. Alami M., Dalal K., Lelj-Garolla B., Sligar S.G. & Duong F. 2007 Nanodiscs unravel the interaction between the SecYEG channel and its cytosolic partner SecA. *EMBO J* **26**, 1995-2004
105. Or E., Navon A. & Rapoport T. 2002 Dissociation of the dimeric SecA ATPase during protein translocation across the bacterial membrane. *EMBO J* **21**, 4470-4479
106. Jilaveanu L.B. & Oliver D. 2006 SecA dimer cross-linked at its subunit interface is functional for protein translocation. *J Bacteriol* **188**, 335-338
107. Wang H., Na B., Yang H. & Tai P.C. 2008 Additional in vitro and in vivo evidence for SecA functioning as dimers in the membrane: dissociation into monomers is not essential for protein translocation in *Escherichia coli*. *J Bacteriol* **190**, 1413-1418
108. Gardel C., Benson S., Hunt J., Michaelis S. & Beckwith J. 1987 secD, a new gene involved in protein export in *Escherichia coli*. *J Bacteriol* **169**, 1286-1290
109. Gardel C., Johnson K., Jacq A. & Beckwith J. 1990 The secD locus of *E.coli* codes for two membrane proteins required for protein export. *EMBO J* **9**, 3209-3216
110. Matsuyama S., Fujita Y. & Mizushima S. 1993 SecD is involved in the release of translocated secretory proteins from the cytoplasmic membrane of *Escherichia coli*. *EMBO J* **12**, 265-270
111. Matsuyama S., Fujita Y., Sagara K. & Mizushima S. 1992 Overproduction, purification and characterization of SecD and SecF, integral membrane components of the protein translocation machinery of *Escherichia coli*. *Biochim Biophys Acta* **1122**, 77-84

-
112. Pogliano J.A. & Beckwith J. 1994 SecD and SecF facilitate protein export in *Escherichia coli*. *EMBO J* **13**, 554-561
113. Kim Y.J., Rajapandi T. & Oliver D. 1994 SecA protein is exposed to the periplasmic surface of the *E. coli* inner membrane in its active state. *Cell* **78**, 845-853
114. Economou A., Pogliano J.A., Beckwith J., Oliver D.B. & Wickner W. 1995 SecA membrane cycling at SecYEG is driven by distinct ATP binding and hydrolysis events and is regulated by SecD and SecF. *Cell* **83**, 1171-1181
115. Eichler J., Brunner J. & Wickner W. 1997 The protease-protected 30 kDa domain of SecA is largely inaccessible to the membrane lipid phase. *EMBO J* **16**, 2188-2196
116. Eichler J. & Wickner W. 1997 Both an N-terminal 65-kDa domain and a C-terminal 30-kDa domain of SecA cycle into the membrane at SecYEG during translocation. *Proc Natl Acad Sci U S A* **94**, 5574-5581
117. Duong F. & Wickner W. 1997 The SecDFyajC domain of preprotein translocase controls preprotein movement by regulating SecA membrane cycling. *EMBO J* **16**, 4871-4879
118. Nouwen N., Piwowarek M., Berrelkamp G. & Driessen A.J. 2005 The large first periplasmic loop of SecD and SecF plays an important role in SecDF functioning. *J Bacteriol* **187**, 5857-5860
119. Tsukazaki T., Mori H., Echizen Y., Ishitani R., Fukai S., Tanaka T., Perederina A., Vassilyev D.G., Kohno T., Maturana A.D., Ito K. & Nureki O. 2011 Structure and function of a membrane component SecDF that enhances protein export. *Nature* **474**, 235-238
120. Bolhuis A., Broekhuizen C.P., Sorokin A., van Roosmalen M.L., Venema G., Bron S., Quax W.J. & van Dijk J.M. 1998 SecDF of *Bacillus subtilis*, a molecular Siamese twin required for the efficient secretion of proteins. *J Biol Chem* **273**, 21217-21224
121. Murakami S., Nakashima R., Yamashita E., Matsumoto T. & Yamaguchi A. 2006 Crystal structures of a multidrug transporter reveal a functionally rotating mechanism. *Nature* **443**, 173-179
122. Tseng T.T., Gratwick K.S., Kollman J., Park D., Nies D.H., Goffeau A. & Saier M.H.Jr. 1999 The RND permease superfamily: an ancient, ubiquitous and diverse family that includes human disease and development proteins. *J Mol Microbiol Biotechnol* **1**, 107-125

123. Schiebel E., Driessen A.J., Hartl F.U. & Wickner W. 1991 Delta mu H⁺ and ATP function at different steps of the catalytic cycle of preprotein translocase. *Cell* **64**, 927-939
124. van der Wolk J.P., de Wit J.G. & Driessen A.J. 1997 The catalytic cycle of the escherichia coli SecA ATPase comprises two distinct preprotein translocation events. *EMBO J* **16**, 7297-7304
125. Driessen A.J. & Wickner W. 1991 Proton transfer is rate-limiting for translocation of precursor proteins by the Escherichia coli translocase. *Proc Natl Acad Sci U S A* **88**, 2471-2475
126. Driessen A.J. 1992 Precursor protein translocation by the Escherichia coli translocase is directed by the protonmotive force. *EMBO J* **11**, 847-853
127. Sagara K., Matsuyama S. & Mizushima S. 1994 SecF stabilizes SecD and SecY, components of the protein translocation machinery of the Escherichia coli cytoplasmic membrane. *J Bacteriol* **176**, 4111-4116
128. Nouwen N. & Driessen A.J. 2002 SecDFyajC forms a heterotetrameric complex with YidC. *Mol Microbiol* **44**, 1397-1405
129. Tornroth-Horsefield S., Gourdon P., Horsefield R., Brive L., Yamamoto N., Mori H., Snijder A. & Neutze R. 2007 Crystal structure of AcrB in complex with a single transmembrane subunit reveals another twist. *Structure* **15**, 1663-1673
130. Fang J. & Wei Y. 2011 Expression, purification and characterization of the Escherichia coli integral membrane protein YajC. *Protein Pept Lett* **18**, 601-608
131. Kol S., Nouwen N. & Driessen A.J. 2008 Mechanisms of YidC-mediated insertion and assembly of multimeric membrane protein complexes. *J Biol Chem* **283**, 31269-31273
132. Economou A. & Wickner W. 1994 SecA promotes preprotein translocation by undergoing ATP-driven cycles of membrane insertion and deinsertion. *Cell* **78**, 835-843
133. Jilaveanu L.B. & Oliver D.B. 2007 In vivo membrane topology of Escherichia coli SecA ATPase reveals extensive periplasmic exposure of multiple functionally important domains clustering on one face of SecA. *J Biol Chem* **282**, 4661-4668
134. Uchida K., Mori H. & Mizushima S. 1995 Stepwise movement of preproteins in the process of translocation across the cytoplasmic membrane of Escherichia coli. *J Biol Chem* **270**, 30862-30868

-
135. Erlandson K.J., Miller S.B., Nam Y., Osborne A.R., Zimmer J. & Rapoport T.A. 2008 A role for the two-helix finger of the SecA ATPase in protein translocation. *Nature* **455**, 984-987
136. Nouwen N., Berrelkamp G. & Driessen A.J. 2009 Charged amino acids in a preprotein inhibit SecA-dependent protein translocation. *J Mol Biol* **386**, 1000-1010
137. Simon S.M., Peskin C.S. & Oster G.F. 1992 What drives the translocation of proteins? *Proc Natl Acad Sci U S A* **89**, 3770-3774
138. Kusters I. & Driessen A.J. 2011 SecA, a remarkable nanomachine. *Cell Mol Life Sci* **68**, 2053-2066
139. Mao C., Hardy S.J. & Randall L.L. 2009 Maximal efficiency of coupling between ATP hydrolysis and translocation of polypeptides mediated by SecB requires two protomers of SecA. *J Bacteriol* **191**, 978-984
140. Randall L.L., Crane J.M., Liu G. & Hardy S.J. 2004 Sites of interaction between SecA and the chaperone SecB, two proteins involved in export. *Protein Sci* **13**, 1124-1133
141. Karamanou S., Gouridis G., Papanikou E., Sianidis G., Gelis I., Keramisanou D., Vrontou E., Kalodimos C.G. & Economou A. 2007 Preprotein-controlled catalysis in the helicase motor of SecA. *EMBO J* **26**, 2904-2914
142. Engelman D.M. & Steitz T.A. 1981 The spontaneous insertion of proteins into and across membranes: the helical hairpin hypothesis. *Cell* **23**, 411-422
143. Joly J.C. & Wickner W. 1993 The SecA and SecY subunits of translocase are the nearest neighbors of a translocating preprotein, shielding it from phospholipids. *EMBO J* **12**, 255-263
144. Natale P., Swaving J., van der Does C., de Keyzer J. & Driessen A.J. 2004 Binding of SecA to the SecYEG complex accelerates the rate of nucleotide exchange on SecA. *J Biol Chem* **279**, 13769-13777
145. Driessen A.J. & Nouwen N. 2008 Protein translocation across the bacterial cytoplasmic membrane. *Annu Rev Biochem* **77**, 643-667
146. Duong F. & Wickner W. 1999 The PrlA and PrlG phenotypes are caused by a loosened association among the translocase SecYEG subunits. *EMBO J* **18**, 3263-3270
147. van der Does C., de Keyzer J., van der Laan M. & Driessen A.J. 2003 Reconstitution of purified bacterial preprotein translocase in liposomes. *Methods Enzymol* **372**, 86-98

148. de Keyzer J., van der Does C. & Driessen A.J. 2002 Kinetic analysis of the translocation of fluorescent precursor proteins into Escherichia coli membrane vesicles. *J Biol Chem* **277**, 46059-46065
149. Hanahan D. 1983 Studies on transformation of Escherichia coli with plasmids. *J Mol Biol* **166**, 557-580
150. Baneyx F. & Georgiou G. 1990 In vivo degradation of secreted fusion proteins by the Escherichia coli outer membrane protease OmpT. *J Bacteriol* **172**, 491-494
151. Nouwen N., van der Laan M. & Driessen A.J. 2001 SecDFyajC is not required for the maintenance of the proton motive force. *FEBS Lett* **508**, 103-106
152. Hess B., Kutzner C., van der Spoel D. & Lindahl E. 2008 GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *Journal of Chemical Theory and Computation* **4**, 435-447
153. Marrink S.J., Risselada H.J., Yefimov S., Tieleman D.P. & de Vries A.H. 2007 The MARTINI force field: coarse grained model for biomolecular simulations. *J Phys Chem B* **111**, 7812-7824
154. Monticelli L., Kandasamy S.K., Periole X., Larson R.G., Tieleman D.P. & Marrink S.J. 2008 The MARTINI Coarse-Grained Force Field: Extension to Proteins. *Journal of Chemical Theory and Computation* **4**, 819-834
155. Sengupta D., Rampioni A. & Marrink S.J. 2009 Simulations of the c-subunit of ATP-synthase reveal helix rearrangements. *Mol Membr Biol* **26**, 422-434
156. Yefimov S., van der Giessen E., Onck P.R. & Marrink S.J. 2008 Mechanosensitive membrane channels in action. *Biophys J* **94**, 2994-3002
157. Green N.S., Reisler E. & Houk K.N. 2001 Quantitative evaluation of the lengths of homobifunctional protein cross-linking reagents used as molecular rulers. *Protein Sci* **10**, 1293-1304
158. Bol R., de Wit J.G. & Driessen A.J. 2007 The active protein-conducting channel of Escherichia coli contains an apolar patch. *J Biol Chem* **282**, 29785-29793
159. Cabelli R.J., Chen L., Tai P.C. & Oliver D.B. 1988 SecA protein is required for secretory protein translocation into E. coli membrane vesicles. *Cell* **55**, 683-692

-
160. Fekkes P., de Wit J.G., van der Wolk J.P., Kimsey H.H., Kumamoto C.A. & Driessen A.J. 1998 Preprotein transfer to the Escherichia coli translocase requires the co-operative binding of SecB and the signal sequence to SecA. *Mol Microbiol* **29**, 1179-1190
161. Rutkowska A., Mayer M.P., Hoffmann A., Merz F., Zachmann-Brand B., Schaffitzel C., Ban N., Deuerling E. & Bukau B. 2008 Dynamics of trigger factor interaction with translating ribosomes. *J Biol Chem* **283**, 4124-4132
162. Klose M., Schimz K.L., van der W.J., Driessen A.J. & Freudl R. 1993 Lysine 106 of the putative catalytic ATP-binding site of the Bacillus subtilis SecA protein is required for functional complementation of Escherichia coli secA mutants in vivo. *J Biol Chem* **268**, 4504-4510
163. Manting E.H., Kaufmann A., van der D.C. & Driessen A.J. 1999 A single amino acid substitution in SecY stabilizes the interaction with SecA. *J Biol Chem* **274**, 23868-23874
164. Schaffitzel C. & Ban N. 2007 Generation of ribosome nascent chain complexes for structural and functional studies. *J Struct Biol* **158**, 463-471
165. Evans M.S., Ugrinov K.G., Frese M.A. & Clark P.L. 2005 Homogeneous stalled ribosome nascent chain complexes produced in vivo or in vitro. *Nat Methods* **2**, 757-762
166. Lakowicz J.R. 1999 Principles of fluorescence spectroscopy.
167. Arkowitz R.A., Joly J.C. & Wickner W. 1993 Translocation can drive the unfolding of a preprotein domain. *EMBO J* **12**, 243-253
168. Nakatogawa H. & Ito K. 2002 The ribosomal exit tunnel functions as a discriminating gate. *Cell* **108**, 629-636
169. Greenough K.P. & Blanchard G.J. 2006 Quantitating the dynamics of NBD hexanoic acid in homogeneous solution and in solutions containing unilamellar vesicles. *J Phys Chem B* **110**, 6351-6358
170. Lin S. & Struve W.S. 1991 Time-resolved fluorescence of nitrobenzoxadiazole-aminohexanoic acid: effect of intermolecular hydrogen-bonding on non-radiative decay. *Photochem Photobiol* **54**, 361-365
171. Ramachandran R., Tweten R.K. & Johnson A.E. 2004 Membrane-dependent conformational changes initiate cholesterol-dependent cytolysin oligomerization and intersubunit beta-strand alignment. *Nat Struct Mol Biol* **11**, 697-705

172. Prinz A., Behrens C., Rapoport T.A., Hartmann E. & Kalies K.U. 2000 Evolutionarily conserved binding of ribosomes to the translocation channel via the large ribosomal RNA. *EMBO J* **19**, 1900-1906
173. Becker T., Bhushan S., Jarasch A., Armache J.P., Funes S., Jossinet F., Gumbart J., Mielke T., Berninghausen O., Schulten K., Westhof E., Gilmore R., Mandon E.C. & Beckmann R. 2009 Structure of monomeric yeast and mammalian Sec61 complexes interacting with the translating ribosome. *Science* **326**, 1369-1373
174. Kontinen V.P., Yamanaka M., Nishiyama K. & Tokuda H. 1996 Roles of the conserved cytoplasmic region and non-conserved carboxy-terminal region of SecE in Escherichia coli protein translocase. *J Biochem* **119**, 1124-1130
175. Hsu H.J., Tsai K.C., Sun Y.K., Chang H.J., Huang Y.J., Yu H.M., Lin C.H., Mao S.S. & Yang A.S. 2008 Factor Xa active site substrate specificity with substrate phage display and computational molecular modeling. *J Biol Chem* **283**, 12343-12353
176. Harris J.L., Backes B.J., Leonetti F., Mahrus S., Ellman J.A. & Craik C.S. 2000 Rapid and general profiling of protease specificity by using combinatorial fluorogenic substrate libraries. *Proc Natl Acad Sci U S A* **97**, 7754-7759
177. Matthews D.J. & Wells J.A. 1993 Substrate phage: selection of protease substrates by monovalent phage display. *Science* **260**, 1113-1117
178. Nishiyama K., Kabuyama Y., Akimaru J., Matsuyama S., Tokuda H. & Mizushima S. 1991 SecY is an indispensable component of the protein secretory machinery of Escherichia coli. *Biochim Biophys Acta* **1065**, 89-97
179. Swaving J., van Wely K.H. & Driessen A.J. 1999 Preprotein translocation by a hybrid translocase composed of Escherichia coli and Bacillus subtilis subunits. *J Bacteriol* **181**, 7021-7027
180. Wu Z.C., de Keyzer J., Kedrov A & Driessen A.J. 2012 Competitive binding of the SecA ATPase and ribosomes to the SecYEG translocon. *J Biol Chem.* **287**, 7885-7895