

University of Groningen

Genetics of asthma and atopy

Koppelman, Gerard Henk

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koppelman, G. H. (2001). *Genetics of asthma and atopy*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 3 | Genetics and environment in asthma: the answer of twin studies

(Eur Respir J 1999;13:2-4)

Gerard H. Koppelman ^{* #}, Henk Los ^{* #} and Dirkje S. Postma ^{*}

[#] Department of Pulmonary Rehabilitation, Beatrixoord Rehabilitation Centre, Haren, the Netherlands

^{*} Department of Pulmonology, University Hospital Groningen, the Netherlands

Table 1 Results of twin studies of asthma

Population [reference]	Number of twin-pairs	MZ correlation+	DZ correlation+	Probandwise concordance MZ/DZ	Heritability	Comments
Swedish [5]	6996	0.65	0.25*			Adult population based study. Correlations calculated by Duffy et al. ²
Australian [2]	3808	0.65	0.24*		0.60-0.75	"Asthma or wheezing" by questionnaire. Adult population.
Finnish [3]	13888	0.43	0.25*	0.13/0.07		Hospitalization, medication or cause of death / adult population based study.
Swedish [6]	434 M 456 F			0.62/0.26* 0.41/0.18*	0.36	Questionnaire (ever wheezing with short-ness of breath, wheezing without a cold, or parental reported asthma) / twins aged 7-9 years.
Norwegian [4]	2570	0.75	0.21*	0.45/0.12*	0.75	Population based study of twins aged 18 - 25 years.
Finnish [7]	1713	0.76	0.45	0.42/0.19*		Population based study of twins aged 16 years.
Danish [1]	1929 M 2131 F 1867 M 2110 F	0.76 0.71 0.81 0.65	0.36 0.47 0.37 0.15	0.48/0.19* 0.42/0.26* 0.51/0.16* 0.38/0.09*		Population based study of twins aged 12 - 26 years. Age 12-26. Age 27-41. Age 27-41.

MZ monozygous; DZ dizygous; M male; F female; + correlation is tetrachoric correlation; * statistically significant differences between MZ and DZ twin pairs

Twin studies have been widely used to estimate the genetic contribution to diseases. In this issue of the *European Respiratory Journal*, Skadhauge *et al.*¹ present the results of a large Danish, population-based twin study on asthma. In this study, the heritability in liability to asthma, i.e. the proportion of variance due to genetic factors, is estimated to be 0.77 for males and 0.68 for females. In other studies, the heritability of asthma is estimated to be between 0.36 and 0.75²⁻⁷ (table 1). Thus, the results of this Danish study are consistent with those of other twin studies and add to the body of evidence indicating that the genetic contribution to asthma is considerable. In addition, the results of this study suggest individual specific, unshared, environmental factors to be important as well. In this editorial, the assumptions and methods of twin studies will be assessed, and the role of genetic and environmental factors in asthma reviewed.

Genetic studies using the twin-design have four major assumptions:⁸

1. Monozygous (MZ) and dizygous(DZ) twins are samples of the same gene pool;
2. twins are representative of the general population;
3. self-reported zygoty is correct in questionnaire based studies;
4. the environment for both MZ and DZ twins is similar.

The first and second assumptions are valid, provided representative or complete samples are taken from the population. The second assumption, the representativity, may not be totally valid because MZ and DZ twins differ from each other and from singletons with respect to their intrauterine environment.⁹ The shared intrauterine environment may have an adverse effect on the growth and organ maturation of the foetus. However, this most likely does not influence the development of asthma, since the prevalence of asthma is comparable in twins and singletons.^{1,2,4} The third assumption has been tested.⁷ In general, self reported zygoty questions are adequate in 95-98 % of cases. Finally, the fourth assumption of an equal environment may not be valid in the case of asthma. For instance, it has been shown that MZ twins have more similar smoking patterns than DZ twins.¹² It is unknown if this higher similarity in MZ twins is also the case for other environmental factors, such as exposure to indoor allergens and viruses. A higher similarity in environment for MZ twins compared to DZ twins may lead to an overestimation of the heritability of asthma.

The method for diagnosing asthma is a self-reported questionnaire in most large scale twin studies. Subjects in these studies are not tested clinically. This method may lead to an under- or overestimation of asthma prevalence.^{13, 14} Overestimation could occur, for instance, if asthma is diagnosed by questions on wheeze. Small children in particular may wheeze during the course of a respiratory infection, but not have asthma.¹² Therefore, studies on the genetics of asthma are currently directed at measurable clinical components of asthma, e.g. airway hyperresponsiveness (AH), reversibility and variability of airway obstruction. If it were known which of these components of asthma have a high genetic contribution, these


components can then be selected for genetic studies to find genes that regulate these components. As an example, one twin study of reasonable size on AH has been published, in which the heritability of AH to methacholine was 0.66.¹⁶ Clearly, more studies are needed.

In general, statistical analyses of twin studies are complicated. In the study of twins, phenotypic similarities and differences are compared between MZ and DZ twins. MZ twins share 100% of their genetic information and DZ twins share on average 50%. If a trait is influenced by genetic factors, MZ twins should resemble each other to a greater extent than DZ twins, and the correlations between MZ and DZ twinpairs may be used to estimate the relative size of genetic and environmental influences.

In biometric modelling, one goes one step further. Since these biometric analyses are not frequently presented in pulmonary journals, we will first address some assumptions and methods before discussing the results of these analyses in the study of asthma. In biometric modelling, a quantitative genetic analysis is performed with dichotomous variables (e.g. asthma/not asthma). To permit these analyses, the first assumption is that disease status is determined by an unobserved continuous variable called the liability. If the liability falls above a threshold, individuals are classified as affected. The second assumption is that the distribution of the liability is normal.¹⁷ The variance of the distribution of the liability is composed of multiple environmental and genetic influences. The environmental component can be dissected in influences shared by both twins and influences not shared. Furthermore, the genetic component consists of an effect of individual alleles on the trait (additive effect) or interaction between alleles at the same locus (dominance effect).⁸ The last possible source of genetic effects, i.e. interaction of alleles at different loci (epistasis) can not be discriminated from dominant genetic effects in twin studies, which is a limitation of this design. Thus, the observed phenotypes P1 of twin 1 and P2 of twin 2 of a twin pair, will be linear functions of the underlying additive genetic influences (A-twin 1, A-twin 2), dominance genetic influences (D-twin 1, D-twin 2), shared environmental influences (C-twin 1, C-twin-2) and specific environmental influences (E-twin 1, E-twin 2). These functions are calculated in a path model by specialized computer programs (figure 1). The results of these calculations are then compared to known models, including a model in which the disease is caused by environmental factors alone, and a model in which the disease is caused by genetic factors, or combinations of these models. In this way, the best fitting model is calculated, i.e. the model that describes the data best. For more background information the reader is referred to the book by Neale and Cardon.¹⁸

By applying the above mentioned methods, Skudhauge *et al.*¹ found evidence for the liability for asthma in a model consisting of additive genetic factors and non-shared environmental influences, with modest evidence of effects of shared environment. Interestingly, other large-scale twin studies in different countries in the world came to the same conclusion.^{2,4,7} The question that arises is: How do these findings relate to current evidence on the role of genetic and environmental factors in asthma?

Figure 1 Path diagram depicting genetic and environmental effects



Twin 1 and twin 2 are the first and second twin in a pair with phenotype P1 and P2, respectively. A (Additive genetic effects), C (shared environmental effects), E (non-shared environmental effects) and D (dominant genetic effects). a, c, d and e are path coefficients which are measures of variance of these genetic and environmental effects. MZ twins share 100% of their genes, therefore the correlation is 1 for additive and 1 for dominant effects. In DZ twins, these correlations are 0.5 for additive, and 0.25 for dominant effects.

Genetic factors

Major susceptibility genes for asthma and atopy have not been determined to date.¹⁹ Several reports indicate a possible role for mutations in a gene on chromosome 11 coding for the β -chain of the high affinity IgE receptor in atopy or asthma, however, this picture is not clear as these mutations seem not to play in role in asthma nor atopy in other populations.²⁰ The gene coding for the β_2 -adrenergic receptor has been studied in more detail. Current data indicate that two common polymorphisms in this gene do not play a role in the causation of asthma. They may, however, modify asthma into a more severe phenotype expressed as more nocturnal complaints, higher use of inhaled corticosteroids in patients with these mutations and a reduced effect of β -mimetics.²¹

Environmental factors

Environmental risk factors in asthma are active or passive smoking, exposure to allergens²², viral respiratory infections, and possibly diet and air pollution.²³ Intuitively, most of these environmental factors appear to be largely shared. However, the twin studies suggest that not the shared, but the unshared individual-specific environment appears to be important. It is a challenge for researchers to assess which factors have these specific effects and to what extent the timing of exposure is relevant. It is of major interest to learn to understand how these factors interact with each other and with genetic factors.

In summary, what answers do twin studies give to the question which genetic and environmental factors cause asthma? First, twin studies have indicated the considerable genetic component of asthma. This component most likely consists of genes of additive effect. Second, twin studies have shown that individual specific environmental factors may be important as well. To further understand the genetics of asthma, we recommend to direct the twin approach at measurable components of asthma, such as airway hyperresponsiveness, reversibility and variability in airway obstruction. Since we currently do not know which genes lead to susceptibility to asthma, the next challenge will be to study interaction of these genes and specific environmental factors in the development of asthma.

Acknowledgement

G. H. Koppelman is supported by a grant from the Dutch Asthma Foundation, AF 95.09.

References

- 1 Skadhauge LR, Christensen K, Kyvik KO, Sigsgaard T. Genetic and environmental influence on asthma. A population-based study of 11,688 Danish twin pairs. *Eur Respir J* 1999;13: 8-14.
- 2 Duffy DL, Martin NG, Battistutta D, Hopper JL, Mathews JD. Genetics of asthma and hay fever in Australian twins. *Am Rev Respir Dis* 1990;142:1351-8.
- 3 Nieminen MM, Kaprio J, Koskenvuo M. A population-based study of bronchial asthma in adult twin pairs. *Chest* 1991;100:70-5.
- 4 Harris JR, Magnus P, Samuelsen SO, Tambs K. No evidence for effects of family environment on asthma. A retrospective study of Norwegian twins. *Am J Respir Crit Care Med* 1997;156:43-9.
- 5 Edfors-Lubs ML. Allergy in 7000 twin pairs. *Acta Allergol* 1971; 26:249-285.
- 6 Lichtenstein P, Svartengren M. Genes, environments, and sex: factors of importance in atopic diseases in 7-9 year-old Swedish twins. *Allergy* 1997; 52: 1079-1086.
- 7 Laitinen T, Rasasen M, Kaprio J, Koskenvuo M, Laitinen LA. Importance of genetic factors in adolescent asthma. *Am J Respir Crit Care Med* 1998; 157: 1073-1078.
- 8 Khoury MJ, Beaty TH, Cohen BH. Fundamentals of genetic epidemiology. New York Oxford: Oxford University Press; 1993; 7, Genetic approaches to familial aggregation. I. Analysis of heritability. p. 200-32.
- 9 Phillips DIW. Twin studies in medical research: can they tell us whether diseases are genetically determined? *Lancet* 1993;341:1008-9.
- 10 Magnus P, Berg K, Nance WE. Predicting zygosity in Norwegian twin pairs born in 1915-1960. *Clin Genet* 1983;24:103-12.
- 11 Boomsma DI, Kaptein A, Kempen A, Gevers-Leuven JA, Princen HMG. Lipoprotein (a): relation to other risk factors and genetic heritability. Results from a Dutch parent-twin study. *Atherosclerosis* 1993;99:23-33.
- 12 Carmelli D, Swan GE, Robinette D, Fabsitz R. Genetic influence on smoking - a study of male twins. *N Engl J Med* 1992;327:829-33.
- 13 Marco de R, Cerveri I, Bugiani M, Ferrari M, Verlato G. An undetected burden of asthma in Italy: the relationship between clinical and epidemiological diagnosis of asthma. *Eur Respir J* 1998;11:599-605.
- 14 Cerveri I, Bruschi C, Ricciardi M, Zocchi L, Zoia MC, Rampulla C. Epidemiological diagnosis of asthma: methodological considerations of prevalence evaluation. *Eur J Epidemiol* 1987;3:202-5.
- 15 Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ, et al. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995;332:133-8.
- 16 Hopp RJ, Bewtra AK, Watt GD, Nair NM, Townley RG. Genetic analysis of allergic disease in twins. *J Allergy Clin Immunol* 1984;73:265-70.
- 17 Falconer DS. The inheritance of liability to certain diseases estimated from the incidence among relatives. *Ann Hum Genet* 1965;29:51-76.
- 18 Neale MC; Cardon LR. Methodology for genetic studies of twins and families. Dordrecht, the Netherlands: Kluwer Academic Publishers; 1992.

- 19 Holgate ST. Asthma genetics: waiting to exhale. *Nat Genet* 1997;15:227-9.
- 20 Thomas NS, Holgate ST. Genes for asthma on chromosome 11: an update (editorial). *Clin Exp Allergy* 1998;28:387-91.
- 21 Liggett SB. Pharmacogenetics of relevant targets in asthma. *Clin Exp Allergy* 1998;28 Supplement 1:77-9.
- 22 Duffy DL, Mitchell CA, Martin NG. Genetic and environmental risk factors for asthma: a cotwin-control study. *Am J Respir Crit Care Med* 1998;157:840-5.
- 23 Sears MR. Descriptive epidemiology of asthma. *Lancet* 1997;350:III1-4.

