

University of Groningen

Visual hallucinations in Parkinson's disease

Meppelink, Anne Marthe

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Meppelink, A. M. (2011). *Visual hallucinations in Parkinson's disease: clinical and fMRI studies*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Bibliography

- Aarsland, D., Andersen, K., Larsen, J. P., Lolk, A., and Kragh-Sorensen, P. (2003). Prevalence and characteristics of dementia in parkinson disease: an 8-year prospective study. *Arch.Neurol.*, 60(3):387–392.
- Aarsland, D., Zaccai, J., and Brayne, C. (2005). A systematic review of prevalence studies of dementia in parkinson’s disease. *Mov Disord.*, 20(10):1255–1263.
- Abe, Y., Kachi, T., Kato, T., Arahata, Y., Yamada, T., Washimi, Y., Iwai, K., Ito, K., Yanagisawa, N., and Sobue, G. (2003). Occipital hypoperfusion in parkinson’s disease without dementia: correlation to impaired cortical visual processing. *J.Neurol.Neurosurg.Psychiatry*, 74(4):419–422.
- Alegret, M., Valdeoriola, F., Marti, M., Pilleri, M., Junque, C., Rumia, J., and Tolosa, E. (2004). Comparative cognitive effects of bilateral subthalamic stimulation and subcutaneous continuous infusion of apomorphine in parkinson’s disease. *Mov Disord.*, 19(12):1463–1469.
- Aleman, A., Laroi, F., and Association, A. P. (2008). *Hallucinations. The science of idiosyncratic perception*. Washington DC.
- Archibald, N. K., Clarke, M. P., Mosimann, U. P., and Burn, D. J. (2009). The retina in parkinson’s disease. *Brain*, 132(Pt 5):1128–1145.
- Arditi, A. (2005). Improving the design of the letter contrast sensitivity test. *Invest Ophthalmol. Vis.Sci.*, 46(6):2225–2229.
- Ashby, F. G., Ennis, J. M., and Spiering, B. J. (2007). A neurobiological theory of automaticity in perceptual categorization. *Psychol.Rev.*, 114(3):632–656.
- Baker, W. L., Silver, D., White, C. M., Kluger, J., Aberle, J., Patel, A. A., and Coleman, C. I. (2009). Dopamine agonists in the treatment of early parkinson’s disease: a meta-analysis. *Parkinsonism.Relat Disord.*, 15(4):287–294.

- Ballanger, B., Strafella, A. P., van Eimeren, T., Zurowski, M., Rusjan, P. M., Houle, S., and Fox, S. H. (2010). Serotonin 2a receptors and visual hallucinations in parkinson disease. *Arch.Neurol.*, 67(4):416–421.
- Ballard, C. G., Aarsland, D., McKeith, I., O'Brien, J., Gray, A., Cormack, F., Burn, D., Cassidy, T., Starfeldt, R., Larsen, J. P., Brown, R., and Tovee, M. (2002). Fluctuations in attention: Pd dementia vs dlb with parkinsonism. *Neurology*, 59(11):1714–1720.
- Bar, M., Kassam, K. S., Ghuman, A. S., Boshyan, J., Schmid, A. M., Dale, A. M., Hamalainen, M. S., Marinkovic, K., Schacter, D. L., Rosen, B. R., and Halgren, E. (2006). Top-down facilitation of visual recognition. *Proc.Natl.Acad.Sci.U.S.A.*, 103(2):449–454.
- Bar, M., Tootell, R. B., Schacter, D. L., Greve, D. N., Fischl, B., Mendola, J. D., Rosen, B. R., and Dale, A. M. (2001). Cortical mechanisms specific to explicit visual object recognition. *Neuron*, 29(2):529–535.
- Barnes, J. and Boubert, L. (2008). Executive functions are impaired in patients with parkinson's disease with visual hallucinations. *J.Neurol.Neurosurg.Psychiatry*, 79(2):190–192.
- Barnes, J., Boubert, L., Harris, J., Lee, A., and David, A. S. (2003). Reality monitoring and visual hallucinations in parkinson's disease. *Neuropsychologia*, 41(5):565–574.
- Barnes, J. and David, A. S. (2001). Visual hallucinations in parkinson's disease: a review and phenomenological survey. *J.Neurol.Neurosurg.Psychiatry*, 70(6):727–733.
- BECK, A. T., WARD, C. H., MENDELSON, M., MOCK, J., and ERBAUGH, J. (1961). An inventory for measuring depression. *Arch.Gen.Psychiatry*, 4:561–571.
- Beck, D. M. and Kastner, S. (2005). Stimulus context modulates competition in human extrastriate cortex. *Nat.Neurosci.*, 8(8):1110–1116.
- Beck, D. M. and Kastner, S. (2009). Top-down and bottom-up mechanisms in biasing competition in the human brain. *Vision Res.*, 49(10):1154–1165.
- Beyer, M. K., Janvin, C. C., Larsen, J. P., and Aarsland, D. (2007). A magnetic resonance imaging study of patients with parkinson's disease with mild cognitive impairment and dementia using voxel-based morphometry. *J.Neurol.Neurosurg.Psychiatry*, 78(3):254–259.

- Binkofski, F., Buccino, G., Stephan, K. M., Rizzolatti, G., Seitz, R. J., and Freund, H. J. (1999). A parieto-premotor network for object manipulation: evidence from neuroimaging. *Exp.Brain Res.*, 128(1-2):210–213.
- Biousse, V., Skibell, B. C., Watts, R. L., Loupe, D. N., Drews-Botsch, C., and Newman, N. J. (2004). Ophthalmologic features of parkinson's disease. *Neurology*, 62(2):177–180.
- Blanke, O., Landis, T., and Seeck, M. (2000). Electrical cortical stimulation of the human prefrontal cortex evokes complex visual hallucinations. *Epilepsy Behav.*, 1(5):356–361.
- Bodis-Wollner, I. (1990). The visual system in parkinson's disease. *Res.Publ.Assoc.Res.Nerv.Ment.Dis.*, 67:297–316.
- Bodis-Wollner, I. (2009). Retinopathy in parkinson disease. *J.Neural Transm.*, 116(11):1493–1501.
- Boecker, H., Ceballos-Baumann, A. O., Volk, D., Conrad, B., Forstl, H., and Haussermann, P. (2007). Metabolic alterations in patients with parkinson disease and visual hallucinations. *Arch.Neurol.*, 64(7):984–988.
- Bohnen, N. I., Kaufer, D. I., Hendrickson, R., Ivanco, L. S., Lopresti, B. J., Constantine, G. M., Mathis, C., Davis, J. G., Moore, R. Y., and Dekosky, S. T. (2006). Cognitive correlates of cortical cholinergic denervation in parkinson's disease and parkinsonian dementia. *J.Neurol.*, 253(2):242–247.
- Bos, E. H., Bouhuys, A. L., Geerts, E., van Os, T. W., Van der Spoel, I. D., Brouwer, W. H., and Ormel, J. (2005). Cognitive, physiological, and personality correlates of recurrence of depression. *J.Affect.Disord.*, 87(2-3):221–229.
- Bosboom, J. L., Stoffers, D., and Wolters, E. C. (2004). Cognitive dysfunction and dementia in parkinson's disease. *J.Neural Transm.*, 111(10-11):1303–1315.
- Braak, H., Ghebremedhin, E., Rub, U., Bratzke, H., and Del Tredici, K. (2004). Stages in the development of parkinson's disease-related pathology. *Cell Tissue Res.*, 318(1):121–134.
- Braak, H., Rub, U., Jansen Steur, E. N., Del Tredici, K., and de Vos, R. A. (2005). Cognitive status correlates with neuropathologic stage in parkinson disease. *Neurology*, 64(8):1404–1410.
- Brainard, D. H. (1997). The psychophysics toolbox. *Spat. Vis.*, 10(4):433–436.

- Buning, H. (1997). Robust analysis of variance. *Journal of Applied Statistics*, 24(3):319–332.
- Burke, W. (2002). The neural basis of charles bonnet hallucinations: a hypothesis. *J.Neurol.Neurosurg.Psychiatry*, 73(5):535–541.
- Burn, D., Emre, M., McKeith, I., De Deyn, P. P., Aarsland, D., Hsu, C., and Lane, R. (2006). Effects of rivastigmine in patients with and without visual hallucinations in dementia associated with parkinson’s disease. *Mov Disord.*, 21(11):1899–1907.
- Burton, E. J., McKeith, I. G., Burn, D. J., Williams, E. D., and O’Brien, J. T. (2004). Cerebral atrophy in parkinson’s disease with and without dementia: a comparison with alzheimer’s disease, dementia with lewy bodies and controls. *Brain*, 127(Pt 4):791–800.
- Buttner, T., Kuhn, W., Muller, T., Patzold, T., Heidbrink, K., and Przuntek, H. (1995). Distorted color discrimination in ‘de novo’ parkinsonian patients. *Neurology*, 45(2):386–387.
- Buttner, T., Kuhn, W., Patzold, T., and Przuntek, H. (1994). L-dopa improves colour vision in parkinson’s disease. *J.Neural Transm.Park Dis.Dement.Sect.*, 7(1):13–19.
- Buttner, T., Muller, T., and Kuhn, W. (2000). Effects of apomorphine on visual functions in parkinson’s disease. *J.Neural Transm.*, 107(1):87–94.
- Catani, M. and Mesulam, M. (2008). The arcuate fasciculus and the disconnection theme in language and aphasia: history and current state. *Cortex*, 44(8):953–961.
- Collerton, D., Perry, E., and McKeith, I. (2005). Why people see things that are not there: A novel perception and attention deficit model for recurrent complex visual hallucinations. *Behav.Brain Sci.*, 28(6):737–757.
- Colosimo, C., Hughes, A. J., Kilford, L., and Lees, A. J. (2003). Lewy body cortical involvement may not always predict dementia in parkinson’s disease. *J.Neurol.Neurosurg.Psychiatry*, 74(7):852–856.
- Corbetta, M., Kincade, J. M., and Shulman, G. L. (2002). Neural systems for visual orienting and their relationships to spatial working memory. *J.Cogn Neurosci.*, 14(3):508–523.

- Corbetta, M., Miezin, F. M., Dobmeyer, S., Shulman, G. L., and Petersen, S. E. (1991). Selective and divided attention during visual discriminations of shape, color, and speed: functional anatomy by positron emission tomography. *J.Neurosci.*, 11(8):2383–2402.
- Coull, J. T., Frackowiak, R. S., and Frith, C. D. (1998). Monitoring for target objects: activation of right frontal and parietal cortices with increasing time on task. *Neuropsychologia*, 36(12):1325–1334.
- Crick, F. (1984). Function of the thalamic reticular complex: the searchlight hypothesis. *Proc.Natl.Acad.Sci.U.S.A.*, 81(14):4586–4590.
- Cunnington, R., Egan, G. F., O’Sullivan, J. D., Hughes, A. J., Bradshaw, J. L., and Colebatch, J. G. (2001). Motor imagery in parkinson’s disease: a pet study. *Mov Disord.*, 16(5):849–857.
- Dagher, A. and Nagano-Saito, A. (2007). Functional and anatomical magnetic resonance imaging in parkinson’s disease. *Mol.Imaging Biol.*, 9(4):234–242.
- Davidson, S., Cronin-Golomb, A., and Lee, A. (2005). Visual and spatial symptoms in parkinson’s disease. *Vision Res.*, 45(10):1285–1296.
- de Jong, B. M. and Paans, A. M. (2007). Medial versus lateral prefrontal dissociation in movement selection and inhibitory control. *Brain Res.*, 1132(1):139–147.
- de Jong, B. M., van der Graaf, F. H., and Paans, A. M. (2001). Brain activation related to the representations of external space and body scheme in visuomotor control. *Neuroimage.*, 14(5):1128–1135.
- de Jong, B. M., van Zomeren, A. H., Willemsen, A. T., and Paans, A. M. (1996). Brain activity related to serial cognitive performance resembles circuitry of higher order motor control. *Exp.Brain Res.*, 109(1):136–140.
- de Jong, B. M., Willemsen, A. T., and Paans, A. M. (1999). Brain activation related to the change between bimanual motor programs. *Neuroimage.*, 9(3):290–297.
- Desimone, R. and Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annu.Rev.Neurosci.*, 18:193–222.
- Di Rosa, A. E., Epifanio, A., Antonini, A., Stocchi, F., Martino, G., Di Blasi, L., Tetto, A., Basile, G., Imbesi, D., La Spina, P., Di Raimondo, G., and

- Morgante, L. (2003). Continuous apomorphine infusion and neuropsychiatric disorders: a controlled study in patients with advanced parkinson's disease. *Neurol.Sci.*, 24(3):174–175.
- Diederich, N. J., Fenelon, G., Stebbins, G., and Goetz, C. G. (2009). Hallucinations in parkinson disease. *Nat.Rev.Neurol.*, 5(6):331–342.
- Diederich, N. J., Goetz, C. G., Raman, R., Pappert, E. J., Leurgans, S., and Piery, V. (1998). Poor visual discrimination and visual hallucinations in parkinson's disease. *Clin.Neuropsycharmacol.*, 21(5):289–295.
- Diederich, N. J., Goetz, C. G., and Stebbins, G. T. (2005). Repeated visual hallucinations in parkinson's disease as disturbed external/internal perceptions: focused review and a new integrative model. *Mov Disord.*, 20(2):130–140.
- Diederich, N. J., Pieri, V., and Goetz, C. G. (2003). Coping strategies for visual hallucinations in parkinson's disease. *Mov Disord.*, 18(7):831–832.
- Dotchin, C. L., Jusabani, A., and Walker, R. W. (2009). Non-motor symptoms in a prevalent population with parkinson's disease in tanzania. *Parkinsonism.Relat Disord.*, 15(6):457–460.
- Downing, P. E., Chan, A. W., Peelen, M. V., Dodds, C. M., and Kanwisher, N. (2006). Domain specificity in visual cortex. *Cereb.Cortex*, 16(10):1453–1461.
- Draganski, B., Gaser, C., Busch, V., Schuierer, G., Bogdahn, U., and May, A. (2004). Neuroplasticity: changes in grey matter induced by training. *Nature*, 427(6972):311–312.
- Draganski, B. and May, A. (2008). Training-induced structural changes in the adult human brain. *Behav.Brain Res.*, 192(1):137–142.
- Dubois, B. and Pillon, B. (1997). Cognitive deficits in parkinson's disease. *J.Neurol.*, 244(1):2–8.
- Dubois, B., Slachevsky, A., Litvan, I., and Pillon, B. (2000). The fab: a frontal assessment battery at bedside. *Neurology*, 55(11):1621–1626.
- Eger, E., Henson, R. N., Driver, J., and Dolan, R. J. (2006). Mechanisms of top-down facilitation in perception of visual objects studied by fmri. *Cereb.Cortex*.
- Ellis, C., Lemmens, G., Parkes, J. D., Abbott, R. J., Pye, I. F., Leigh, P. N., and Chaudhuri, K. R. (1997). Use of apomorphine in parkinsonian patients with neuropsychiatric complications to oral treatment. *Parkinsonism.Relat Disord.*, 3(2):103–107.

- Epstein, J., Stern, E., and Silbersweig, D. (1999a). Mesolimbic activity associated with psychosis in schizophrenia. symptom-specific pet studies. *Ann.N.Y.Acad.Sci.*, 877:562–574.
- Epstein, R., Harris, A., Stanley, D., and Kanwisher, N. (1999b). The parahippocampal place area: recognition, navigation, or encoding? *Neuron*, 23(1):115–125.
- Esselink, R. A., de Bie, R. M., de Haan, R. J., Lenders, M. W., Nijssen, P. C., Staal, M. J., Smeding, H. M., Schuurman, P. R., Bosch, D. A., and Speelman, J. D. (2004). Unilateral pallidotomy versus bilateral subthalamic nucleus stimulation in pd: a randomized trial. *Neurology*, 62(2):201–207.
- Farnsworth, D. and Society, T. P. (1947). *The Farnsworth Dichotomous Test for Color Blindness*. New York.
- Faul, F., Erdfelder, E., Lang, A. G., and Buchner, A. (2007). G*power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav.Res.Methods*, 39(2):175–191.
- Feldmann, A., Illes, Z., Kosztolanyi, P., Illes, E., Mike, A., Kover, F., Balas, I., Kovacs, N., and Nagy, F. (2008). Morphometric changes of gray matter in parkinson’s disease with depression: a voxel-based morphometry study. *Mov Disord.*, 23(1):42–46.
- Fenelon, G., Goetz, C. G., and Karenberg, A. (2006). Hallucinations in parkinson disease in the prelevodopa era. *Neurology*, 66(1):93–98.
- Fenelon, G., Mahieux, F., Huon, R., and Ziegler, M. (2000). Hallucinations in parkinson’s disease: prevalence, phenomenology and risk factors. *Brain*, 123 (Pt 4):733–745.
- Fenelon, G., Souals, T., de Langavant L., C., and C, B.-L. A. (2008). Sense of presence in parkinson’s disease: A prospective phenomenological study.
- Ffytche, D. H., Howard, R. J., Brammer, M. J., David, A., Woodruff, P., and Williams, S. (1998). The anatomy of conscious vision: an fmri study of visual hallucinations. *Nat.Neurosci.*, 1(8):738–742.
- Fink, G. R., Dolan, R. J., Halligan, P. W., Marshall, J. C., and Frith, C. D. (1997). Space-based and object-based visual attention: shared and specific neural domains. *Brain*, 120 (Pt 11):2013–2028.

- Flowers, K. A. and Robertson, C. (1995). Perceptual abnormalities in parkinsons-disease - top-down or bottom-up processes. *Perception*, 24(10):1201–1221.
- Folstein, M. F., Folstein, S. E., and McHugh, P. R. (1975). "mini-mental state". a practical method for grading the cognitive state of patients for the clinician. *J.Psychiatr.Res.*, 12(3):189–198.
- Fox, P. T. and Raichle, M. E. (1986). Focal physiological uncoupling of cerebral blood flow and oxidative metabolism during somatosensory stimulation in human subjects. *Proc.Natl.Acad.Sci.U.S.A*, 83(4):1140–1144.
- Francis, P. T. and Perry, E. K. (2007). Cholinergic and other neurotransmitter mechanisms in parkinson's disease, parkinson's disease dementia, and dementia with lewy bodies. *Mov Disord.*, 22 Suppl 17:S351–S357.
- Frankel, J. P., Lees, A. J., Kempster, P. A., and Stern, G. M. (1990). Subcutaneous apomorphine in the treatment of parkinson's disease. *J.Neurol.Neurosurg.Psychiatry*, 53(2):96–101.
- Frederick, J. M., Rayborn, M. E., Laties, A. M., Lam, D. M., and Hollyfield, J. G. (1982). Dopaminergic neurons in the human retina. *J.Comp Neurol.*, 210(1):65–79.
- Friston, K. J., Fletcher, P., Josephs, O., Holmes, A., Rugg, M. D., and Turner, R. (1998). Event-related fmri: characterizing differential responses. *Neuroimage.*, 7(1):30–40.
- Friston, K. J., Holmes, A. P., Poline, J. B., Grasby, P. J., Williams, S. C., Frackowiak, R. S., and Turner, R. (1995). Analysis of fmri time-series revisited. *Neuroimage.*, 2(1):45–53.
- Geerligs, L., Meppelink, A. M., Brouwer, W. H., and van Laar, T. (2009). The effects of apomorphine on visual perception in patients with parkinson disease and visual hallucinations: a pilot study. *Clin.Neuropharmacol.*, 32(5):266–268.
- Goetz, C. G., Pappert, E. J., Blasucci, L. M., Stebbins, G. T., Ling, Z. D., Nora, M. V., and Carvey, P. M. (1998). Intravenous levodopa in hallucinating parkinson's disease patients: high-dose challenge does not precipitate hallucinations. *Neurology*, 50(2):515–517.
- Goetz, C. G. and Stebbins, G. T. (1993). Risk factors for nursing home placement in advanced parkinson's disease. *Neurology*, 43(11):2227–2229.

- Goetz, C. G., Wu, J., Curgian, L. M., and Leurgans, S. (2005). Hallucinations and sleep disorders in pd: six-year prospective longitudinal study. *Neurology*, 64(1):81–86.
- Goodale, M. A., Westwood, D. A., and Milner, A. D. (2004). Two distinct modes of control for object-directed action. *Prog.Brain Res.*, 144:131–144.
- Grill-Spector, K. (2003). The neural basis of object perception. *Curr.Opin.Neurobiol.*, 13(2):159–166.
- Grill-Spector, K., Kushnir, T., Hendler, T., and Malach, R. (2000). The dynamics of object-selective activation correlate with recognition performance in humans. *Nat.Neurosci.*, 3(8):837–843.
- Groenewegen, H. J. (2009). The basal ganglia.
- Grossi, D., Trojano, L., Pellecchia, M. T., Amboni, M., Fragassi, N. A., and Barone, P. (2005). Frontal dysfunction contributes to the genesis of hallucinations in non-demented parkinsonian patients. *Int.J.Geriatr.Psychiatry*, 20(7):668–673.
- Hajee, M. E., March, W. F., Lazzaro, D. R., Wolintz, A. H., Shrier, E. M., Glazman, S., and Bodis-Wollner, I. G. (2009). Inner retinal layer thinning in parkinson disease. *Arch.Ophthalmol.*, 127(6):737–741.
- Harding, A. J., Broe, G. A., and Halliday, G. M. (2002). Visual hallucinations in lewy body disease relate to lewy bodies in the temporal lobe. *Brain*, 125(Pt 2):391–403.
- Harnois, C. and Di Paolo, T. (1990). Decreased dopamine in the retinas of patients with parkinson’s disease. *Invest Ophthalmol.Vis.Sci.*, 31(11):2473–2475.
- Harris, J. P., Calvert, J. E., and Phillipson, O. T. (1992). Processing of spatial contrast in peripheral vision in parkinson’s disease. *Brain*, 115 (Pt 5):1447–1457.
- Hattori, N., Shibasaki, H., Wheaton, L., Wu, T., Matsushashi, M., and Hallett, M. (2009). Discrete parieto-frontal functional connectivity related to grasping. *J.Neurophysiol.*, 101(3):1267–1282.
- Hazlett, E. A., Buchsbaum, M. S., Kemether, E., Bloom, R., Platholi, J., Brickman, A. M., Shihabuddin, L., Tang, C., and Byne, W. (2004). Abnormal glucose metabolism in the mediodorsal nucleus of the thalamus in schizophrenia. *Am.J.Psychiatry*, 161(2):305–314.

- Hendry, S. H. and Yoshioka, T. (1994). A neurochemically distinct third channel in the macaque dorsal lateral geniculate nucleus. *Science*, 264(5158):575–577.
- Hilker, R., Thomas, A. V., Klein, J. C., Weisenbach, S., Kalbe, E., Burghaus, L., Jacobs, A. H., Herholz, K., and Heiss, W. D. (2005). Dementia in parkinson disease: functional imaging of cholinergic and dopaminergic pathways. *Neurology*, 65(11):1716–1722.
- Holroyd, S., Currie, L., and Wooten, G. F. (2001). Prospective study of hallucinations and delusions in parkinson’s disease. *J.Neurol.Neurosurg.Psychiatry*, 70(6):734–738.
- Hopfinger, J. B. and West, V. M. (2006). Interactions between endogenous and exogenous attention on cortical visual processing. *Neuroimage.*, 31(2):774–789.
- Horowitz, T. S., Choi, W. Y., Horvitz, J. C., Cote, L. J., and Mangels, J. A. (2006). Visual search deficits in parkinson’s disease are attenuated by bottom-up target salience and top-down information. *Neuropsychologia*, 44(10):1962–1977.
- Huang, C., Mattis, P., Tang, C., Perrine, K., Carbon, M., and Eidelberg, D. (2007). Metabolic brain networks associated with cognitive function in parkinson’s disease. *Neuroimage.*, 34(2):714–723.
- Huettel, S. A., Song, A. W., and McCarthy, G. (2004). Functional magnetic resonance imaging.
- Ibarretxe-Bilbao, N., Ramirez-Ruiz, B., Junque, C., Marti, M. J., Valldeoriola, F., Bargallo, N., Juanes, S., and Tolosa, E. (2009). Differential progression of brain atrophy in parkinson disease with and without visual hallucinations. *J.Neurol.Neurosurg.Psychiatry*.
- Imamura, K., Wada-Isoe, K., Kitayama, M., and Nakashima, K. (2007). Executive dysfunction in non-demented parkinson’s disease patients with hallucinations. *Acta Neurol.Scand*.
- James, T. W., Humphrey, G. K., Gati, J. S., Menon, R. S., and Goodale, M. A. (2000). The effects of visual object priming on brain activation before and after recognition. *Current Biology*, 10(17):1017–1024.
- Jancke, L., Koeneke, S., Hoppe, A., Rominger, C., and Hanggi, J. (2009). The architecture of the golfer’s brain. *PLoS.One.*, 4(3).

- Jellinger, K. A. (2009a). A critical evaluation of current staging of alpha-synuclein pathology in lewy body disorders. *Biochim.Biophys.Acta*, 1792(7):730–740.
- Jellinger, K. A. (2009b). Formation and development of lewy pathology: a critical update. *J.Neurol.*, 256 Suppl 3:270–279.
- Johannsen, P., Jakobsen, J., Bruhn, P., Hansen, S. B., Gee, A., Stodkilde-Jorgensen, H., and Gjedde, A. (1997). Cortical sites of sustained and divided attention in normal elderly humans. *Neuroimage.*, 6(3):145–155.
- Kandel, E. (2000). *Principles of Neural Science*. McGraw-Hill.
- Kanwisher, N., McDermott, J., and Chun, M. M. (1997). The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J.Neurosci.*, 17(11):4302–4311.
- Kastner, S., De Weerd, P., Desimone, R., and Ungerleider, L. G. (1998). Mechanisms of directed attention in the human extrastriate cortex as revealed by functional mri. *Science*, 282(5386):108–111.
- Kastner, S., O'Connor, D. H., Fukui, M. M., Fehd, H. M., Herwig, U., and Pinsk, M. A. (2004). Functional imaging of the human lateral geniculate nucleus and pulvinar. *J.Neurophysiol.*, 91(1):438–448.
- Kataoka, H., Furiya, Y., Morikawa, M., Ueno, S., and Inoue, M. (2008). Increased temporal blood flow associated with visual hallucinations in parkinson's disease with dementia. *Mov Disord.*, 23(3):464–465.
- Kempster, P. A., Williams, D. R., Selikhova, M., Holton, J., Revesz, T., and Lees, A. J. (2007). Patterns of levodopa response in parkinson's disease: a clinico-pathological study. *Brain*, 130(Pt 8):2123–2128.
- Klein, R. C., de Jong, B. M., de Vries, J. J., and Leenders, K. L. (2005). Direct comparison between regional cerebral metabolism in progressive supranuclear palsy and parkinson's disease. *Mov Disord.*, 20(8):1021–1030.
- Kleinschmidt, A., Buchel, C., Hutton, C., Friston, K. J., and Frackowiak, R. S. J. (2002). The neural structures expressing perceptual hysteresis in visual letter recognition. *Neuron*, 34(4):659–666.
- Kobayashi, Y. and Isa, T. (2002). Sensory-motor gating and cognitive control by the brainstem cholinergic system. *Neural Netw.*, 15(4-6):731–741.

- Koerts, J., Borg, M. A., Meppelink, A. M., Leenders, K. L., van Beilen, M., and van Laar, T. (2010). Attentional and perceptual impairments in parkinson's disease with visual hallucinations. *Parkinsonism.Relat Disord.*
- Ku, J., Kim, J. J., Jung, Y. C., Park, I. H., Lee, H., Han, K., Yoon, K. J., Kim, I. Y., and Kim, S. I. (2008). Brain mechanisms involved in processing unreal perceptions. *Neuroimage.*, 43(4):793–800.
- Kveraga, K., Boshyan, J., and Bar, M. (2007). Magnocellular projections as the trigger of top-down facilitation in recognition. *J.Neurosci.*, 27(48):13232–13240.
- La Vega-Talbot, M., Duchowny, M., and Jayakar, P. (2006). Orbitofrontal seizures presenting with ictal visual hallucinations and interictal psychosis. *Pediatr.Neurol.*, 35(1):78–81.
- Laatu, S., Revonsuo, A., Pihko, L., Portin, R., and Rinne, J. O. (2004). Visual object recognition deficits in early parkinson's disease. *Parkinsonism.Relat Disord.*, 10(4):227–233.
- LaBerge, D. and Buchsbaum, M. S. (1990). Positron emission tomographic measurements of pulvinal activity during an attention task. *J.Neurosci.*, 10(2):613–619.
- Lang, A. E. and Lozano, A. M. (1998). Parkinson's disease. first of two parts. *N.Engl.J.Med.*, 339(15):1044–1053.
- Leh, S. E., Chakravarty, M. M., and Ptito, A. (2008). The connectivity of the human pulvinal: a diffusion tensor imaging tractography study. *Int.J.Biomed.Imaging*, 2008.
- Lieb, K., Brucker, S., Bach, M., Els, T., Lucking, C. H., and Greenlee, M. W. (1999). Impairment in preattentive visual processing in patients with parkinson's disease. *Brain*, 122 (Pt 2):303–313.
- Litvinenko, I. V., Odinak, M. M., Mogil'naya, V. I., and Perstnev, S. V. (2010). Use of memantine (akatinol) for the correction of cognitive impairments in parkinson's disease complicated by dementia. *Neurosci.Behav.Physiol.*, 40(2):149–155.
- Liu, T., Pestilli, F., and Carrasco, M. (2005). Transient attention enhances perceptual performance and fmri response in human visual cortex. *Neuron*, 45(3):469–477.

- Lo, R. Y., Tanner, C. M., Albers, K. B., Leimpeter, A. D., Fross, R. D., Bernstein, A. L., McGuire, V., Quesenberry, C. P., Nelson, L. M., and Van Den Eeden, S. K. (2009). Clinical features in early parkinson disease and survival. *Arch.Neurol.*, 66(11):1353–1358.
- Ma, Y., Tang, C., Spetsieris, P. G., Dhawan, V., and Eidelberg, D. (2007). Abnormal metabolic network activity in parkinson’s disease: test-retest reproducibility. *J.Cereb.Blood Flow Metab*, 27(3):597–605.
- Malach, R., Reppas, J. B., Benson, R. R., Kwong, K. K., Jiang, H., Kennedy, W. A., Ledden, P. J., Brady, T. J., Rosen, B. R., and Tootell, R. B. (1995). Object-related activity revealed by functional magnetic resonance imaging in human occipital cortex. *Proc.Natl.Acad.Sci.U.S.A*, 92(18):8135–8139.
- Manford, M. and Andermann, F. (1998). Complex visual hallucinations. clinical and neurobiological insights. *Brain*, 121 (Pt 10):1819–1840.
- Manganelli, F., Vitale, C., Santangelo, G., Pisciotta, C., Iodice, R., Cozzolino, A., Dubbioso, R., Picillo, M., Barone, P., and Santoro, L. (2009). Functional involvement of central cholinergic circuits and visual hallucinations in parkinson’s disease. *Brain*, 132(Pt 9):2350–2355.
- Marinus, J., Visser, M., Stiggelbout, A. M., Rabey, J. M., Martinez-Martin, P., Bonuccelli, U., Kraus, P. H., and van Hilten, J. J. (2004). A short scale for the assessment of motor impairments and disabilities in parkinson’s disease: the spes/scopa. *J.Neurol.Neurosurg.Psychiatry*, 75(3):388–395.
- Marinus, J., Visser, M., Verwey, N. A., Verhey, F. R., Middelkoop, H. A., Stiggelbout, A. M., and van Hilten, J. J. (2003). Assessment of cognition in parkinson’s disease. *Neurology*, 61(9):1222–1228.
- Matsui, H., Nishinaka, K., Oda, M., Hara, N., Komatsu, K., Kubori, T., and Udaka, F. (2006a). Hypoperfusion of the visual pathway in parkinsonian patients with visual hallucinations. *Mov Disord.*, 21(12):2140–2144.
- Matsui, H., Udaka, F., Tamura, A., Oda, M., Kubori, T., Nishinaka, K., and Kameyama, M. (2006b). Impaired visual acuity as a risk factor for visual hallucinations in parkinson’s disease. *J.Geriater.Psychiatry Neurol.*, 19(1):36–40.
- Mayo, J. P. (2009). Intrathalamic mechanisms of visual attention. *J.Neurophysiol.*, 101(3):1123–1125.

- McAlonan, K., Cavanaugh, J., and Wurtz, R. H. (2008). Guarding the gateway to cortex with attention in visual thalamus. *Nature*, 456(7220):391–394.
- Meppelink, A. M., de Jong, B. M., Renken, R., Leenders, K. L., Cornelissen, F. W., and van Laar, T. (2009). Impaired visual processing preceding image recognition in parkinson’s disease patients with visual hallucinations. *Brain*, 132(Pt 11):2980–2993.
- Meppelink, A. M., de Jong, B. M., van der Hoeven, J. H., and van Laar, T. (2010). Lasting visual hallucinations in visual deprivation; fmri correlates and the influence of rtms. *J.Neurol.Neurosurg.Psychiatry*.
- Meppelink, A. M., Koerts, J., Borg, M., Leenders, K. L., and van Laar, T. (2008). Visual object recognition and attention in parkinson’s disease patients with visual hallucinations. *Mov Disord*.
- Merabet, L. B., Kobayashi, M., Barton, J., and Pascual-Leone, A. (2003). Suppression of complex visual hallucinatory experiences by occipital transcranial magnetic stimulation: a case report. *Neurocase.*, 9(5):436–440.
- Merims, D., Shabtai, H., Korczyn, A. D., Peretz, C., Weizman, N., and Giladi, N. (2004). Antiparkinsonian medication is not a risk factor for the development of hallucinations in parkinson’s disease. *J.Neural Transm.*, 111(10-11):1447–1453.
- Mesulam, M. M. (2004). The cholinergic innervation of the human cerebral cortex. *Prog.Brain Res.*, 145:67–78.
- Mesulam, M. M. and Geula, C. (1994). Chemoarchitectonics of axonal and perikaryal acetylcholinesterase along information processing systems of the human cerebral cortex. *Brain Res.Bull.*, 33(2):137–153.
- Middleton, F. A. and Strick, P. L. (1996). The temporal lobe is a target of output from the basal ganglia. *Proc.Natl.Acad.Sci.U.S.A*, 93(16):8683–8687.
- Mink, J. W. (1996). The basal ganglia: focused selection and inhibition of competing motor programs. *Prog.Neurobiol.*, 50(4):381–425.
- Morgante, L., Basile, G., Epifanio, A., Spina, E., Antonini, A., Stocchi, F., Di Rosa, E., Martino, G., Marconi, R., La Spina, P., Nicita-Mauro, V., and Di Rosa, A. E. (2004). Continuous apomorphine infusion (cai) and neuropsychiatric disorders in patients with advanced parkinson’s disease: a follow-up of two years. *Arch.Gerontol.Geriatr.Suppl*, (9):291–296.

- Mosimann, U. P., Rowan, E. N., Partington, C. E., Collerton, D., Littlewood, E., O'Brien, J. T., Burn, D. J., and McKeith, I. G. (2006). Characteristics of visual hallucinations in parkinson disease dementia and dementia with lewy bodies. *American Journal of Geriatric Psychiatry*, 14(2):153–160.
- Muller, T., Benz, S., and Przuntek, H. (2002). Apomorphine delays simple reaction time in parkinsonian patients. *Parkinsonism.Relat Disord.*, 8(5):357–360.
- Muslimovic, D., Post, B., Speelman, J. D., and Schmand, B. (2005). Cognitive profile of patients with newly diagnosed parkinson disease. *Neurology*, 65(8):1239–1245.
- Nagano-Saito, A., Washimi, Y., Arahata, Y., Iwai, K., Kawatsu, S., Ito, K., Nakamura, A., Abe, Y., Yamada, T., Kato, T., and Kachi, T. (2004). Visual hallucination in parkinson's disease with fdg pet. *Mov Disord.*, 19(7):801–806.
- Nagano-Saito, A., Washimi, Y., Arahata, Y., Kachi, T., Lerch, J. P., Evans, A. C., Dagher, A., and Ito, K. (2005). Cerebral atrophy and its relation to cognitive impairment in parkinson disease. *Neurology*, 64(2):224–229.
- Newell, F. N., Wallraven, C., and Huber, S. (2004). The role of characteristic motion in object categorization. *J. Vis.*, 4(2):118–129.
- Oishi, N., Udaka, F., Kameyama, M., Sawamoto, N., Hashikawa, K., and Fukuyama, H. (2005). Regional cerebral blood flow in parkinson disease with nonpsychotic visual hallucinations. *Neurology*, 65(11):1708–1715.
- Okada, K., Suyama, N., Oguro, H., Yamaguchi, S., and Kobayashi, S. (1999). Medication-induced hallucination and cerebral blood flow in parkinson's disease. *J.Neurol.*, 246(5):365–368.
- Parkkinen, L., Pirttila, T., and Alafuzoff, I. (2008). Applicability of current staging/categorization of alpha-synuclein pathology and their clinical relevance. *Acta Neuropathol.*, 115(4):399–407.
- Pauling, L. and Coryell, C. D. (1936). The magnetic properties and structure of hemoglobin, oxyhemoglobin and carbonmonoxyhemoglobin. *Proc.Natl.Acad.Sci.U.S.A*, 22(4):210–216.
- Pelli, D. G. (1997). The videotoolbox software for visual psychophysics: transforming numbers into movies. *Spat. Vis.*, 10(4):437–442.

- Pereira, J. B., Junque, C., Marti, M. J., Ramirez-Ruiz, B., Bargallo, N., and Tolosa, E. (2009). Neuroanatomical substrate of visuospatial and visuoperceptual impairment in parkinson's disease. *Mov Disord.*, 24(8):1193–1199.
- Perry, E. K., Curtis, M., Dick, D. J., Candy, J. M., Attack, J. R., Bloxham, C. A., Blessed, G., Fairbairn, A., Tomlinson, B. E., and Perry, R. H. (1985). Cholinergic correlates of cognitive impairment in parkinson's disease: comparisons with alzheimer's disease. *J.Neurol.Neurosurg.Psychiatry*, 48(5):413–421.
- Perry, E. K. and Perry, R. H. (1995). Acetylcholine and hallucinations: disease-related compared to drug-induced alterations in human consciousness. *Brain Cogn*, 28(3):240–258.
- Pieri, V., Diederich, N. J., Raman, R., and Goetz, C. G. (2000). Decreased color discrimination and contrast sensitivity in parkinson's disease. *J.Neurol.Sci.*, 172(1):7–11.
- Pitzalis, S., Sereno, M. I., Committeri, G., Fattori, P., Galati, G., Patria, F., and Galletti, C. (2010). Human v6: the medial motion area. *Cereb.Cortex*, 20(2):411–424.
- Playford, E. D., Jenkins, I. H., Passingham, R. E., Nutt, J., Frackowiak, R. S., and Brooks, D. J. (1992). Impaired mesial frontal and putamen activation in parkinson's disease: a positron emission tomography study. *Ann.Neurol.*, 32(2):151–161.
- Quene, H. and van den Bergh, H. (2004). On multi-level modeling of data from repeated measures designs: A tutorial. *Speech Communication*, 43:103–121.
- Ramirez-Ruiz, B., Junque, C., Marti, M. J., Valldeoriola, F., and Tolosa, E. (2006). Neuropsychological deficits in parkinson's disease patients with visual hallucinations. *Mov Disord.*, 21(9):1483–1487.
- Ramirez-Ruiz, B., Junque, C., Marti, M. J., Valldeoriola, F., and Tolosa, E. (2007a). Cognitive changes in parkinson's disease patients with visual hallucinations. *Dement.Geriatr.Cogn Disord.*, 23(5):281–288.
- Ramirez-Ruiz, B., Marti, M. J., Tolosa, E., Falcon, C., Bargallo, N., Valldeoriola, F., and Junque, C. (2008). Brain response to complex visual stimuli in parkinson's patients with hallucinations: A functional magnetic resonance imaging study. *Mov Disord.*

- Ramirez-Ruiz, B., Marti, M. J., Tolosa, E., Gimenez, M., Bargallo, N., Valldeoriola, F., and Junque, C. (2007b). Cerebral atrophy in parkinson's disease patients with visual hallucinations. *Eur.J.Neurol.*, 14(7):750–756.
- Redgrave, P., Prescott, T. J., and Gurney, K. (1999). The basal ganglia: a vertebrate solution to the selection problem? *Neuroscience*, 89(4):1009–1023.
- Reinders, A. A. T. S., Glascher, J., de Jong, J. R., Willemsen, A. T. M., den Boer, J. A., and Buchel, C. (2006). Detecting fearful and neutral faces: Bold latency differences in amygdala-hippocampal junction. *Neuroimage*, 33(2):805–814.
- Robinson, D. L. and Petersen, S. E. (1992). The pulvinar and visual salience. *Trends Neurosci.*, 15(4):127–132.
- Roselli, F., Pisciotta, N. M., Perneckzy, R., Pennelli, M., Aniello, M. S., De Caro, M. F., Ferrannini, E., Tartaglione, B., Defazio, G., Rubini, G., and Livrea, P. (2009). Severity of neuropsychiatric symptoms and dopamine transporter levels in dementia with lewy bodies: a 123i-fp-cit spect study. *Mov Disord.*, 24(14):2097–2103.
- Rye, D. B. (1997). Contributions of the pedunculopontine region to normal and altered rem sleep. *Sleep*, 20(9):757–788.
- Saalmann, Y. B. and Kastner, S. (2009). Gain control in the visual thalamus during perception and cognition. *Curr.Opin.Neurobiol.*, 19(4):408–414.
- Sanchez-Ramos, J. R., Ortoll, R., and Paulson, G. W. (1996). Visual hallucinations associated with parkinson disease. *Arch.Neurol.*, 53(12):1265–1268.
- Santangelo, G., Trojano, L., Vitale, C., Ianniciello, M., Amboni, M., Grossi, D., and Barone, P. (2007). A neuropsychological longitudinal study in parkinson's patients with and without hallucinations. *Mov Disord.*, 22(16):2418–2425.
- Sarter, M., Givens, B., and Bruno, J. P. (2001). The cognitive neuroscience of sustained attention: where top-down meets bottom-up. *Brain Res.Brain Res.Rev.*, 35(2):146–160.
- Schapira, A. H., Mann, V. M., Cooper, J. M., Krige, D., Jenner, P. J., and Marsden, C. D. (1992). Mitochondrial function in parkinson's disease. the royal kings and queens parkinson's disease research group. *Ann.Neurol.*, 32 Suppl:S116–S124.

- Schmeichel, A. M., Buchhalter, L. C., Low, P. A., Parisi, J. E., Boeve, B. W., Sandroni, P., and Benarroch, E. E. (2008). Mesopontine cholinergic neuron involvement in lewy body dementia and multiple system atrophy. *Neurology*, 70(5):368–373.
- Schneider, E., Fischer, P. A., Jacobi, P., Becker, H., and Hacker, H. (1979). The significance of cerebral atrophy for the symptomatology of parkinson's disease. *J.Neurol.Sci.*, 42(2):187–197.
- Schotte, A., Janssen, P. F., Megens, A. A., and Leysen, J. E. (1993). Occupancy of central neurotransmitter receptors by risperidone, clozapine and haloperidol, measured ex vivo by quantitative autoradiography. *Brain Res.*, 631(2):191–202.
- Schufried, G. and Midling (1992). *Test Management Program: Reaction Test - S7*.
- Seger, C. A. and Cincotta, C. M. (2005). The roles of the caudate nucleus in human classification learning. *J.Neurosci.*, 25(11):2941–2951.
- Selby, G. (1968). Cerebral atrophy in parkinsonism. *J.Neurol.Sci.*, 6(3):517–559.
- Serences, J. T. and Yantis, S. (2006). Selective visual attention and perceptual coherence. *Trends Cogn Sci.*, 10(1):38–45.
- Shimada, H., Hirano, S., Shinotoh, H., Aotsuka, A., Sato, K., Tanaka, N., Ota, T., Asahina, M., Fukushi, K., Kuwabara, S., Hattori, T., Suhara, T., and Irie, T. (2009). Mapping of brain acetylcholinesterase alterations in lewy body disease by pet. *Neurology*.
- Silbersweig, D. A., Stern, E., Frith, C., Cahill, C., Holmes, A., Grootoink, S., Seaward, J., McKenna, P., Chua, S. E., Schnorr, L., and . (1995). A functional neuroanatomy of hallucinations in schizophrenia. *Nature*, 378(6553):176–179.
- Silva, M. F., Faria, P., Regateiro, F. S., Forjaz, V., Januario, C., Freire, A., and Castelo-Branco, M. (2005). Independent patterns of damage within magno-, parvo- and koniocellular pathways in parkinson's disease. *Brain*, 128(Pt 10):2260–2271.
- Simons, J. S., Henson, R. N., Gilbert, S. J., and Fletcher, P. C. (2008). Separable forms of reality monitoring supported by anterior prefrontal cortex. *J.Cogn Neurosci.*, 20(3):447–457.

- Sincich, L. C., Park, K. F., Wohlgenuth, M. J., and Horton, J. C. (2004). Bypassing v1: a direct geniculate input to area mt. *Nat.Neurosci.*, 7(10):1123–1128.
- Smith, A. T., Cotton, P. L., Bruno, A., and Moutsiana, C. (2009). Dissociating vision and visual attention in the human pulvinar. *J.Neurophysiol.*, 101(2):917–925.
- Snow, J. C., Allen, H. A., Rafal, R. D., and Humphreys, G. W. (2009). Impaired attentional selection following lesions to human pulvinar: evidence for homology between human and monkey. *Proc.Natl.Acad.Sci.U.S.A.*, 106(10):4054–4059.
- Sossi, V., Fuente-Fernandez, R., Schulzer, M., Troiano, A. R., Ruth, T. J., and Stoessl, A. J. (2007). Dopamine transporter relation to dopamine turnover in parkinson’s disease: a positron emission tomography study. *Ann.Neurol.*, 62(5):468–474.
- Stebbins, G. T., Goetz, C. G., Carrillo, M. C., Bangen, K. J., Turner, D. A., Glover, G. H., and Gabrieli, J. D. (2004). Altered cortical visual processing in pd with hallucinations: an fmri study. *Neurology*, 63(8):1409–1416.
- Steriade, M. (2004). Acetylcholine systems and rhythmic activities during the waking–sleep cycle. *Prog.Brain Res.*, 145:179–196.
- Summerfield, C., Junque, C., Tolosa, E., Salgado-Pineda, P., Gomez-Anson, B., Marti, M. J., Pastor, P., Ramirez-Ruiz, B., and Mercader, J. (2005). Structural brain changes in parkinson disease with dementia: a voxel-based morphometry study. *Arch.Neurol.*, 62(2):281–285.
- Summerfield, C. and Koechlin, E. (2008). A neural representation of prior information during perceptual inference. *Neuron*, 59(2):336–347.
- Tagliati, M., Bodis-Wollner, I., and Yahr, M. D. (1996). The pattern electroretinogram in parkinson’s disease reveals lack of retinal spatial tuning. *Electroencephalogr.Clin.Neurophysiol.*, 100(1):1–11.
- Tan, E. K. and Skipper, L. M. (2007). Pathogenic mutations in parkinson disease. *Hum.Mutat.*, 28(7):641–653.
- Teunisse, R. J., Cruysberg, J. R., Hoefnagels, W. H., Verbeek, A. L., and Zitman, F. G. (1996). Visual hallucinations in psychologically normal people: Charles bonnet’s syndrome. *Lancet*, 347(9004):794–797.

- Thannickal, T. C., Lai, Y. Y., and Siegel, J. M. (2007). Hypocretin (orexin) cell loss in parkinson's disease. *Brain*, 130(Pt 6):1586–1595.
- Thannickal, T. C., Moore, R. Y., Nienhuis, R., Ramanathan, L., Gulyani, S., Aldrich, M., Cornford, M., and Siegel, J. M. (2000). Reduced number of hypocretin neurons in human narcolepsy. *Neuron*, 27(3):469–474.
- Thomas, C., Moya, L., Avidan, G., Humphreys, K., Jung, K. J., Peterson, M. A., and Behrmann, M. (2008). Reduction in white matter connectivity, revealed by diffusion tensor imaging, may account for age-related changes in face perception. *J. Cogn. Neurosci.*, 20(2):268–284.
- Tir, M., Delmaire, C., le Thuc, V., Duhamel, A., Destee, A., Pruvo, J. P., and Defebvre, L. (2009). Motor-related circuit dysfunction in msa-p: Usefulness of combined whole-brain imaging analysis. *Mov Disord.*, 24(6):863–870.
- True, S. (2003). Climbing the cortical ladder from sensation to perception. *Trends Cogn. Sci.*, 7(11):469–471.
- Ungerleider, L. G. and Haxby, J. V. (1994). 'what' and 'where' in the human brain. *Curr. Opin. Neurobiol.*, 4(2):157–165.
- Verleden, S., Vingerhoets, G., and Santens, P. (2007). Heterogeneity of cognitive dysfunction in parkinson's disease: a cohort study. *Eur. Neurol.*, 58(1):34–40.
- von Campenhausen, S., Bornschein, B., Wick, R., Botzel, K., Sampaio, C., Poewe, W., Oertel, W., Siebert, U., Berger, K., and Dodel, R. (2005). Prevalence and incidence of parkinson's disease in europe. *Eur. Neuropsychopharmacol.*, 15(4):473–490.
- Walker, M. P., Ayre, G. A., Cummings, J. L., Wesnes, K., McKeith, I. G., O'Brien, J. T., and Ballard, C. G. (2000). Quantifying fluctuation in dementia with lewy bodies, alzheimer's disease, and vascular dementia. *Neurology*, 54(8):1616–1625.
- Warrington (1991). *The visual object and space perception battery*. Thames Valley Test Company, Bury St. Edmonds UK.
- Wesnes, K. A., McKeith, I., Edgar, C., Emre, M., and Lane, R. (2005). Benefits of rivastigmine on attention in dementia associated with parkinson disease. *Neurology*, 65(10):1654–1656.

- Westerink, B. H. (2002). Can antipsychotic drugs be classified by their effects on a particular group of dopamine neurons in the brain? *Eur.J.Pharmacol.*, 455(1):1–18.
- Wheaton, L. A. and Hallett, M. (2007). Ideomotor apraxia: a review. *J.Neurol.Sci.*, 260(1-2):1–10.
- Williams, D. R. and Lees, A. J. (2005). Visual hallucinations in the diagnosis of idiopathic parkinson's disease: a retrospective autopsy study. *Lancet Neurol.*, 4(10):605–610.
- Wolters, E., van Laar, T., and Berendse, H. (2007). Parkinsonism and related disorders.
- Zeki, S. (2001). Localization and globalization in conscious vision. *Annu.Rev.Neurosci.*, 24:57–86.
- Zweig, R. M., Jankel, W. R., Hedreen, J. C., Mayeux, R., and Price, D. L. (1989). The pedunclopontine nucleus in parkinson's disease. *Ann.Neurol.*, 26(1):41–46.

