

University of Groningen

Parental divorce and adolescent excessive drinking

Tomcikova, Zuzana

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Tomcikova, Z. (2011). *Parental divorce and adolescent excessive drinking: role of parent-adolescent relationship and other social and psychosocial factors*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Characteristics of adolescent excessive drinkers compared with consumers and abstainers

Zuzana Tomcikova, Andrea Madarasova Geckova, Jitse P. van Dijk,
Sijmen A. Reijneveld

Published: Drug and Alcohol Review 2010
DOI: 10.1111/j.1465-3362.2010.00209.x.

Abstract

This study aimed at comparing adolescent abstainers, consumers and excessive drinkers in terms of family characteristics (structure of family, socioeconomic factors), perceived social support, personality characteristics (extraversion, self-esteem, aggression) and well-being. Cross-sectional data were obtained from 3694 elementary school students in the 8th and 9th grades from several cities in Slovakia (mean age 14.5, 49.0% men; response rate 93%). Respondents completed questions on the use of alcohol and on family structure (parental divorce), the socioeconomic position of the family (parents' education and family affluence), perceived social support, extraversion, self-esteem, aggression and psychological well-being. They were split into three groups based on the pattern of alcohol use – abstainers, consumers and excessive drinkers (i.e. being drunk at least once during the past 4 weeks). The results showed significant differences between abstainers, consumers and excessive drinkers in almost every characteristic explored. A risky pattern of alcohol consumption occurs more frequently among adolescents who have divorced parents, higher socioeconomic position, higher scores for perceived social support from friends, extraversion, negative self-esteem and aggression, and lower scores for social support from family and for well-being. A risky pattern of alcohol consumption is more likely among relatively easily identifiable groups of adolescents from high socioeconomic position and divorced families. Their personalities and social networks have characteristics that could be accommodated in preventive interventions as well.

Introduction

Excessive drinking is a relatively common behaviour, particularly among adolescents, and also has become a major public health concern. The results of the European school survey on alcohol and other drugs (ESPAD) (Hibbel, 2004) indicate that more than half of all students have consumed alcohol at the age of 13 years or younger. The proportion of students who reported having been drunk at the age of 13 or younger varies greatly across countries (Hibbel, 2004). Slovak participants in this study placed approximately in the middle: 27% of boys and 17% of girls reported having been drunk at this age. A wide variety of factors that may play a role as possible risk factors of hazardous alcohol drinking in adolescence could be divided into three groups: (1) factors related to family and social background of the adolescent—within this group we distinguish structural characteristics (e.g. structure of family) and psychosocial characteristics (e.g. social support); (2) individual personality factors; and (3) factors proximal to behaviour, such as immediate intentions, reasons or expectations related to alcohol drinking, but also one's immediate condition (e.g. well-being) (Petraitis et al., 1995). Besides these groups of factors genetic and biological factors play an important role as well (Jessor, 1991).

Family factors

Undoubtedly, family is one of the most significant contexts that determine the development of children and adolescents. Basic patterns of behaviour are formed in the family, as well as values, norms and attitudes. When a family is not complete, this may lead to developmental disturbances, including risk behaviour (Eickhoff, 2001; Currie et al., 2004). Many studies (Miller, 1997; Blum et al., 2000; Ledoux et al., 2002; Kuntsche & Kuendig, 2006; Fisher et al., 2007) have found that living in a single-parent family increases the risk of adolescent alcohol use. This may be explained by the decreased parental control in one-parent families (Freeman & Newland, 2002; Marsden et al., 2005), by the fact that single parents tend to have fewer financial resources and to suffer from greater social isolation (Griffin et al., 2000) or through the immediate consequences of the disruption of the family structure (divorce) on adolescents (e.g. lowered self-esteem, increased levels of depression and anxiety) (Storksens et al., 2005). The socioeconomic position (SEP) of the family seems to play a role in adolescent alcohol use as well, although contradictory results about this association could be found. On one hand, the more financial resources are available to adolescents, the higher the rates of excessive drinking; but on the other hand, low levels of education are associated with more excessive drinking (Kuntsche et al., 2004).

Social support

Besides the above-mentioned effects of the family environment, social support from the family is an important buffer against stressful life events and plays a considerable role in coping with demanding life situations (Bal et al., 2003; Geckova et al., 2003; Murberg & Bru, 2004). This also holds true, to a lesser degree, for social support from other sources. Concerning excessive drinking, several studies have found that less parental support (support from family) is associated with greater alcohol use in adolescents (Stice et al., 1993; Shucksmith et al., 1997; Windle & Miller-Tutzauer, 1997), although not all studies have confirmed this association (Lifrak et al., 1997). Besides the social support gained from the family, adolescents can receive support from peers (friends) and significant others as well (Blumenthal et al., 1987). Higher perceived social support from peer friends seems to be associated with greater alcohol use (Engels & Ter Bogt, 2001).

Personality factors

Although many studies have confirmed the impact of social and environmental factors on alcohol use in adolescents, there is a growing body of evidence that personality highly determines someone's vulnerability to excessive drinking (Merenakk et al., 2003). One personality trait that is of great importance in explaining hazardous drinking is extraversion, which is defined as gregariousness and sociability (Kutsche et al., 2006). Some studies have found that people scoring higher in extraversion are at higher risk to drink more frequently or more hazardously (Martsh & Miller, 1997; Merenakk et al., 2003) and to have more tolerant attitudes towards alcohol use (Francis, 1996). Another important personality factor regarding excessive drinking is self-esteem, typically defined as one's overall sense of worthiness as a person (Rosenberg, 1979). The role of self-esteem in alcohol use among adolescents is not clear. On one hand, it is known that positive self-esteem may function as a buffer against deviant behaviour by facilitating better psychological adjustment (Schweitzer et al., 1992). On the other hand, here are some inconsistent results from studies showing both abstainers and high/ excessive users having higher levels of self-esteem (Pandina & Schuele, 1983; Lifrak et al., 1997; Freeman & Newland, 2002). A final personality trait that plays a role in excessive drinking is aggressiveness. Aggressive behaviour is, on one hand, a common result of problematic drinking (Swahn & Donovan, 2004), but on the other hand, aggressive tendencies in behaviour also predict excessive alcohol use (Gerra et al., 2004).

Well-being

It is well-known that problem drinking is associated with lower states of psychological well-being (Pitkanen, 1999), meaning that in some cases,

drinking alcohol (and particularly hazardous drinking) might function as a coping mechanism, as an example of an avoidance strategy (Unger et al., 2001; Catanzaro & Laurent, 2004; Marsden et al., 2005), especially among women (Pitkanen, 1999). Research indicates that each pattern or stage of drinking may have its own predictors (Power et al., 2005). The movement from abstaining to 'non-risk' drinking may thus be influenced by different factors from the movement from 'non-risk' to 'risk' drinking. Therefore, we decided to examine three patterns of alcohol use in adolescence – abstainers, consumers ('non-risk') and excessive drinkers ('risk').

The aim of our study was to compare adolescent abstainers, consumers and excessive drinkers with regard to family characteristics (socioeconomic factors, structure of family), perceived social support, personality characteristics (extraversion, self-esteem, aggression) and well-being.

Methods

Sample and procedure

The total sample of our study consisted of 3694 elementary school students from 8th and 9th grades from three cities in Slovakia Bratislava (600 000 inhabitants, Western Slovakia), Zilina (156 000 inhabitants, Northern Slovakia) and Kosice (240 000 inhabitants, Eastern Slovakia), and several smaller towns in the Kosice region (10 000–40 000 inhabitants). The age range was from 13 to 16; mean age was 14.5 (SD=0.5). The sample was randomly selected after stratification by region and gender (49.0% men, 51.0% women). The representation of the regions was as follows: 24.6% of the participants lived in Bratislava, 21.3% in Zilina, 32.1% in Kosice and 22.0% in several smaller towns in the Kosice region. This reflects the distribution of these types of areas across Slovakia, so that the sample can be considered to be representative for this country. Data were collected in autumn 2006 by a team of trained researchers and their assistants. The schools and classes were selected in every mentioned region or city randomly. We asked the directors of the schools for participation, and after their approval and approval from parents, we performed the data collection. Respondents filled in the questionnaire during two regular school lessons (45 min each) on a voluntary and anonymous basis, without the presence of the teacher. Response rate was 93.0%, with non-response due mainly to illness.

Measures

Questions concerning alcohol drinking: Drinking alcohol: 'How many times in the last 4 weeks have you drunk alcohol?' – I haven't drunk during the last 4 weeks/1–2 times/3 and more times. Being drunk: 'In the last 4 weeks have you been drunk?' – no/1–2 times/3 and more times. Both

questions were dichotomised, and based on the results; we divided the respondents into three groups: (1) total abstainers (had neither drunk alcohol nor been drunk); (2) consumers (had drunk alcohol without being drunk during last 4 weeks); and (3) excessive drinkers (had been drunk at least once during the last 4 weeks).

As we already mentioned in the Introduction, three main groups of factors that may play a role as possible risk factors of hazardous alcohol drinking in adolescence can be found in literature – social/family factors, personality factors and factors related to immediate condition of adolescents. In our study we explore following factors representing each of these groups:

Family structure: Respondents were asked to answer a question about whether their parents are divorced (legally), with the responses: no/yes, less than 12 months ago/yes, more than 12 months ago, but less than 3 years ago/yes, more than 3 years ago. A dichotomised variable was then constructed for the analysis – no/yes (any period since divorce).

SEP of the family: Two indicators of family SEP were used: the parents' education level and the family affluence. Parents' education level was defined as the highest level of education attained by the parents of the respondents: as high (university), medium (secondary school) or low (apprenticeship or primary school only). Family affluence was measured using the Family Affluence Scale (Currie et al., 2004), which consists of four questions concerning possession of a car and computer in the family, the family going on holiday (longer than 5 days) during the past year and respondents having their own room. Possible answers were: no/yes, one/yes, two or more for the question about the car; none/one/two/three or more for the question about the computer; no/once/ twice/three or more times for the question about the holiday and yes/no for the question about their own room. The score ranges from 0 to 7; the sum score was computed, and for the analysis we used a 3-point ordinal scale: low affluence (score = 0–3), middle affluence (score = 4–5) and high affluence (score = 6–7).

Perceived social support: Social support was measured using the Perceived Social Support Scale (Blumenthal et al., 1987), which is a 12-item self-reported questionnaire assessing perceived social support in three dimensions (from the family, friends and significant others). A 7-point Likert-type format was used ranging from totally disagree (1) to totally agree (7). The score for each of the 4-item subscales ranges from 4 to 28, with a higher score indicating a higher level of perceived social support. Internal reliability was satisfactory; Cronbach's alpha coefficient for the social support from family dimension was 0.91, for the social support from friends dimension 0.91 and for the social support from significant others dimension 0.85.

Extraversion: Extraversion was assessed with the Ten Item Personality Inventory (Gosling, 2003), a brief measure of the Big-Five personality

dimensions. For the purposes of this study, we used the extraversion dimension saturated by two items. A 7-point Likert-type format was used, ranging from strongly disagree (1) to strongly agree (7). The score ranges from 2 to 14, with a higher score indicating a higher level of extraversion. Cronbach's alpha coefficient was 0.31, and the mean inter-item correlation was 0.19. According to the guidelines of Briggs and Cheek (Clark & Watson, 1995; Parker et al., 2003), the mean inter-item correlation should range around 0.20, but not be less than 0.15 (Clark & Watson, 1995; Parker et al., 2003).

Self-esteem: Self-esteem was measured with the Rosenberg Self-Esteem Scale (Rosenberg, 1965), a widely used measure of global self-esteem in adolescents. The scale consists of 10 items rated on a 4-point scale, with responses ranging from strongly agree (4) to strongly disagree (1). The Rosenberg Self-Esteem Scale could be divided into an equal number of positively and negatively worded items measuring positive and negative self-esteem (Sarkova et al., 2006). Items were standardised and summed for the two subscales (positive and negative self-esteem), with the range of the sum score from 5 to 20 for each subscale. A higher score indicates higher positive or negative self-esteem. Cronbach's alpha coefficient for the positive self-esteem subscale was 0.73 and for the negative self-esteem subscale 0.64.

Aggression: Aggression was measured with the Aggression Questionnaire (Nakano, 2001), which is a 29-item self reported measure of four dimensions of aggression— physical aggression (nine items), verbal aggression (five items), anger (seven items) and hostility (eight items). We used a 5-point Likert-type score ranging from extremely uncharacteristic of me (1) to extremely characteristic of me (7), with a higher score indicating a higher level of aggression. The internal reliability coefficient for the physical aggression dimension was 0.80, for verbal aggression 0.64, for anger 0.64 and for hostility 0.75.

Psychological well-being: Psychological well-being was measured using the 12-item version of General Health Questionnaire (Goldberg & Williams, 1988). The General Health Questionnaire-12 is a widely used self-reported questionnaire assessing psychological illness. It has been divided into two subscales: social dysfunction and depression/anxiety. The factor 'depression/anxiety' consists of items about loss of sleep, being under strain, overcoming difficulties, feelings of unhappiness and a loss of self-confidence. Items about concentration, playing a useful part, making decisions, enjoying activities, facing up to problems and feeling happy are components of the 'social dysfunction' factor (Sarkova et al., 2006). We used a 4-point Likert score to score the items, which were then summed for the two subscales (depression/anxiety and social dysfunction), with the range of the sum score from 6 to 24 for each subscale. A higher score indicates higher levels of depression/anxiety and social dysfunction, thus

worse well-being. Cronbach's alpha coefficient for depression/anxiety was 0.82 and for social dysfunction 0.65.

Statistical analysis

Data were analysed using SPSS, version 14. We first assessed the characteristics of the sample. To compare adolescent abstainers, consumers and excessive drinkers regarding family characteristics, social support, personality traits and well-being we compared means and proportions, depending on the measurement scale. Differences were then tested using *F*-tests and χ^2 -tests, respectively. Additionally, post-hoc tests were computed to determine which means differ significantly. In the next step we compared the two most extreme groups—abstainers and excessive drinkers ($n = 2565$)—regarding consumption pattern, leaving out the consumers group. We analysed, using logistic regression, the degree to which excessive drinking was more likely among specific groups of adolescents. We only included characteristics that showed statistically significant differences in the bivariate analyses. Five models were constructed, all adjusted for gender. In the first model we analysed the effect of family characteristics that were significant in previous analyses (affluence and divorce). In the second step we analysed the effect of perceived social support from family and friends. In the third model we analysed the effect of the personality characteristics (extraversion, self-esteem and aggression) and in the fourth we analysed the effect of well-being. In the last model we analysed the effect of all characteristics that were significant in previous steps, simultaneously. To acquire the information on the group of consumers as well, we repeated the analyses comparing them with abstainers, constructing the same five models. Because the data were collected in entire school classes, a clustering of the students' outcomes per class might affect our findings. To account for this clustering, we performed all binary logistic analyses using MIWin 2.02 (Rasbash et al., 2005).

Results

Table 3.1 shows the differences between the three groups in social support, extraversion, self-esteem, aggression and psychological well-being. The higher the score in aggression, extraversion, perceived social support from friends and negative self-esteem, and the lower the scoring in social support from family and well-being, the more risky the pattern of alcohol consumption. Furthermore, Table 3.1 shows the proportion of highest family education, family affluence and family structure in the three explored groups. Adolescents from divorced families and those from families with higher affluence are significantly more likely to be excessive drinkers.

The results of multilevel logistic regression comparing the groups of abstainers and excessive drinkers are shown in Table 3.2. In the first four models we analysed separately the effect of four groups of factors (family characteristics, social support, personality characteristics, and well-being) on excessive drinking. Low family affluence, parental divorce, social support from family, social support from friends, extraversion, negative self-esteem, physical aggression, anger, hostility, depression/anxiety and social dysfunction all showed to have an effect on the probability of excessive drinking. In the final model we analysed the effect of all these significant characteristics together. All of them except negative self-esteem, anger and depression/anxiety remained significant. In general, the multilevel analyses showed a significant clustering of the students' outcomes per class, as shown by the random variances that are indicated in the bottom row of Table 3.2. This clustering hardly affected the estimates concerned, however. For instance, the odds ratio (95% confidence interval) for the effect of social support from family in the final model 5 in Table 3.2 was 0.93 (0.90–0.96), compared with 0.93 (0.91–0.96) for the ordinary logistic regression. The additional analyses comparing the group of abstainers with the group of consumers showed that most of the variables that were associated with excessive drinking were associated with consuming, although associations were mostly slightly weaker. One important exception was parental divorce, however. This had hardly any association with consuming, whereas it had with excessive drinking (both compared with abstaining).

Table 3.1 Family factors, social support, personality factors and well-being of adolescents, split by drinking behaviour

	Descriptive ^{a)}			Chi square / F-value	Differences Post hoc ^{b)}
	1 abstainers	2 consumers	3 excessive drinkers		
Highest educ. of parents ^{a)}					
low	362 (20)	235 (21)	101 (16)	12.0 ^{ns}	
medium	984 (53)	579 (51)	347 (53)		
high	462 (25)	296 (26)	193(30)		
Family affluence ^{a)}					
low	787 (43)	406 (36)	230 (36)	21.3 ^{***}	
medium	740 (41)	485 (44)	277 (43)		
high	292 (16)	218 (20)	135 (21)		
Family structure ^{a)}					
Divorced	335 (18)	209 (19)	180 (28)	30.3 ^{***}	
Not divorced	1483 (82)	906 (81)	458 (72)		
Social support family ^{b)}	22.1 (5.3)	21.5 (5.3)	20.9 (6.1)	11.1 ^{***}	1-2*, 1-3 ^{***}
Social support friends ^{b)}	21.5 (5.4)	21.5 (5.4)	22.4 (5.7)	6.0 *	1-3 ^{**} , 2-3*
Social support others ^{b)}	22.1 (5.2)	22.0 (5.2)	22.2 (5.7)	0.3 ^{ns}	ns
Extraversion ^{b)}	9.1 (2.8)	9.6 (2.8)	9.7 (2.9)	13.5 ^{***}	1-2 ^{***} , 1-3 ^{***}
Positive self-esteem ^{b)}	15.1 (2.3)	15.0 (2.4)	15.1 (2.6)	0.1 ^{ns}	ns
Negative self-esteem ^{b)}	11.8 (2.8)	12.0 (2.8)	12.4 (2.9)	11.5 ^{***}	1-3 ^{***} , 2-3*
Physical aggression ^{b)}	22.2 (6.6)	24.4 (7.18)	28.1 (7.7)	156.7 ^{***}	1-2 ^{***} , 1-3 ^{***} , 2-3 ^{***}
Verbal aggression ^{b)}	14.7 (3.8)	15.4 (3.9)	16.2 (4.2)	36.7 ^{***}	1-2 ^{***} , 1-3 ^{***} , 2-3 ^{***}
Anger ^{b)}	17.1 (4.8)	18.1 (4.9)	19.7 (5.1)	61.4 ^{***}	1-2 ^{***} , 1-3 ^{***} , 2-3 ^{***}
Hostility ^{b)}	21.1 (6.1)	22.0 (6.0)	22.8 (6.6)	18.2 ^{***}	1-2 ^{**} , 1-3 ^{***} , 2-3*
Depression/anxiety ^{b)}	11.5 (4.2)	11.8 (4.1)	12.6 (4.6)	16.9 ^{***}	1-3 ^{***} , 2-3 ^{***}
Social dysfunction ^{b)}	11.6 (2.5)	11.7 (2.5)	12.0 (3.0)	5.2 [*]	1-3 ^{**} , 2-3*

^{a)} descriptives in column 1, 2 and 3 concern the number of respondents (percentage of the sample), ^{b)} post hoc; 1 = abstainers, 2 = consumers, 3 = excessive drinkers, *** p < 0.001; ** p < 0.01, * p < 0.05; ns=not significant

Table 3.2 Associations of family factors, perceived social support, personality factors and psychological well-being with excessive drinking

	Excessive drinking OR (95% CI)				
	Model 1 Family factors	Model 2 Social support	Model 3 Personality traits	Model 4 Well-being	Model 5 Final model
Family affluence					
high	1 (Ref)				1 (Ref)
medium	0.89 (0.64 – 1.24)				0.87 (0.60 – 1.25)
low	0.66 (0.47 – 0.94)***				0.66 (0.45 – 0.97)**
Family structure					
intact	1 (Ref)				1 (Ref)
divorced	1.95 (1.47 – 2.58)***				1.65 (1.21 – 2.26)***
Social support from family		0.91 (0.88 – 0.93)***			0.93 (0.90 – 0.96)***
Social support from friends		1.10 (1.07 – 1.13)***			1.10 (1.06 – 1.13)***
Extraversion			1.08 (1.03 – 1.13)***		1.06 (1.01 – 1.11)***
Negative self-esteem			1.08 (1.03 – 1.13)**		1.02 (0.96 – 1.08)
Physical aggression			1.11 (1.09 – 1.13)***		1.12 (1.09 – 1.14)***
Verbal aggression			1.00 (0.96 – 1.04)		
Anger			1.04 (1.01 – 1.08)**		
Hostility			0.97 (0.94 – 0.99)**		1.03 (0.988 – 1.07)
Depression / anxiety				1.05 (1.02 – 1.08)***	0.97 (0.94 – 0.99)**
Social dysfunction				1.06 (1.003 – 1.12)***	1.03 (0.99 – 1.07)
Random variation at class level (standard error) #	0.427 (0.124)	0.423 (0.297)	0.404 (0.129)	0.395 (0.119)	1.06 (1.003 – 1.13)*** 0.400 (0.132)

** p < 0.01, *** p < 0.001; Ref=reference category; OR/CI values are based on multilevel logistic regression

For the empty model, this was 0.415 (0.121)

Discussion

The current study explored the differences between adolescent abstainers, consumers and excessive drinkers in regards to their family characteristics, social support, personality characteristics and well-being. We found differences between all three explored groups. Adolescent abstainers and excessive drinkers differed in every explored characteristic except for positive self-esteem and social support from others. Moreover, we found differences between consumers and abstainers in extraversion, aggression and social support from family; and between consumers and excessive drinkers in negative self-esteem, aggression, well-being and social support from friends. After mutual adjustment, eight differences remained statistically significant between excessive drinkers and abstainers, the former being more likely to have divorced parents, to be from families with higher affluence, to perceive less social support from family but more social support from friends, to report higher levels of extraversion, physical aggression, hostility and social dysfunction. Our finding regarding family structure confirms the findings of several other studies that explored this issue (Miller, 1997; Blum et al., 2000; Kuntsche & Kuendig, 2006; Paxton et al., 2007) – adolescents living in divorced families are at higher risk of excessive drinking. One of the explanations for this fact might be an often decreased parental control after divorce, but this hypothesis needs to be proved by further research. Findings in the literature about the association between SEP and hazardous drinking among adolescents are contradictory (Osler et al., 2001; Littlejohn, 2006; Zambon et al., 2006; Arvanitidou et al., 2007). In our study we assumed that lower SEP would be associated with a higher probability of excessive drinking. However, our results do not support this assumption: excessive drinking respondents were those with higher family affluence; parents' education did not show a significant association with excessive drinking. A speculation might be that the roots of the association between higher SEP and the higher probability of excessive drinking might be found in the particular youth subculture related to high SEP (particularly the attitude to drinking alcohol) rather than in the possession of more financial resources available for buying alcohol. In Slovakia, alcohol is very cheap, much cheaper than soft drinks [e.g. typical price of a beer (0.3 L) is €0.50 and of a soft drink (0.3 L) is €1 in a pub], making it rather cheap to get drunk. However, we cannot fully exclude that financial means plays a role. Our results concerning perceived social support from family are in line with studies that have found an association between low support from family and alcohol use in adolescents (Stice et al., 1993; Shucksmith et al., 1997; Windle & Miller-Tutzauer, 1997). We also confirmed the association between high perceived support from friends and excessive drinking (Engels & Ter Bogt, 2001). However, this does not necessarily mean that

the relationships among peers themselves are risky; such relationships are an essential part of healthy socialisation during adolescence (Kerr et al., 2003). Places where alcohol is sold (bars, pubs, discos, etc.) are where these relationships with peers take place, so maintaining a social network in adolescence is strongly connected with places or situations in which alcohol is easily obtainable. Extraversion as personality trait is often found to be associated with risk behaviours, including hazardous alcohol drinking (Martsh & Miller, 1997; Merenakk et al., 2003). Our results supported this assumption only partly – extraversion makes one more likely to be a consumer, but not an excessive drinker. This means that extraversion stimulates the participation in social activities, but as we stated above, the real risk of excessive drinking is more related to the context in which these activities are taking place. Drunkenness was found to cluster per class, but it has hardly an effect on model outcomes. This may be interpreted as meaning that classroom-bound factors do not affect drunkenness in an important way, but children in a given class share common background characteristics like family support and divorce background to some degree. The present study has several strengths and limitations. Its main strengths are the size of the study sample, the high response rate and the proportional representation from several different regions of Slovakia. A main limitation of our study is that it relies on the self-report of our respondents. However, the answers were filled out anonymously, which has been shown to lead to rather valid self-reports (Del Boca & Noll, 2000). Another limitation that has to be taken into account is that although parents are not legally divorced, it does not necessarily mean they still live together and this might have the same impact on their children as divorce. And, finally, cross-sectional data may not provide us with sufficient information about the causal mechanisms. As the design of this study was cross-sectional, the implication for further research might be to examine longitudinal data to confirm the hypothesised causal mechanisms with regard to hazardous drinking. Two main targets for practice could be tackled in this study. Our results show that particular groups (children of divorced parents, adolescents from families with higher affluence, those with lower social support from family but higher from friends, those with higher levels of physical aggression and hostility and those with lower well-being) run a higher risk of becoming excessive drinkers and thus need particular attention in prevention. The second implication arises from the results on peer support. We have already mentioned above that peers usually meet in an environment that is not alcohol-free. The prevention strategy might be to support alcohol-free, safe environments for these peer interactions on one hand and to limit the availability of alcoholic drinks in environments that are frequented by young adolescents (e.g. to increase the age limit for selling alcohol to adolescents in public places) on the other.

References

1. Arvanitidou M, Tirodimos I, Kyriakidis I, Tsinaslanidou Z, Seretopoulos D. Decreasing prevalence of alcohol consumption among Greek adolescents. *Am J Drug Alcohol Ab* 2007; 33: 411–17.
2. Bal S, Crombez G, van Oost P, Debourdeaudhuij I. The role of social support in well-being and coping with self-reported stressful events in adolescents. *Child Abuse Negl* 2003; 7: 1377–95.
3. Blum RW, Buehring T, Shew ML, Bearinger LH, Sieving RE, Resnick MD. The effects of race/ethnicity, income, and family structure on adolescent risk behaviors. *Am J Public Health* 2000; 90: 1879–84.
4. Blumenthal JA, Burg MM, Barefoot J, Williams RB, Haney T, Zimet G. Social support, type A behavior, and coronary artera disease. *Psychosom Med* 1987; 49: 331–40.
5. Catanzaro SJ, Laurent J. Perceived family support, negative mood regulation expectancies, coping, and adolescent alcohol use: evidence of mediation and moderation effects. *Addict Behav* 2004; 29: 1779–97.
6. Clark LA, Watson D. Constructing validity: basic issues in objective scale development. *Psychol Assess* 1995; 7: 309–19.
7. Currie C, Roberts Ch, Morgan A, Smith R, Settertobulte W, Samdal O, Barnekow Ramussen V. Young people's health in context. Health behaviour in school-aged children (HBSC) study: International report from the 2001/2002 survey. Health Policy for Children and Adolescents; Report No. 4, 2004.
8. Del Boca FK, Noll JA. Truth or consequences: the validity of self-report data in health services research on addictions. *Addiction* 2000; 95: 347–60.
9. Eickhoff C. Protection or risk? Family environments reflected in communication between parents and their children. In: Research and Practice of Health Promotion, Volume 10. Cologne: Federal Centre for Health Education, 2001: 34–39.
10. Engels RCME, Ter Bogt T. Influences of risk behaviors on the quality of peer relations in adolescence. *J Youth Adolesc* 2001; 30: 675–95.
11. Fisher LB, Williams Miles I, Austin SB, Camargo CA, Colditz GA. Predictors of initiation of alcohol use among US adolescents. *Arch Pediat Adol Med* 2007; 161: 959–66.
12. Francis LJ. The relationship between Eysenck's personality factors and attitude towards substance use among 13–15- year-olds. *Pers Individ Dif* 1996; 21: 633–40.
13. Freeman HS, Newland LA. Family transitions during the adolescent transition: implications for parenting. *Adolescence* 2002; 37: 457–75.
14. Geckova A, van Dijk JP, Stewart R, Groothoff JW, Post D. Influence of social support on health among gender and socio-economic groups of adolescents. *Eur J Public Health* 2003; 13: 44–50.

15. Gerra G, Angioni L, Zaimovic A, Moj G, Bussandri M, Bertacca S, Santoro G, Gardini S, Caccavari R, Nicoli MA: Substance use among high-school students: relationship with temperament, personality traits and parental care perception. *Subst Use Misuse* 2004; 39: 345–67.
16. Goldberg D, Williams P. A user's guide to the General Health Questionnaire. Windsor: NFER—Nelson, 1988.
17. Gosling SD, Renfrow PJ, Swann WB Jr. A very brief measure of Big-Five personality domains. *J Res Pers* 2003; 37: 504–28.
18. Griffin KW, Botvin GJ, Scheier LM, Diaz T, Miller NL. Parenting practices as predictors of substance use, delinquency, and aggression among urban minority youth: moderating effects of family structure and gender. *Psychol Addict Behav* 2000; 14: 174–84.
19. Hibbel B. The ESPAD Report 2003: Alcohol and other drug use among students in 35 European countries. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs, 2004.
20. Jessor R. Risk behavior in adolescence: a psychosocial framework for understanding and action. *J Adolesc Health* 1991; 12: 597–605.
21. Kerr M, Stattin H, Biesecker G, Ferrer-Wreder L. Relationships with parents and peers in adolescence. In: Lerner RM, Easterbrooks MA, Mistry J, eds. *Handbook of psychology, volume 6, developmental psychology*. Hoboken: John Wiley & Sons, Inc, 2003:395–422.
22. Kuntsche E, Kuendig H. What is worse? A hierarchy of family-related risk factors predicting alcohol use in adolescents. *Subst Use Misuse* 2006; 41: 71–86.
23. Kuntsche E, Rehm J, Gmel G. Characteristics of binge drinkers in Europe. *Soc Sci Med* 2004; 59: 113–27.
24. Ledoux S, Miller P, Choquet M, Plant M. Family structure, parent-child relationships, and alcohol and other drug use among teenagers in France and the United Kingdom. *Alcohol Alcoholism* 2002; 37: 52–60.
25. Lifrak PD, McKay J, Rostain A, Alterman AI, O'Brien CHP. Relationship of perceived competencies, perceived social support, and gender to substance use in young adolescents. *J Am Acad Child Psychiatry* 1997; 36: 933–40.
26. Littlejohn C. Does socio-economic status influence the acceptability of, attendance for, and outcome of, screening and brief interventions for alcohol misuse: a review. *Alcohol Alcoholism* 2006; 41: 540–5.
27. Marsden J, Boys A, Farrell M, et al. Personal and social correlates of alcohol consumption among mid-adolescents. *Br J Dev Psychol* 2005; 23: 427–50.
28. Martsh CT, Miller WR. Extraversion predicts heavy drinking in college students. *Pers Individ Dif* 1997; 23: 153–5.

29. Merenakk L, Harro M, Kiive E, Laidra K, Eensoo D, Allik J, Oreland L, Harro J: Association between substance use, personality traits, and platelet MAO activity in preadolescents and adolescents. *Addict Behav* 2003; 28: 1507–14.
30. Miller P. Family structure, personality, drinking, smoking and illicit drug use: a study of UK teenagers. *Drug Alcohol Depend* 1997; 45: 121–9.
31. Murberg TA, Bru E. Social support, negative life events and emotional problems among Norwegian adolescents. *Sch Psychol Int* 2004; 25: 387–403.
32. Nakano K. Psychometric evaluation on the Japanese adaptation of the aggression questionnaire. *Behav Res Ther* 2001; 39: 853–8.
33. Osler M, Jorgensen T, Davidsen M, et al. Socioeconomic status and trends in alcohol drinking in the Danish MONICA population, 1982–92. *Scand J Public Health* 2001; 29: 40–3.
34. Pandina RJ, Schuele JA. Psychosocial correlates of alcohol and drug use of adolescent students and adolescents in treatment. *J Stud Alcohol* 1983; 44: 950–73.
35. Parker JDA, Taylor GJ, Bagby RM. The 20-item Toronto Alexithymia Scale III. Reliability and factorial validity in a community population. *J Psychosom Res* 2003; 55: 269–75.
36. Paxton RJ, Valoism RE, Dranem JW. Is there a relationship between family structure and substance use among public middle school students? *J Child Fam Stud* 2007; 16:593–605.
37. Petraitis J, Flay BR, Miller TQ. Reviewing theories of adolescent substance use: organizing pieces in the puzzle. *Psychol Bull* 1995; 11: 67–86.
38. Pitkanen T. Problem drinking and psychological wellbeing: a five-year follow-up study from adolescence to young adulthood. *Scand J Psychol* 1999; 40: 197–207.
39. Power TG, Stewart CD, Hughes SO, Arbona C. Predicting patterns of adolescents alcohol use: a longitudinal study. *J Stud Alcohol* 2005; 66: 74–81.
40. Rasbash J, Charlton C, Browne WJ, Healy M, Cameron B. MLwiN Version 2.02. Centre for Multilevel Modelling. Bristol: University of Bristol, 2005. Available at: <http://www.cmm.bristol.ac.uk/MLwiN/index.shtml> (accessed May 2010).
41. Rosenberg M. Society and the adolescent self-image. Princeton: Princeton University Press, 1965.
42. Rosenberg M. Conceiving the self. New York: Basic Books, 1979.
43. Sarkova M, Nagyova I, Katreniakova Z, Geckova AM, Orosova O, Middel B, van Dijk JP, van den Heuvel W. Psychometric evaluation of the General Health Questionnaire-12 and Rosenberg Self-esteem Scale in Hungarian and Slovak early adolescents. *Stud Psychol* 2006; 48: 69–79.

44. Schweitzer RD, Seth-Smith M, Callan V. The relationship between self-esteem and psychological adjustment in young adolescents. *J Adolesc* 1992; 15: 83–97.
45. Shucksmith J, Glendinning A, Hendry L. Adolescents' drinking behaviour and the role of family life: a Scottish perspective. *J Adolesc* 1997; 20: 85–101.
46. Stice E, Barrera M, Chassin L. Relation of parental support and control to adolescents' externalizing symptomatology and substance use: a longitudinal examination of curvilinear effects. *J Abnorm Child Psychol* 1993; 21: 609–29.
47. Storksen I, Roysamb E, Moum T, Tambs K. Adolescents with a childhood experience of parental divorce: a longitudinal study of mental health and adjustment. *J Adolesc* 2005; 28: 725–39.
48. Swahn MH, Donovan JE. Correlates and predictors of violent behavior among adolescent drinkers. *J Adolesc Health* 2004; 34: 480–92.
49. Unger JB, Li Y, Johnson CA, et al. Stressful life events among adolescents in Wuhan, China: associations with smoking, alcohol use, and depressive symptoms. *Int J Behav Med* 2001; 8: 1–18.
50. Windle M, Miller-Tutzauer C. Confirmatory factor analysis and concurrent validity of the perceived social support— family measure among adolescents. *J Marriage Fam* 1997; 54: 777–87.
51. Zambon A, Lemma P, Borraccino A, Dalmaso P, Cavallo F. Socio-economic position and adolescents' health in Italy: the role of the quality of social relations. *Eur J Public Health* 2006; 16: 627–32.