

University of Groningen

## Of Stalks and Diamonds. Simulation Studies of Membrane Fusion and the Role of Fusion Peptides

Fuhrmans, Marc

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Fuhrmans, M. (2010). *Of Stalks and Diamonds. Simulation Studies of Membrane Fusion and the Role of Fusion Peptides*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Bibliography

- [1] Berendsen HJC. *Simulating the physical world: hierarchical modeling from quantum mechanics to fluid dynamics*. Cambridge University Press, 2007
- [2] Frenkel D, Smit B. *Understanding molecular simulation: from algorithms to applications*. Academic Press, 2002
- [3] Hadwiger H. *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*. Springer, 1957
- [4] Michielsen K, De Raedt H. Integral-geometry morphological image analysis. *Phys Rep*, 347:461, 2001
- [5] Seddon JM, Templer RH. Polymorphism of lipid-water systems. In Lipowsky R, Sackmann E, eds., *Handbook of biological physics*, 97–160. Elsevier, 1995
- [6] Israelachvili JN, Mitchell DJ, Ninham BW. Theory of self-assembly of hydrocarbon amphiphiles into micelles and bilayers. *J Chem Soc, Faraday Trans 2*, 72:1525, 1976
- [7] Hyde S, Andersson S, Larsson K, Blum Z, Landh T, Lidin S, Ninham BW. *The language of shape*. Elsevier, 1997
- [8] Helfrich W. Elastic properties of lipid bilayers: theory and possible experiments. *Z Naturforsch*, 28c:693, 1973
- [9] Siegel DP, Kozlov MM. The Gaussian curvature elastic modulus of N-monomethylated dioleoylphosphatidylethanolamine: relevance to membrane fusion and lipid phase behavior. *Biophys J*, 87:366, 2004
- [10] Cullis PR, Hope MJ, de Kruijff B, Verkleij AJ, Tilcock CPS. Structural properties and functional roles of phospholipids in biological membranes. In Kuo JF, ed., *Phospholipids and Cellular Regulation*, 1–59. CRC Press, 1985
- [11] Hui SW. Non-bilayer-forming lipids: Why are they necessary in biomembranes? *Comments Mol Cell Biophys*, 4:233, 1987
- [12] Markin VS, Albanesi JP. Membrane fusion: stalk model revisited. *Biophys J*, 82:693, 2002
- [13] Kozlovsky Y, Kozlov MM. Stalk model of membrane fusion: solution of energy crisis. *Biophys J*, 82:882, 2002
- [14] Siegel DP. The Gaussian curvature elastic energy of intermediates in membrane fusion. *Biophys J*, 95:5200, 2008
- [15] Siegel DP. The modified stalk mechanism of lamellar/inverted phase transitions and its implications for membrane fusion. *Biophys J*, 76:291, 1999
- [16] Kozlov MM, Leikin SL, Chernomordik LV, Markin VS, Chizmadzhev YA. Stalk mechanism of vesicle fusion. *Eur Biophys J*, 17:121, 1989
- [17] Lentz BR, Lee J. Poly(ethylene glycol)(PEG)-mediated fusion between pure lipid bilayers: a mechanism in common with viral fusion and secretory vesicle release? *Mol Membr Biol*, 16:279, 1999

- [18] Pantazatos DP, MacDonald RC. Directly observed membrane fusion between oppositely charged phospholipid bilayers. *J Membr Biol*, 170:27, 1999
- [19] Leikina E, Mittal A, Cho MS, Melikov K, Kozlov MM, Chernomordik LV. Influenza hemagglutinins outside of the contact zone are necessary for fusion pore expansion. *J Biol Chem*, 279:26526, 2004
- [20] Jahn R, Scheller RH. SNAREs – engines for membrane fusion. *Nat Rev Mol Cell Biol*, 7:631, 2006
- [21] Martens S, McMahon HT. Mechanisms of membrane fusion: disparate players and common principles. *Nat Rev Mol Cell Biol*, 9:543, 2008
- [22] Armstrong RT, Kushnir AS, White JM. The transmembrane domain of influenza hemagglutinin exhibits a stringent length requirement to support the hemifusion to fusion transition. *J Cell Biol*, 151:425, 2000
- [23] Siegel DP, Cherezov V, Greathouse DV, Koeppe RE, Killian JA, Caffrey M. Transmembrane peptides stabilize inverted cubic phases in a biphasic length-dependent manner: implications for protein-induced membrane fusion. *Biophys J*, 90:200, 2006
- [24] Han X, Tamm LK. A host–guest system to study structure–function relationships of membrane fusion peptides. *Proc Natl Acad Sci U S A*, 97:13097, 2000
- [25] Yeagle PL, Epanand RM, Richardson CD, Flanagan TD. Effects of the 'fusion peptide' from measles virus on the structure of N-methyl dioleoylphosphatidylethanolamine membranes and their fusion with Sendai virus. *Biochim Biophys Acta*, 1065:49, 1991
- [26] Epanand RM, Epanand RF. Relationship between the infectivity of influenza virus and the ability of its fusion peptide to perturb bilayers. *Biochem Biophys Res Commun*, 202:1420, 1994
- [27] Epanand RF, Martin I, Ruyschaert JM, Epanand RM. Membrane orientation of the SIV fusion peptide determines its effect on bilayer stability and ability to promote membrane fusion. *Biochem Biophys Res Commun*, 205:1938, 1994
- [28] Davies SM, Epanand RF, Bradshaw JP, Epanand RM. Modulation of lipid polymorphism by the feline leukemia virus fusion peptide: implications for the fusion mechanism. *Biochem*, 37:5720, 1998
- [29] Peisajovich SG, Epanand RF, Pritsker M, Shai Y, Epanand RM, et al. The polar region consecutive to the HIV fusion peptide participates in membrane fusion. *Biochem*, 39:1826, 2000
- [30] Aranda FJ, Teruel JA, Ortiz A. Interaction of a synthetic peptide corresponding to the N terminus of canine distemper virus fusion protein with phospholipid vesicles: a biophysical study. *Biochim Biophys Acta, Biomembr*, 1618:51, 2003
- [31] Bertocco A, Formaggio F, Toniolo C, Broxterman QB, Epanand RF, Epanand RM. Design and function of a conformationally restricted analog of the influenza virus fusion peptide. *J Pept Res*, 62:19, 2003
- [32] Epanand RM, Epanand RF, Martin I, Ruyschaerts JM. Membrane interactions of mutated forms of the influenza fusion peptide. *Biochem*, 40:8800, 2001
- [33] Colotto A, Martin I, Ruyschaert JM, Sen A, Hui SW, Epanand RM. Structural study of the interaction between the SIV fusion peptide and model membranes. *Biochem*, 35:980, 1996
- [34] Siegel DP, Epanand RM. Effect of influenza hemagglutinin fusion peptide on lamellar/inverted phase transitions in dipalmitoleoylphosphatidylethanolamine: implications for membrane fusion mechanisms. *Biochim Biophys Acta, Biomembr*, 1468:87, 2000

- [35] Harroun TA, Balali-Mood K, Gourlay I, Bradshaw JP. The fusion peptide of simian immunodeficiency virus and the phase behaviour of N-methylated dioleoylphosphatidylethanolamine. *Biochim Biophys Acta, Biomembr*, 1617:62, 2003
- [36] Egberts E, Berendsen HJC. Molecular dynamics simulation of a smectic liquid crystal with atomic detail. *J Chem Phys*, 89:3718, 1988
- [37] Lindahl E, Edholm O. Mesoscopic undulations and thickness fluctuations in lipid bilayers from molecular dynamics simulations. *Biophys J*, 79:426, 2000
- [38] Marrink SJ, Lindahl E, Edholm O, Mark AE. Simulation of the spontaneous aggregation of phospholipids into bilayers. *J Am Chem Soc*, 123:8638, 2001
- [39] Leontiadou H, Mark AE, Marrink SJ. Antimicrobial peptides in action. *J Am Chem Soc*, 128:12156, 2006
- [40] Periole X, Huber T, Marrink SJ, Sakmar TP. G protein-coupled receptors self-assemble in dynamics simulations of model bilayers. *J Am Chem Soc*, 129:10126, 2007
- [41] Marrink SJ, de Vries AH, Tieleman DP. Lipids on the move: Simulations of membrane pores, domains, stalks and curves. *Biochim Biophys Acta, Biomembr*, 1788:149, 2009
- [42] Berger O, Edholm O, Jähnig F. Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophys J*, 72:2002, 1997
- [43] Elezgaray J, Laguerre M. A systematic method to derive force fields for coarse-grained simulations of phospholipids. *Comp Phys Comm*, 175:264, 2006
- [44] Izvekov S, Violi A, Voth GA. Systematic coarse-graining of nanoparticle interactions in molecular dynamics simulation. *J Phys Chem B*, 109:17019, 2005
- [45] Müller M, Katsov K, Schick M. Biological and synthetic membranes: What can be learned from a coarse-grained description? *Phys Rep*, 434:113, 2006
- [46] Marrink SJ, Risselada HJ, Yefimov S, Tieleman DP, de Vries AH. The MARTINI force field: coarse grained model for biomolecular simulations. *J Phys Chem B*, 111:7812, 2007
- [47] Marrink SJ, Mark AE. The mechanism of vesicle fusion as revealed by molecular dynamics simulations. *J Am Chem Soc*, 125:11144, 2003
- [48] Kasson PM, Kelley NW, Singhal N, Vrljic M, Brunger AT, Pande VS. Ensemble molecular dynamics yields submillisecond kinetics and intermediates of membrane fusion. *Proc Natl Acad Sci U S A*, 103:11916, 2006
- [49] Knecht V, Marrink SJ. Molecular dynamics simulations of lipid vesicle fusion in atomic detail. *Biophys J*, 92:4254, 2007
- [50] Smeijers AF, Markvoort AJ, Pieterse K, Hilbers PAJ. A detailed look at vesicle fusion. *J Phys Chem B*, 110:13212, 2006
- [51] Risselada HJ, Mark AE, Marrink SJ. Application of mean field boundary potentials in simulations of lipid vesicles. *J Phys Chem B*, 112:7438, 2008
- [52] Ingebrigtsen L, Brandl M. Determination of the size distribution of liposomes by sec fractionation, and pcs analysis and enzymatic assay of lipid content. *AAPS PharmSciTech*, 3:9, 2002
- [53] Berendsen HJC, Postma JPM, DiNola A, Haak JR. Molecular dynamics with coupling to an external bath. *J Chem Phys*, 81:3684, 1984
- [54] Müller M, Katsov K, Schick M. New mechanism of membrane fusion. *J Chem Phys*, 116:2342, 2002
- [55] Noguchi H, Takasu M. Fusion pathways of vesicles: a brownian dynamics simulation. *J Chem Phys*, 115:9547, 2001

- [56] Katsov K, Müller M, Schick M. Field theoretic study of bilayer membrane fusion: II. mechanism of a stalk-hole complex. *Biophys J*, 90:915, 2006
- [57] Bonnafous P, Stegmann T. Membrane perturbation and fusion pore formation in influenza hemagglutinin-mediated membrane fusion. A new model for fusion. *J Biol Chem*, 275:6160, 2000
- [58] Cross KJ, Langley WA, Russell RJ, Skehel JJ, Steinhauer DA. Composition and functions of the influenza fusion peptide. *Protein Pept Lett*, 16:766, 2009
- [59] Steinhauer DA, Wharton SA, Skehel JJ, Wiley DC. Studies of the membrane fusion activities of fusion peptide mutants of influenza virus hemagglutinin. *J Virol*, 69:6643, 1995
- [60] Carr CM, Chaudhry C, Kim PS. Influenza hemagglutinin is spring-loaded by a metastable native conformation. *Proc Natl Acad Sci U S A*, 94:14306, 1997
- [61] Lai AL, Park H, White JM, Tamm LK. Fusion peptide of influenza hemagglutinin requires a fixed angle boomerang structure for activity. *J Biol Chem*, 281:5760, 2006
- [62] Han X, Bushweller JH, Cafiso DS, Tamm LK. Membrane structure and fusion-triggering conformational change of the fusion domain from influenza hemagglutinin. *Nat Struct Biol*, 8:715, 2001
- [63] Huang Q, Chen CL, Herrmann A. Bilayer conformation of fusion peptide of influenza virus hemagglutinin: a molecular dynamics simulation study. *Biophys J*, 87:14, 2004
- [64] Lagüe P, Roux B, Pastor RW. Molecular dynamics simulations of the influenza hemagglutinin fusion peptide in micelles and bilayers: conformational analysis of peptide and lipids. *J Mol Biol*, 354:1129, 2005
- [65] Marrink SJ, Mark AE. Molecular view of hexagonal phase formation in phospholipid membranes. *Biophys J*, 87:3894, 2004
- [66] Monticelli L, Kandasamy SK, Periole X, Larson RG, Tieleman DP, Marrink SJ. The MARTINI coarse-grained force field: extension to proteins. *J Chem Theory Comput*, 4:819, 2008
- [67] van der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC. GROMACS: fast, flexible, and free. *J Comput Chem*, 26:1701, 2005
- [68] Hess B. Determining the shear viscosity of model liquids from molecular dynamics simulations. *J Chem Phys*, 116:209, 2002
- [69] Fuhrmans M, Knecht V, Marrink SJ. A single bicontinuous cubic phase induced by fusion peptides. *J Am Chem Soc*, 131:9166, 2009
- [70] Rand RP, Fuller NL. Structural dimensions and their changes in a reentrant hexagonal-lamellar transition of phospholipids. *Biophys J*, 66:2127, 1994
- [71] Yang L, Ding L, Huang HW, et al. New phases of phospholipids and implications to the membrane fusion problem. *Biochem*, 42:6631, 2003
- [72] Siegel DP. Determining the ratio of the Gaussian curvature and bending elastic moduli of phospholipids from QII phase unit cell dimensions. *Biophys J*, 91:608, 2006
- [73] Siegel DP. The relationship between bicontinuous inverted cubic phases and membrane fusion. In Lynch ML, Spicer PT, eds., *Bicontinuous Liquid Crystals*, 59–98. Taylor and Francis, 2005
- [74] Marrink SJ, Fuhrmans M, Risselada HJ, Periole X. The MARTINI forcefield. In Voth G, ed., *Coarse-Graining of Condensed Phase and Biomolecular Systems*. CRC Press, 2008

- [75] Loison C, Mareschal M, Schmid F. Pores in bilayer membranes of amphiphilic molecules: Coarse-grained molecular dynamics simulations compared with simple mesoscopic models. *J Chem Phys*, 121:1890, 2004
- [76] Marrink SJ, Tieleman DP. Molecular dynamics simulation of spontaneous membrane fusion during a cubic-hexagonal phase transition. *Biophys J*, 83:2386, 2002
- [77] Sevink GJA, Zvelindovsky AV. Self-assembly of complex vesicles. *Macromol*, 38:7502, 2005
- [78] Edelsbrunner H, Mücke E. 3D alpha shapes. *Assoc Comput Mach Trans Graphics*, 13:43, 1994
- [79] Zomorodian A, Carlsson G. Computing persistent homology. *Discrete Comput Geom*, 33:249, 2005
- [80] Kasson PM, Zomorodian A, Park S, Singhal N, Guibas LJ, Pande VS. Persistent voids: a new structural metric for membrane fusion. *Bioinformatics*, 23:1753, 2007
- [81] Hyde ST, Barnes IS, Ninham BW. Curvature energy of surfactant interfaces confined to the plaquettes of a cubic lattice. *Langmuir*, 6:1055, 1990
- [82] Delano WL. The PyMOL molecular graphics system, 2002
- [83] Voth G, ed. *Coarse-Graining of Condensed Phase and Biomolecular Systems*. CRC Press, 2008
- [84] Ayton GS, Noid WG, Voth GA. Multiscale modeling of biomolecular systems: in serial and in parallel. *Curr Opin Struct Biol*, 17:192, 2007
- [85] Ensing B, Nielsen SO, Moore PB, Klein ML, Parrinello M. Energy conservation in adaptive hybrid atomistic/coarse-grain molecular dynamics. *J Chem Theory Comput*, 3:1100, 2007
- [86] Christen M, van Gunsteren WF. Multigraining: an algorithm for simultaneous fine-grained and coarse-grained simulation of molecular systems. *J Chem Phys*, 124:154106, 2006
- [87] Berendsen HJC, Postma JPM, van Gunsteren WF, Hermans J. Interaction models for water in relation to protein hydration. In Pullman B, ed., *Intermolecular Forces: Proc. Fourteenth Jerusalem Symp. on Quantum Chemistry and Biochemistry*. Reidel, 1981
- [88] Oostenbrink C, Villa A, Mark AE, Van Gunsteren WF. A biomolecular force field based on the free enthalpy of hydration and solvation: the GROMOS force-field parameter sets 53A5 and 53A6. *J Comput Chem*, 25:1656, 2004
- [89] Tironi IG, Sperb R, Smith PE, van Gunsteren WF. A generalized reaction field method for molecular dynamics simulations. *J Chem Phys*, 102:5451, 1995
- [90] Miyamoto S, Kollman PA. SETTLE: an analytical version of the SHAKE and RATTLE algorithm for rigid water models. *J Comput Chem*, 13:952, 1992
- [91] Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. LINCS: a linear constraint solver for molecular simulations. *J Comput Chem*, 18:1463, 1997
- [92] de Vries AH. 53A6 for lipids. In preparation, topology available on request.
- [93] de Vries AH, Chandrasekhar I, van Gunsteren WF, Hünenberger PH. Molecular dynamics simulations of phospholipid bilayers: Influence of artificial periodicity, system size, and simulation time. *J Phys Chem B*, 109:11643, 2005
- [94] Artymiuk PJ, Blake CCF, Rice DW, Wilson KS. The structures of the monoclinic and orthorhombic forms of hen egg-white lysozyme at 6 Å resolution. *Acta Crystallogr, Sect B: Struct Crystallogr Cryst Chem*, 38:778, 1982
- [95] Guo GJ, Zhang YG, Zhao YJ. Comment on computation of the viscosity of a liquid from time averages of stress fluctuations. *Phys Rev E: Stat, Nonlinear, Soft Matter Phys*, 67:43101, 2003

- [96] van Gunsteren WF, Daura X, Mark AE. Computation of free energy. *Helv Chim Acta*, 85:3113, 2002
- [97] Baron R, Trzesniak D, de Vries AH, Elsener A, Marrink SJ, van Gunsteren WF. Comparison of thermodynamic properties of coarse-grained and atomic-level simulation models. *ChemPhysChem*, 8:452, 2007
- [98] Krynicki K, Green CD, Sawyer DW. Pressure and temperature dependence of self-diffusion in water. *Faraday Discuss Chem Soc*, 66:199, 1978
- [99] Weast RC. *Handbook of Chemistry and Physics*. CRC Press, 1986
- [100] Gildnyi T, Stergiopoulos C, Wolfram E. Equilibrium surface tension of aqueous surfactant solutions. *Colloid Polym Sci*, 254:1018, 1976
- [101] Groot RD, Rabone KL. Mesoscopic simulation of cell membrane damage, morphology change and rupture by nonionic surfactants. *Biophys J*, 81:725, 2001
- [102] van der Spoel D, van Maaren PJ, Berendsen HJC. A systematic study of water models for molecular simulation: Derivation of water models optimized for use with a reaction field. *J Chem Phys*, 108:10220, 1998
- [103] Halle B, Wennerström H. Interpretation of magnetic resonance data from water nuclei in heterogeneous systems. *J Chem Phys*, 75:1928, 1981
- [104] Cabani S, Gianni P, Mollica V, Lepori L. Group contributions to the thermodynamic properties of non-ionic organic solutes in dilute aqueous solution. *J Solution Chem*, 10:563, 1981
- [105] Nagle JF, Tristram-Nagle S. Structure of lipid bilayers. *Biochim Biophys Acta, Biomembr*, 1469:159, 2000
- [106] Kabsch W, Sander C. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers*, 22:2577, 1983
- [107] Chu JW, Ayton GS, Izvekov S, Voth GA. Emerging methods for multiscale simulation of biomolecular systems. *Mol Phys*, 105:167, 2007
- [108] Marrink SJ, de Vries AH, Mark AE. Coarse grained model for semiquantitative lipid simulations. *J Phys Chem B*, 108:750, 2004
- [109] Bennett WFD, MacCallum JL, Hinner MJ, Marrink SJ, Tieleman DP. Molecular view of cholesterol flip-flop and chemical potential in different membrane environments. *J Am Chem Soc*, 131:12714, 2009
- [110] Risselada HJ, Marrink SJ. The molecular face of lipid rafts in model membranes. *Proc Natl Acad Sci U S A*, 105:17367, 2008
- [111] Marrink SJ, Risselada J, Mark AE. Simulation of gel phase formation and melting in lipid bilayers using a coarse grained model. *Chem Phys Lip*, 135:223, 2005
- [112] Radzicka A, Wolfenden R. Comparing the polarities of the amino acids: side-chain distribution coefficients between the vapor phase, cyclohexane, 1-octanol, and neutral aqueous solution. *Biochem*, 27:1664, 1988
- [113] Bond PJ, Sansom MSP. Insertion and assembly of membrane proteins via simulation. *J Am Chem Soc*, 128:2697, 2006
- [114] Shih AY, Arkhipov A, Freddolino PL, Schulten K. Coarse grained protein-lipid model with application to lipoprotein particles. *J Phys Chem B*, 110:3674, 2006
- [115] Yefimov S, Van Der Giessen E, Onck PR, Marrink SJ. Mechanosensitive membrane channels in action. *Biophys J*, 94:2994, 2008