

University of Groningen

Delayed presentation to regular Dutch paediatric care in COVID-19 times: a national survey

Jansen, Danielle; Illy, Karoly

Published in:
BMJ Paediatrics Open

DOI:
[10.1136/ bmjpo-2020-000834](https://doi.org/10.1136/bmjpo-2020-000834)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Jansen, D., & Illy, K. (2020). Delayed presentation to regular Dutch paediatric care in COVID-19 times: a national survey. *BMJ Paediatrics Open*, 1-3. [https://doi.org/10.1136/ bmjpo-2020-000834](https://doi.org/10.1136/bmjpo-2020-000834)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Delayed presentation to regular Dutch paediatric care in COVID-19 times: a national survey

Danielle E M C Jansen ,^{1,2} Károly E Illy^{3,4}

To cite: Jansen DEMC, Illy KE. Delayed presentation to regular Dutch paediatric care in COVID-19 times: a national survey. *BMJ Paediatrics Open* 2020;**4**:e000834. doi:10.1136/bmjpo-2020-000834

Received 8 August 2020

Accepted 20 September 2020

ABSTRACT

We explored the collateral harm in Dutch children and adolescents during the COVID-19 pandemic from experience of paediatricians via an open question distributed via the website of the Dutch Paediatric Society. From the end of March till the first week of July, we received 51 reports of collateral harm involving mostly very young children with mainly acute physical problems but also social problems. In older children, several cases of diabetic ketoacidosis were reported. Our results show that delaying care can lead to seriously ill children, life-threatening situations and that in some cases it can even lead to death. If we want to avoid such a delay at a possible second peak of Corona, general care providers and paediatricians have to join forces and find new ways of working. Systematic data collection of collateral harm in children is needed to be able to intervene adequately.

In their report on the consequences of the corona crisis for regular care, the Dutch Healthcare Authority showed that of all specialisms, paediatrics has experienced the

largest decline in the number of urgent referrals in the first weeks of the COVID-19 crisis. In addition, paediatrics was the specialty that showed the least signs of a recovery in referrals (it even fell back slightly) while all other specialties showed a steady recovery from early April onwards.¹ The decrease in referrals might be explained by a reduced incidence of, for example, traffic and sports injuries as a result of social distancing and school closures. However, the decline might also indicate a delay in care and subsequently in collateral harm. Although some studies report on the consequences of delayed presentations to emergency departments, insight in the nature and severity of delayed presentation to regular paediatric care is missing.^{2,3} Our objective was to explore the collateral harm in Dutch children and adolescents during the COVID-19 pandemic from the experience of paediatricians.

Table 1 Background characteristics of cases (n=51)

Background characteristics	
Gender	Male: 9 Female: 11 Unknown: 29
Age of reported cases	Neonate (under 28 days of age): 9 (18%) Infant (under 1 year of age): 10 (20%) Aged 1–4 years: 8 (16%) Aged 4–12 years: 5 (10%) Aged 12–16 years: 6 (12%) Unknown/not reported: 13 (26%)
Province of report (location of province and no/% of registered patients with COVID-19 on 24 March 2020)*	South Holland (West Netherlands; 758 (13.6%)): 10 North Holland (West Netherlands; 740 (13.3%)): 9 North Brabant (South Netherlands; 1739 (31.3%)): 9 Gelderland (East Netherlands; 593 (10.7%)): 6 Utrecht (West Netherlands; 478 (8.6%)): 4 Limburg (South Netherlands; 690 (12.4%)): 3 Drenthe (North Netherlands; 55 (1%)): 2 Friesland (North Netherlands; 42 (0.8%)): 2 Zeeland (West Netherlands; 63 (1.1%)): 2 Groningen (North Netherlands; 71 (1.3%)): 1 Overijssel (East Netherlands; 257 (4.4%)): 1 Flevoland (East Netherlands; 74 (1.3%)): 0

*https://www.rivm.nl/sites/default/files/2020-03/Epidemiologische_situatie_COVID-19_24_maart_2020.pdf.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

¹Department of Health Sciences, University Medical Centre Groningen, Groningen, The Netherlands

²Sociology, University of Groningen Faculty of Behavioural and Social Sciences, Groningen, The Netherlands

³Ziekenhuis Rivierenland, Tiel, Gelderland, The Netherlands

⁴NVK, Utrecht, The Netherlands

Correspondence to

Dr Danielle E M C Jansen; d.e.m.c.jansen@umcg.nl

**Table 2** Reports of collateral harm**Paediatric subspecialty**

↓	Collateral harm →
Cardiology	<ul style="list-style-type: none"> ▶ Broadened mediastinum with a vena cava superior syndrome, due to a lymphoma ▶ Complex cor vitium ▶ Critical pulmonary stenosis and right ventricular hypertrophy with poor right ventricle dysfunction ▶ Congenital cyanotic heart disease
Child abuse paediatrics	<ul style="list-style-type: none"> ▶ Died due to serious abuse ▶ Impending out of home placement ▶ Oppression of the brains due to subdural hematoma
Ear, nose and throat	<ul style="list-style-type: none"> ▶ Extensive soft-tissue swelling in the mouth due to abscess
Endocrinology	<ul style="list-style-type: none"> ▶ Diabetes mellitus de novo (1x) ▶ Diabetes mellitus de novo with severe diabetic ketoacidosis 3x ▶ Diabetic ketoacidosis (5x)
Gastroenterology and nutrition	<ul style="list-style-type: none"> ▶ Abscess in the abdomen after appendicitis (3x) ▶ Developmental delay due to carnitine deficiency ▶ Low birth weight (2x) ▶ Oral aversion ▶ Severe weight loss ▶ Severe dehydration with hypochloraemic alkalosis, hypokalemia and hyponatremia ▶ Vitamin B₁₂ and folic acid deficiency
Genetics and metabolic diseases	<ul style="list-style-type: none"> ▶ Long-term breathing stop and diarrhoea in child with Cockayne syndrome
Haematology	<ul style="list-style-type: none"> ▶ Anaemia with signs of impending circulatory insufficiency ▶ Hyperbilirubinaemia (2x)
Infectious diseases	<ul style="list-style-type: none"> ▶ A-typical COVID-19 symptoms ▶ Impetigo bullosa and suspected Staphylococcal Scalded Skin Syndrome (SSSS) ▶ Infected, necrotic varicella lesions ▶ Kawasaki-like symptoms ▶ Mastoiditis ▶ Sepsis ▶ Shock due to group B streptococcal septicaemia
Neurology	<ul style="list-style-type: none"> ▶ Severe neurological complication after manual therapy
Oncology	<ul style="list-style-type: none"> ▶ Leukaemia ▶ Persistent fever and suspicion of lymphoma ▶ Possible benign tumour from tonsil/uvula
Pulmonology	<ul style="list-style-type: none"> ▶ Asphyxia ▶ Asthma ▶ Respiratory failure with respiratory infection ▶ Subglottic stenosis, due to haemangioma
Psychiatry	<ul style="list-style-type: none"> ▶ Anorexia ▶ Severe weight loss, a. mesenteric superior syndrome, acute renal failure, ulcers, leucopenia with fever, traumatic injury after attempted suicide
Other	<ul style="list-style-type: none"> ▶ Unexplained clinical deterioration ▶ Insufficient follow-up ex-premature

Via the website of the Dutch Paediatric Society, we requested all 1400 paediatricians affiliated with the professional association (93% of all Dutch paediatricians) to report on collateral harm in children and adolescents, from 2 weeks since the initiation of the Dutch 'intelligent lockdown',⁴ a lighter version of a full lockdown⁵ (end of March) to the first week of July. The question was: 'We ask you to report if, in your opinion, a child was presented too

late to acute, regular or chronic care due to parental or healthcare provider concerns about corona, and which resulted in unnecessary harm'.

The results of this inventory showed 51 reports (from 38 respondents divided over 31 hospitals) of collateral harm since the end of March. Although the majority of reports of harm were received in the first 4 weeks of the intelligent lockdown (n=27), there were still 24 reports in

the months after, up to and including the second week of July. The reports came from all over the Netherlands, but most reports were received from the west and south-west of the Netherlands, the regions where the corona crisis was more severe and, in all likelihood, experienced the highest pressure on healthcare.

Most reports (54%) of collateral harm involved young children: neonates, infants and children aged 1–4 years (table 1). The symptoms with which the children were presented—too late—to the paediatrician varied widely, including mainly acute physical problems but also social problems. The delay in neonatal care was mainly related to hyperbilirubinaemia and weight-related problems (low birth weight and severe weight loss). Spread across multiple age groups, several children presented with diabetic ketoacidosis (table 2).

Although the results of this exploration among paediatricians shows an alarming situation, this is probably only the tip of the iceberg since it is an exploration in which the data were not collected systematically. Our results show that delaying care can lead to seriously ill children, life-threatening situations and that in some cases it can even lead to death. If we want to avoid such a delay in providing the right care at the right place by the right person, at a possible second peak of Corona, action must be taken in which general care providers and paediatricians have to join forces, in particular regarding triage. We need to find new ways of working for unusual times like this so that the delay in care is avoided at all times. Finally, there should be systematic data collection of collateral harm in children; this is the only way to clarify its causes so that targeted interventions can be made.

Contributors DJ drafted the initial manuscript. KI provided significant contributions to the data collection that led to this manuscript and critically reviewed and revised the manuscript. Both authors provided significant contributions to the data analysis and presentation used in this manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Danielle E M C Jansen <http://orcid.org/0000-0002-2654-1038>

REFERENCES

- 1 Dutch Healthcare Authority. Analysis of the consequences of the corona crisis for regular care, 2020. Available: https://puc.overheid.nl/nza/doc/PUC_307166_22/1/ [Accessed 13 Jul 2020].
- 2 Ciacchini B, Tonioli F, Marciano C, *et al*. Reluctance to seek pediatric care during the COVID-19 pandemic and the risks of delayed diagnosis. *Ital J Pediatr* 2020;46:87.
- 3 Roland D, Harwood R, Bishop N, *et al*. Children's emergency presentations during the COVID-19 pandemic. *Lancet Child Adolesc Health* 2020;4:e32–3.
- 4 COVID-19 pandemic in the Netherlands. Available: https://en.wikipedia.org/wiki/COVID-19_pandemic_in_the_Netherlands
- 5 de Haas M, Faber R, Hamersma M. How COVID-19 and the Dutch 'intelligent lockdown' change activities, work and travel behaviour: Evidence from longitudinal data in the Netherlands. *Transportation Research Interdisciplinary Perspectives* 2020;6:100150.