

University of Groningen

Epidemiology of Erosive Tooth Wear, Dental Fluorosis and Molar Incisor Hypomineralization in the American Continent

Martignon, Stefania; Bartlett, David; Manton, David J; Martinez-Mier, E Angeles; Splieth, Christian; Avila, Viviana

Published in:
Caries Research

DOI:
[10.1159/000512483](https://doi.org/10.1159/000512483)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Martignon, S., Bartlett, D., Manton, D. J., Martinez-Mier, E. A., Splieth, C., & Avila, V. (2021). Epidemiology of Erosive Tooth Wear, Dental Fluorosis and Molar Incisor Hypomineralization in the American Continent. *Caries Research*, 55(1), 1-11. <https://doi.org/10.1159/000512483>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Epidemiology of Erosive Tooth Wear, Dental Fluorosis and Molar Incisor Hypomineralization in the American Continent

Stefania Martignon^a David Bartlett^b David J. Manton^{c, d}
E. Angeles Martinez-Mier^e Christian Splieth^f Viviana Avila^a

^aUNICA – Caries Unit Research, Research Department, Universidad El Bosque, Bogotá, Colombia; ^bCentre for Oral, Clinical and Translational Sciences, Faculty of Dentistry, Oral & Craniofacial Sciences, King's College London, London, UK; ^cCentrum voor Tandheelkunde en Mondzorgkunde, UMCG, University of Groningen, Groningen, The Netherlands; ^dMelbourne Dental School, University of Melbourne, Melbourne, VIC, Australia; ^eDepartment of Cariology, Operative Dentistry and Dental Public Health, Indiana University School of Dentistry, Indianapolis, IN, USA; ^fSection of Preventive and Pediatric Dentistry, University of Greifswald, Greifswald, Germany

Keywords

Epidemiology · Tooth erosion · Tooth wear · Dental fluorosis · Molar incisor hypomineralization

Abstract

Non-carious dental lesions such as developmental defects of enamel (DDE) and erosive tooth wear (ETW) are the subject of intensive research. This paper aims to give perspectives on both DDE, including dental fluorosis and molar incisor hypomineralization (MIH), and ETW, presenting epidemiological data from the Americas and associated diagnostic aspects. Besides, it is important to present evidence to guide the clinical assessment process, supporting the clinicians' management decisions towards better oral health of their patients. The overall increase in the worldwide prevalence of non-carious lesions discussed in this paper may reflect the need of perceptual changes. Although the number of publications related to these conditions has been increasing in the last years, there is still a need for clinical diagnostic and management awareness to include these conditions in routine dental practice. Besides, it is important to provide recommendations for standardized clinical assessment criteria,

improving the process and helping clinicians' adherence. In this sense, this paper discusses the most commonly implemented indices for each condition. Thus, despite the wide range of diagnostic indices, BEWE is proposed to be the index recommended for ETW assessment, Dean or Thylstrup & Fejerskov indices for fluorosis and preferably the EAPD criteria (or modified DDE index) for MIH. Overall, non-carious lesions are a growing concern, and it is important to implement preventive measures that control their severity and progression, and accurate diagnosis by the dental clinician.

© 2021 S. Karger AG, Basel

Introduction

This paper is a product of the Saturday Afternoon Symposium (SAS) that took place last July during the 66th ORCA Congress in Cartagena, Colombia and offers valuable information for recognizing dental conditions (apart from dental caries) that constitute clinical, epidemiological, research and education challenges. Thus, a perspective based on best available evidence, on erosive tooth wear (ETW) and developmental defects of enamel

(including molar incisal hypomineralization (MIH) and dental fluorosis) (DDE) is given here, emphasizing the American continent epidemiological information in relation to the context of the ORCA meeting held in a Latin American country.

In this sense, non-carious dental conditions such as ETW, MIH and dental fluorosis are subject of extensive research as well as growing concern to dentists around the world. The aim of this symposium paper is to give a perspective on the aforementioned conditions presenting, in addition to the epidemiological data, the most frequently reported indices for each condition and supporting guidance to clinicians regarding the assessment process in order to inform best practice management decisions, thus leading to better oral health outcomes for their patients.

The understanding of these dental conditions is also relevant to increase dental practitioners' awareness and skills regarding differential diagnosis of non-carious lesions for routine practice.

Erosive Tooth Wear

General Aspects

ETW has received increasing attention over the past 15 years [Schlueter and Luka, 2018]. This is in no small part to the data derived from global epidemiological studies, but most has been published in Europe. In 2009 [Van't Spijker et al., 2009] and 2010 [Kreulen et al., 2010] published two systematic reviews reporting higher levels in children and adolescents up to 82% in primary teeth and 5% in permanent teeth and for adults up to 17%. However, comparison between studies was challenging, because different indices were used to assess the severity of wear. In part, to overcome this the Basic Erosive Tooth Wear Examination (BEWE) was devised as part of a consensus meeting [Bartlett et al., 2008]. Although primarily designed for screening in general dental practice it has now been used regularly in prevalence studies and has since its introduction has been used in 58 clinical and epidemiological studies on 36,982 participants in 35 countries. Our team at KCL has reported prevalence rates for over 10,000 adults in 14 developed and developing countries with reported prevalence rates of 30% in adults aged 18–30 years [Bartlett et al., 2013] and up to 60% in developing countries such as Oman [Awad et al., 2019]. Overall, the prevalence rate is reported to be over 30%.

In calculating data, the convention is to accept the highest value from a single tooth or surface in a dentition. This gives an estimation of the impact of the condition,

which is different to a patient's perspective, particularly for those with a mouth destroyed by tooth wear. However, using these methods data can be compared between countries with some confidence.

Epidemiology

European data suggests that the condition is common. One study from seven European countries reported BEWE scores of 2 with 29% of 3100 participants and 3% with BEWE 3, the higher score representing more severe wear [Bartlett et al., 2013]. Typically, with large multi-country studies there was variation between countries at different BEWE scores. This might reflect variation in risk factors between countries or differences in assessment processes which are inevitable with large population studies. A similar methodological process was used in an Arab study across 6 countries, which also reported variation between 60% in Oman to 15% in Jordan [Awad et al., 2019]. Training is essential to reduce variation between examiners, but some differences are inevitable. The UK has reported a comparatively large number of studies [Smith et al., 1997; Bartlett et al., 1998; Fares et al., 2009], but many have used the Smith and Knight [Smith and Knight, 1984] index which calculates tooth wear by dentine exposure. These data, based on dentine exposure, give a range between 2–5% in younger adults and up to 10% for older age groups [Van't Spijker et al., 2009]. But direct comparison to the BEWE is not possible as they use different criteria. But overall, the evidence suggests that ETW is common and increases severity with age (Table 1).

A recent paper by Schlueter and Luka [2018] reported prevalence data of 22% in Europe and 21% in America for children and adolescents. The number of studies reported in Europe was far higher than in other countries with 23 in Europe and 9 in the America's. Amongst other countries there were around ten. This paper also reported that the condition was age related, generally more common in males than females and children had higher levels. Their analysis of the data reported that in Europe the condition in adults varied between countries with Germany reporting 6- to 7-year-olds with 72%, the Netherlands 15- to 16-year-olds 30%, Sweden 18–19 years 22%. Very little data is available from South America or the USA.

The clinical finding that dentine is exposed causes some controversy partly due to a study on extracted teeth where dentine was observed to be exposed but on microscopic examination enamel remained visible [Ganss et al., 2006]. However, from a clinical perspective exposed dentine is a useful predictor of the depth of ETW whereas the

Table 1. The prevalence of erosive tooth wear in the Americas

| Country | Authors | Year | Age, years | Sample size | ETW prevalence | Criteria |
|----------|--------------------------------|------|------------|-------------|----------------|-----------------------------------|
| Brazil | Nahás-Pires et al. | 2011 | 2–20 | 232 | 25.4 | O'Brien index |
| Brazil | De Carvalho Sales-Peres et al. | 2013 | 4 & 12 | 2,731 | 51.7 | Erosion wear index (EWI) |
| Brazil | Moimaz et al. | 2013 | 6–7 | 2,759 | 0.6 | Tooth Wear Index (TWI) |
| Brazil | Frazao et al. | 2018 | 6–10 | 239 | 11.7 | BEWE |
| Brazil | Mangueira et al. | 2009 | 6–12 | 983 | 19.9 | O'Sullivan index |
| Brazil | Salas et al. | 2014 | 8–12 | 1,211 | 25.0 | O'Sullivan index |
| Brazil | Vargas-Ferreira et al. | 2011 | 11–14 | 944 | 7.2 | O'Sullivan index |
| Brazil | Peres et al. | 2005 | 12 | 499 | 13.0 | O'Sullivan index |
| Brazil | Correr et al. | 2009 | 12 | 389 | 26.0 | O'Sullivan index |
| Brazil | Alves et al. | 2015 | 12 | 1,528 | 15.0 | BEWE |
| Brazil | Gurgel et al. | 2011 | 12 & 16 | 414 | 20.0 | O'Brien index |
| Brazil | Auad et al. | 2007 | 13–14 | 458 | 34.1 | O'Sullivan-Index |
| Brazil | Pedraño et al. | 2018 | 35–74 | 207 | 58.9 | BEWE |
| Chile | Dugmore et al. | 2003 | 12 | 1,753 | 56.3 | O'Brien index |
| Chile | Marró et al. | 2020 | 18–46 | 535 | 97.9 | BEWE |
| Colombia | Mafla et al. | 2017 | 10–15 | 384 | 57.3 | O'Sullivan index |
| Colombia | Avila et al. | 2019 | 12–15 | 176 | 73.9 | BEWE |
| Colombia | Martignon et al. | 2019 | 18–25 | 601 | 73.0 | BEWE |
| Ecuador | Martínez et al. | 2018 | 8–12 | 175 | 53.1 | O'Brien index |
| Mexico | Pineda et al. | 2019 | 11–14 | 512 | 63.9 | BEWE |
| México | Pineda et al. | 2016 | 14–19 | 417 | 31.7 | Lussi index |
| Peru | Torres-Vargas et al. | 2012 | Adults | 150 | 30.0 | Lussi index |
| Uruguay | Alvarez Loureiro et al. | 2015 | 12 | 1,136 | 52.9 | BEWE |
| USA | Habib M et al. | 2013 | 2–4 & 12 | 243 | 12.0 | Tooth surface loss |
| USA | Deery et al. | 2000 | 11–13 | 129 | 41.0 | O'Brien index |
| USA | Mungia et al. | 2009 | 12–17 | 307 | 5.5 | Tooth Wear Index |
| USA | Okunseri et al. | 2011 | 13–19 | 1,314 | 45.0 | Smith and Knight tooth wear index |
| USA | McGuire et al. | 2009 | 13–19 | 1,962 | 45.9 | Smith and Knight tooth wear index |

BEWE is predominantly a screening tool. Dentine exposure is calculated in a similar way to the highest BEWE score with one surface with dentine exposed used to calculate the prevalence. Using this method dentine exposure ranges from 2–45% and mostly on the occlusal and incisal surfaces.

The most common tooth surfaces are consistently the occlusal surfaces of the lower molars and the buccal surfaces of the upper central incisors [Schlueter and Luka, 2018]. This could be considered key teeth and may in large population studies reduce the time needed to screen for the condition. There has not been any comprehensive epidemiological study to analyze the prevalence across large populations. Most studies are based on convenience studies but even with this proviso the indication from multiple studies across different countries is that ETW is a common clinical finding.

The clinical management of ETW is wide ranging and complex depending on its associated origin and severity,

emphasizing on its regular routine examination and recording using the BEWE [Bartlett et al., 2008] and involving from avoiding acidic foods and drinks, to self- and professionally application of fluoride to restorative care [O'Toole et al., 2015, 2017, 2018].

Developmental Defects of Enamel

There are two major types of DDE – hypoplasia and hypomineralization. Hypoplasia occurs when the ameloblast's secretion of enamel matrix is interrupted, creating a quantitative defect – thin enamel. This normally occurs as pits or horizontal lines in the enamel – indicating the age at which the insult to the ameloblast occurred. Hypomineralization is a qualitative defect – lower mineral density occurs due to an insult to the ameloblast during the calcification or maturation phase, however the enamel is usually full thickness, although hypoplasia and hypomin-

Table 2. The prevalence of fluorosis in the Americas national reports

| Country | Authors | Year | Age, years | Sample size | Defect prevalence | Criteria |
|-------------|---|------|--------------|-------------------------|---------------------|--------------|
| Bolivia | Ministry of Health and Social Welfare | 1997 | 6–15 | 2,707 | 40.8 | Dean |
| Canada | Canadian Health Measures Survey (CHMS) | 2010 | <12 | 5,604 | 37.1 | Dean |
| Colombia | Ministry of Health and Social Protection | 2014 | 5, 12 & 15 | 2,276 2,183 2,046 | 8.4 65.1 56.1 | Dean |
| Costa Rica | Ministry of Health | 2002 | 6–8, 12 & 15 | 3,780 | 5.7 31.9 24.4 | Dean |
| Ecuador | Ministry of Public Health | 2009 | 12 & 15 | 700 | 5.0 | Dean |
| El Salvador | Ministry of Public Health and Social Assistance | 2000 | 12–15 | 2,000 | 5.2 | Dean |
| Mexico | Secretary of Health | 2017 | >6 | 307,993 | 2.6 | Not reported |
| Nicaragua | Pan American Health Organization – a | 1999 | 12–15 | 292 | 16.7 | Dean |
| Paraguay | Ministry of Public Health and Social Welfare | 2017 | 12 & 15 | 1,340 | 14.4 | Dean |
| USA | National Center for Health Statistics | 2019 | 6–19 | 7,158 | 85.7 | Dean |
| Venezuela | Pan American Health Organization – b | 1998 | 6–8, 12 & 15 | 4,462 | 15.0 | Dean |

eralization can occur in the same lesion. Hypomineralized lesions can be diffuse (such as fluorosis) or demarcated (such as MIH). Both primary and permanent dentitions can be affected [Wright et al., 2015].

Dental Fluorosis

General Aspects

At appropriate levels, fluoride has been established as an effective agent in the prevention of dental caries [ADA, 2019]. However, excessive intake of fluoride during tooth formation may result in the development of fluorosis. Histologically, enamel fluorosis presents as a subsurface hypomineralization with various degrees of subsurface porosity. There is contradictory evidence on the relationship between the concentration of fluoride in enamel and fluorosis severity [Richards et al., 1989, 1992; Vieira et al., 2004, 2005]. However, more recent research has indicated that a positive linear relationship does exist [Martinez-Mier et al., 2016].

Clinically, the first signs of enamel fluorosis present as thin white striae across teeth surfaces, which follow the perikymata pattern. In mild fluorosis, the cusps, tips, incisal edges, and marginal ridges may be completely

opaque, a sign called “snow capping.” In moderate cases, the white lines appear more pronounced and may merge to produce cloudy areas scattered over the surface. As severity increases, the entire tooth surface exhibits opaque cloudy areas mixed with areas of brownish discoloration and there may be pitting [Thylstrup and Fejerskov, 1978].

The differential diagnosis of enamel fluorosis is complex and includes caries, MIH, and other non-fluoride opacities. Multiple indices have been developed for the visual examination of enamel fluorosis. In 1934, Dean developed the first index to describe the condition [Dean, 1934]. In 1978, the Thylstrup and Fejerskov Index (TFI) was introduced; and, in 1984, the Tooth Surface Index of Fluorosis was developed [Thylstrup and Fejerskov, 1978; Horowitz et al., 1984]. Since then, multiple efforts have been focused on the development of methods to identify and quantify fluorosis using imaging technology [Pretty et al., 2012; Cuevas-Espinoza et al., 2017].

Epidemiology

Table 2 summarizes selected national reports made in American countries. In the United States, assessments of changes in the prevalence of enamel fluorosis have generally found increases at all levels of severity over time [Clark, 1994; Beltrán-Aguilar et al., 2010; Wiener et al.,

2018]. Larger increases in severity and prevalence have been found in the most recent national surveys. Total fluorosis prevalence increased from 22 to 41 to 65% in the 1986–1987, 1999–2004, and 2011–2012 National Health and Nutrition Examination Survey (NHANES) surveys, respectively [Neurath et al., 2019]. However, a quality assessment of the enamel fluorosis clinical data concluded that factors other than a true change in prevalence may explain the observed differences, including potential examiner subjectivity [National Center for Health Statistics, 2019]. In Mexico, clinical examinations for enamel fluorosis are not included in national health surveys; however, two review articles presented prevalence data. These reviews included data from naturally fluoridated areas where water fluoride was above optimal levels as well as areas where water has negligible, unknown or optimal fluoride levels and where fluoridated salt was distributed [Soto et al., 2004; Aguilar-Díaz et al., 2017]. Studies span over two decades and the prevalence ranged from 40 to 100%. These studies point towards a potential increase in both the prevalence and severity of fluorosis over time.

In Colombia, the Community Fluorosis Index (CFI) has been used to measure the burden of dental fluorosis in more than 70 communities as part of a sentinel monitoring program for the national salt fluoridation program. The reported CFIs have ranged from 0.60 to 3.0 [Tovar & Misnaza, 2016]. A scoping review of fluorosis prevalence in Colombia reported a range in prevalence from 0% to 98% [Agudelo-Suárez et al., 2013] and an epidemiological study conducted in a fluorosis endemic Colombian region included besides dental fluorosis with the TFI the assessment of caries lesions with the ICDAS criteria [Tellez et al., 2012].

Enamel fluorosis development has been linked to prolonged, excessive ingestion from varied sources of fluoride [Bronckers et al., 2009]. Potential sources of fluoride include water for drinking and cooking, fluoridated milk and salt, dentifrice ingestion, environmental pollution, pesticides, and professionally applied fluoride products, among others. In the United States, Health and Human Services (HHS) and the Environmental Protection Agency (EPA) conducted a review of standards and guidelines on fluoride in drinking water. As part of this review, the EPA completed and peer-reviewed a study on environmental exposure to fluoride and the relative source contribution of known sources of fluoride [EPA, 2010]. Water, food, beverages, sulfuryl fluoride, toothpaste and soils were identified as major sources of fluoride. In Mexico and Colombia, no national reviews on relative source contribution have been conducted thus far. Selected stud-

ies on fluoride intake have identified diet and dental products as major sources of fluoride. Toothpaste ingestion in young children has been identified as a major source of fluoride in both countries [Martinez-Mier et al., 2003; Franco et al., 2005].

The clinical management of fluorosis depends on its severity [Alhawij et al., 2015; Marín et al., 2016], the individual's caries risk and the esthetic patient's perspective. While mild cases do not require treatment, more severe forms of fluorosis would, including preventive, microinvasive and operative care [Shahroom et al., 2019].

A dental epidemiological transition is underway at the global level and research is needed to describe it. There is a need to characterize the prevalence of enamel fluorosis and to identify current sources of fluoride intake and their relative source contributions in the Americas.

The clinical management of fluorosis depends on its severity [Alhawij et al., 2015; Marín et al., 2016], the individual's caries risk and the esthetic patient's perspective. While mild cases do not require treatment, more severe forms of fluorosis would, including preventive, microinvasive and operative care [Shahroom et al., 2019].

Molar-Incisor Hypomineralization

General Aspects

In the past 20 years, increased interest has been engendered in the dental research community regarding the DDE termed molar incisor hypomineralization (MIH) and defined as a demarcated hypomineralized defect on at least one first permanent molar, often also affecting the permanent incisors [Weerheijm et al., 2001]. Other less-used terms with respect to MIH include cheesy molars, idiopathic enamel hypomineralization and non-endemic mottling [Weerheijm et al., 2001].

There is variability in the number of teeth affected in MIH, with a tendency for incisors to be affected more frequently when the molars are more severely affected [Jälevik et al., 2001]. Severity can vary both within the individual but also within the tooth. This feature is confusing with respect to the pathogenesis of the lesion. It has been hypothesized that MIH is influenced by multiple genes and the environment within a developmental system that can be affected in various ways which allows the variable phenotype to occur [Vieira and Manton, 2019].

Although MIH is the most prevalent presentation of demarcated hypomineralized defects, all teeth in both primary and permanent dentitions can be affected by similar lesions [Kühnisch et al., 2014], highlighting the

Table 3. The prevalence of molar incisor hypomineralization and demarcated opacities in the Americas

| Country | Authors | Year | Age, years | Sample size | Defect prevalence, % | Criteria |
|-----------|---------------------------------|------|------------|-------------|----------------------|-----------------------|
| Argentina | López et al. | 2014 | 7–17 | 1,090 | 16.1 | DDE |
| Argentina | Biondi et al. | 2011 | 11.3 | 1,098 | 15.9 | DDE |
| Argentina | Biondi et al. | 2012 | 11.6 | 512 | 6.4 | DDE |
| Brazil | da Silva Junior et al. | 2015 | 5–17 | 260 | 8.8 | EAPD |
| Brazil | Sé | 2017 | 6–11 | 858 | 14.7 | EAPD |
| Brazil | da Costa-Silva et al. | 2010 | 6–12 | 918 | 19.8 | EAPD |
| Brazil | Souza et al. | 2012 | 6–12 | 903 | 19.8 | EAPD |
| Brazil | Jeremias et al. | 2013 | 6–12 | 1,157 | 12.3 | EAPD |
| Brazil | Souza et al. | 2013 | 7–12 | 1,151 | 12.3 | EAPD |
| Brazil | Soviero et al. | 2009 | 7–13 | 249 | 40.2 | EAPD |
| Brazil | Rodrigues et al. | 2015 | 7–14 | 1,179 | 2.5 | mDDE* |
| Brazil | Portella et al. | 2019 | 8 | 728 | 12.1 | EAPD |
| Brazil | Tourino et al. | 2016 | 8–9 | 1,181 | 20.4 | EAPD |
| Brazil | Dantas-Neta et al. | 2018 | 8–10 | 744 | 25.0 | EAPD |
| Brazil | Vargas-Ferreira et al. | 2018 | 8–12 | 1,206 | 26.4 | mDDE |
| Brazil | De Lima et al. | 2015 | 11–14 | 594 | 18.4 | EAPD |
| Canada | Sidhu et al. | 2019 | 0–17 | 233 | 12.4 | EAPD |
| Chile | Corral-Núñez et al. | 2016 | 6–12 | 851 | 12.7 | EAPD |
| Chile | Gambetta et al. | 2019 | 6–12 | 577 | 15.8 | EAPD |
| Chile | Jans-Muñoz et al. | 2011 | 6–13 | 334 | 16.8 | Weerheijm |
| Colombia | Mejía et al. | 2019 | 6–15 | 1,075 | 11.2 | EAPD |
| Colombia | Beltran et al. | 2019 | 12–15 | 176 | 13.6 | EAPD |
| Mexico | Gurrusquieta et al. | 2017 | 6–12 | 1,156 | 15.8 | EAPD |
| Mexico | Villanueva-Gutiérrez et al. | 2019 | 7–12 | 686 | 35.4 | EAPD |
| Peru | Vásquez-Muñoz & Aguilar-Gálvez. | 2014 | 6–13 | 970 | 10.0 | EAPD |
| USA | Davenport et al. | 2019 | 6–14 | 375 | 9.6 | EAPD |
| USA | Tagelsir et al. | 2019 | 6–15 | 266 | 12.0 | EAPD |
| Uruguay | López et al. | 2014 | 7–17 | 626 | 12.3 | Mathu-Muju and Wright |
| Uruguay | Biondi et al. | 2012 | 11 | 463 | 7.1 | Mathu-Muju and Wright |

* mDDE will have demarcated opacities on first permanent molars and/or incisors.

importance of hypomineralized second primary molars (HSPM) [Elfrink et al., 2012].

Hypomineralized enamel can have markedly altered physical characteristics, with up to 70% mineral density deficit, increased protein content (mainly serum albumin), decreased hardness and lower fracture resistance, more carbonate, and an altered elemental profile [Robinson et al., 1992; Mangum et al., 2010; Crombie et al., 2013; Elhennawy et al., 2017].

Approximately one third of MIH lesions are severe, with increasing severity being associated with increased caries experience and burden, as well as rapid lesion breakdown, especially in high-caries risk individuals [Jälevik et al., 2001; Kuhnisch et al., 2018; Paixão-Gonçalves et al., 2019].

The etiological factors involved in MIH (and HSPM) are environmental and genetic in nature, including infan-

tile illnesses, especially hyper-pyrexia, asthma and pneumonia [Crombie et al., 2009; Alaluusua, 2010]. In a recent twins study, after potential confounding factors were controlled for, vitamin D levels at birth, in vitro fertilization, infantile eczema, socioeconomic position, dizygosity and maternal smoking beyond the first trimester of pregnancy demonstrated the strongest associations with HSPM [Silva et al., 2019].

Epidemiology

The prevalence of MIH in the Americas varies from 2.5% to 40.2%, with most studies finding between 10 and 20% prevalence; however, as different indices have been used, direct comparison of data must be done with caution (Table 3). The majority of data come from Brazil, and until recently, no USA data was published, however, several studies are currently underway. The more prevalent

indices being the modified DDE index and the EAPD MIH index [Clarkson and O'Mullane, 1992; Ghanim et al., 2019]. Worldwide prevalence figures of 13.1% and 14.2% have been recently reported [Schwendicke et al., 2018; Zhao et al., 2018].

MIH affected teeth place a great burden on individuals, requiring nearly 10 times as much intervention than teeth in non-affected individuals by the age of 9 years [Jälevik and Klingberg, 2002] and by 18 years, affected molars had been treated 4.2 times as often as the controls [Jälevik and Klingberg, 2012]. There is a direct association between MIH presence and increased caries experience and treatment burden [Elhennawy and Schwendicke, 2016; Americano et al., 2017; Kuhnisch et al., 2018]. Hypomineralized teeth can be increasingly sensitive with increased severity, also leading to poor oral hygiene and increased risk of breakdown [Raposo et al., 2019].

Finally, we can hope for early identification of MIH/HSPM, possibly through genetic means – even pre-eruptive diagnosis, which could allow support of affected ameloblasts to minimize the mineral deficit; and development of remineralizing techniques to prevent lesion breakdown after eruption.

Clinical management options for MIH-affected teeth vary considerably, depending on severity and whether they are molars or incisors. Treatment options include from non-operative to operative care and extraction [Baroni et al., 2014; Crombie et al., 2014].

Conclusion

The growing worldwide interest in non-carious lesions such as dental fluorosis, MIH and ETW leads to changes in their approach, both at the clinical and at the community level. Although the number of studies related to these conditions in the American continent has been increasing lately, there is still a need to increase awareness of their accurate diagnosis and management in order to include non-carious lesions in routine dental practice to ensure the maintenance of oral health. Additionally, correct diagnosis is crucial for systematic data collection favoring epidemiological surveillance reliability. Recommendations regarding diagnostic indices to be used during assessment are given, improving this process and helping clinicians' adherence. The most commonly used indices are – BEWE for ETW assessment and quantification, Dean or Thylstrup & Fejerskov indices for dental fluorosis and the EAPD criteria for MIH. Besides, since a wider range of indices has been used, this per se can lead to variation of prevalence data. Thus, a consensus for

diagnostic criteria in these three conditions is needed, as already proposed by international groups such as the Erosive Tooth Wear Foundation [Erosive Tooth Wear Foundation, 2020] or the D3 Group [The D3 Group, 2020]. Finally, it is important for dental practitioners to develop skills and knowledge in the diagnosis, quantification and management of non-carious dental disorders, taking into account that their clinical management should be focused on prevention and mitigation of associated consequences as well as control of progression, in order to preserve tooth structure.

Acknowledgements

We acknowledge LAOHA (Latin American Oral Health Association) – Colgate Palmolive for funding the SAS, the ORCA Board for its approval and support, and UNICA for the logistic and local support.

Conflict of Interest Statement

No potential conflicts of interest exist for any author regarding this symposium paper.

Author Contributions

Stefania Martignon: Conception and planning, analysis and interpretation of the epidemiological data, drafting the article, revising critically the article, approval of the version to be published.

David Bartlett: Analysis and interpretation of the Erosive Tooth Wear data, drafting the article, revising critically the article, approval of the version to be published.

David J. Manton: Analysis and interpretation of the Molar Incisor Hypomineralization data, drafting the article, revising critically the article, approval of the version to be published.

E. Angeles Martinez-Mier: Analysis and interpretation of the Dental Fluorosis data, drafting the article, revising critically the article, approval of the version to be published.

Christian Splieth: Revising critically the article, approval of the version to be published.

Viviana Avila: Conception and planning, analysis and interpretation of the epidemiological data, drafting the article, revising critically the article, approval of the version to be published.

References

- ADA - American Dental Association. [Internet]. Council on Access, Prevention and Interpersonal Affairs: Fluoridation Facts 2019. [cited 2019 October 10]. Available from: www.ada.org
- Agudelo-Suárez A, Martínez-Flórez L, Madrid-Gutiérrez L, Vivares-Builes A, Rocha-Buelvas A. Panorama de la fluorosis dental en Colombia: una revisión exploratoria de la literatura. *Univ Odontol*. 2013;32:133–45.

- Aguilar-Díaz FD, Morales-Corona F, Cintra-Vieiro AC, Fuente-Hernández J. Prevalence of dental fluorosis in Mexico 2005-2015: a literature review. *Salud Publica Mex.* 2017 May-Jun;59(3):306-13.
- Alaluusua S. Aetiology of Molar-Incisor Hypomineralisation: A systematic review. *Eur Arch Paediatr Dent.* 2010 Apr;11(2):53-8.
- Alhawij H, Lippert F, Martínez-Mier EA. Relative fluoride response of caries lesions created in fluorotic and sound teeth studied under remineralizing conditions. *J Dent.* 2015 Jan;43(1):103-9.
- Alvarez Loureiro L, Fabruccini Fager A, Alves LS, Alvarez Vaz R, Maltz M. Erosive tooth wear among 12-year-old schoolchildren: a population-based cross-sectional study in Montevideo, Uruguay. *Caries Res.* 2015;49(3):216-25.
- Alves LS, Brusius CD, Damé-Teixeira N, Maltz M, Susin C. Dental erosion among 12-year-old schoolchildren: a population-based cross-sectional study in South Brazil. *Int Dent J.* 2015 Dec;65(6):322-30.
- Americano GC, Jacobsen PE, Soviero VM, Haubek D. A systematic review on the association between molar incisor hypomineralization and dental caries. *Int J Paediatr Dent.* 2017 Jan;27(1):11-21.
- Auad SM, Waterhouse PJ, Nunn JH, Steen N, Moynihan PJ. Dental erosion amongst 13- and 14-year-old Brazilian schoolchildren. *Int Dent J.* 2007 Jun;57(3):161-7.
- Avila V, Beltran E, Castellanos J, Lara J, Martignon S. Erosive tooth wear and risk factors in 12-15 years old schoolchildren in Bogota, Colombia. *Caries Res.* 2019;53:357-410.
- Awad MA, El Kassas D, Al Harthi L, Abraham SB, Al-Khalifa KS, Khalaf ME, et al. Prevalence, severity and explanatory factors of tooth wear in Arab populations. *J Dent.* 2019 Jan;80:69-74.
- Baroni C, Bazzocchi M, Marchionni S, Bedini R, Pecci R, Manton D. MIH supplementation strategies. *Dent Mater.* 2014;30:e87-8.
- Bartlett D, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Invest.* 2008 Mar;12(S1 Suppl 1):S65-8.
- Bartlett DW, Coward PY, Nikkah C, Wilson RF. The prevalence of tooth wear in a cluster sample of adolescent schoolchildren and its relationship with potential explanatory factors. *Br Dent J.* 1998 Feb;184(3):125-9.
- Bartlett DW, Lussi A, West NX, Bouchard P, Sanz M, Bourgeois D. Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. *J Dent.* 2013 Nov;41(11):1007-13.
- Beltrán E, Avila V, Úsuga-Vacca M, Cortés A, Martignon S. Prevalence of MIH and dental fluorosis in 12-15 years old schoolchildren from Bogotá, Colombia. *Caries Res.* 2019;53:357-410.
- Beltrán-Aguilar ED, Barker L, Dye BA. Prevalence and severity of dental fluorosis in the United States, 1999-2004. *NCHS Data Brief.* 2010 Nov;53(53):1-8.
- Biondi AM, Cortese SG, Martínez K, Ortolani AM, Sebelli PM, Ienco M, et al. Prevalence of molar incisor hypomineralization in the city of Buenos Aires. *Acta Odontol Latinoam.* 2011;24(1):81-5.
- Biondi AM, López Jordi MC, Cortese SG, Alvarez L, Salveraglio I, Ortolani AM. Prevalence of molar-incisor hypomineralization (MIH) in children seeking dental care at the Schools of Dentistry of the University of Buenos Aires (Argentina) and University of la Republica (Uruguay). *Acta Odontol Latinoam.* 2012; 25(2):224-30.
- Bronckers AL, Lyaruu DM, DenBesten PK. The impact of fluoride on ameloblasts and the mechanisms of enamel fluorosis. *J Dent Res.* 2009 Oct;88(10):877-93.
- Canadian Health Measures Survey (CHMS). Statistics Canada. [Internet]. 2010. Canadian Health Measures Survey (CHMS) Cycle 1: Data Dictionary. [cited 2019 Oct 30]. Available from: https://library.carleton.ca/sites/default/files/find/data/surveys/pdf_files/chms-c1-07-09-dic.pdf
- CCI - CariesCare International. [Internet]. Official website 2020. [cited 2020 April 23]. Available from: <https://cariescareinternational.com/>
- Clark DC. Trends in prevalence of dental fluorosis in North America. *Community Dent Oral Epidemiol.* 1994 Jun;22(3):148-52.
- Clarkson JJ, O'Mullane DM. Prevalence of enamel defects/fluorosis in fluoridated and non-fluoridated areas in Ireland. *Community Dent Oral Epidemiol.* 1992 Aug;20(4):196-9.
- Corral-Núñez C, Rodríguez H, Cabello R, Bersezio-Miranda C, Cordeiro R, Fresno-Rivas M. Impacto de la hipomineralización incisivo molar en la experiencia de caries en escolares de 6-12 años en Santiago, Chile. *Rev Clín Periodoncia Implantol Rehabil Oral.* 2016; 9(3):277-83.
- Correr GM, Alonso RC, Correa MA, Campos EA, Baratto-Filho F, Puppim-Rontani RM. Influence of diet and salivary characteristics on the prevalence of dental erosion among 12-year-old schoolchildren. *J Dent Child (Chic).* 2009 Sep-Dec;76(3):181-7.
- Crombie F, Manton D, Kilpatrick N. Aetiology of molar-incisor hypomineralization: a critical review. *Int J Paediatr Dent.* 2009 Mar;19(2):73-83.
- Crombie F, Manton D, Palamara J, Reynolds E. Resin infiltration of developmentally hypomineralised enamel. *Int J Paediatr Dent.* 2014 Jan;24(1):51-5.
- Crombie FA, Manton DJ, Palamara JE, Zalzniak I, Cochrane NJ, Reynolds EC. Characterisation of developmentally hypomineralised human enamel. *J Dent.* 2013 Jul;41(7):611-8.
- Cuevas-Espinosa DM, Martínez-Mier EA, Ando M, Castiblanco GA, Cortes F, Rincon-Bermudez CM, et al. In vitro Validation of Quantitative Light-Induced Fluorescence for the Diagnosis of Enamel Fluorosis in Permanent Teeth. *Caries Res.* 2017;51(5):515-26.
- da Costa-Silva CM, Jeremias F, de Souza JF, Cordeiro RC, Santos-Pinto L, Zuanon AC. Molar incisor hypomineralization: prevalence, severity and clinical consequences in Brazilian children. *Int J Paediatr Dent.* 2010 Nov;20(6):426-34.
- Dantas-Neta N, Soares-Figueiredo M, Lima C, Bendo C, Matos de Andrade E, Lima M. Factors associated with molar-incisor hypomineralisation in schoolchildren aged 8-10 years: a case-control study. *Int J Paediatr Dent* 2018; 28:570-577.
- Davenport M, Welles AD, Angelopoulou MV, Gonzalez C, Okunseri C, Barbeau L, et al. Prevalence of molar-incisor hypomineralization in Milwaukee, Wisconsin, USA: a pilot study. *Clin Cosmet Investig Dent.* 2019 May; 11:109-17.
- De Carvalho Sales-Peres SH, De Carvalho Sales-Peres A, Marsicano JA, De Moura-Grec PG, De Carvalho CA, De Freitas AR, et al. An epidemiological scoring system for tooth wear and dental erosive wear. *Int Dent J.* 2013 Jun; 63(3):154-60.
- de Lima MD, Andrade MJ, Dantas-Neta NB, Andrade NS, Teixeira RJ, de Moura MS, et al. Epidemiologic study of molar-incisor hypomineralization in schoolchildren in Northeastern Brazil. *Pediatr Dent.* 2015 Nov-Dec; 37(7):513-9.
- Dean H. Classification of mottled enamel diagnosis. *J Am Dent Assoc.* 1934;21:1421-6.
- Deery C, Wagner ML, Longbottom C, Simon R, Nugent ZJ. The prevalence of dental erosion in a United States and a United Kingdom sample of adolescents. *Pediatr Dent.* 2000 Nov-Dec;22(6):505-10.
- Dugmore CR, Rock WP. The progression of tooth erosion in a cohort of adolescents of mixed ethnicity. *Int J Paediatr Dent.* 2003 Sep;13(5):295-303.
- Elfrink ME, ten Cate JM, Jaddoe VW, Hofman A, Moll HA, Veerkamp JS. Deciduous molar hypomineralization and molar incisor hypomineralization. *J Dent Res.* 2012 Jun;91(6):551-5.
- Elhennawy K, Manton DJ, Crombie F, Zaslansky P, Radlanski RJ, Jost-Brinkmann PG, et al. Structural, mechanical and chemical evaluation of molar-incisor hypomineralization-affected enamel: A systematic review. *Arch Oral Biol.* 2017 Nov;83:272-81.
- Elhennawy K, Schwendicke F. Managing molar-incisor hypomineralization: A systematic review. *J Dent.* 2016 Dec;55:16-24.
- EPA - Environmental Protection Agency. [Internet]. Health and Ecological Criteria Division Office of Water: Fluoride: Exposure and Relative Source Contribution Analysis. 820-R-10-015. 2010. [cited 2019 October 10]. Available from: <https://www.epa.gov/sites/production/files/2019-03/documents/fluoride-exposure-relative-report.pdf>
- Erosive Tooth Wear Foundation. [Internet]. Official website. 2020. [cited 2020 March 23]. Available from: <https://www.erosivetoothwear.com/>

- Fares J, Shirodaria S, Chiu K, Ahmad N, Sherriff M, Bartlett D. A new index of tooth wear. Reproducibility and application to a sample of 18- to 30-year-old university students. *Caries Res*. 2009;43(2):119–25.
- Franco AM, Martignon S, Saldarriaga A, González MC, Arbeláez MI, Ocampo A, et al. Total fluoride intake in children aged 22-35 months in four Colombian cities. *Community Dent Oral Epidemiol*. 2005 Feb;33(1):1–8.
- Frazao JB, Machado LG, Ferreira MC. Dental erosion in schoolchildren and associated factors: A cross-sectional study. *J Indian Soc Pedod Prev Dent*. 2018 Apr-Jun;36(2):113–9.
- Gambetta-Tessini K, Mariño R, Ghanim A, Calache H, Manton DJ. The impact of MIH/HSPM on the carious lesion severity of schoolchildren from Talca, Chile. *Eur Arch Paediatr Dent*. 2019 Oct;20(5):417–23.
- Ganss C, Klimek J, Lussi A. Accuracy and consistency of the visual diagnosis of exposed dentine on worn occlusal/incisal surfaces. *Caries Res*. 2006;40(3):208–12.
- Ghanim A, Mariño R, Manton DJ. Validity and reproducibility testing of the Molar Incisor Hypomineralisation (MIH) Index. *Int J Paediatr Dent*. 2019 Jan;29(1):6–13.
- González-Aragón Pineda ÁE, Borges-Yáñez SA, Irigoyen-Camacho ME, Lussi A. Relationship between erosive tooth wear and beverage consumption among a group of schoolchildren in Mexico City. *Clin Oral Investig*. 2019 Feb; 23(2):715–23.
- González-Aragón Pineda ÁE, Borges-Yáñez SA, Lussi A, Irigoyen-Camacho ME, Angeles Medina F. Prevalence of erosive tooth wear and associated factors in a group of Mexican adolescents. *J Am Dent Assoc*. 2016 Feb;147(2): 92–7.
- Gurgel CV, Rios D, Buzalaf MA, da Silva SM, Araújo JJ, Pauletto AR, et al. Dental erosion in a group of 12- and 16-year-old Brazilian schoolchildren. *Pediatr Dent*. 2011 Jan-Feb; 33(1):23–8.
- Gurrusquieta BJ, Núñez VM, López ML. Prevalence of molar incisor hypomineralization in Mexican children. *J Clin Pediatr Dent*. 2017; 41(1):18–21.
- Habib M, Hottel TL, Hong L. Prevalence and risk factors of dental erosion in American children. *J Clin Pediatr Dent*. 2013;38(2):143–8.
- Horowitz HS, Driscoll WS, Meyers RJ, Heifetz SB, Kingman A. A new method for assessing the prevalence of dental fluorosis—the Tooth Surface Index of Fluorosis. *J Am Dent Assoc*. 1984 Jul;109(1):37–41.
- ICCMSTM - International Caries Classification and Management System. [Internet]. Maintaining and preserving tooth structure. Official website 2020. [cited 2020 April 23]. Available from: <https://www.iccms-web.com/>
- Jälevik B, Klingberg G, Barregård L, Norén JG. The prevalence of demarcated opacities in permanent first molars in a group of Swedish children. *Acta Odontol Scand*. 2001 Oct; 59(5):255–60.
- Jälevik B, Klingberg G. Treatment outcomes and dental anxiety in 18-year-olds with MIH, comparisons with healthy controls - a longitudinal study. *Int J Paediatr Dent*. 2012 Mar; 22(2):85–91.
- Jälevik B, Klingberg GA. Dental treatment, dental fear and behaviour management problems in children with severe enamel hypomineralization of their permanent first molars. *Int J Paediatr Dent*. 2002 Jan;12(1):24–32.
- Jans Muñoz A, Díaz Meléndez J, Vergara González C, Zaror Sánchez C. Frecuencia y severidad de la hipomineralización molar incisal en pacientes atendidos en las clínicas odontológicas de la Universidad de La Frontera, Chile. *Int J Odontostomatol*. 2011;5(2):133–40.
- Jeremias F, de Souza JF, Silva CM, Cordeiro RC, Zuanon AC, Santos-Pinto L. Dental caries experience and Molar-Incisor Hypomineralization. *Acta Odontol Scand*. 2013 May-Jul; 71(3-4):870–6.
- Kreulen CM, Van 't Spijker A, Rodriguez JM, Bronkhorst EM, Creugers NH, Bartlett DW. Systematic review of the prevalence of tooth wear in children and adolescents. *Caries Res*. 2010;44(2):151–9.
- Kühnisch J, Heitmüller D, Thiering E, Brockow I, Hoffmann U, Neumann C, et al. Proportion and extent of manifestation of molar-incisor-hypomineralizations according to different phenotypes. *J Public Health Dent*. 2014;74(1): 42–9.
- Kühnisch J, Kabary L, Malyk Y, Rothmaier K, Metz I, Hickel R, et al. Relationship between caries experience and demarcated hypomineralised lesions (including MIH) in the permanent dentition of 15-year-olds. *Clin Oral Investig*. 2018 Jun;22(5):2013–9.
- López Jordi MC, Cortese SG, Álvarez L, Salveraglio I, Ortolani AM, Biondi AM. Comparación de la prevalencia de hipomineralización molar incisiva en niños con diferente cobertura asistencial en las ciudades de Buenos Aires (Argentina) y Montevideo (Uruguay). *Salud Colect*. 2014 Aug;10(2):243–51.
- Mafla AC, Cerón-Bastidas XA, Muñoz-Ceballos ME, Vallejo-Bravo DC, Fajardo-Santacruz MC. Prevalence and Extrinsic Risk Factors for Dental Erosion in Adolescents. *J Clin Pediatr Dent*. 2017;41(2):102–11.
- Mangueira DF, Sampaio FC, Oliveira AF. Association between socioeconomic factors and dental erosion in Brazilian schoolchildren. *J Public Health Dent*. 2009;69(4):254–9.
- Mangum JE, Crombie FA, Kilpatrick N, Manton DJ, Hubbard MJ. Surface integrity governs the proteome of hypomineralized enamel. *J Dent Res*. 2010 Oct;89(10):1160–5.
- Marín LM, Cury JA, Tenuta LM, Castellanos JE, Martignon S. Higher fluorosis severity makes enamel less resistant to demineralization. *Caries Res*. 2016;50(4):407–13.
- Marró ML, Aránguiz V, Ramirez V, Lussi A. Prevalence of erosive tooth wear in Chilean adults, 2016: A cross-sectional study. *J Oral Rehabil*. 2020 Apr;47(4):467–72.
- Martignon S, López-Macias AM, Bartlett D, Pitts N, Usuga-Vacca M, Gamboa LF, et al. The use of index teeth vs. full mouth in erosive tooth wear to assess risk factors in the diet: A cross-sectional epidemiological study. *J Dent*. 2019 Sep;88:103164.
- Martínez J, Armas A, Aguilera F, Tello G. Prevalencia y factores asociados del desgaste dental erosivo en niños de 8-12 años del norte de Quito, Ecuador. *Rev Odontol*. 2018;20:61–74.
- Martinez-Mier EA, Shone DB, Buckley CM, Ando M, Lippert F, Soto-Rojas AE. Relationship between enamel fluorosis severity and fluoride content. *J Dent*. 2016 Mar;46:42–6.
- Martínez-Mier EA, Soto-Rojas AE, Ureña-Cirett JL, Stookey GK, Dunipace AJ. Fluoride intake from foods, beverages and dentifrice by children in Mexico. *Community Dent Oral Epidemiol*. 2003 Jun;31(3):221–30.
- McGuire J, Szabo A, Jackson S, Bradley TG, Okunseri C. Erosive tooth wear among children in the United States: relationship to race/ethnicity and obesity. *Int J Paediatr Dent*. 2009 Mar;19(2):91–8.
- Mejía JD, Restrepo M, González S, Álvarez LG, Santos-Pinto L, Escobar A. Molar Incisor Hypomineralization in Colombia: Prevalence, Severity and Associated Risk Factors. *J Clin Pediatr Dent*. 2019;43(3):185–9.
- Ministry of Health - Ministerio de Salud. Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud. Caja Costarricense de Seguro Social. OPS. [Internet]. Encuesta nacional de salud oral: fluorosis de esmalte, 1999. Série de documentos técnicos del INCIENSA No 7. Tres Ríos, Costa Rica 2002. [cited 2019 Nov 10]. Available from: https://www.paho.org/hq/dmdocuments/2009/OH_COR_EncNacFluoresmal1999.pdf
- Ministry of Health and Social Protection - Ministerio de Salud y Protección Social. [Internet]. IV Estudio Nacional de Salud Bucal. ENSAB IV: Situación en Salud Bucal. Para Saber cómo estamos y saber qué hacemos. Colombia 2014. [cited 2019 Nov 10]. Available from: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/PP/ENSAB-IV-Situacion-Bucal-Actual.pdf>
- Ministry of Health and Social Welfare - Ministerio de Salud y Previsión Social Bolivia. [Internet]. Estudio Epidemiológico de Salud Bucal 1995. Bolivia 1997. [cited 2019 Nov 8]. Available from: https://www.paho.org/hq/dmdocuments/2009/OH_BOL_EstudEpidemSalud1995.pdf
- Ministry of Public Health - Ministerio de Salud Pública Del Ecuador. OPS. [Internet]. Programa Nacional de Fluoruración de la Sal. Estudio epidemiológico de salud bucal en escolares fiscales menores de 15 años del Ecuador. I parte- Estudio Descriptivo. Quito- Ecuador-1995-1996. Ecuador 2009. [cited 2019 Nov 10]. Available from: https://www.paho.org/hq/dmdocuments/2009/OH_ECU_EpidemEscolDesc1996.pdf

- Ministry of Public Health and Social Assistance - Ministerio de Salud Pública y Asistencia Social de El Salvador. [Internet]. Estudio epidemiológico de caries y fluorosis dental en escolares de 6, 7-8, 12 y 15 años de centros de enseñanza pública de El Salvador. Unidad de Salud Bucal. El Salvador, 2000. [cited 2019 Nov 08]. Available from: https://www.paho.org/hq/dmdocuments/2009/OH_ELS_EpidemCariesFluorEscol2000.pdf
- Ministry of Public Health and Social Welfare - Ministerio de Salud Pública y Bienestar Social. Encuesta Nacional de Salud Bucodental del Paraguay. [Internet]. ENSABUD PY 2017: Del diagnóstico al diseño y la ejecución de políticas públicas de salud bucodental. Asunción: OPS/OMS Paraguay 2018. [cited 2019 Nov 10]. Available from: http://portal.mspbs.gov.py/bucodental/wp-content/uploads/2018/09/BUCODENTAL-PY-_2017_FINAL_AGOSTO-2018.pdf
- Moimaz SA, Araújo PC, Chiba FY, Garbín CA, Saliba NA. Prevalence of deciduous tooth erosion in childhood. *Int J Dent Hyg*. 2013 Aug; 11(3):226-30.
- Mungia R, Zarzabal LA, Dang SC, Baez M, Stookley GK, Brown JP. Epidemiologic survey of erosive tooth wear in San Antonio, Texas. *Tex Dent J*. 2009 Nov;126(11):1097-109.
- Nahás Pires Corrêa MS, Nahás Pires Corrêa F, Nahás Pires Corrêa JP, Murakami C, Mendes FM. Prevalence and associated factors of dental erosion in children and adolescents of a private dental practice. *Int J Paediatr Dent*. 2011 Nov;21(6):451-8.
- National Center for Health Statistics, National Center for Chronic Disease Prevention and Health Promotion. [Internet]. Data quality evaluation of the dental fluorosis clinical assessment data from the National Health and Nutrition Examination Survey, 1999-2004 and 2011-2016. National Center for Health Statistics. *Vital Health Stat* 2(183). [cited 2019 Nov 20]. Available from: United States 2019. https://www.cdc.gov/nchs/data/series/sr_02/sr02_183-508.pdf
- Neurath C, Limeback H, Osmunson B, Connett M, Kanter V, Wells C. Enamel fluorosis Trends in US Oral Health Surveys: 1986 to 2012. *JDR Clin Trans Res*. 2019;4(4):298-308.
- O'Toole S, Bernabé E, Moazzez R, Bartlett D. Timing of dietary acid intake and erosive tooth wear: A case-control study. *J Dent*. 2017 Jan;56:99-104.
- O'Toole S, Mistry M, Mutahar M, Moazzez R, Bartlett D. Sequence of stannous and sodium fluoride solutions to prevent enamel erosion. *J Dent*. 2015 Dec;43(12):1498-503.
- O'Toole S, Pennington M, Varma S, Bartlett DW. The treatment need and associated cost of erosive tooth wear rehabilitation - a service evaluation within an NHS dental hospital. *Br Dent J*. 2018 Jun;224(12):957-61.
- Okunseri C, Okunseri E, Gonzalez C, Visotcky A, Szabo A. Erosive tooth wear and consumption of beverages among children in the United States. *Caries Res*. 2011;45(2):130-5.
- Paixão-Gonçalves S, Corrêa-Faria P, Ferreira FM, Ramos-Jorge ML, Paiva SM, Pordeus IA. Risk of Dental Caries in Primary Teeth with Developmental Defects of Enamel: A Longitudinal Study with a Multilevel Approach. *Caries Res*. 2019;53(6):667-74.
- Pan American Health Organization-a. World Health Organization. Ministerio de Salud. [Internet]. Estudio Epidemiológico de Salud Bucal en Niños de 6, 7, 8 12 y 15 Años de Escuelas y Colegios Públicos de Nicaragua. Nicaragua 1999. [cited 2019 Oct 30]. Available from: https://www.paho.org/hq/dmdocuments/2009/OH_NIC_EpidemSaludBucNin1999.pdf
- Pan American Health Organization-b. [Internet]. Estudio Basal de Prevalencia de caries y fluorosis dental en niños escolarizados. Venezuela 1997. Maracaibo, Venezuela 1998. [cited 2019 Oct 30]. Available from: http://new.paho.org/hq/dmdocuments/2009/OH_VENpreval_caries_fluorosis1997.pdf
- Pedrao AM, Andrews Portes L, Padilha Gomes E, Figueira Teixeira FC, da Costa Pereira A, de Oliveira NC. Erosive Tooth Wear and Dietary Patterns: A Clinical Study. *Oral Health Prev Dent*. 2018;16(2):145-51.
- Peres KG, Armênio MF, Peres MA, Traebert J, De Lacerda JT. Dental erosion in 12-year-old schoolchildren: a cross-sectional study in Southern Brazil. *Int J Paediatr Dent*. 2005 Jul; 15(4):249-55.
- Portella PD, Menoncin BL, de Souza JF, de Menezes JV, Fraiz FC, Assunção LR. Impact of molar incisor hypomineralization on quality of life in children with early mixed dentition: A hierarchical approach. *Int J Paediatr Dent*. 2019 Jul;29(4):496-506.
- Pretty IA, McGrady M, Zakian C, Ellwood RP, Taylor A, Sharif MO, et al. Quantitative light fluorescence (QLF) and polarized white light (PWL) assessments of dental fluorosis in an epidemiological setting. *BMC Public Health*. 2012 May;12(1):366.
- Raposo F, de Carvalho Rodrigues AC, Lia ÉN, Leal SC. Prevalence of Hypersensitivity in Teeth Affected by Molar-Incisor Hypomineralization (MIH). *Caries Res*. 2019;53(4):424-30.
- Richards A, Fejerskov O, Baelum V. Enamel fluoride in relation to severity of human dental fluorosis. *Adv Dent Res*. 1989 Sep;3(2):147-53.
- Richards A, Likimani S, Baelum V, Fejerskov O. Fluoride concentrations in unerupted fluorotic human enamel. *Caries Res*. 1992;26(5): 328-32.
- Robinson C, Kirkham J, Brookes SJ, Shore RC. The role of albumin in developing rodent dental enamel: a possible explanation for white spot hypoplasia. *J Dent Res*. 1992 Jun; 71(6):1270-4.
- Rodrigues F, Ribeiro P, Thomaz É, Lima G, Neves P, Ribeiro C. Molar-Incisor hypomineralization in schoolchildren of Sao Luis, Brazil Maranhao: prevalence and associated factors. *Pesq Bras Odontoped Clín Integr*. 2015;15(1): 271-8.
- Salas M, Dantas R, Sarmento H, Vargas-Ferreira F, Torriani D, Demarco F. Tooth erosion and dental caries in schoolchildren: is there a relationship between them? *Braz J Oral Sci*. 2014; 13(1):12-6.
- Schlueter N, Luka B. Erosive tooth wear - a review on global prevalence and on its prevalence in risk groups. *Br Dent J*. 2018 Mar;224(5):364-70.
- Schwendicke F, Elhennawy K, Reda S, Bekes K, Manton DJ, Krois J. Global burden of molar incisor hypomineralization. *J Dent*. 2018 Jan; 68:10-8.
- Sé M. [Internet]. Cárie dentária, hipomineralização molar incisivo, hipomineralização em deciduos e dor em escolares do Paranoá-DF. [PhD]. Universidade de Brasília, Universidade de Brasília 2017. [cited 2019 Nov 10]. Available from: [https://repositorio.unb.br/bitstream/10482/34963/1/2017_MariaJos% c3%a9daSilvaFigueir% c3%aadoS% c3%a9.pdf](https://repositorio.unb.br/bitstream/10482/34963/1/2017_MariaJos%c3%a9daSilvaFigueir%c3%aadoS%c3%a9.pdf)
- Secretary of Health - Secretaria de Salud. [Internet]. Resultados del Sistema de Vigilancia Epidemiológica de Patologías Bucales SIVEPAB 2017. Mexico 2017. [cited 2019 Nov 10]. Available from: https://www.gob.mx/cms/uploads/attachment/file/459279/SIVEPAB_2017.pdf
- Shahroom NS, Mani G, Ramakrishnan M. Interventions in management of dental fluorosis, an endemic disease: A systematic review. *J Family Med Prim Care*. 2019 Oct;8(10):3108-13.
- Sidhu N, Wang Y, Barrett E, Casas M. Prevalence and presentation patterns of enamel hypomineralisation (MIH and HSPM) among paediatric hospital dental patients in Toronto, Canada: a cross-sectional study. *Eur Arch Paediatr Dent*. 2020;21:263-270. <https://doi.org/10.1007/s40368-019-00477-x>.
- Silva MJ, Kilpatrick NM, Craig JM, Manton DJ, Leong P, Burgner D, et al. Etiology of Hypomineralized Second Primary Molars: A Prospective Twin Study. *J Dent Res*. 2019 Jan; 98(1):77-83.
- Silva Júnior I, Aguiar N, Barros W, da Silva L, Arantes D, Nascimento L. Prevalence and severity of molar incisor hypomineralization in students of Belém, Brazil. *Pesq Bras Odontoped Clin Integr*. 2015;15(1):377-85.
- Smith BG, Bartlett DW, Robb ND. The prevalence, etiology and management of tooth wear in the United Kingdom. *J Prosthet Dent*. 1997 Oct;78(4):367-72.
- Smith BG, Knight JK. An index for measuring the wear of teeth. *Br Dent J*. 1984 Jun;156(12): 435-8.
- Soto-Rojas AE, Ureña-Cirett JL, Martínez-Mier EA. A review of the prevalence of dental fluorosis in Mexico. *Rev Panam Salud Publica*. 2004 Jan;15(1):9-18.
- Souza JF, Costa-Silva CM, Jeremias F, Santos-Pinto L, Zuanon AC, Cordeiro RC. Molar incisor hypomineralisation: possible aetiological factors in children from urban and rural areas. *Eur Arch Paediatr Dent*. 2012 Aug; 13(4):164-70.

- Souza JF, Jeremias F, Costa-Silva CM, Santos-Pinto L, Zuanon AC, Cordeiro RC. Aetiology of molar-incisor hypomineralisation (MIH) in Brazilian children. *Eur Arch Paediatr Dent*. 2013 Jun;14(4):233–8.
- Soviero V, Haubek D, Trindade C, Da Matta T, Poulsen S. Prevalence and distribution of demarcated opacities and their sequelae in permanent 1st molars and incisors in 7 to 13-year-old Brazilian children. *Acta Odontol Scand*. 2009;67(3):170–5.
- Tagelsir A, Soto-Rojas J, Dean G, Eckert E, Martinez-Mier A. Prevalence of Molar Hypomineralization in Indiana, USA. In: 66th ORCA Congress. Cartagena, Colombia, 3–6 July, 2019. *Caries Res*. 2019;53:357–410.
- Tellez M, Santamaria RM, Gomez J, Martignon S. Dental fluorosis, dental caries, and quality of life factors among schoolchildren in a Colombian fluorotic area. *Community Dent Health*. 2012 Mar;29(1):95–9.
- The D3 Group. [Internet]. The D3 Group for developmental dental defects. Official website 2020. [cited 2020 March 23]. Available from: <https://www.thed3group.org/>
- Thylstrup A, Fejerskov O. Clinical appearance of dental fluorosis in permanent teeth in relation to histologic changes. *Community Dent Oral Epidemiol*. 1978 Nov;6(6):315–28.
- Torres Vargas L, Torres Vargas N, Vargas Cárdenas G. Erosiones dentales en pacientes con diagnóstico de enfermedad por refl ujo gastroesofágico en el Hospital Nacional Arzobispo Loayza. *Rev Gastroenterol Peru*. 2012 Oct-Dec;32(4):343–50.
- Tourino LF, Corrêa-Faria P, Ferreira RC, Bendo CB, Zarzar PM, Vale MP. Association between molar incisor hypomineralization in schoolchildren and both prenatal and postnatal factors: a population-based study. *PLoS One*. 2016 Jun;11(6):e0156332.
- Tovar S, Misnaza S. [Internet]. Documento Técnico - Perspectiva del Uso del Fluor vs Caries y Fluorosis Dental en Colombia. Ministerio de Salud y Protección Social 2016. [cited 2019 October 2020]. Available from: <https://www.minsalud.gov.co/sites/rid/Lists/Biblioteca-Digital/RIDE/VS/PP/ENT/perspectiva-uso-fluor.pdf>
- Van't Spijker A, Rodriguez JM, Kreulen CM, Bronkhorst EM, Bartlett DW, Creugers NH. Prevalence of tooth wear in adults. *Int J Prosthodont*. 2009 Jan-Feb;22(1):35–42.
- Vargas-Ferreira F, Peres MA, Dumith SC, Thomson WM, Demarco FF. Association of preperi- and postnatal factors with developmental defects of enamel in schoolchildren. *J Clin Pediatr Dent*. 2018;42(2):125–34.
- Vargas-Ferreira F, Praetzel JR, Ardenghi TM. Prevalence of tooth erosion and associated factors in 11-14-year-old Brazilian schoolchildren. *J Public Health Dent*. 2011;71(1):6–12.
- Vásquez-Muñoz E, Aguilar-Gálvez D. Prevalencia y severidad del síndrome de hipomineralización incisivo molar en niños de 6 a 13 años de edad que asistieron a la institución educativa Lee de Forest, Lima 2012. *Rev Cient Odontol*. 2014;2:75–81.
- Vieira AP, Hancock R, Limeback H, Maia R, Grynpsas MD. Is fluoride concentration in dentin and enamel a good indicator of dental fluorosis? *J Dent Res*. 2004 Jan;83(1):76–80.
- Vieira AR, Manton DJ. On the Variable Clinical Presentation of Molar-Incisor Hypomineralization. *Caries Res*. 2019;53(4):482–8.
- Villanueva-Gutiérrez T, Irigoyen-Camacho ME, Castaño-Seiquer A, Zepeda-Zepeda MA, Sanchez-Pérez L, Frechero NM. Prevalence and Severity of Molar-Incisor Hypomineralization, Maternal Education, and Dental Caries: A Cross-Sectional Study of Mexican Schoolchildren with Low Socioeconomic Status. *J Int Soc Prev Community Dent*. 2019 Sep;9(5):513–21.
- Weerheijm KL, Jälevik B, Alaluusua S. Molar-incisor hypomineralisation. *Caries Res*. 2001 Sep-Oct;35(5):390–1.
- Wiener RC, Shen C, Findley P, Tan X, Sambamoorthi U. Dental fluorosis over time: A comparison of National Health and Nutrition Examination Survey data from 2001–2002 and 2011–2012. *J Dent Hyg*. 2018 Feb;92(1):23–9.
- Wright JT, Carrion IA, Morris C. The molecular basis of hereditary enamel defects in humans. *J Dent Res*. 2015 Jan;94(1):52–61.
- Zhao D, Dong B, Yu D, Ren Q, Sun Y. The prevalence of molar incisor hypomineralization: evidence from 70 studies. *Int J Paediatr Dent*. 2018 Mar;28(2):170–9.