

University of Groningen

Novel Approaches for Developing Small Molecules to Target Histone Deacetylases

Cao, Fangyuan

DOI:
[10.33612/diss.157448844](https://doi.org/10.33612/diss.157448844)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Cao, F. (2021). *Novel Approaches for Developing Small Molecules to Target Histone Deacetylases*. University of Groningen. <https://doi.org/10.33612/diss.157448844>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

1. Histone deacetylases (HDACs) are better called lysine deacetylases (KDAC), to describe their biological function rather than their protein target, since they also deacetylate non-histone proteins. 2. HDAC isoenzymes may not only be involved in regulation of inflammatory gene transcription through their lysine deacetylase catalytic activity but also through non-catalytic interactions. 3. Proteolysis targeting chimera (PROTAC) molecules are different from the traditional occupancy-driven inhibitors. PROTACs act in an event-driven manner and degrade the targeted protein from the cells. 4. The side effects of Cereblon-recruiting PROTACs should be assessed carefully, because the E3-ligase binding components, alone or incorporated into PROTACs, can induce downregulation of NF- κ B signaling pathway. 5. For multifactorial diseases such as cancer and inflammation, approaches to manipulate multiple targets might be more beneficial than approaches directed at a single target. Sometimes, one plus one equals more than two. 6. All men by nature desire to know. —Aristotle 7. When you are lost during a project, think about its beginning and recapitulate your original aims. Stay true to your original intentions.