

University of Groningen

Detection of autism in childhood

van 't Hof, Maarten

DOI:
[10.33612/diss.159007941](https://doi.org/10.33612/diss.159007941)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
van 't Hof, M. (2021). *Detection of autism in childhood*. [Thesis fully internal (DIV), University of Groningen]. Parnassia Groep. <https://doi.org/10.33612/diss.159007941>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 7

General discussion

The objective of this thesis was to identify factors to enhance the detection of autism spectrum disorder (ASD) in childhood. Three aspects of detection of ASD were addressed:

- Age at ASD diagnosis: *what is the current global age at ASD diagnosis?*
- Early detection of ASD by preventive care physicians: *can we improve early detection by healthcare professionals?*
- Eating problems and ASD: *can the assessment of eating behavior potentially be useful in the (early) detection of ASD?*


The strengths and shortcomings of the individual studies have been discussed in chapters 2-6. This final chapter offers a broader overview of the early detection of ASD and suggestions for future research.

AGE AT ASD DIAGNOSIS

In chapter 2 we discussed the results of a meta-analysis on data from 2012–2019 that indicated a mean age at ASD diagnosis of 60.5 months (range 30.9–234.6 months).¹ Map presentations of the meta-analysis results are provided in figures 7.1 and 7.2. This update was needed as the previous literature review used studies up to 2012.² However, the ultimate objective is to convert these results into healthcare measures to improve the early detection of ASD.

For people with ASD, early detection and a younger age at diagnosis can lead to earlier treatment,³ which has been shown to improve later language and cognitive abilities, and help improve the core symptoms in children and adults with ASD.^{4,5} Although an ASD diagnosis can be established as early as age 18 months,⁶ the current global mean age at diagnosis is 43 months for children up to 10 years old;¹ this makes a huge difference, especially for young children – at age 18 months they are in the toddler development phase, while at age 43 months they are in the late early-childhood development phase. A solid and comprehensive evaluation of the age at diagnosis can contribute to reducing the local or national age at ASD diagnosis and thereby increasing the quality of healthcare offered to children with potential ASD. Such an evaluation can: 1) indicate factors affecting age at ASD diagnosis, and 2) refine and evaluate the efficacy of a national/local healthcare system in detecting and diagnosing ASD (and evaluate the effects of policy changes).

Our literature review and meta-analysis identified several limitations that make it difficult to compare the different studies and guide policy-making effectively. The largest limitations in the current literature are: 1) study heterogeneity (large variation in study characteristics, e.g. type of study sample, sample size, age of study sample, location of study), 2) an insufficient evaluation of factors affecting age at ASD diagnosis (e.g. ASD type or severity), and 3) incomplete reports on study results (e.g. missing SD, gender ratio).



Global mean age (95%CI) = 60.5 (50.1;70.8) months, range = 30.9–234.6 months

Figure 7.1 Global mean age at ASD diagnosis.




Figure 7.2 Global mean age at ASD diagnosis for children up to 10 years of age.

Factors affecting age at ASD diagnosis: the need for an integral perspective

One challenge in putting the age at ASD diagnosis literature into practice and policy change is in defining the concept of age at diagnosis. Research has indicated that a multitude of clinical, sociodemographic, parental, healthcare, educational, geographical, cohort and period factors affect the reported age at ASD diagnosis.^{1,2} These factors must therefore be included in evaluating age at diagnosis.

The effect of some factors on age at diagnosis appears to be clear. For example, including a large sample with Asperger's Disorder (AD) increases the age at diagnosis, as children with AD were diagnosed 8 to 80 months later than children with ASD in general.^{34,41-46} Including children with more severe ASD symptoms lowers the reported age at diagnosis¹⁰ by 1.9–4.1 years,⁴⁷ while using a younger study sample lowers the reported age at diagnosis.^{10,17,32,48} However, numerous factors have conflicting effects on the age at ASD diagnosis (e.g. co-morbid disorders, gender, parental age), and national or regional differences in age at diagnosis are also found,^{23,41,46} raising questions about the accessibility to healthcare in different countries.

Another challenge is that many influencing factors are intertwined. For example, conflicting results are found for the effect of ethnicity on the age at ASD diagnosis.^{9,18,46,49} However, research indicates that cultural and contextual factors affect the identification and diagnosis of ASD and whether parents seek help⁵⁰: i.e. ethnic minority families with a child with (possible) ASD experience more difficulties in gaining access to healthcare and an ASD diagnosis and they are less likely to contact a healthcare provider with their concerns.⁵¹ A major challenge is that ethnicity is also related to other characteristics like parental education, poverty status and residential area,⁵¹ which all affect access to healthcare services and increase the age at ASD diagnosis. Ethnicity is therefore not an isolated construct, but a factor intertwined with several other factors.

A more integral perspective towards the age at diagnosis and influencing factors is needed to standardize future research and enable inter-study comparisons. Figure 7.3 displays a proposal for a multi-level model providing an integral perspective on age at ASD diagnosis.

Based on results from this thesis, we recommend that future research on age at ASD diagnosis should: 1) include more general population samples instead of localized and/or specific population studies (e.g. tertiary mental healthcare clinics), 2) include more influencing factors in their study design, 3) report more completely the study results, so that data can be used in meta-analyses, and 4) use meta-analyses to adjust for a wide range of influencing factors.

A more comprehensive meta-analysis that adjusts for a wide variety of personal, socio-environmental, and national or local influencing factors would yield a more standardized current age at ASD diagnosis. This could then be used to compare results and identify weak points in local or national healthcare systems and improve the early detection of ASD.



Figure 7.3 Multi-level model for gaining an integral perspective on the age at ASD diagnosis

EARLY DETECTION OF ASD BY PREVENTIVE CARE PHYSICIANS

It is necessary to establish the current global mean age at ASD diagnosis to evaluate the barriers to and possible facilitators of an earlier detection of ASD. The current burden on healthcare makes it important to create an efficient system and develop methods to identify children at risk for ASD.⁵² Such a system should include developmental surveillance and screening to identify conditions that affect children's development and achievements.⁵³ As preventive care professionals hold a key position in child healthcare systems, one way of facilitating earlier detection would be to optimize the knowledge and skills of this group of professionals. In chapters 3 and 4 we have evaluated the early detection of ASD by preventive care physicians working in Dutch Youth and Family Healthcare Centers (YFC).

Educational program to enhance early detection of ASD

Chapter 3 indicated that Dutch preventive care physicians have low levels of specific ASD knowledge. These results are in line with previous studies indicating misconceptions about ASD held by Iranian healthcare workers and paediatricians⁵⁴ and low levels of ASD training in well-child care professionals in Iceland.⁵⁵ Our results in chapter 3 show that we should be aware that some healthcare

professionals who screen for ASD in the general population may not have sufficient knowledge about ASD. In agreement with our findings, the literature emphasizes the need for global ASD-specific training in African healthcare workers,⁵⁶ Iranian healthcare workers and paediatricians,⁵⁴ and Canadian physicians.⁵⁷ In chapter 4 we describe how a live online educational program can be used to increase the level of ASD knowledge and self-confidence in detecting ASD among preventive care physicians, highlighting the importance of such educational programs.

Live online educational program

Face-to-face educational programs are traditionally used to enhance the early detection of ASD by well-child care professionals.⁵⁵ However, online inter-professional learning offers multiple benefits in enhancing the educational experience: it is time-efficient, overcomes geographic limitations, and offers greater flexibility than face-to-face programs.⁵⁸ Online programs can be especially beneficial in low-resource countries, in which the high cost of training is one of the largest barriers in ASD diagnosis and healthcare,⁵⁹ or in a context where face-to-face training is not feasible, for example, in conflict areas or in the current COVID-19 pandemic.

Training materials for ASD detection are mostly standardized but do not include advice, practical cases, or supervision that can offer advice and tips based on specific local issues (e.g. cultural differences or available materials). Yet the method used for the educational program described in this thesis made it possible to hold live video-based sessions between tutor and students in an online school platform. This online live contact enabled not only knowledge transfer, but also discussion of case studies and responses to students' questions. The relevance of this type of interaction is supported by the results of a study on ASD distance learning for parents and professionals that showed the efficacy of a self-directed online educational program.⁶⁰ However, Wainer and Ingersoll⁶⁰ also mentioned that the added value of a more supportive and interactive educational program, including feedback and coaching, may be beneficial for some trainees.

To increase ASD knowledge and reduce stigma in low- and middle-income countries, a basic, easy-to-implement, live online ASD educational program needs to be developed for healthcare professionals who screen children. It should be implemented worldwide and made freely available for countries and organizations to use, just like the *Caregiver Skill Training* developed by WHO and *Autism Speaks* for carers of children with developmental delays and disorders.^{61,62} The program developed for this thesis could be the basis for a generic global training program that could be made accessible for everyone to use. For a good uptake of the training program in a new setting, it is important to collaborate and engage with local stakeholders and experts to ensure buy-in and that the program's contents and delivery strategy fits into the local culture and context.^{63,64} Implementing such a program might also reduce the higher level of stigma towards ASD and other mental illnesses that is

seen in non-Western countries.⁶⁵ The level of stigma towards autism could be evaluated internationally with the Autism Stigma and Knowledge Questionnaire (ASK-Q), which was developed by Harrison et al.^{66,67} during the writing of this thesis.

Other methods to enhance global ASD training should also be further explored, such as the remote machine-learning models for detecting developmental delays and autism that showed promising results in US⁶⁸ and Bangali⁶⁹ settings.

Preventive healthcare systems

Besides evaluating individual factors that could decrease the age at ASD diagnosis, health system factors should also be considered. Preventive healthcare (e.g. well-baby/child clinics) plays a vital role in the national and local healthcare system and in detecting ASD. There is consensus that early identification of children with ASD and appropriate intervention are a public health priority and that universal screening is an essential tool for the early detection of ASD.⁷⁰ However, the accessibility and efficiency of well-baby clinics depend on a variety of factors, such as professionals' level of ASD knowledge, availability of ASD screening questionnaires, national/local presence of well-baby clinics, and financial consequences.

Differences in the various healthcare systems can also explain possible barriers to earlier detection or facilitators. Well-baby clinics are the core of public health services for infants and preschool children; they are present in several European countries (Sweden, Norway, Finland, Belgium, Iceland, United Kingdom, Italy and the Netherlands),^{71,72} and in the United States,⁷³ Australia,⁷² Taiwan,⁷⁴ Turkey,⁷⁵ Nigeria,⁷⁶ Ghana,⁷⁷ and South Africa.⁷⁸ However, there are major differences in the purpose and activities of well-child/well-baby visits amongst countries.⁷⁹ Where some clinics offer universal surveillance of the complete child population, others provide well-child checks that are driven by parental request. The differences between clinic surveillance and well-child checks is summarized by The European Observatory on Health Systems and Policies⁸⁰ and described in Table 7.1.

There is a limited amount of information regarding the specific details (e.g. presence, procedures) of national well-child clinics and their role in the national system of infant and child healthcare.

Table 7.1 Differences between surveillance and well-child check⁸⁰

Surveillance	Well-child checks
Clinician driven	Parent or caregiver driven
Occurs at regular or routine intervals	At the parent or caregiver's initiative
Expensive in health care provider time	Less costly
Early detection of disease or risk factors	Promoting a healthy lifestyle to prevent disease or risk factors
Universal for all children	May exclude the children most in need
May pick up too many false positives	May pick up fewer deviations from normal (false negatives)
Provides support for parents (including those with normal children)	Puts responsibility with parents and may widen inequalities

Table 7.2 provides an overview of well-child clinics and screening capacities in several counties, based on literature/internet searches and personal inquiry among international colleagues. The results are similar to a 2010 study that indicated a wide range in the number of preventive child health examinations between European countries.⁷⁹

Table 7.2 Screening capacities of well-child clinics in selected countries around the world

	Well-child clinics present	Surveillance or well-child check	Screening for developmental delays	Additional costs	National coverage	Number of visits
The Netherlands ⁵⁵	Yes	Surveillance	Yes	No	Yes	22 (first 18 years) ^a
Iceland ⁵⁵	Yes	Surveillance	Yes	No	Yes	11 (first 4 years)
Belgium ⁸¹	Yes	Surveillance	Yes	No	Yes	10 (first 2.5 years)
Germany ⁸²	No ^b	Surveillance	Yes	No ^c	Yes	13 (first 18 years)
Norway ⁸³	Yes	Surveillance	Yes	No	Yes	14 (first 5 years) ^d

^a 13 individual-, 6 collective-, and 3 optional individual consultations (programs vary slightly between centers)

^b provided by pediatrician

^c covered by German health insurance

^d After the 5th year, health surveillance is organized by the schooling system.

Although there are reports of the early detection of ASD in well-baby clinics⁸⁴ or by related professionals⁸⁵ (e.g. pediatricians), the number of studies is limited and their results are difficult to compare because the healthcare systems differ and the precise role of the professionals is uncertain. In general, although an international report stated that most countries (91%) provided some child preventive care, this was only seen in 64% of low-income countries.⁸⁶ A 2019 report on the International Child Neurology Association Meeting on ASD in Africa indicated a huge variation in available services amongst 14 African countries, with only small numbers of specialists assessing and managing ASD compared to the size of population served.⁵⁶

Evaluating local and national healthcare systems: development of an international rating scale

Certain healthcare system factors are known to improve the early detection of ASD. For example, the availability of suitable ASD screening questionnaires has a significant effect on the early detection of ASD. However, although a review in 2019 identified 37 different ASD screening tools, many of these are not freely available, nor available in multiple languages.⁸⁷ But another 2019 study identified ten ASD screening tools that show promise for effective use in low- and middle-income countries; the ten tools are brief, low-cost and can be administered by paraprofessionals or community health workers.⁸⁸ The literature thus highlights the importance of identifying such universal tools for more general use in national healthcare systems to enhance the early detection of ASD.

In order to expand the facilities to detect and diagnose ASD at an early age, the accessibility and efficiency of many (child) healthcare systems needs to be improved. This means it is first necessary to assess the current situation to see what is needed in a specific context. A global rating tool to perform such an assessment would be very helpful and we feel future research should focus on its development. The tool should cover the factors that affect the quality of an early detection system for ASD but that are often not evaluated or reported in the literature (see below). A study on vaccination coverage has already demonstrated the relevance of rating healthcare systems; it categorized European and Australian healthcare systems based on freedom of primary care provider (hierarchical vs non-hierarchical organization) and whether preventive care services were provided by a separate organization (e.g. well-baby clinics).⁷²

Box 7.1 Factors in a healthcare system that have a positive effect on the (early) detection and diagnosis of ASD^{2,89,90}

- (free) well-baby/child healthcare centers with multiple screening moments early in life
- national guidelines on the early detection of ASD
- training for professionals on the early signs of ASD
- availability of validated (translated) screening tools for ASD
- availability of validated (translated) diagnostic tools for ASD
- availability of an early intervention service/referral options for ASD cases
- working in multidisciplinary teams, adopting appropriate working procedures

The above items (Box 7.1) can be used to develop a rating system, which would make it easier to compare regional and national differences in the early detection and age at ASD diagnosis, and the rating could then serve to highlight points needing improvement.

Box 7.2 How could Bram have benefited from these insights?

The general introduction of this thesis described the case histories of Bram and Mara. What could the insights in this thesis offer them?

Bram

Bram's signs of ASD were recognized by when he was 6 years old and adequate steps were taken to establish a diagnosis. However, it is likely that more subtle, early signs were present but missed during previous visits to the well-child clinic. An educational program on the early signs of ASD could probably have given the preventive care professional more knowledge, potentially leading to earlier detection, improving treatment outcome, and sparing child and parental distress.

EATING PROBLEMS AND ASD

Identifying the early signs of ASD can enhance the timely detection of ASD in a child. This thesis (chapters 5 and 6) has explored eating problems as an early ASD sign. Many children (and adults) with ASD struggle with eating-related problems, but the general public and many healthcare professionals are unaware of the association between ASD and eating problems. This illustrates a big gap in useful knowledge that would aid the detection of ASD in infancy and middle childhood.

Infant eating behavior as an early sign of ASD

The results reported in this thesis (chapter 5) indicate an association between infant eating behavior (formula feeding and eating problems) and later autistic traits. Our results are in line with previous studies suggesting that children with ASD are less likely to have been breastfed than their non-ASD peers.⁹¹ We hypothesize that a child's ASD characteristics complicate breastfeeding, for example, sensory problems⁹²⁻⁹⁴ (e.g. elevated oral stimuli due to flavors in breastmilk⁹⁵), skin-to-skin contact,⁹⁶ or dysregulated breastfeeding patterns⁹⁷ could negatively affect the duration of breastfeeding. It is also possible that maternal factors (general maternal psychopathology,^{98,99} elevated depressive symptomology¹⁰⁰, socio-affective impairment¹⁰¹ related to the broader autism phenotype,^{102,103} and under- or over-responsiveness to sensory stimuli¹⁰⁴⁻¹⁰⁶) could negatively affect breastfeeding. Our work also indicated an association between infant eating problems (drinking only small quantities and being hungry/not satisfied) and later autistic traits. The literature shows multiple associations between infant feeding practices and later health issues, for example, an increased risk of obesity.¹⁰⁷ An evaluation of childhood eating trajectories indicated associations between overeating and later binge eating/binge eating disorder, and between persistent undereating (in girls only) and fussy eating and later anorexia nervosa.¹⁰⁸ In addition, previous research in the same general population sample that we studied found that persistent picky eating during childhood (1.5-6 years) predicted more autistic traits at 7 years old.¹⁰⁹ Studies often focus on how caregivers affect their infants eating behavior, but it has also been suggested that infant eating behavior can be seen as a dyadic model in which the behavior is modified by the relational dynamics between an infant's characteristics (e.g. temperament) and its parents (e.g. food type choice, frequency, duration, amount, method, response to feeding cues).¹¹⁰ Although further studies are needed, our results indicate that evaluation of infant eating behavior could improve the early detection of ASD.

Sex-specific association between eating behavior in middle childhood

The results reported in chapter 6 show an association between autistic traits and eating behavior

problems (picky eating and food responsiveness) in middle childhood. Our results are in line with previous studies indicating an association between eating behavior and autistic traits.¹⁰⁹ However, we were also able to show that autistic traits were associated with girls' emotionally based eating problems (emotional over- and undereating).

The literature emphasizes the need for girl-specific ASD screening to prevent under- or misdiagnosis in females.¹¹¹ Although it is still being debated, the girl-specific ASD phenotype likely includes social imitation or the camouflaging of social difficulties with less evident impairments.¹¹² A recent report indicated that girls and women with ASD are not only significantly underdiagnosed but they also often struggle with internalizing problems such as anxiety, depression, and eating disorders, and in addition, they are regularly mislabeled and criticized in a social context.¹¹³

A recent study by Wallace et al.¹¹⁴ partly confirms our results as they found that children (4-17 years) with ASD exhibited more emotional over- and undereating than their non-ASD peers. In addition, girls were more inclined towards emotional overeating than boys with ASD, resulting in a higher consumption of sweet foods and fewer vegetables. This study found no differences in the emotional eating behavior of non-ASD children,¹¹⁴ although it included children with only an ASD diagnosis so that girls with high autistic traits and another diagnosis – e.g. anorexia nervosa, often showing high levels of emotional undereating – were excluded, which would explain the lack of gender difference. Although much about the underlying mechanisms is unknown, this thesis indicates that emotional eating problems may complement the girl-specific ASD phenotype.

Eating behavior as an (early) sign of ASD: from research to clinical practice and future research

This thesis has made several recommendations on improving the (early) detection of ASD in clinical practice and the aims of future research into associations between ASD and eating behavior.

Future research: adding eating behavior to ASD screening tools

The reported association between infant eating behavior and later autistic traits indicates that the inclusion of eating behavior in ASD screening tools could enhance the early detection of ASD. Widely used screening instruments like the Modified Checklist for Autism in Toddlers (M-CHAT(R/F)),^{115,116} the Social Communication Questionnaire (SCQ)¹¹⁷ and the Early Screening of Autistic Traits Questionnaire (ESAT/CoSoS in Dutch)¹¹⁸ often include items specific to ASD-related deficiencies, such as having an unusual response to sensory stimuli,^{115,117,118} but none of them evaluates eating behavior specifically. Yet problems with sensory stimuli may well appear while eating; if healthcare professionals are more aware of the relationship between eating behavior and ASD they will notice this when problems are being reported by parents. Indeed, the Copenhagen Infant Mental Health

Screening (CIMHS), used at age 9-10 months, includes eating behavior and shows potential for infant mental health screening based on community health nurses.¹¹⁹

Future research should evaluate if the sensitivity and/or specificity of ASD screening questionnaires increases when eating behavior questions are included. We recommend a study be performed in collaboration with well-child clinics (e.g. Youth and Family Centre in the Netherlands), in which children who are suspected of ASD are screened by the ESAT with additional questions on eating behavior. These children should be followed through their diagnostic process to confirm ESAT screening results and evaluate whether the results improve when eating behavior is included.

Future research: association of ASD with eating behavior

We recommend that the association between infant eating behavior and later ASD is further evaluated. More specific evaluation of infant eating behaviors with the Baby Eating Behaviour Questionnaire (BEBQ)¹²⁰ and the Child Eating Behaviour Questionnaire (CEBQ)¹²¹ in a general population cohort would provide insight into which type(s) of infant eating behavior is associated with ASD. It would be beneficial to evaluate eating behavior at multiple moments throughout infancy and early childhood (e.g. 2 months, 6 months, 1 year, 2 years) to see if specific eating behavior trajectories are associated with later diagnosis of ASD and if any gender differences are present.

Future research: gender differences and overlap with other disorders

Firstly, although our work found gender differences were present in the association between eating behavior and ASD in early childhood,¹²² we did not find any in infancy.¹²³ This may be due to the fact that gender differences were found in emotional eating – these behaviors are shaped by shared environmental factors rather than genetic risk,¹²⁴ and may develop only later during childhood. Nonetheless, previous research has demonstrated an association between the parental use of food to soothe (emotional feeding strategies) and the later development of emotional eating problems,¹²⁵ indicating that eating-related behavior is already being shaped as early as infancy and suggesting it can be observed during infancy. However, there were no gender differences found in the effects of parental use of food to soothe during infancy.¹²⁶ More precise and repeated evaluations of eating behaviors in infancy may reveal any gender differences.

Secondly, we must continue to follow the Generation R population to evaluate how eating behavior and ASD traits develop into pre-adolescence and adolescence, the period in which most eating disorders start to manifest.¹²⁷ Studies should focus especially on the development of eating behavior (especially emotional eating) in girls with ASD and the possible overlap with anorexia nervosa,^{128–134} to enhance early identification and referral to suitable treatment.

Finally, future research should evaluate how the inclusion of (emotional) eating behavior can enhance the diagnosis of ASD in girls. There is ongoing debate and research on the girl-specific ASD phenotype but there is a clinical relevance to enhancing the detection/diagnosis of ASD in girls to ensure they have appropriate support and treatment.

Box 7.3 How could Mara have benefited from these insights?

The general introduction of this thesis described the case histories of Bram and Mara. How could Mara have benefited from the insights provided in this thesis?

Mara

Although Mara had some early signs of ASD at age 4 years, e.g. playing alone and butterflying in the nursery school yard, her eating problems were the main cause of concern at that time and not linked to her ASD. If the preventive care professional had had more knowledge of eating problems as an early sign of ASD and, in Mara's case, of emotional eating problems (not eating after getting upset), this would have raised the possibility of her having ASD and led to further screening and an earlier ASD diagnosis.

CONCLUSIONS

The objective of this thesis was to identify factors which can enhance the detection of autism spectrum disorder (ASD) in childhood. Three aspects of the detection of ASD were addressed:

- **Age at ASD diagnosis: *what is the current global age at ASD diagnosis?*** (Chapter 2)
We found that the current global mean age at ASD diagnosis is 60.48 months (range 30.9–234 months). Efforts are needed to evaluate the age at ASD diagnosis by a meta-analysis that includes a wide variety of influencing factors in order to advance early detection of ASD in healthcare systems.
- **Early detection of ASD by preventive care physicians: *can we improve early detection by healthcare professionals?*** (Chapters 3 & 4)
Our work indicates that a live online ASD educational program for preventive care professionals contributes to the early detection of ASD. Efforts are needed to evaluate global well-baby facilities and to implement ASD educational programs to enhance the early detection by healthcare professionals screening for ASD in the child population.
- **Eating problems and ASD: *can the assessment of eating behavior potentially be useful in the (early) detection of ASD?*** (Chapters 5 & 6)
Our work indicates an association between eating behavior and ASD in early and middle childhood; it could therefore be beneficial to include questions on eating behavior in ASD

screening tools. Future research should further specify the association between eating behavior and ASD, and determine its usefulness as a screening item.

This thesis describes improvements that can be made to improve global ASD preventive healthcare. We want to encourage researchers and clinicians to further explore the possibilities of detecting ASD at a younger age in order to enhance the quality of care and life for children with ASD.

REFERENCES

1. Van 't Hof M, Tisseur C, van Berckeleer-Onnes I, et al. Age at autism spectrum disorder diagnosis: A systematic review and meta-analysis from 2012 to 2019. *Autism*. In press. doi:10.1177/1362361320971107
2. Daniels AM, Mandell DS. Explaining differences in age at autism spectrum disorder diagnosis: A critical review. *Autism*. 2014;18(5):583-597. doi:10.1177/1362361313480277
3. Rogers SJ, Vismara L, Wagner AL, McCormick C, Young G, Ozonoff S. Autism treatment in the first year of life: A pilot study of infant start, a parent-implemented intervention for symptomatic infants. *J Autism Dev Disord*. 2014;44(12):2981-2995. doi:10.1007/s10803-014-2202-y
4. Dawson G, Burner K. Behavioral interventions in children and adolescents with autism spectrum disorder. *Curr Opin Pediatr*. 2011;23(6):616-620. doi:10.1097/MOP.0b013e32834cf082
5. Clark MLE, Vinen Z, Barbaro J, Dissanayake C. School age outcomes of children diagnosed early and later with autism spectrum disorder. *J Autism Dev Disord*. 2018;48(1):92-102. doi:10.1007/s10803-017-3279-x
6. Hyman SL, Levy SE, Myers SM. Identification, evaluation, and management of children with autism spectrum disorder. *Pediatrics*. 2020;145(1). doi:10.1542/peds.2019-3447
7. Coo H, Ouellette-Kuntz H, Lam M, et al. Correlates of age at diagnosis of autism spectrum disorders in six Canadian regions. *Chronic Dis Inj Can*. 2012;32(2):90-100.
8. Valicenti-McDermott M, Hottinger K, Seijo R, Shulman L. Age at Diagnosis of Autism Spectrum Disorders. *J Pediatr*. 2012;161(3):554-556. doi:10.1016/j.jpeds.2012.05.012
9. Magaña S, Lopez K, Aguinaga A, Morton H. Access to diagnosis and treatment services among latino children with autism spectrum disorders. *Intellect Dev Disabil*. 2013;51(3):141-153. doi:10.1352/1934-9556-51.3.141
10. Mazurek MO, Handen BL, Wodka EL, Nowinski L, Butter E, Engelhardt CR. Age at first autism spectrum disorder diagnosis: The role of birth cohort, demographic factors, and clinical features. *J Dev Behav Pediatr*. 2014;35(9):561-569. doi:10.1097/DBP.0000000000000097
11. Hausman-Kedem M, Kosofsky BE, Ross G, et al. Accuracy of reported community diagnosis of autism spectrum disorder. *J Psychopathol Behav Assess*. 2018;40(3):367-375. doi:10.1007/s10862-018-9642-1
12. Martinez M, Thomas KC, Williams CS, et al. Family experiences with the diagnosis of autism spectrum disorder: System barriers and facilitators of efficient diagnosis. *J Autism Dev Disord*. 2018;48(7):2368-2378. doi:10.1007/s10803-018-3493-1
13. Miodovnik A, Harstad E, Sideridis G, Huntington N. Timing of the diagnosis of attention-deficit/hyperactivity disorder and autism spectrum disorder. *Pediatrics*. 2015;136(4):e830-e837. doi:10.1542/peds.2015-1502
14. Darcy-Mahoney A, Minter B, Higgins M, Guo Y, Zauche LH, Hirst J. Maternal and neonatal birth factors affecting the age of asd diagnosis. *Newborn Infant Nurs Rev*. 2016;16(4):340-347. doi:10.1053/j.nainr.2016.09.033
15. Emerson ND, Morrell HER, Neece C. Predictors of age of diagnosis for Children with Autism Spectrum Disorder: The Role of a Consistent Source of Medical Care, Race, and Condition Severity. *J Autism Dev Disord*. 2016;46(1):127-138. doi:10.1007/s10803-015-2555-x
16. Sicherman N, Loewenstein G, Tavassoli T, Buxbaum JD. Grandma knows best: Family structure and age of diagnosis of autism spectrum disorder. *Autism*. 2017;136236131667963. doi:10.1177/1362361316679632
17. Zablotsky B, Colpe LJ, Pringle BA, Kogan MD, Rice C, Blumberg SJ. Age of parental concern, diagnosis, and service initiation among children with autism spectrum disorder. *Am J Intellect Dev Disabil*. 2017;122(1):49-61. doi:10.1352/1944-7558-122.1.49
18. Becerra-Culqui TA, Lynch FL, Owen-Smith AA, Spitzer J, Croen LA. Parental first concerns and

- timing of autism spectrum disorder diagnosis. *J Autism Dev Disord.* 2018;48(10):3367-3376. doi:10.1007/s10803-018-3598-6
19. Berg KL, Acharya K, Shiu C-S, Msall ME. Delayed diagnosis and treatment among children with autism who experience adversity. *J Autism Dev Disord.* 2017. doi:10.1007/s10803-017-3294-y
 20. Hall-Lande J, Esler AN, Hewitt A, Gunty AL. Age of initial identification of autism spectrum disorder in a diverse urban sample. *J Autism Dev Disord.* 2018. doi:10.1007/s10803-018-3763-y
 21. Ribeiro SHB, de Paula CS, Bordini D, Mari JJ, Caetano SC. Barriers to early identification of autism in Brazil. *Rev Bras Psiquiatr.* 2017;39(4):352-354. doi:10.1590/1516-4446-2016-2141
 22. Lagunju IA, Bella-Awusah TT, Omigbodun OO. Autistic disorder in Nigeria: Profile and challenges to management. *Epilepsy Behav.* 2014;39:126-129. doi:10.1016/j.yebeh.2014.08.020
 23. Bello-Mojeed MA, Omigbodun OO, Bakare MO, Adewuya AO. Pattern of impairments and late diagnosis of autism spectrum disorder among a sub-Saharan African clinical population of children in Nigeria. *Glob Ment Heal.* 2017;4:e5. doi:10.1017/gmh.2016.30
 24. Mpaka DM, Okitundu DLEA, Ndjukendi AO, et al. Prevalence and comorbidities of autism among children referred to the outpatient clinics for neurodevelopmental disorders. *Pan Afr Med J.* 2016;25:82. doi:10.11604/pamj.2016.25.82.4151
 25. Daniels AM, Como A, Hergüner S, Kostadinova K, Stosic J, Shih A. Autism in southeast europe: A survey of caregivers of children with autism spectrum disorders. *J Autism Dev Disord.* 2017;47(8):2314-2325. doi:10.1007/s10803-017-3145-x
 26. Salomone E, Charman T, McConachie H, Warreyn P. Child's verbal ability and gender are associated with age at diagnosis in a sample of young children with ASD in Europe. *Child Care Health Dev.* 2016;42(1):141-145. doi:10.1111/cch.12261
 27. Brett D, Warnell F, McConachie H, Parr JR. Factors affecting age at asd diagnosis in uk: No evidence that diagnosis age has decreased between 2004 and 2014. *J Autism Dev Disord.* 2016;46(6):1974-1984. doi:10.1007/s10803-016-2716-6
 28. Hrdlicka M, Vacova M, Oslejskova H, et al. Age at diagnosis of autism spectrum disorders: Is there an association with socioeconomic status and family self-education about autism? *Neuropsychiatr Dis Treat.* 2016;12:1639-1644. doi:10.2147/NDT.S107239
 29. Höfer J, Hoffmann F, Kamp-Becker I, et al. Pathways to a diagnosis of autism spectrum disorder in Germany: A survey of parents. *Child Adolesc Psychiatry Ment Health.* 2019;13(1):1-10. doi:10.1186/s13034-019-0276-1
 30. Ververi A, Vargiami E, Papadopoulou V, Tryfonas D, Zafeiriou DI. Clinical and laboratory data in a sample of greek children with autism spectrum disorders. *J Autism Dev Disord.* 2012;42(7):1470-1476. doi:10.1007/s10803-011-1414-7
 31. Larsen K. The early diagnosis of preschool children with autism spectrum disorder in Norway: A study of diagnostic age and its associated factors. *Scand J Child Adolesc Psychiatry Psychol.* 2015;3(2):136-145.
 32. Rutherford M, McKenzie K, Johnson T, et al. Gender ratio in a clinical population sample, age of diagnosis and duration of assessment in children and adults with autism spectrum disorder. *Autism.* 2016;20(5):628-634. doi:10.1177/1362361315617879
 33. Garrido D, Carballo G, Artis J, Garcia-Retamero R. Timing of parents' concerns related to autism spectrum disorder and its diagnosis: A mediation analysis. *Span J Psychol.* 2018;(2018):1-8. doi:10.1017/sjp.2018.64
 34. Begeer S, Mandell D, Wijnker-Holmes B, et al. Sex differences in the timing of identification among children and adults with autism spectrum disorders. *J Autism Dev Disord.* 2013;43(5):1151-1156. doi:10.1007/s10803-012-1656-z
 35. Kentrou V, de Veld DMJ, Mataw KJK, Begeer S. Delayed autism spectrum disorder recognition in children and adolescents previously diagnosed with attention-deficit/hyperactivity disorder. *Autism.* 2018. doi:10.1177/1362361318785171
 36. Mishaal RA, Ben-Itzhak E, Zachor DA. Age of autism spectrum disorder diagnosis is associated

- with child's variables and parental experience. *Res Autism Spectr Disord*. 2014;8(7):873-880. doi:10.1016/j.rasd.2014.04.001
37. Manohar H, Kandasamy P, Chandrasekaran V, Rajkumar RP. Early diagnosis and intervention for autism spectrum disorder: Need for pediatrician–child psychiatrist liaison. *Indian J Psychol Med*. 2019;41(1):87-90. doi:10.4103/IJPSYM.IJPSYM_154_18
 38. Shrestha R, Dissanayake C, Barbaro J. Age of diagnosis of autism spectrum disorder in Nepal. *J Autism Dev Disord*. 2019;49(6):2258-2267. doi:10.1007/s10803-019-03884-7
 39. Kurasawa S, Tateyama K, Iwanaga R, Ohtoshi T, Nakatani K, Yokoi K. The age at diagnosis of autism spectrum disorder in children in Japan. *Int J Pediatr*. 2018;2018:1-5. doi:10.1155/2018/5374725
 40. Wei HT, Hsu JW, Huang KL, et al. Timing of the diagnoses of attention deficit hyperactivity disorder and autism spectrum disorder in Taiwan. *J Autism Dev Disord*. 2018. doi:10.1007/s10803-018-3655-1
 41. Bent, Catherine A; Dissanayake, Cheryl; Barbaro J. Mapping the diagnosis of autism spectrum disorders in children aged under 7 years in Australia, 2010–2012. *Med J Aust*. 2015;202(6):317-320. doi:10.5694/mja14.00328
 42. Bravo Oro A, Vázquez Briseño J, Cuello García CA, Calderón Sepúlveda RF, Hernández Villalobos AM, Esmer Sánchez C. Early manifestations of autism spectrum disorders. Experience of 393 cases in a paediatric neurology. *Neurol (Barcelona, Spain)*. 2012;27(7):414-420. doi:10.1016/j.nrl.2011.09.011
 43. Hinkka-Yli-Salomäki S, Banerjee PN, Gissler M, et al. The incidence of diagnosed autism spectrum disorders in Finland. *Nord J Psychiatry*. 2014;68(7):472-480. doi:10.3109/08039488.2013.861017
 44. Baio J, Wiggins L, Christensen DL, et al. Prevalence of autism spectrum disorder among children aged 8 years - Autism and developmental disabilities monitoring network, 11 sites, United States, 2014. *MMWR Surveill Summ*. 2018;67(6):1-23. doi:10.15585/mmwr.ss6706a1
 45. U.S. Department of Health and Human Services. Prevalence of autism spectrum disorder among children aged 8 years - autism and developmental disabilities monitoring network, 11 sites, United States, 2010. *MMWR Surveill Summ*. 2014;63(2):1-21. doi:24670961
 46. Christensen DL, Baio J, Braun KVN, et al. Prevalence and characteristics of autism spectrum disorder among children aged 8 years - Autism and developmental disabilities monitoring network, 11 sites, United States, 2012. *MMWR Surveill Summ*. 2016;65(3):1-23. doi:10.15585/mmwr.ss6503a1
 47. Sheldrick RC, Maye MP, Carter AS. Age at first identification of autism spectrum disorder: An analysis of two US surveys. *J Am Acad Child Adolesc Psychiatry*. 2017;56(4):313-320. doi:10.1016/j.jaac.2017.01.012
 48. Jo H, Schieve LA, Rice CE, et al. Age at autism spectrum disorder (asd) diagnosis by race, ethnicity, and primary household language among children with special health care needs, United States, 2009-2010. *Matern Child Health J*. 2015;1687-1697. doi:10.1007/s10995-015-1683-4
 49. Emerson ND, Morrell HER, Neece C. Predictors of age of diagnosis for children with autism spectrum disorder: The role of a consistent source of medical care, race, and condition severity. *J Autism Dev Disord*. 2016;46(1):127-138. doi:10.1007/s10803-015-2555-x
 50. de Leeuw A, Happé F, Hoekstra RA. A conceptual framework for understanding the cultural and contextual factors on autism across the globe. *Autism Res*. 2020;13(7):1029-1050. doi:10.1002/aur.2276
 51. Zeleke WA, Hughes TL, Drozda N. Disparities in diagnosis and service access for minority children with ASD in the United States. *J Autism Dev Disord*. 2019;49(10):4320-4331. doi:10.1007/s10803-019-04131-9
 52. McCarty P, Frye RE. Early detection and diagnosis of autism spectrum disorder: Why is it so difficult? *Semin Pediatr Neurol*. 2020;35:100831. doi:10.1016/j.spen.2020.100831
 53. Lipkin PH, Macias MM. Promoting optimal development: Identifying infants and young

- children with developmental disorders through developmental surveillance and screening. *Pediatrics*. 2020;145(1). doi:10.1542/peds.2019-3449
54. Effatpanah M, Shariatpanahi GH, Ramaghi R, Tavakolizadeh R. A preliminary survey of autism knowledge and attitude among health care workers and pediatricians in Tehran, Iran. *Iran J child Neurol*. 2019;13(2):29-35.
 55. Jonsdottir SL, Saemundsen E, Gudmundsdottir S, Haraldsdottir GS, Palsdottir AH, Rafnsson V. Implementing an early detection program for autism in primary healthcare: Screening, education of healthcare professionals, referrals for diagnostic evaluation, and early intervention. *Res Autism Spectr Disord*. 2020;77:101616. doi:10.1016/j.rasd.2020.101616
 56. Ruparelia K, Abubakar A, Badoo E, et al. Autism spectrum disorders in Africa: Current challenges in identification, assessment, and treatment. *J Child Neurol*. 2016;31(8):1018-1026. doi:10.1177/08830738166635748
 57. Ghaderi G, Watson SL. Autism spectrum disorder knowledge, training and experience: Ontario physicians' perspectives about what helps and what does not. *J Dev Disabil*. 2019;24(2):51-60.
 58. Reeves S, Fletcher S, Mcloughlin C, Yim A, Patel KD. Interprofessional online learning for primary healthcare: Findings from a scoping review. *BMJ Open*. 2017;7(8). doi:10.1136/bmjopen-2017-016872
 59. Durkin MS, Elsabbagh M, Barbaro J, et al. Autism screening and diagnosis in low resource settings: Challenges and opportunities to enhance research and services worldwide. *Autism Res*. 2015;8(5):473-476. doi:10.1002/aur.1575
 60. Wainer AL, Ingersoll BR. Disseminating asd interventions: A pilot study of a distance learning program for parents and professionals. *J Autism Dev Disord*. 2013;43:11-24. doi:10.1007/s10803-012-1538-4
 61. WHO. The WHO Caregiver Skills Training programme. https://www.who.int/mental_health/maternal-child/PST/en/. Accessed April 21, 2020.
 62. Autism Speaks. Autism caregiver skills training: Helping families at home and abroad. <https://www.autismspeaks.org/science-news/autism-caregiver-skills-training-helping-families-home-and-abroad>. Accessed November 5, 2020.
 63. Tekola B, Girma F, Kinfe M, et al. Adapting and pre-testing the World Health Organization's Caregiver Skills Training programme for autism and other developmental disorders in a very low-resource setting: Findings from Ethiopia. *Autism*. 2020;24(1):51-63. doi:10.1177/1362361319848532
 64. Hoekstra RA, Girma F, Tekola B, Yenus Z. Nothing about us without us: the importance of local collaboration and engagement in the global study of autism. *BJPsycho Int*. 2018;15(2):40-43. doi:10.1192/bji.2017.26
 65. van 't Hof M, van Berckelaer-Onnes I, Deen M, et al. Novel insights into autism knowledge and stigmatizing attitudes toward mental illness in dutch youth and family center physicians. *Community Ment Health J*. 2020. doi:10.1007/s10597-020-00568-w
 66. Harrison AJ, Paff ML, Kaff MS. Examining the psychometric properties of the autism stigma and knowledge questionnaire (ASK-Q) in multiple contexts. *Res Autism Spectr Disord*. 2019;57:28-34. doi:10.1016/j.rasd.2018.10.002
 67. Harrison AJ, Bradshaw LP, Naqvi NC, Paff ML, Campbell JM. Development and psychometric evaluation of the autism stigma and knowledge questionnaire (ASK-Q). *J Autism Dev Disord*. 2017;47(10):3281-3295. doi:10.1007/s10803-017-3242-x
 68. Tariq Q, Daniels J, Schwartz JN, Washington P, Kalantarian H, Wall DP. Mobile detection of autism through machine learning on home video: A development and prospective validation study. *PLoS Med*. 2018;15(11):1-20. doi:10.1371/journal.pmed.1002705
 69. Tariq Q, Fleming SL, Schwartz JN, et al. Detecting developmental delay and autism through machine learning models using home videos of bangladeshi children: Development and validation study. *J Med Internet Res*. 2019;21(4):1-15. doi:10.2196/13822
 70. Pierce K, Courchesne E, Bacon E. To screen or not to screen universally for autism is not the question: Why the task force got it wrong. *J Pediatr*. 2016;176:182-194.

- doi:10.1016/j.jpeds.2016.06.004
71. Sundelin C, Magnusson M, Lagerberg D. Child health services in transition: I. Theories, methods and launching. *Acta Paediatr Int J Paediatr*. 2005. doi:10.1080/08035250410023412
 72. Arat A, Burström B, Östberg V, Hjern A. Social inequities in vaccination coverage among infants and pre-school children in Europe and Australia - A systematic review. *BMC Public Health*. 2019;19(1):1-10. doi:10.1186/s12889-019-6597-4
 73. Kumaraswami T, Rankin KM, Lunde B, Cowett A, Caskey R, Harwood B. Acceptability of postpartum contraception counseling at the well baby visit. *Matern Child Health J*. 2018;22(11):1624-1631. doi:10.1007/s10995-018-2558-2
 74. Wu SC, Lue HC, Tseng LL. A pediatric clinic-based approach to early literacy promotion - experience in a well-baby clinic in Taiwan. *J Formos Med Assoc*. 2012;111(5):258-264. doi:10.1016/j.jfma.2011.05.008
 75. Çelik SB, Şahin F, Beyazova U, Can H. Growth status of children in well-baby outpatient clinics and related factors. *Turk Pediatr Ars*. 2014;49(2):104-110. doi:10.5152/tpa.2014.1145
 76. Adam V, Abhulimhen-Iyoha B. Teething: Beliefs and behaviors of mothers attending well baby clinics in Benin City, Nigeria. *African J Med Heal Sci*. 2015;14(1):8-12. doi:10.4103/2384-5589.153377
 77. Morhe ESK, Ankobe F, Asubonteng GO, Opoku B, Turpin CA, Dalton VK. Postpartum contraceptive choices among women attending a well-baby clinic in Ghana. *Int J Gynecol Obstet*. 2017;138(2):219-224. doi:10.1002/ijgo.12216
 78. Sokhela DG, Sibiyi MN, Gwele NS. Monitoring well-baby visits in primary healthcare facilities in a middle-income country. *SAJCH South African J Child Heal*. 2018;12(2):9-12. doi:10.7196/SAJCH.2018.v12i2.1262
 79. Van Esso D, Del Torso S, Hadjipanayis A, et al. Paediatric primary care in Europe: Variation between countries. *Arch Dis Child*. 2010;95(10):791-795. doi:10.1136/adc.2009.178459
 80. European Observatory on health system and policies. *European Child Health Services and Systems Lessons: Without Borders.*; 2013. http://www.euro.who.int/__data/assets/pdf_file/0003/254928/European-Child-Health-Services-and-Systems-Lessons-without-borders.pdf.
 81. Gezin K en. Consulten. <https://www.kindengezin.be/>. Accessed October 15, 2020.
 82. Ehrich J, Grote U, Gerber-Grote A, Strassburg M. The child health care system of Germany. *J Pediatr*. 2016;177:S71-S86. doi:10.1016/j.jpeds.2016.04.045
 83. Norwegian Institute of Public Health. High coverage for the childhood immunisation programme in Norway. <https://www.fhi.no/en/news/2020/high-coverage-for-the-childhood-immunisation-programme-in-norway/>. Accessed November 5, 2020.
 84. Suma K, Adamson LB, Bakeman R, Robins DL, Abrams DN. After early autism diagnosis: Changes in intervention and parent-child interaction. *J Autism Dev Disord*. 2016;46(8):2720-2733. doi:10.1007/s10803-016-2808-3
 85. Pijl MKJ, Buitelaar JK, de Korte MWP, Rommelse NNJ, Oosterling IJ. Sustainability of an early detection program for autism spectrum disorder over the course of 8 years. *Autism*. 2018;22(8):1018-1024. doi:10.1177/1362361317717977
 86. Harper BD, Nganga W, Armstrong R, et al. Where are the paediatricians? An international survey to understand the global paediatric workforce. *BMJ Paediatr Open*. 2019;3(1):1-9. doi:10.1136/bmjpo-2018-000397
 87. Thabtah F, Peebles D. Early autism screening: A comprehensive review. *Int J Environ Res Public Health*. 2019;16(18):3502. doi:10.3390/ijerph16183502
 88. Marlow M, Servili C, Tomlinson M. A review of screening tools for the identification of autism spectrum disorders and developmental delay in infants and young children: recommendations for use in low- and middle-income countries. *Autism Res*. 2019;12(2):176-199. doi:10.1002/aur.2033
 89. NICE. *Autism Spectrum Disorder in under 19s: Recognition, Referral and Diagnosis.*; 2011. <https://www.nice.org.uk/guidance/cg128>.

90. Centers for Disease Control and Prevention (CDC). Autism spectrum disorder - recommendations and guidelines for healthcare providers. <https://www.cdc.gov/ncbddd/autism/hcp-recommendations.html>. Accessed October 12, 2020.
91. Tseng P-T, Chen Y-W, Stubbs B, et al. Maternal breastfeeding and autism spectrum disorder in children: A systematic review and meta-analysis. *Nutr Neurosci*. 2019;22(5):354-362. doi:10.1080/1028415X.2017.1388598
92. Bryson SE, Zwaigenbaum L, Brian J, et al. A prospective case series of high-risk infants who developed autism. *J Autism Dev Disord*. 2007;37(1):12-24. doi:10.1007/s10803-006-0328-2
93. Mulligan S, White BP. Sensory and motor behaviors of infant siblings of children with and without autism. *Am J Occup Ther*. 2012;66(5):556-566. doi:10.5014/ajot.2012.004077
94. Germani T, Zwaigenbaum L, Bryson S, et al. Brief report: Assessment of early sensory processing in infants at high-risk of autism spectrum disorder. *J Autism Dev Disord*. 2014;44(12):3264-3270. doi:10.1007/s10803-014-2175-x
95. Beauchamp GK, Mennella JA. Flavor perception in human infants: Development and functional significance. *Digestion*. 2011;83(SUPPL.1):1-6. doi:10.1159/000323397
96. Liu J, Leung P, Yang A. Breastfeeding and active bonding protects against children's internalizing behavior problems. *Nutrients*. 2013;6(1):76-89. doi:10.3390/nu6010076
97. Lucas J, Cutler. Autism: Dysregulated breastfeeding behaviors? *J Perinat Educ*. 2015;24(3):171-180. doi:10.1891/1058-1243.24.3.171
98. Boucher O, Julvez J, Guxens M, et al. Association between breastfeeding duration and cognitive development, autistic traits and ADHD symptoms: A multicenter study in Spain. *Pediatr Res*. 2017;81(3):434-442. doi:10.1038/pr.2016.238
99. Tavoulari E-F, Benetou V, Vlastarakos P V, et al. Factors affecting breastfeeding duration in Greece: What is important? *World J Clin Pediatr*. 2016;5(3):349. doi:10.5409/wjcp.v5.i3.349
100. Ingersoll B, Hambrick DZ. The relationship between the broader autism phenotype, child severity, and stress and depression in parents of children with autism spectrum disorders. *Res Autism Spectr Disord*. 2011;5(1):337-344. doi:10.1016/j.rasd.2010.04.017
101. Berthoz S, Lalanne C, Crane L, Hill EL. Investigating emotional impairments in adults with autism spectrum disorders and the broader autism phenotype. *Psychiatry Res*. 2013;208(3):257-264. doi:10.1016/j.psychres.2013.05.014
102. Henderson JJ, Evans SF, Straton JAY, Priest SR, Hagan R. Impact of postnatal depression on breastfeeding duration. *Birth*. 2003;30(3):175-180. doi:10.1046/j.1523-536X.2003.00242.x
103. Falsett CF, Santos IMM dos, Vasconcellos AM. Interfering factors of the breastfeeding process in children bearing various health needs: Contributions to nursing / Fatores que interferem no processo de aleitamento materno de crianças com necessidades de saúde variadas: Contribuições para a enfermagem. *Rev Pesqui Cuid é Fundam Online*. 2019;11(5):1278. doi:10.9789/2175-5361.2019.v11i5.1278-1285
104. Britton JR, Britton HL, Gronwaldt V. Breastfeeding, sensitivity, and attachment. *Pediatrics*. 2006;118(5):e1436-e1443. doi:10.1542/peds.2005-2916
105. Tharner A, Luijk MPCM, Raat H, et al. Breastfeeding and its relation to maternal sensitivity and infant attachment. *J Dev Behav Pediatr*. 2012;33(5):396-404. doi:10.1097/DBP.0b013e318257fac3
106. Lai CYY, Chung JCC, Chan CCH, Li-Tsang CWP. Sensory Processing Measure-HK Chinese version: Psychometric properties and pattern of response across environments. *Res Dev Disabil*. 2011;32(6):2636-2643. doi:10.1016/j.ridd.2011.06.010
107. Matvienko-Sikar K, Griffin C, McGrath N, et al. Developing a core outcome set for childhood obesity prevention: A systematic review. *Matern Child Nutr*. 2019;15(1). doi:10.1111/mcn.12680
108. Herle M, Stavola B De, Hübel C, et al. A longitudinal study of eating behaviours in childhood and later eating disorder behaviours and diagnoses. *Br J Psychiatry*. 2020;216(2):113-119. doi:10.1192/bjp.2019.174

109. Cardona Cano S, Hoek HW, van Hoeken D, et al. Behavioral outcomes of picky eating in childhood: A prospective study in the general population. *J Child Psychol Psychiatry*. 2016;57(11):1239-1246. doi:10.1111/jcpp.12530
110. Hodges EA, Propper CB, Estrem H, Schultz MB. Feeding during infancy: interpersonal behavior, physiology, and obesity risk. *Child Dev Perspect*. 2020;14(3):185-191. doi:10.1111/cdep.12376
111. Kirkovski M, Enticott PG, Fitzgerald PB. A review of the role of female gender in autism spectrum disorders. *J Autism Dev Disord*. 2013;43(11):2584-2603. doi:10.1007/s10803-013-1811-1
112. Allely CS. Understanding and recognising the female phenotype of autism spectrum disorder and the “camouflage” hypothesis: a systematic PRISMA review. *Adv Autism*. 2019;5(1):14-37. doi:10.1108/AIA-09-2018-0036
113. Eckerd M. Detection and Diagnosis of ASD in Females. *J Heal Serv Psychol*. 2020;46(1):37-47. doi:10.1007/s42843-020-00006-1
114. Wallace GL, Richard E, Wolff A, Nadeau M, Zucker N. Increased emotional eating behaviors in children with autism: Sex differences and links with dietary variety. *Autism*. 2020. doi:10.1177/1362361320942087
115. Robins DL, Fein D, Barton ML, Green JA. The modified checklist for autism in toddlers: An initial study investigating the early detection of autism and pervasive developmental disorders. *J Autism Dev Disord*. 2001;31(2):131-144. doi:10.1023/A:1010738829569
116. Robins DL, Casagrande K, Barton M, Chen C-MA, Dumont-Mathieu T, Fein D. Validation of the modified checklist for autism in toddlers, revised with follow-up (M-CHAT-R/F). *Pediatrics*. 2014;133(1):37-45. doi:10.1542/peds.2013-1813
117. Rutter M, Bailey A, Lord C. *The Social Communication Questionnaire*. Torrance, CA: Western Psychological Services; 2003.
118. Swinkels SHN, Dietz C, Daalen E Van, Kerkhof IHGM, Engeland H Van, Buitelaar JK. Screening for autistic spectrum in children aged 14 to 15 months. I : The development of the early screening of autistic traits questionnaire (ESAT). *J Autism Dev Disord*. 2006;36:723-732. doi:10.1007/s10803-006-0115-0
119. Ammitzbøll J, Holstein BE, Wilms L, Andersen A, Skovgaard AM. A new measure for infant mental health screening: Development and initial validation. *BMC Pediatr*. 2016;16(1):1-10. doi:10.1186/s12887-016-0744-1
120. Llewellyn CH, van Jaarsveld CHM, Johnson L, Carnell S, Wardle J. Development and factor structure of the baby eating behaviour questionnaire in the gemini birth cohort. *Appetite*. 2011;57(2):388-396. doi:10.1016/j.appet.2011.05.324
121. Wardle J, Guthrie CA, Sanderson S, Rapoport L. Development of the children’s eating behaviour questionnaire. *J Child Psychol Psychiatry Allied Discip*. 2001;42(7):963-970. doi:10.1111/1469-7610.00792
122. van ’t Hof M, Ester WA, Serdarevic F, et al. The sex-specific association between autistic traits and eating behavior in childhood: An exploratory study in the general population. *Appetite*. 2020;147. doi:10.1016/j.appet.2019.104519
123. van ’t Hof M, Ester WA, van Berckelaer-Onnes I, Hillegers MHJ, Hoek HW, Jansen PW. Do early-life eating habits predict later autistic traits? Results from a population-based study. *Appetite*. 2021;156:104976. doi:10.1016/j.appet.2020.104976
124. Herle M, Fildes A, Llewellyn CH. Emotional eating is learned not inherited in children, regardless of obesity risk. *Pediatr Obes*. 2018;13(10):628-631. doi:10.1111/ijpo.12428
125. Hardman CA, Christiansen P, Wilkinson LL. Using food to soothe: Maternal attachment anxiety is associated with child emotional eating. *Appetite*. 2016;99:91-96. doi:10.1016/j.appet.2016.01.017
126. Jansen PW, Derks IPM, Batenburg A, et al. Using food to soothe in infancy is prospectively associated with childhood BMI in a population- based cohort. *J Nutr*. 2019;149(5):788-794. doi:10.1093/jn/nxy277
127. Galmiche M, Déchelotte P, Lambert G, Tavolacci MP. Prevalence of eating disorders over the

- 2000-2018 period: A systematic literature review. *Am J Clin Nutr.* 2019;109(5):1402-1413. doi:10.1093/ajcn/nqy342
128. Harrison A, Sullivan S, Tchanturia K, Treasure J. Emotional functioning in eating disorders: attentional bias, emotion recognition and emotion regulation. *Psychol Med.* 2010;40(11):1887-1897. doi:10.1017/S0033291710000036
129. Ricca V, Castellini G, Fioravanti G, et al. Emotional eating in anorexia nervosa and bulimia nervosa. *Compr Psychiatry.* 2012;53(3):245-251. doi:10.1016/j.comppsy.2011.04.062
130. Samson AC, Hardan AY, Podell RW, Phillips JM, Gross JJ. Emotion regulation in children and adolescents with autism spectrum disorder. *Autism Res.* 2015;8(1):9-18. doi:10.1002/aur.1387
131. Huke V, Turk J, Saeidi S, Kent A, Morgan JF. Autism spectrum disorders in eating disorder populations: a systematic review. *Eur Eat Disord Rev.* 2013;21(5):345-351. doi:10.1002/erv.2244
132. Zucker NL, Losh M, Bulik CM, Labar KS, Piven J. Anorexia nervosa and autism spectrum disorders : Guided investigation of social cognitive endophenotypes. *Psychol Bull.* 2007;133(6):976-1006. doi:10.1037/0033-2909.133.6.976
133. Kalyva E. Comparison of eating attitudes between adolescent girls with and without asperger syndrome: Daughters' and mothers' reports. *J Autism Dev Disord.* 2009;39(3):480-486. doi:10.1007/s10803-008-0648-5
134. Renwick B, Musiat P, Lose A, et al. Neuro- and social-cognitive clustering highlights distinct profiles in adults with anorexia nervosa. *Int J Eat Disord.* 2015;48(1):26-34. doi:10.1002/eat.22366

