

University of Groningen

Management of heavy menstrual bleeding

van den Brink, Marian

DOI:
[10.33612/diss.160486947](https://doi.org/10.33612/diss.160486947)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
van den Brink, M. (2021). *Management of heavy menstrual bleeding: Towards a patient-centred approach*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.160486947>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Levonorgestrel-releasing intrauterine system (Mirena) versus endometrial ablation (Novasure) in women with heavy menstrual bleeding: a multicentre randomised controlled trial (*Study protocol*)

Malou C Herman, Marian J van den Brink, Peggy M Geomini, Hannah S van Meurs, Judith A Huirne, Heleen P Eising, Anne Timmermans, Johanna MA Pijnenborg, Ellen R Klinkert, Sjors F Coppus, Theodoor E Nieboer, Ruby Catshoek, Lucet F van der Voet, Hugo WF van Eijndhoven, Giuseppe CM Graziosi, Sebastiaan Veersema, Paul J van Kesteren, Josje Langenveld, Nicol AC Smeets, Huib AAM van Vliet, Jan Willem van der Steeg, Yvonne Lisman-van Leeuwen, Janny H Dekker, Ben W Mol, Marjolein Y Berger, Marlies Y Bongers

BMC Women's Health. 2013;13:32.

ABSTRACT

Background

Heavy menstrual bleeding is an important health problem. Two frequently used therapies are the levonorgestrel intrauterine system (LNG-IUS) and endometrial ablation. The LNG-IUS can be applied easily by the general practitioner, which saves costs, but has considerable failure rates. As an alternative, endometrial ablation is also very effective, but this treatment has to be performed by a gynaecologist. Due to lack of direct comparison of LNG-IUS with endometrial ablation, there is no evidence based preferred advice for the use of one of these treatment possibilities.

Method

A multicentre randomised controlled trial, organised in a network infrastructure in the Netherlands in which general practitioners and gynaecologists collaborate.

Women ≥ 34 years with heavy menstrual bleeding, a Pictorial Blood Assessment Chart (PBAC) score exceeding 150 points and no future child wish can participate in the trial. After informed consent, women will be randomised to a strategy starting with the LNG-IUS or a strategy starting with endometrial ablation.

The primary outcome is the PBAC-score at 24 months of follow-up. Secondary outcomes are patient satisfaction, complications, number of reinterventions, menstrual bleeding pattern, quality of life, sexual function, sick leave and costs. As predictors of effect of intervention we also measure level of coagulation factors.

Discussion

This study, considering both effectiveness and cost effectiveness of LNG-IUS versus endometrial ablation may well improve care for women with heavy menstrual bleeding.

Trial registration: Dutch trial register, number NTR2984

Background

Heavy menstrual bleeding is a frequent problem that affects many premenopausal women in the Netherlands and is the most important reason for a visit to the outpatient department of gynaecology.^{1,2} Each year one in 20 women between 30 to 49 years of age consult their general practitioner (GP) with heavy menstrual bleeding.³ Hormonal treatment with the oral contraceptive pill or the levonorgestrel-releasing intrauterine system (LNG-IUS) or non-hormonal treatment with tranexamic acid or non-steroidal anti-inflammatory drugs (NSAIDs) are advised as treatments of first choice. Nevertheless 77% of the women are not willing to continue their treatment and often end up undergoing other treatment or even surgery.^{3,4}

Hysterectomy is a definitive solution for the treatment of heavy menstrual bleeding, and in 2010 11,038 women underwent a hysterectomy due to bleeding disorders in the Netherlands.⁵ Recently it was reported that hysterectomy should be considered the preferred strategy for the treatment of heavy menstrual bleeding based on cost-effectiveness.⁶ Nevertheless, it is a major surgical procedure and has significant physical complications and social and economic costs.⁷

A significant number of women with heavy menstrual bleeding who seek treatment will not benefit from, or will not wish to continue the medical treatment and are keen to preserve their uterus.^{3,8} Many women opt for a less invasive treatment, even when they are informed of the fact that success is not always assured.^{1,9}

Two frequently used minimally invasive treatment options for heavy menstrual bleeding are the LNG-IUS and endometrial ablation. Intrauterine devices were initially introduced as contraceptives, but after the addition of progestogen (LNG-IUS) these devices also reduce menstrual bleeding effectively. The local release of levonorgestrel in the uterine cavity suppresses endometrial growth. A systematic review on the effectiveness of the LNG-IUS in heavy menstrual bleeding concluded that the reduction of menstrual blood loss was 79-96% in the LNG-IUS group.¹⁰⁻¹³ In women with heavy menstrual bleeding who presented to primary care providers, the LNG-IUS was more effective than usual medical treatment in reducing the effect of heavy menstrual bleeding on quality of life.¹⁴ However, up to 60% of women discontinue LNG-IUS within 5 years because of unscheduled bleeding, pain, and/or systemic progestogenic side-effects.¹⁵ In the Royal College of Obstetricians and Gynaecologists (RCOG) guideline on heavy menstrual

bleeding, the use of the LNG-IUS is the first therapeutic option when drug treatment has failed. This is not based on proven cost-effectiveness.¹⁵ Endometrial destruction techniques, which aim to destroy or remove the endometrial tissue, have become alternatives to hysterectomies. In 1991 in the Netherlands 21,433 hysterectomies were performed, a number that was reduced to 16,320 in 1998 and just above 11,000 in 2010, which was partly due to the start of using endometrial ablation.¹⁶ Approximately 47-58% percent of women report an amenorrhea after treatment with the bipolar ablation device.^{5,17-20} The satisfaction rates of endometrial ablation have been evaluated by several randomised controlled trials and is showing a rate of about 90%.^{19,21,22} Ablation techniques result in shorter duration of surgery, shorter hospital stay and quicker recovery time compared to hysterectomy.

Seven trials compared the LNG- IUS with transcervical resection of the endometrium or balloon ablation. In the meta-analysis a significant reduction in mean pictorial blood assessment chart (PBAC) score was reported for all women and a significant lower PBAC-score was reported in the surgical group. One study showed a significantly lower median PBAC-score in the LNG-IUS group at nine months and one year follow-up.²³ Further outcomes showed no difference in satisfaction rates, amenorrhea rates, duration of menstruation, further surgical treatment or quality of life (QoL). Nevertheless, the studies are small, most studies have a short period of follow-up and contain a lot of non-compliance, which makes interpretation of outcomes difficult.^{12,20,23-27}

Consequently, usual care in the Netherlands implies one of these two strategies for the treatment of heavy menstrual bleeding if drug therapy fails. The strategy LNG-IUS means prescription and placement by the GP or gynaecologist without anaesthesia. The strategy endometrial ablation implies performing an endometrial ablation by a gynaecologist in day-care or outpatient clinic with or without general anaesthesia. Due to lack of sufficiently powered studies directly comparing LNG-IUS with endometrial ablation, there is currently no evidence based advice for the use of one of these treatment possibilities. We propose a randomised controlled trial in which these two strategies are compared in the treatment of heavy menstrual bleeding. The study will focus on (cost-) effectiveness, patient satisfaction and QoL.

METHODS

Objective

The aim of this study is to assess the effects and cost-effectiveness of a strategy starting with LNG-IUS versus a strategy starting with endometrial ablation in women with heavy menstrual bleeding. This study will also evaluate QoL, sexual function and as predictors of effect of the intervention we also measure level of coagulation factors.

Trial design

This study is a multicentre randomised controlled trial with an economic evaluation alongside it. It will be performed by the Dutch Consortium for Studies in Women's Health and the department of General Practice of the University Medical Centre Groningen. The study will be organised in a network infrastructure in which GPs and gynaecologists collaborate. Participating hospitals can be district, teaching, third referral or university hospitals. The study is conducted according to the principles of the Declaration of Helsinki and in accordance with the Medical Research Involving Human Subjects Act (WMO) and has been approved by the ethics committee of the Academic Medical Centre Amsterdam (ref. no MEC 10/183). The protocol is registered in the Dutch trial register, number NTR2984.

Eligibility criteria

Women ≥ 34 years suffering from heavy menstrual bleeding, with a PBAC-score exceeding 150 points can participate in the trial.

Women who might want to get pregnant in the future will be excluded as an endometrial ablation is an absolute contraindication for pregnancies. Other exclusion criteria are abnormal cervix cytology in the past 5 years, a sounding length of more than 10 cm, intracavitary fibroids or polyps or large intramural fibroids determined by either a transvaginal ultrasound (TVU) or a bimanual vaginal examination.

Patient recruitment, randomization and collection of data

Eligible patients are identified by the gynaecologist in the participating hospitals or by participating GPs. Before entry into the study, inclusion and exclusion criteria are

checked by research nurses and if women are eligible they will also be counselled by these experienced research nurses. They will be informed about the aims, methods, reasonably anticipated benefits and potential hazards of the study. We will ask separate informed consent for taking and analysing blood on levels of coagulation factors. After given written informed consent women will be randomised to a strategy starting with a LNG-IUS or a strategy starting with endometrial ablation. Randomisation is performed by accessing a web-based randomisation program. We will stratify for inclusion by GP or by gynaecologist. Patients will be randomised into two groups in a ratio of 1:1, using permuted block randomisation with a variable block size.

Participants will be given a computer generated numeric code. Data handling will be done anonymously, with the patient code only available to the local investigator and the research nurse working in the local centre. At the local centres, baseline data and follow-up data collection is the responsibility of the specialised research nurse. Data will be collected using a website dedicated to studies in the Dutch Consortium for women's health and reproductive medicine studies (www.studies-obsgyn.nl). In accordance with guidelines of the Dutch Federation of University Medical Centres (NFU) the data will be kept for 20 years.

Interventions

The LNG-IUS can be placed either by a GP or by a gynaecologist. In the endometrial ablation group, a second generation bipolar ablation technique (Novasure®) will be performed by a gynaecologist.

Measurements and follow-up

Before randomisation patients are asked to fill out a PBAC. Prior to start of the treatment, participating patients will fill in two questionnaires measuring health-related quality of life: the Short Form-36 (SF-36), a validated general questionnaire and the Shaw heavy menstrual bleeding questionnaire, a disease specific questionnaire, which we validated linguistically. Furthermore, patients are asked to complete questionnaires to measure sexual functioning: the Female Sexual Function Index (FSFI), Female Sexual Distress Scale (FSDS) and a questionnaire to measure (indirect) costs. Baseline demographic

characteristics, medical history and menstruation pattern are recorded in a Case Record Form (CRF).

Patients will fill in the PBAC and the same questionnaires used at baseline at 3, 6, 12 and 24 months after randomisation. Since we compare strategies starting with LNG-IUS or endometrial ablation, patients who are not satisfied with the (randomised) treatment effect can choose another treatment, or opt for hysterectomy. Nevertheless, they will be asked to complete follow-up questionnaires.

Outcome measures

Primary outcome is the PBAC-score at 24 months of follow-up. Secondary outcomes are patient satisfaction, complications, number of reinterventions, menstrual bleeding pattern, including rates of amenorrhea, QoL, sexual function, sick leave and costs and as predictors of effect of intervention we also measure level of coagulation factors.

Statistical analysis

Sample size

The study is designed as a noninferiority study. Based on previous studies, we estimate the mean PBAC-score at 24 months of follow-up to be 50 points in the LNG-IUS group and 40 points in the endometrial ablation group, with a SD of 40.²³⁻²⁶ Previous studies in women with heavy menstrual bleeding have shown that a 50 point difference in PBAC-score between treatments represents a clinically meaningful difference in reduction of menstrual bleeding. In our study we use a noninferiority margin of 25 points. With this margin and estimated mean PBAC-scores of 50 and 40 points in resp. the LNG-IUS and endometrial ablation group, the estimated scores in both treatment groups will be far below the upper range of normal menstrual blood loss. Using a dropout rate of 15%, an alpha error of 2.5% and Beta error of 20%, we need to include 266 patients (133 patients per treatment group).

Data analysis

Data will be analysed according to the intention to treat principle. Statistical analysis will be performed using the software Statistical Package for the Social Sciences (SPSS, Inc., Chicago, IL, USA).

Point estimates and 95% confidence intervals will be calculated for different time points. First, the treatment effect at 24 months of follow-up of a strategy starting with LNG-IUS compared to a strategy starting with endometrial ablation will be investigated. For not normally distributed continuous variables, differences between groups will be tested with the Mann–Whitney U test. Differences between categorical variables will be tested with the Chi-square test or Fisher’s exact test. Secondary, to assess the treatment effect over time, and the interaction of the intervention effect, we will use a mixed effects model. Time to reintervention will be compared with Kaplan Meier analysis and Cox regression. The analysis of the influence of lower levels of coagulation on the treatment effect of LNG-IUS and endometrial ablation will be explorative.

Economic evaluation

The economic analysis will be conducted from a societal perspective including direct medical and direct and indirect non-medical costs. Relevant direct costs components that will be taken into account are costs of the LNG-IUS and endometrial ablation, interventions for complications, hospital admission and home care, consisting of both professional care as well as informal care. Sick leave and loss of productivity at work will be taken into account as indirect non-medical costs.

The economic evaluation will be designed as a cost-effectiveness analysis, with the costs per treatment resulting in a reduction of menstrual blood loss as an outcome measure. We will also perform a cost-utility analysis expressing the incremental costs per Quality Adjusted Life Years based on the SF-36. Robustness of the results (costs and health outcomes) for various assumptions and parameter estimates will be explored in sensitivity analyses and visualised in ICER-graphs and cost-effectiveness acceptability curves.

DISCUSSION

In view of its high prevalence, an optimal treatment for heavy menstrual bleeding is of utmost importance. Usual care in The Netherlands implies two strategies for the treatment of heavy menstrual bleeding if drug therapy fails. First, there is the LNG-IUS that can be applied easily by the GP, which saves costs, but has considerable failure

rates. As an alternative, endometrial ablation is also very effective, but this treatment has to be performed by a gynaecologist. Large randomised controlled trials comparing LNG-IUS with endometrial ablation are lacking and hence no preferred advice for the use of one of these treatment possibilities is available. When we estimate that treatment for heavy menstrual bleeding involves 10,000 women annually, there can be a potential saving from implementing the best strategy as appears from the proposed study. Therefore, a direct comparison of both strategies on cost-effectiveness is essential to determine which strategy should be advocated.

This is the first study comparing LNG-IUS and endometrial ablation which will be organised in a network infrastructure in which both GP's and gynaecologists collaborate. It has got the largest study population compared to other trials and has a long-term follow-up of 2 years. Besides the objective outcome PBAC we also measure patient satisfaction, QoL and sexual function. Due to this study design and outcome measures, the results will be applicable for a large group of women suffering from heavy menstrual bleeding.

Acknowledgements

This study is funded by The Dutch organisation for Health Research and Development (ZonMw) grant, number 171202001.

REFERENCES

1. Palep-Singh M, Prentice A: Epidemiology of abnormal uterine bleeding. *Best Pract Res Clin Obstet Gynaecol* 2007, 21(6):887–890.
2. Shapley M, Jordan K, Croft PR: An epidemiological survey of symptoms of menstrual loss in the community. *Br J Gen Pract* 2004, 54(502):359–363.
3. Lethaby AE, Cooke I, Rees M: Progesterone or progestogen-releasing intrauterine systems for heavy menstrual bleeding. *Cochrane Database Syst Rev* 2005, 4, CD002126.
4. Cooper K, Lee A, Chien P, Raja E, Timmaraju V, Bhattacharya S: Outcomes following hysterectomy or endometrial ablation for heavy menstrual bleeding: Retrospective analysis of hospital episode statistics in scotland. *BJOG* 2011, 118(10):1171–1179.
5. Prisma: Dutch Hospital Data; 2009.
6. Roberts TE, Tsourapas A, Middleton LJ, et al: Hysterectomy, endometrial ablation, and levonorgestrel releasing intrauterine system (mirena) for treatment of heavy menstrual bleeding: Cost effectiveness analysis. *BMJ* 2011, 342:d2202.
7. Abbott JA, Garry R: The surgical management of menorrhagia. *Hum Reprod Update* 2002, 8(1):68–78.
8. Shaw RW: Assessment of medical treatments for menorrhagia. *Br J Obstet Gynaecol* 1994, 101(Suppl 11):15–18.
9. Nagele F, Rubinger T, Magos A: Why do women choose endometrial ablation rather than hysterectomy? *Fertil Steril* 1998, 69(6):1063–1066.
10. Lahteenmaki P, Haukkamaa M, Puolakka J, et al: Open randomised study of use of levonorgestrel releasing intrauterine system as alternative to hysterectomy. *BMJ* 1998, 316(7138):1122–1126.
11. Milsom I, Andersson K, Andersch B, Rybo G: A comparison of flurbiprofen, tranexamic acid, and a levonorgestrel-releasing intrauterine contraceptive device in the treatment of idiopathic menorrhagia. *Am J Obstet Gynecol* 1991, 164(3):879–883.
12. Crosignani PG, Vercellini P, Mosconi P, Oldani S, Cortesi I, De Giorgi O: Levonorgestrel-releasing intrauterine device versus hysteroscopic endometrial resection in the treatment of dysfunctional uterine bleeding. *Obstet Gynecol* 1997, 90(2):257–263.
13. Irvine GA, Campbell-Brown MB, Lumsden MA, Heikkila A, Walker JJ, Cameron IT: Randomised comparative trial of the levonorgestrel intrauterine system and

- norethisterone for treatment of idiopathic menorrhagia. *Br J Obstet Gynaecol* 1998, 105(6):592–598.
14. Gupta J, Kai J, Middleton L, et al: Levonorgestrel intrauterine system versus medical therapy for menorrhagia. *N Engl J Med* 2013, 368(2):128–137.
 15. Ewies AA: Levonorgestrel-releasing intrauterine system—the discontinuing story. *Gynecol Endocrinol* 2009, 25(10):668–673.
 16. Middleton LJ, Champaneria R, Daniels JP, et al: Hysterectomy, endometrial destruction, and levonorgestrel releasing intrauterine system (mirena) for heavy menstrual bleeding: Systematic review and meta-analysis of data from individual patients. *BMJ* 2010, 341:c3929.
 17. Gallinat A, Nugent W: NovaSure impedance-controlled system for endometrial ablation. *J Am Assoc Gynecol Laparosc* 2002, 9(3):283–289.
 18. Clark TJ, Samuels N, Malick S, Middleton L, Daniels J, Gupta J: Bipolar radiofrequency compared with thermal balloon endometrial ablation in the office: a randomised controlled trial. *Obstet Gynecol* 2011, 117(5):1228.
 19. Bongers MY, Bourdrez P, Mol BW, Heintz AP, Brodmann HA: Randomised controlled trial of bipolar radio-frequency endometrial ablation and balloon endometrial ablation. *BJOG* 2004, 111(10):1095–1102.
 20. Marjoribanks J, Lethaby A, Farquhar C: Surgery versus medical therapy for heavy menstrual bleeding. *Cochrane Database Syst Rev* 2006, 2, CD003855.
 21. Penninx JP, Herman MC, Mol BW, Bongers MY: Five-year follow-up after comparing bipolar endometrial ablation with hydrothermablation for menorrhagia. *Obstet Gynecol* 2011, 118(6):1287–1292.
 22. Bongers MY, Mol BW: Thermal balloon ablation versus endometrial resection for treatment of abnormal uterine bleeding. *Hum Reprod* 2000, 15(6):1424–1425.
 23. Shaw RW, Symonds IM, Tamizian O, Chaplain J, Mukhopadhyay S: Randomised comparative trial of thermal balloon ablation and levonorgestrel intrauterine system in patients with idiopathic menorrhagia. *Aust N Z J Obstet Gynaecol* 2007, 47(4):335–340.
 24. Istre O, Trolle B: Treatment of menorrhagia with the levonorgestrel intrauterine system versus endometrial resection. *Fertil Steril* 2001, 76(2):304–309.
 25. Barrington JW, Arunkalaivanan AS, Abdel-Fattah M: Comparison between the levonorgestrel intrauterine system (LNG-IUS) and thermal balloon ablation in the treatment of menorrhagia. *Eur J Obstet Gynecol Reprod Biol* 2003, 108(1):72–74.

26. Busfield RA, Farquhar CM, Sowter MC, et al: A randomised trial comparing the levonorgestrel intrauterine system and thermal balloon ablation for heavy menstrual bleeding. *BJOG* 2006, 113(3):257-263.
27. Soysal M, Soysal S, Ozer S: A randomised controlled trial of levonorgestrel releasing IUD and thermal balloon ablation in the treatment of menorrhagia. *Zentralbl Gynakol* 2002, 124(4):213-219.

