

University of Groningen

## Understanding How Asthma Starts

Koppelman, Gerard H; Kersten, Elin T G

*Published in:*  
 American Journal of Respiratory and Critical Care Medicine

*DOI:*  
[10.1164/rccm.202102-0443ED](https://doi.org/10.1164/rccm.202102-0443ED)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
 Publisher's PDF, also known as Version of record

*Publication date:*  
 2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Koppelman, G. H., & Kersten, E. T. G. (2021). Understanding How Asthma Starts: Longitudinal Patterns of Wheeze and the Chromosome 17q Locus. *American Journal of Respiratory and Critical Care Medicine*, 203(7), 793-795. <https://doi.org/10.1164/rccm.202102-0443ED>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

found no association between nasal ORM DL3 gene expression and 17q21 genotype. Nevertheless, these findings should inform further studies on the regulation and role of SPT subunits in asthma.

Overall, this study lends evidence supporting the concept that genetically altered sphingolipid metabolism in children who carry 17q21 asthma-risk genotypes may lead to functional consequences on airway resistance, acting as a predisposing factor for the development of asthma. Although the manifestations of classic inborn disorders of sphingolipid metabolism mainly result from the accumulation of toxic products affecting the nervous system and skin, decreased synthesis of bioactive lipids such as sphinganine-1-phosphate and sphingosine-1-phosphate may also have specific functional consequences (10). We may also learn from other manifestations of decreased SPT activity, such as in hereditary sensory autonomic neuropathy caused by a loss of function mutation of the SPT subunit *Sptlc2*, which has recently been shown to have relevant consequences on immune cell function (11). Although the exact role of sphingolipids in asthma remains enigmatic, Rago and colleagues have opened the door a little further, providing another glimpse of how this class of lipids is involved in the complex pathogenesis of childhood asthma. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Jennie G. Ono, M.D., M.S.  
Department of Pediatrics  
Weill Cornell Medicine  
New York, New York

Stefan Worgall, M.D., Ph.D.  
Department of Pediatrics  
Department of Genetic Medicine  
and  
Druker Institute for Children's Health  
Weill Cornell Medicine  
New York, New York

## References

1. Moffatt MF, Kabesch M, Liang L, Dixon AL, Strachan D, Heath S, *et al*. Genetic variants regulating ORM DL3 expression contribute to the risk of childhood asthma. *Nature* 2007;448:470–473.
2. Miller M, Tam AB, Cho JY, Doherty TA, Pham A, Khorram N, *et al*. ORM DL3 is an inducible lung epithelial gene regulating metalloproteases, chemokines, OAS, and ATF6. *Proc Natl Acad Sci USA* 2012;109:16648–16653.
3. Breslow DK, Collins SR, Bodenmiller B, Aebersold R, Simons K, Shevchenko A, *et al*. Orm family proteins mediate sphingolipid homeostasis. *Nature* 2010;463:1048–1053.
4. Rago D, Pedersen C-ET, Huang M, Kelly RS, Gürdeniz G, Brustad N, *et al*. Characteristics and mechanisms of a sphingolipid-associated childhood asthma endotype. *Am J Respir Crit Care Med* 2021;203:853–863.
5. Ono JG, Kim BI, Zhao Y, Christos PJ, Tesfaigzi Y, Worgall TS, *et al*. Decreased sphingolipid synthesis in children with 17q21 asthma-risk genotypes. *J Clin Invest* 2020;130:921–926.
6. Huang M, Kelly RS, Chu SH, Kachroo P, Gürdeniz G, Chawes BL, *et al*. Maternal metabolome in pregnancy and childhood asthma or recurrent wheeze in the Vitamin D Antenatal Asthma Reduction Trial. *Metabolites* 2021;11:65.
7. Worgall TS, Veerappan A, Sung B, Kim BI, Weiner E, Bholah R, *et al*. Impaired sphingolipid synthesis in the respiratory tract induces airway hyperreactivity. *Sci Transl Med* 2013;5:186ra67.
8. Das S, Miller M, Broide DH. Chromosome 17q21 genes ORM DL3 and GSDMB in asthma and immune diseases. *Adv Immunol* 2017;135:1–52.
9. Siow D, Sunkara M, Dunn TM, Morris AJ, Wattenberg B. ORM DL3/serine palmitoyltransferase stoichiometry determines effects of ORM DL3 expression on sphingolipid biosynthesis. *J Lipid Res* 2015;56:898–908.
10. Dunn TM, Tiff CJ, Proia RL. A perilous path: the inborn errors of sphingolipid metabolism. *J Lipid Res* 2019;60:475–483.
11. Wu J, Ma S, Sandhoff R, Ming Y, Hotz-Wagenblatt A, Timmerman V, *et al*. Loss of neurological disease HSN-1-associated gene SPTLC2 impairs CD8<sup>+</sup> T cell responses to infection by inhibiting T cell metabolic fitness. *Immunity* 2019;50:1218–1231, e5.

Copyright © 2021 by the American Thoracic Society



## Understanding How Asthma Starts: Longitudinal Patterns of Wheeze and the Chromosome 17q Locus

Most childhood asthma starts in the preschool years. Symptoms such as wheeze, cough, and dyspnea are not specific to asthma and can also represent transient symptoms due to viral respiratory tract infections. Triggers of preschool wheeze can change over time (1) and therefore are not reliable predictors of asthma. No valid, reproducible diagnostic or predictive test of preschool asthma is

currently available (2). This inability to diagnose preschool asthma has seriously impeded better understanding of childhood-onset asthma and the ability to design targeted, early-life interventions.

In this issue of the *Journal*, Hallmark and colleagues (pp. 864–870) combine two strategies to better understand the development of childhood wheeze: the description of longitudinal patterns of wheeze in seven U.S. birth cohorts participating in the Children's Respiratory Research and Environment Workgroup, and the investigation of the association of these longitudinal wheezing phenotypes with the 17q12–21 locus ("17q") (3). 17q is the most replicated childhood-onset asthma locus (4). Using data from birth until age 11 years, their latent class modeling revealed four classes of wheeze: infrequent (no or low presence of wheeze;

Ⓒ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Originally Published in Press as DOI: 10.1164/rccm.202102-0443ED on February 23, 2021

estimated prevalence 62% based on *post hoc* assignment to most probable class), transient (wheeze in the first years of life, but absent at school age; 17%), late-onset (10%), and persistent (wheeze persisting from preschool to school age; 11%). These patterns were similar in children of European American (EA) and African American (AA) descent. Notably, AA children had higher probability to be assigned to the persistent wheeze group, indicating a higher burden of wheeze in these children. This clearly calls for further research to better understand this higher burden in AA children.

Are four wheezing patterns all there are? Whereas many longitudinal birth cohorts have identified these four patterns (5, 6), two additional wheezing patterns have been described by latent class analysis of UK and Dutch birth cohorts, with an additional transient wheeze group (prolonged early wheeze) and an additional late-onset group, the intermediate-onset wheeze group, which is strongly associated with atopy (7). This illustrates the strength but also the limitation of data-driven methods: the ability to define wheezing phenotypes is indeed strongly driven by the data, such as the number of assessments of wheeze and number of children studied.

How does the description of longitudinal wheezing phenotypes advance our understanding of asthma development? No wheezing pattern was uniquely predictive of asthma; in fact, all wheezing patterns were associated with a higher chance of ever developing asthma (3). Interpretation of this analysis may be complicated by the fact that a doctor's diagnosis of asthma at any age was taken as an outcome, as many children obtained their asthma diagnosis in the first 3 years of life. However, other studies also indicate that there is no wheezing pattern that uniquely leads to asthma, defined by asthma at school age, reduced lung function, or airway hyperresponsiveness (7). Thus, the pattern itself has limited ability to predict asthma, and the fact that these wheezing patterns can only be defined retrospectively limits their clinical use.

Do the longitudinal wheezing phenotypes represent unique pathophysiological mechanisms and thus represent endotypes of asthma (8)? Genetic studies may reveal this, and Hallmark and coworkers chose to investigate SNPs at the 17q locus. Recent analysis has shown that this locus is strongly related to the age of onset of asthma but not to the onset of eczema or hay fever (9). 17q SNPs were associated with any of the three wheezing phenotypes in both AA and EA children, suggesting that these phenotypes have a shared genetic origin and, thus, that differences may actually be influenced by other factors, such as the environment. Fine mapping of this locus has been challenging because of extensive linkage disequilibrium (LD). By investigating associations between wheezing phenotypes and 17q SNPs in both EA and AA children, in whom the LD blocks extend over shorter distances, they aimed to pinpoint potential causal variants. The same technique recently allowed Ober and coworkers (10) to narrow down the association between 17q SNPs and childhood-onset asthma (<6 yr) in AA children to two SNPs: rs2305480 and rs8076131. In a subsequent expression quantitative trait loci analysis, rs2305480, an SNP in the coding region for *GSDMB* (gasdermin B), was strongly associated with expression of *GSDMB* in airway epithelial cells. Similarly, Gui and coworkers (11) used next-generation DNA sequencing data from AA individuals in three cohorts to evaluate the association between 17q and childhood-onset asthma (<5 yr). The lead association from their meta-analysis, rs11078928, is located in a

4-kb haplotype block containing four potentially functional polymorphisms in very strong LD, including rs2305480. rs11078928 affects splicing of *GSDMB* transcripts, and expression levels of these transcript isoforms in whole blood RNA sequencing were associated with asthma status, making this a likely causal candidate mechanism of childhood-onset asthma (12).

The authors propose that the 17q locus may be considered a "wheezing locus" instead of an "asthma locus." However, this statement may need support from future studies. For most of the tested SNPs, the odd ratios are higher in wheezing classes with higher proportions of children with doctor-diagnosed asthma. These results are in line with previous observations by Granell and coworkers (13) who describe a strong association of 17q SNPs, including rs2305480, with persistent and intermediate-onset childhood wheezing phenotypes, but a much weaker association with transient wheeze. Similarly, Sordillo and coworkers (14) observed an increased odds for persistent wheeze, but no association with transient or late-onset wheeze, for the 17q SNP rs12603332. In fact, in one of the first studies to assess the association between rs2305480 and preschool wheeze, the association was strongest for children wheezing in the third and fourth year of life (likely indicating persistent wheeze) and nonsignificant for wheeze in the first year of life (with the highest prevalence of the transient wheezing phenotype) (15).

Hallmark and coworkers (3) have shown that it is possible to harmonize and jointly analyze multiple birth cohorts. Investigating the genetics of longitudinal wheezing phenotypes, in the future at a genome-wide basis, may help to disentangle mechanisms of early life wheeze and subsequent asthma development. ■

---

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Gerard H. Koppelman, M.D., Ph.D.  
Elin T. G. Kersten, Ph.D.  
*Department of Pediatric Pulmonology and Pediatric Allergology  
University Medical Center Groningen  
Groningen, the Netherlands*  
and  
*Groningen Research Institute for Asthma and COPD  
University Medical Center Groningen  
Groningen, the Netherlands*

ORCID IDs: 0000-0001-8567-3252 (G.H.K.); 0000-0001-7423-860X (E.T.G.K.).

---

## References

1. Brand PLP, Caudri D, Eber E, Gaillard EA, Garcia-Marcos L, Hedlin G, *et al*. Classification and pharmacological treatment of preschool wheezing: changes since 2008. *Eur Respir J* 2014;43: 1172–1177.
2. Kothalawala DM, Kadalayil L, Weiss VBN, Kyaly MA, Arshad SH, Holloway JW, *et al*. Prediction models for childhood asthma: a systematic review. *Pediatr Allergy Immunol* 2020;31:616–627.
3. Hallmark B, Wegienka G, Havstad S, Billheimer D, Ownby D, Mendonca EA, *et al*. ECHO-CREW. Chromosome 17q12-21 variants are associated with multiple wheezing phenotypes in childhood. *Am J Respir Crit Care Med* 2021;203:864–870.
4. Stein MM, Thompson EE, Schoettler N, Helling BA, Magnaye KM, Stanhope C, *et al*. A decade of research on the 17q12-21 asthma locus: piecing together the puzzle. *J Allergy Clin Immunol* 2018;142: 749–764, e3.

5. Belgrave DCM, Custovic A, Simpson A. Characterizing wheeze phenotypes to identify endotypes of childhood asthma, and the implications for future management. *Expert Rev Clin Immunol* 2013;9: 921–936.
6. Oksel C, Granell R, Haider S, Fontanella S, Simpson A, Turner S, *et al.*; STELAR investigators, breathing Together investigators. Distinguishing wheezing phenotypes from infancy to adolescence: a pooled analysis of five birth cohorts. *Ann Am Thorac Soc* 2019;16:868–876.
7. Savenije OE, Granell R, Caudri D, Koppelman GH, Smit HA, Wijga A, *et al.* Comparison of childhood wheezing phenotypes in 2 birth cohorts: ALSPAC and PIAMA. *J Allergy Clin Immunol* 2011;127: 1505–1512, e14.
8. Lötvaall J, Akdis CA, Bacharier LB, Bjermer L, Casale TB, Custovic A, *et al.* Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011;127: 355–360.
9. Ferreira MAR, Vonk JM, Baurecht H, Marenholz I, Tian C, Hoffman JD, *et al.*; 23andMe Research Team; collaborators of the SHARE study. Age-of-onset information helps identify 76 genetic variants associated with allergic disease. *PLoS Genet* 2020;16:e1008725.
10. Ober C, McKennan CG, Magnaye KM, Altman MC, Washington C III, Stanhope C, *et al.*; Environmental Influences on Child Health Outcomes-Children's Respiratory Research Workgroup. Expression quantitative trait locus fine mapping of the 17q12-21 asthma locus in African American children: a genetic association and gene expression study. *Lancet Respir Med* 2020;8:482–492.
11. Gui H, Levin AM, Hu D, Sleiman P, Xiao S, Mak ACY, *et al.* Mapping the 17q12-21.1 locus for variants associated with early-onset asthma in African Americans. *Am J Respir Crit Care Med* 2021;203: 424–436.
12. Raby BA, Weiss ST. Diversity and the splice of life: mapping the 17q12-21.1 locus for variants associated with early-onset asthma in African American individuals. *Am J Respir Crit Care Med* 2021;203:401–403.
13. Granell R, Henderson AJ, Timpson N, St Pourcain B, Kemp JP, Ring SM, *et al.* Examination of the relationship between variation at 17q21 and childhood wheeze phenotypes. *J Allergy Clin Immunol* 2013; 131:685–694.
14. Sordillo JE, Coull BA, Rifas-Shiman SL, Wu AC, Lutz SM, Hivert MF, *et al.* Characterization of longitudinal wheeze phenotypes from infancy to adolescence in Project Viva, a prebirth cohort study. *J Allergy Clin Immunol* 2020;145:716–719, e8.
15. van der Valk RJP, Duijts L, Kerkhof M, Willemssen SP, Hofman A, Moll HA, *et al.* Interaction of a 17q12 variant with both fetal and infant smoke exposure in the development of childhood asthma-like symptoms. *Allergy* 2012;67:767–774.

Copyright © 2021 by the American Thoracic Society



## Oral Corticosteroids Tapering in Severe Asthma

First marketed 70 years ago, corticosteroids transformed the life of patients suffering from asthma and quickly became the mainstay of treatment for this condition. Despite major developments in therapeutic options since, particularly with the use of inhaled corticosteroids more than 40 years ago, the powerful antiinflammatory effects of oral corticosteroids (OCS) are as of yet impossible to replace completely, explaining their persistent use in asthma management. Most of the time, they are prescribed intermittently to treat severe exacerbations, although some patients require them chronically to achieve asthma control (1). However, OCS are associated with well-recognized long-term side effects and an increase in mortality (2, 3). Recent evidence suggests that this risk is related to the cumulative lifetime exposure to OCS (4, 5), implying that even repeated short courses may have a significant impact on their associated morbidity.

More recently, monoclonal antibodies brought the first real long-term alternative to OCS in severe asthma since the 1950s. They are powerful antiinflammatory agents targeting T-helper cell type 2 (Th2) inflammation with minimal side effects and with corticosteroid-sparing properties (6–9). Their availability provoked a change in OCS perception in severe asthma, from a necessary evil to an increasingly avoidable one. With the increasing use of biologics, tapering and cessation of maintenance OCS has become much more common and feasible, but specific guidance on how to proceed is lacking.

Numerous studies exploring steroid-sparing drugs have reported their OCS weaning protocols. The OCS tapering regimens used were quite variable, as were the assessments of asthma control, biomarker use, and screening for adrenal insufficiency, thus making generalization difficult. Many of them also lacked the details needed to be efficiently implemented in clinical practice. An exception would be the ongoing PONENTE trial, investigating the safety and efficacy of OCS tapering after initiation of benralizumab. Although not evidenced-based, it provides a detailed OCS reduction algorithm with systematic assessment of adrenal insufficiency that could be used by clinicians.

Research and guidelines have recognized the need to reach the minimal effective dose when OCS are needed for long-term treatment of severe asthma. To achieve this, their focus and advice has been to optimize asthma control strategies and use of OCS-sparing drugs without clear guidance on how to actually proceed with weaning. Hence, there are currently no standardized guidelines on how and when to safely perform OCS tapering. A recent review identified this lack of clear recommendations as a clinical barrier to reduce OCS exposure in severe asthma (10).

In this issue of the *Journal*, Suehs and colleagues (pp. 871–881) provide an expert consensus report on the important topic of OCS use and tapering in patients with asthma, including statements on less frequent conditions such as eosinophilic granulomatosis with polyangiitis and allergic bronchopulmonary aspergillosis (11). A modified Delphi method was used to develop a consensus (>70% agreement) among 131 experts from different specialties, mostly pulmonologists (73%) and allergists (18%), in addition to patient advocacy organization representatives. Although opinions sometimes differed, some general principles for use and reduction of OCS were agreed on.

This study is a first major attempt to provide clinicians with guidelines based on expert opinion specifically on OCS use for

©This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Originally Published in Press as DOI: 10.1164/rccm.202010-4001ED on November 19, 2020