

University of Groningen

The role of E-cadherin/ β -catenin signalling in the development of an asthmatic airway epithelial phenotype

Kuchibhotla, Virinchi

DOI:
[10.33612/diss.172561514](https://doi.org/10.33612/diss.172561514)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Kuchibhotla, V. (2021). *The role of E-cadherin/ β -catenin signalling in the development of an asthmatic airway epithelial phenotype*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.172561514>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER 6

Inhibition of β -catenin/CREB binding protein signaling attenuates house dust mite-induced goblet cell metaplasia in mice

Virinchi N. S. Kuchibhotla^{1,2,3*}, Malcolm R. Starkey^{1,4,5*}, Andrew T. Reid^{6,7}, Irene H. Heijink^{2,3,8}, Martijn C. Nawijn^{2,3}, Philip M. Hansbro^{1,7,9*}, Darryl A. Knight^{1,10,11*}

¹School of Biomedical Sciences and Pharmacy, University of Newcastle, Callaghan, New South Wales, Australia.

²University of Groningen, University Medical Center Groningen, Department of Pathology & Medical Biology, laboratory of Experimental Pulmonology and Inflammation Research (EXPIRE), Groningen, The Netherlands.

³University of Groningen, University Medical Center Groningen, GRIAC Research Institute, Groningen, the Netherlands.

⁴Department of Immunology and Pathology, Central Clinical School, Monash University, Melbourne, Victoria, Australia

⁵Priority Research Centre GrowUpWell, The Faculty of Health and Medicine and Hunter Medical Research Institute, The University of Newcastle, Newcastle, New South Wales, Australia

⁶School of Medicine and Public Health, University of Newcastle, Callaghan, New South Wales, Australia.

⁷Priority Research Centre for Healthy Lungs, Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia.

⁸University of Groningen, University Medical Center Groningen, Department of Pulmonology, Groningen, the Netherlands

⁹Centre for Inflammation, Centenary Institute and University of Technology Sydney, School of Life Science, Faculty of Science, Sydney, NSW, Australia.

¹⁰Providence Health Care Research Institute, Vancouver, BC, Canada.

¹¹Department of Anesthesiology, Pharmacology and Therapeutics, University of British Columbia, Vancouver, BC, Canada.

*** Authors contributed equally and are co-first and senior authors**

This chapter is under revision at *Frontiers in Physiology*, 2021

Abstract

Excessive mucus production is a major feature of allergic asthma. Disruption of epithelial junctions by allergens such as house dust mite (HDM) results in the activation of β -catenin signaling, which has been reported to stimulate goblet cell differentiation. β -catenin interacts with various co-activators including CREB binding protein (CBP) and p300, thereby regulating the expression of genes involved in cell proliferation and differentiation, respectively. We specifically investigated the role of the β -catenin/CBP signaling pathway in goblet cell metaplasia in a HDM-induced allergic airway disease model in mice using a small molecule inhibitor ICG-001. Female 6-8-week-old BALB/c mice were sensitized to HDM/saline on days 0, 1 and 2, followed by intranasal challenge with HDM/saline with or without subcutaneous ICG-001/vehicle treatment from days 14-17, and samples harvested 24 hours after the last challenge/treatment. Differential inflammatory cells in bronchoalveolar lavage (BAL) fluid were enumerated. Periodic acid–Schiff (PAS)/alcian blue (AB) staining was used to identify goblet cells/mucus production, and airway hyperresponsiveness (AHR) was assessed using invasive plethysmography. Exposure to HDM induced airway inflammation, goblet cell metaplasia and increased AHR, with increased airway resistance in response to the non-specific spasmogen methacholine. Inhibition of the β -catenin/CBP pathway using treatment with ICG-001 significantly attenuated the HDM-induced goblet cell metaplasia and infiltration of macrophages, but had no effect on eosinophils, neutrophils, lymphocytes or AHR. Increased β -catenin/CBP signaling may promote HDM-induced goblet cell metaplasia in mice.

Key words: asthma, β -catenin, goblet cell metaplasia, ICG-001

Introduction

Asthma is broadly characterized by chronic inflammation and remodeling of the airways, excessive airway mucus production, and reversible airflow obstruction with loss of lung function (1). The airway epithelium is the first line of defense against inhaled allergens and is more susceptible to damage by allergens such as house dust mite (HDM) in asthma. This results in the deterioration of cellular junctions (2), increased epithelial growth factor receptor (EGFR) signaling (3), extracellular matrix proteins (4) and expression of mesenchymal markers such as α -smooth muscle actin (α -SMA), fibronectin and vimentin (5), and airway hyperresponsiveness (AHR) (4,6). Damaged epithelial cells release pro-inflammatory

cytokines like Chemokine (C-C motif) ligand 20 (CCL20), CCL17, interleukin (IL)-25, IL-33 and thymic stromal lymphopoietin (TSLP), which activate and attract immune cells like dendritic cells (DCs) and macrophages, promoting the differentiation of type-2 cells (3,7,8). T helper 2 (Th2) cells and innate lymphoid cells (ILCs) release various cytokines including IL-4, IL-5 and IL-13, which activate B-lymphocytes, induce the infiltration of eosinophils, and enhance the differentiation of goblet cells (9–11). Besides IL-13, EGF also enhances goblet cell differentiation (12,13). Goblet cells are secretory cells that produce mucus, which consists of polypeptides, water, DNA, enzymes and high molecular weight glycoproteins called mucins (14). MUC5AC and MUC5B are the major mucins secreted by goblet cells, and their expression is increased in asthma and is associated with goblet cell metaplasia (15,16). MUC5AC production is regulated by the transcription factors SAM pointed domain containing ETS transcription factor (SPDEF) and forkhead box a2 (FOXA2), which activate and repress the expression of *MUC5AC*, respectively (15,17). In the asthmatic airway, goblet cell hyperplasia is accompanied by loss of ciliated cells concomitant with higher expression of basal cell markers cytokeratin 5 and p63 compared to healthy individuals, resembling undifferentiated epithelium undergoing aberrant repair responses that is unable to re-differentiate into a pseudostratified layer (18).

Adherens junctions, which mainly constitute the transmembrane protein E-cadherin, help in maintaining the integrity and function of the airway epithelium. Reduced expression of E-cadherin has been observed in airway epithelial cells from asthmatic donors along with decreased epithelial barrier function (19). We have previously shown that allergens like HDM disrupt E-cadherin at the cell junctions and induce goblet cell metaplasia in mice (20). In mice, embryonic knock-out of E-cadherin in airway epithelial cells induced club cell hyperplasia (21) and excessive mucus production (22). The loss of E-cadherin leads to the delocalization of its intracellular binding partner β -catenin into the cytoplasm, where its levels are regulated by a destruction complex consisting of axin, adenomatous polyposis coli APC and GSK-3 β . Activation of Wnt and/or growth factor signaling prevents phosphorylation and degradation of β -catenin by the destruction complex in the cytoplasm. The non-phosphorylated β -catenin, which is the transcriptionally active form of β -catenin, translocates to the nucleus and acts as a transcription factor to regulate the expression of various genes involved in cellular development and regulate cell fate decisions (23). Increased transcriptional activity of β -catenin has been shown to induce goblet cell

metaplasia in mice (24). The regulation of a variety of genes by β -catenin is possible because of its ability to bind to various transcriptional coactivators including CREB binding protein (CBP) and p300 leading to divergent cellular processes of cell migration and proliferation versus differentiation, respectively (25). Specifically, β -catenin/CBP signaling has been shown to regulate epithelial-to-mesenchymal transition of primary airway epithelial cells, leading to loss of epithelial markers such as E-cadherin, possibly through transcriptional regulation of E-cadherin repressors such as SNAIL and SLUG in addition to EGFR, whose downstream signaling results in the loss of E-cadherin (26,27). In addition, inhibition of β -catenin/CBP pathway by the small molecule inhibitor ICG-001 improved airway epithelial barrier function *in vitro* upon Ca^{+2} signaling-induced damage by stabilizing E-cadherin at cell junctions (28). We hypothesized that inhibition of the β -catenin/CBP pathway upon HDM-induced E-cadherin loss using ICG-001 would attenuate HDM-induced goblet cell metaplasia and AHR in a mouse model of allergic asthma.

Materials and Methods

Mice

6-8 week-old female BALB/c mice were obtained from the University of Newcastle's central animal house and housed at the Hunter Medical Research Institute animal facility in individually ventilated cages in an SPF PC2 facility. Mice were provided standard rodent chow *ad libitum*. All protocols were approved by the University of Newcastle Animal Care and Ethics Committee.

HDM model of allergic airway disease

Mice (n=6-8/group) were sensitized to HDM (*Dermatophagoides pteronyssinus*) extract (intranasal (i.n.): days 0, 1 and 2; 50 μg ; Greer Labs, North Carolina, USA) in sterile saline (50 μl) and challenged with HDM (i.n.: day 14–17; 5 μg in 50 μl saline) with/without ICG-001 (subcutaneous: day 14–17; 5mg/Kg) under isoflurane anesthesia and euthanized on day 18 (Figure 1A) as previously described (6,29).

Quantitative polymerase chain reaction (qPCR)

Total RNA was extracted from lung homogenates using TRIZOL (Sigma, Missouri, USA) as previously described (30–32). Synthesis of cDNA was performed using high-capacity cDNA reverse transcription kit with RNase inhibitor (Thermo Fisher Scientific,

Massachusetts, USA) and T100 Thermal Cycler (Bio-Rad Laboratories, California, USA). Quantitative real time PCR (qRT-PCR) targeting *Cdh1* (Mm01247357_m1, Thermo Fisher Scientific), *Muc5ac* (Mm01276718_m1, Thermo Fisher Scientific), *Muc5b* (Mm00466391_m1, Thermo Fisher Scientific), *Spdef* (Hs00171942_m1, Thermo Fisher Scientific), and *Foxa2* (Mm01976556_s1, Thermo Fisher Scientific) was performed on individual biological replicates, which were normalized to *Hprt* (Mm03024075_m1, Thermo Fisher Scientific) and presented as gene expression ($2^{-\Delta Ct}$) as previously described (33).

Histological analysis

Lung sections of 5 μm thickness were obtained for all the experimental mice and stained for goblet cells/mucins using standard Alcian blue pH 2.5 (AB) followed by periodic acid and Schiff's reagent (PAS) as previously described (30,34,35). A minimum of 8 different airways per section were imaged at 10x magnification and quantified using color deconvolution algorithm in ImageJ (National institute of Health) as previously described (30,34,35). Each data point represents the percentage of total staining intensity of mucus in the whole section per total area of the airway epithelial cells in that section. After importing the AB/PAS staining image of the section into ImageJ, the airway epithelial cells were selected by drawing a region of interest (ROI) using the 'free hand selection tool'. The area outside the airway(s) was deleted using 'clear outside' function. The 'Threshold color' was adjusted to select all airway epithelial cells and the total area of the airway epithelial cells was measured using the 'Measure' function, which was exported to an excel file. Next, by selecting the "color deconvolution" function with H-PAS vector, the image was split into three different color channels. The blue/magenta color channel was selected to quantify the AB/PAS staining intensity using the 'Threshold' function. By changing the value of the 'threshold' function, the AB/PAS staining area was selected. Setting a very high threshold value results in an increase of incorrect selection of unstained areas in the image and setting a very low threshold may not sufficiently select the AB/PAS staining area. So, the threshold was manually adjusted to a value which best covers the whole area of the AB/PAS staining. Importantly, the same threshold value was used for the selection of AB/PAS staining for all the images for consistency. After selecting the area of AB/PAS staining, the image was 'flattened' and 'inverted'. The area of the AB/PAS staining was selected using 'Threshold color' and the staining intensity was measured using 'Measure' function, which were also exported to the excel file. All the values of AB/PAS staining intensities and area of airway epithelial cells

from different airways of a single image section were separately added to get the total AB/PAS staining intensity and total area of airway epithelial cells, respectively. Finally, the % AB/PAS staining was calculated using the following equation: (total AB/PAS staining intensity/total area of airway epithelial cells) x 100%. Each data point represents the % AB/PAS staining of mucins per total area of the airway epithelial cells in a section from a single mouse.

Immune cell quantification

BAL fluid (2 ml) was prepared, and total cell numbers were determined with a hemocytometer. Cells prepared by cytocentrifugation (Shandon Cytospin; Thermo Fisher Scientific, Waltham, MA) were stained with May-Grünwald-Giemsa and leukocytes were enumerated on the basis of morphologic criteria (200 cells by light microscopy [$\times 40$]) as previously described (30,36,37). Further, eosinophils, neutrophils, lymphocytes and macrophages were calculated and represented as the percentage of the total leukocytes in BAL fluid.

Airway hyperresponsiveness (AHR)

Mice were anesthetized (ketamine and xylazine [80–100 and 10 mg/kg, respectively]; Troy Laboratories, Smithfield, Australia) and the tracheas were cannulated. Each cannula was connected to an inline aerosol administrator and ventilator, which were attached to a preamplifier and computer (Buxco, Wilmington, NC) to analyze pressure and flow waveforms and to determine airway resistance and dynamic compliance. Mice were nebulized with saline followed by increasing doses of methacholine (Sigma) as previously described (30,32,38).

Statistical analysis

All statistical analyses were performed using GraphPad Prism (Graphpad software, San Diego, USA). Non-parametric Mann–Whitney U test was performed to assess for significant differences in gene expression, staining quantification, and the infiltration of immune cells between different groups. For AHR, two-way analysis of variance (ANOVA) was used to compare different groups and multiple comparisons were done by uncorrected Dunn's test. $P < 0.05$ was considered statistically significant. Outliers were identified by Grubb's test. We detected one outlier in the HDM + ICG-001 group for the *Spdef* gene expression analysis

(Figure 1F) and one outlier in the HDM + ICG-001 group for the AB/PAS staining quantification (Figure 1H) and removed these from the specific analyses. In addition, we also detected and removed one outlier each in the Saline + ICG-001 group for the measurement of total leucocytes, eosinophils, neutrophils, lymphocytes, and macrophages (Figure 2A-E and Figure S1A-E), which are from a single mouse.

Results

ICG-001 decreases HDM-induced goblet cell metaplasia in vivo

Sensitization of mice for 3 days from day 0 to 3 followed by exposure to HDM for 4 days from day 14-17 showed a strong trend ($p=0.06$) towards a decrease in the mRNA expression of E-cadherin (*Cdh1*) compared to PBS-sensitized and PBS-challenged mice. In mice treated with ICG-001, no differences were observed in the *Cdh1* mRNA expression levels after HDM exposure (Figure 1B). HDM significantly increased the mRNA expression of secretory mucins *Muc5ac* and *Muc5b* (Figure 1C, D). Treatment of mice with ICG-001 resulted in a trend ($p=0.09$) towards the suppression of HDM-induced mRNA expression of *Muc5ac*, but had no effect on *Muc5b* expression (Figure 1C, D). The *Muc5ac* transcriptional activator *Spdef* was significantly increased (Figure 1E) and the *Muc5ac* transcriptional repressor *Foxa2* was significantly decreased (Figure 1F) following HDM exposure, which was not affected by ICG-001 treatment (Figure 1E, F). The increase in *Muc5ac* and *Muc5b* expression was validated at the protein level by AB/PAS staining, revealing that HDM exposure promoted goblet cell metaplasia and mucus production. Notably, HDM-induced goblet cell metaplasia and mucus production was significantly repressed by ICG-001 treatment (Figure 1G, H).

ICG-001 inhibits HDM-induced infiltration of macrophages, but does not have an effect on HDM-induced AHR

Airway inflammation with increases in leukocytes and notably eosinophils is a hallmark of the HDM-induced asthma model in mice. HDM significantly increased the total leukocyte counts. The majority of leukocytes in the BAL fluid of PBS exposed mice were macrophages ($98.3 \pm 1.6\%$) and the remaining fraction included eosinophils, neutrophils and lymphocytes (Figure S1A-E). Exposure to HDM significantly increased the total number of eosinophils, neutrophils, lymphocytes, and to a lesser extent, macrophages in BAL fluid (Figure 2A-E), resulting in a decrease in the percentage of macrophages (Figure S1E). ICG-001 did not have




Figure 1: ICG-001 decreases HDM-induced goblet cell metaplasia. (A) HDM-induced allergic airway disease model in mice treated with the ICG-001 inhibitor. (B-F) Gene expression of *Cdh1*, *Muc5ac*, *Muc5b*, *Spdef* and *Foxa2* normalized to *Hprt*. (G) Alcian Blue (AB)/ Periodic acid-Schiff (PAS) staining of lung sections from mice for mucus, scale = 100 μ m. (H) Semi-quantification of AB-PAS staining for mucus using color deconvolution. Data is presented as median \pm IQR; n=4-8, *p<0.05, **p<0.01, Mann-Whitney U test.

any effect on HDM-induced increases in the absolute number of total leukocytes including eosinophils, neutrophils, and lymphocytes (Figure 2A-D). ICG-001 treatment did result in a significant reduction in HDM-induced macrophages (Figure 2E). Methacholine induced a dose-dependent increase in airway resistance, which was significantly elevated by HDM sensitization and challenge (Figure 2F). In line with the lack of effect on inflammation, ICG-001 treatment did not suppress HDM-induced increase in the airway resistance (Figure 2F).




Figure 2: ICG-001 inhibits HDM-induced infiltration of macrophages, but does not have an effect on HDM-induced AHR. Numbers of (A) total leukocytes, (B) eosinophils, (C) neutrophils, (D) lymphocytes, and (E) macrophages present in BAL fluid of mice. Data is presented as median \pm IQR; * p <0.05, ** p <0.01, Mann-Whitney U test. (F) Change in the methacholine-induced airway resistance (%) in mice following exposure to HDM and treated with ICG-001 or vehicle-treated controls. Data is presented as median \pm IQR; n =5-6, * p <0.05, two-way ANOVA.

Discussion

In this study, we investigated the role of β -catenin/CBP signaling in allergen-induced manifestations of asthma. HDM exposure showed a trend towards a decrease in E-cadherin

expression, which was accompanied by increases in *Muc5ac*, *Muc5b* and *Spdef* expression, decreased *Foxa2* expression, the induction of goblet cell metaplasia and the infiltration of inflammatory cells, thereby representing the phenotype of asthmatic airway epithelium. Specific inhibition of β -catenin/CBP signaling using ICG-001 significantly decreased goblet cell metaplasia and mucus production. This effect of ICG-001 on mucus production was in line with a trend towards a decrease in HDM-induced *Muc5ac* expression with ICG-001 treatment, which likely failed to reach statistical significance due to high variation between mice. Our findings are in line with a previous study in which ICG-001 was shown to inhibit goblet cell metaplasia in a toluene diisocyanate treated mouse model of asthma (39). Also, we have recently shown that ICG-001 significantly inhibited IL-13-induced MUC5AC gene expression and production in an air-liquid interface (ALI) culture model of human primary bronchial epithelial cells (Chapter 5), which was accompanied by higher *FOXA2* expression, while leaving *SPDEF* expression unaffected. The discrepancy of the effects of ICG-001 on *MUC5AC* expression between *in vitro* and *in vivo* models could be due to differences in the cellular composition of airway epithelium between human (our *in vitro* data) and mice (this study). Indeed, lineage tracing studies revealed that IL-13-induced goblet cells were derived from FOXJ1-expressing ciliated cells in human primary airway epithelial cells in ALI (40), but this was not demonstrated *in vivo* in an Ovalbumin-induced mouse model of allergic airway disease (41). Newly differentiated goblet cells generated in response to HDM in mice may originate predominantly from club cells that are present in the pseudostratified layer of airway epithelium (42). Also, RNA analysis of lung tissue exhibits greater within-group variance than PAS staining, as the former is dependent on the relative proportion of RNA coming from the airway epithelial cells in the total lung tissue RNA pool. Furthermore, the design of our study may explain the discrepancy between the effect of ICG-001 on HDM-induced *Muc5ac* expression and production. We treated mice with ICG-001 after the initial HDM sensitization and we cannot be sure that the changes in *Muc5ac* gene expression were already induced during the initial sensitization phase.

ICG-001 treatment also had no effect on the gene expression of *Foxa2*, which was previously identified to be regulated by the β -catenin pathway (24). Of note, *Foxa2* is a transcriptional activator of E-cadherin (43,44), and the HDM-induced downregulation of *Foxa2* may have resulted in reduced expression of E-cadherin. In line with the lack of effect on *Foxa2*, we did not observe an effect of ICG-001 on E-cadherin expression. ICG-001 was able to

significantly decrease the HDM-induced infiltration of macrophages, suggesting the important role of β -catenin/CBP signaling in HDM-induced pro-inflammatory responses in mice. Indeed, we have previously shown that ICG-001 inhibits the HDM-induced granulocyte macrophage-colony stimulating factor (GM-CSF) (28), a cytokine that stimulates the production of macrophages. Alternatively, ICG-001 was not able to prevent the infiltration of eosinophils, neutrophils, and lymphocytes. This is in contrast to our *in vitro* data where we showed that ICG-001 reduced HDM-induced CCL20 levels in primary bronchial epithelial cells (28), as neutrophils and T cells (predominantly Th17) are known to express CCL20 receptor CCR6. Moreover, ICG-001 was previously shown to be able to significantly decrease eosinophil and neutrophil influx into the airways induced by toluene diisocyanate in mice (39). These contrasting findings could be due to the difference in the mechanism of action of HDM and toluene diisocyanate. HDM activates pathogen recognition receptors (PRRs) like dectin-1 and Toll-like receptor (TLR)-4 resulting in the release of pro-inflammatory cytokines like CCL17, CCL20, IL-5 and IL-13 (45,46), whereas toluene diisocyanate acts on transient receptor potential melastatin 8 (TRPM8) resulting in the release of IL-25, IL-4 and IL-13 (47). This suggests that TRPM8, but not PRR activation is regulated by β -catenin/CBP signaling. As HDM is a major allergen responsible for airway inflammation in Type-2 driven, atopic asthma, our model may be mainly relevant for better understanding of the role of β -catenin/CBP signaling in allergic airway inflammation.

Here, we show that ICG-001 attenuates HDM-induced goblet cell metaplasia independent of HDM-induced airway inflammation. Inhibition of β -catenin/CBP pathway could be an alternative strategy to regulate mucus hypersecretion in asthma. Future studies should be directed towards more targeted delivery of the ICG-001 to specific airway epithelial cell types for increased efficiency of the drug.

Conflicts of interest

Dr. Knight reports grant from National Health and Medical Research Council (NHMRC #1064405) of Australia. Prof. Hansbro was funded by a Fellowship and grants from the NHMRC of Australia (#1079187, #1175134). Dr Starkey was funded by fellowships from the NHMRC (#1072000) and Australian Research Council (#170100226).

References

1. Fehrenbach H, Wagner C, Wegmann M. Airway remodeling in asthma: what really matters. Vol. 367, *Cell and Tissue Research*. Springer Verlag; 2017. p. 551–69.
2. Heijink IH, Nawijn MC, Hackett T-L. Airway epithelial barrier function regulates the pathogenesis of allergic asthma. *Clin Exp Allergy*. 2014;44(5):620–30.
3. Heijink IH, Kies PM, Kauffman HF, Postma DS, van Oosterhout AJM, Vellenga E. Down-Regulation of E-Cadherin in Human Bronchial Epithelial Cells Leads to Epidermal Growth Factor Receptor-Dependent Th2 Cell-Promoting Activity. *J Immunol*. 2007;178(12):7678–85.
4. Liu G, Cooley MA, Nair PM, Donovan C, Hsu AC, Jarnicki AG, et al. Airway remodelling and inflammation in asthma are dependent on the extracellular matrix protein fibulin-1c. *J Pathol*. 2017 Dec 1;243(4):510–23.
5. Hackett TL. Epithelial-mesenchymal transition in the pathophysiology of airway remodelling in asthma. *Curr Opin Allergy Clin Immunol*. 2012;12(1):53–9.
6. Thorburn AN, Foster PS, Gibson PG, Hansbro PM. Components of *Streptococcus pneumoniae* Suppress Allergic Airways Disease and NKT Cells by Inducing Regulatory T Cells. *J Immunol*. 2012 May 1;188(9):4611–20.
7. Hallstrand TS, Hackett TL, Altemeier WA, Matute-Bello G, Hansbro PM, Knight DA. Airway epithelial regulation of pulmonary immune homeostasis and inflammation. *Clin Immunol*. 2014 Mar 1;151(1):1–15.
8. Hansbro PM, Scott G V, Essilfie A-T, Kim RY, Starkey MR, Nguyen DH, et al. Th2 cytokine antagonists: potential treatments for severe asthma. *Expert Opin Investig Drugs*. 2013 Jan 5;22(1):49–69.
9. Hammad H, Lambrecht BN. Barrier Epithelial Cells and the Control of Type 2 Immunity. *Immunity*. 2015;43(1):29–40.
10. Reid AT, Veerati PC, Gosens R, Bartlett NW, Wark PA, Grainge CL, et al. Persistent induction of goblet cell differentiation in the airways: Therapeutic approaches. *Pharmacol Ther*. 2018 May 1;185:155–69.
11. Starkey MR, McKenzie AN, Belz GT, Hansbro PM. Pulmonary group 2 innate lymphoid cells: surprises and challenges. *Mucosal Immunol*. 2019 Mar 1;12(2):299–311.
12. Perrais M, Pigny P, Copin MC, Aubert JP, Van Seuningen I. Induction of MUC2 and MUC5AC mucins by factors of the epidermal growth factor (EGF) family is mediated by EGF receptor/Ras/Raf/extracellular signal-regulated kinase cascade and Sp1. *J Biol Chem*. 2002 Aug 30;277(35):32258–67.
13. Parker JC, Douglas I, Bell J, Comer D, Bailie K, Skibinski G, et al. Epidermal Growth Factor Removal or Tyrophostin AG1478 Treatment Reduces Goblet Cells & Mucus Secretion of Epithelial Cells from Asthmatic Children Using the Air-Liquid Interface Model. Ahmad S, editor. *PLoS One*. 2015 Jun 9;10(6):e0129546.
14. Williams OW, Sharafkhaneh A, Kim V, Dickey BF, Evans CM. Airway mucus: From production to secretion. *Am J Respir Cell Mol Biol*. 2006 May;34(5):527–36.
15. Bonser LR, Erle DJ. Airway Mucus and Asthma: The Role of MUC5AC and MUC5B. *J Clin Med*. 2017 Nov 29;6(12).
16. Vieira Braga FA, Kar G, Berg M, Carpaij OA, Polanski K, Simon LM, et al. A cellular census of human lungs identifies novel cell states in health and in asthma. *Nat Med*. 2019 Jul 1;25(7):1153–63.
17. Evans CM, Kim K, Tuvim MJ, Dickey BF. Mucus hypersecretion in asthma: Causes and effects. *Curr Opin Pulm Med*. 2009 Jan;15(1):4–11.

18. Hackett T-L, Warner SM, Stefanowicz D, Shaheen F, Pechkovsky D V., Murray LA, et al. Induction of Epithelial–Mesenchymal Transition in Primary Airway Epithelial Cells from Patients with Asthma by Transforming Growth Factor- β 1. *Am J Respir Crit Care Med.* 2009;180(2):122–33.
19. Post S, Nawijn MC, Jonker MR, Kliphuis N, Van Den Berge M, Van Oosterhout AJM, et al. House dust mite-induced calcium signaling instigates epithelial barrier dysfunction and CCL20 production. *Allergy Eur J Allergy Clin Immunol.* 2013;68(9):1117–25.
20. Post S, Nawijn MC, Hackett TL, Baranowska M, Gras R, Van Oosterhout AJM, et al. The composition of house dust mite is critical for mucosal barrier dysfunction and allergic sensitisation. *Thorax.* 2012;67(6):488–95.
21. Ceteci F, Ceteci S, Zanucco E, Thakur C, Becker M, El-Nikhely N, et al. E-cadherin controls bronchiolar progenitor cells and onset of preneoplastic lesions in mice. *Neoplasia (United States).* 2012;14(12):1164–77.
22. Post S, Heijink IH, Hesse L, Koo HK, Shaheen F, Fouadi M, et al. Characterization of a lung epithelium specific E-cadherin knock-out model: Implications for obstructive lung pathology. *Sci Rep.* 2018 Sep 5;8(1):13275.
23. Valenta T, Hausmann G, Basler K. The many faces and functions of β -catenin. *EMBO J.* 2012;31(12):2714–36.
24. Mucenski ML, Nation JM, Thitoff AR, Besnard V, Xu Y, Wert SE, et al. β -Catenin regulates differentiation of respiratory epithelial cells in vivo. *Am J Physiol Cell Mol Physiol.* 2005 Dec;289(6):L971–9.
25. Ma H, Nguyen C, Lee K-S, Kahn M. Differential roles for the coactivators CBP and p300 on TCF/ β -catenin-mediated survivin gene expression. *Oncogene.* 2005;24:3619–31.
26. Moheimani F, Roth HM, Cross J, Reid AT, Shaheen F, Warner SM, et al. Disruption of β -catenin/CBP signaling inhibits human airway epithelial-mesenchymal transition and repair. *Int J Biochem Cell Biol.* 2015;68:59–69.
27. Nawijn MC, Hackett TL, Postma DS, van Oosterhout AJM, Heijink IH. E-cadherin: Gatekeeper of airway mucosa and allergic sensitization. *Trends Immunol.* 2011;32(6):248–55.
28. Kuchibhotla VNS, Jonker MR, Bruin HG, Noordhoek JA, Knight DA, Nawijn MC, et al. Inhibition of β -catenin/CBP signalling improves airway epithelial barrier function and suppresses CCL20 release. *Allergy.* 2020 Feb 24;
29. Thorburn AN, McKenzie CI, Shen S, Stanley D, MacIa L, Mason LJ, et al. Evidence that asthma is a developmental origin disease influenced by maternal diet and bacterial metabolites. *Nat Commun.* 2015 Jun 23;6(1):1–13.
30. Horvat JC, Beagley KW, Wade MA, Preston JA, Hansbro NG, Hickey DK, et al. Neonatal chlamydial infection induces mixed T-cell responses that drive allergic airway disease. *Am J Respir Crit Care Med.* 2007 Sep 15;176(6):556–64.
31. Essilfie AT, Simpson JL, Horvat JC, Preston JA, Dunkley ML, Foster PS, et al. Haemophilus influenzae infection drives IL-17-mediated neutrophilic Allergic airways disease. *PLoS Pathog.* 2011 Oct;7(10):1002244.
32. Kim RY, Horvat JC, Pinkerton JW, Starkey MR, Essilfie AT, Mayall JR, et al. MicroRNA-21 drives severe, steroid-insensitive experimental asthma by amplifying phosphoinositide 3-kinase-mediated suppression of histone deacetylase 2. *J Allergy Clin Immunol.* 2017 Feb 1;139(2):519–32.

33. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc.* 2008 May 5;3(6):1101–8.
34. Starkey MR, Essilfie AT, Horvat JC, Kim RY, Nguyen DH, Beagley KW, et al. Constitutive production of IL-13 promotes early-life Chlamydia respiratory infection and allergic airway disease. *Mucosal Immunol.* 2013 May 7;6(3):569–79.
35. Reid AT, Nichol KS, Veerati PC, Moheimani F, Kicic A, Stick SM, et al. Blocking notch3 signaling abolishes MUC5AC production in airway epithelial cells from individuals with asthma. *Am J Respir Cell Mol Biol.* 2020 Apr 1;62(4):513–23.
36. Kaiko GE, Phipps S, Hickey DK, Lam CE, Hansbro PM, Foster PS, et al. Chlamydia muridarum Infection Subverts Dendritic Cell Function to Promote Th2 Immunity and Airways Hyperreactivity . *J Immunol.* 2008 Feb 15;180(4):2225–32.
37. Thorburn AN, O’Sullivan BJ, Thomas R, Kumar RK, Foster PS, Gibson PG, et al. Pneumococcal conjugate vaccine-induced regulatory T cells suppress the development of allergic airways disease. *Thorax.* 2010 Dec 1;65(12):1053–60.
38. Kim RY, Pinkerton JW, Essilfie AT, Robertson AAB, Baines KJ, Brown AC, et al. Role for NLRP3 inflammasome-mediated, IL-1 β -dependent responses in severe, steroid-resistant asthma. *Am J Respir Crit Care Med.* 2017 Aug 1;196(3):283–97.
39. Yao L, Zhao H, Tang H, Xiong J, Zhao W, Liu L, et al. Blockade of β -catenin signaling attenuates toluene diisocyanate-induced experimental asthma. *Allergy.* 2017 Apr 1;72(4):579–89.
40. Turner J, Roger J, Fitau J, Combe D, Giddings J, Van Heeke G, et al. Goblet cells are derived from a FOXJ1-expressing progenitor in a human airway epithelium. *Am J Respir Cell Mol Biol.* 2011 Mar 1;44(3):276–84.
41. Pardo-Saganta A, Law BM, Gonzalez-Celeiro M, Vinarsky V, Rajagopal J. Ciliated cells of pseudostratified airway epithelium do not become mucous cells after ovalbumin challenge. *Am J Respir Cell Mol Biol.* 2013 Mar;48(3):364–73.
42. Park KS, Korfhagen TR, Bruno MD, Kitzmiller JA, Wan H, Wert SE, et al. SPDEF regulates goblet cell hyperplasia in the airway epithelium. *J Clin Invest.* 2007 Apr 2;117(4):978–88.
43. Song Y, Washington MK, Crawford HC. Loss of FOXA1/2 is essential for the epithelial-to-mesenchymal transition in pancreatic cancer. *Cancer Res.* 2010 Mar 1;70(5):2115–25.
44. Zhang Z, Yang C, Gao W, Chen T, Qian T, Hu J, et al. FOXA2 attenuates the epithelial to mesenchymal transition by regulating the transcription of E-cadherin and ZEB2 in human breast cancer. *Cancer Lett.* 2015 Jun 1;361(2):240–50.
45. Gregory LG, Lloyd CM. Orchestrating house dust mite-associated allergy in the lung. *Trends Immunol.* 2011 Sep;32(9):402–11.
46. Post S, Heijink IH, Petersen AH, De Bruin HG, Van Oosterhout AJM, Nawijn MC. Protease-activated receptor-2 activation contributes to house dust mite-induced IgE responses in mice. *PLoS One.* 2014;9(3):1–9.
47. Kim JH, Jang YS, Jang SH, Jung KS, Kim SH, Ye YM, et al. Toluene diisocyanate exposure induces airway inflammation of bronchial epithelial cells via the activation of transient receptor potential melastatin 8. *Exp Mol Med.* 2017 Mar 3;49(3):e299.

Supplementary Data




Figure S1: Effect of ICG-001 on HDM-induced airway inflammation and AHR in mice. Numbers of (A) total leukocytes, and percentage of (B) eosinophils, (C) neutrophils, (D) lymphocytes, and (E) macrophages present in BAL fluid of mice. Data is presented as median \pm IQR; n=5-6, **p<0.01, Mann-Whitney U test.

