

University of Groningen

**Erratum: Effect of Nonelastic Compression With an Adjustable Wrap After Total Knee Arthroplasty (vol 39, pg 377, 2020)**

Hendrickx, A. A.; Krijnen, W. P.; Bimmel, R.; van der Schans, C. P.; Damstra, R. J.

*Published in:*  
Orthopaedic nursing

*DOI:*  
[10.1097/NOR.0000000000000736](https://doi.org/10.1097/NOR.0000000000000736)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Hendrickx, A. A., Krijnen, W. P., Bimmel, R., van der Schans, C. P., & Damstra, R. J. (2021). Erratum: Effect of Nonelastic Compression With an Adjustable Wrap After Total Knee Arthroplasty (vol 39, pg 377, 2020). *Orthopaedic nursing*, 40(1), 32-32. <https://doi.org/10.1097/NOR.0000000000000736>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

- from the perspective of people living with rheumatoid arthritis. *Arthritis Care & Research*, 66(11), 1662–1672. <https://doi.org/10.1002/acr.22374>
- Phillips, S., & Dow, L. (2012). Impact of impaired morning function on quality of life in rheumatoid arthritis: Results of an exploratory patient survey. *International Journal of Clinical Rheumatology*, 7, 597–606. <https://doi.org/10.2217/IJR.12.62>
- Rhind, V. M., Unsworth, A., & Haslock, I. (1987). Assessment of stiffness in rheumatology: The use of rating scales. *Rheumatology*, 26(2), 126–130. <https://doi.org/10.1093/rheumatology/26.2.126>
- Sacks, J. J., Luo, Y. H., & Helmick, C. G. (2010). Prevalence of specific types of arthritis and other rheumatic conditions in the ambulatory health care system in the United States, 2001–2005. *Arthritis Care & Research*, 62, 460–464. <https://doi.org/10.1002/acr.20041>
- Schlaeger, J. M., Roach, K., Golas, M., Takayama, M., & Wilkie, D. J. (2018). Treatment-seeking behaviors of persons with rheumatoid arthritis. *Journal of Holistic Nursing*, 36(2), 179–191. <https://doi.org/10.1177/0898010117693539>
- Sierakowski, S., & Cutolo, M. (2011). Morning symptoms in rheumatoid arthritis: A defining characteristic and marker of active disease. *Scandinavian Journal of Rheumatology, Supplement*, 40(125), 1–5. <https://doi.org/10.3109/03009742.2011.566433>
- Soubrier, M., Zerkak, D., Gossec, L., Ayral, X., Roux, C., & Dougados, M. (2006). Which variables best predict change in rheumatoid arthritis therapy in daily clinical practice? *The Journal of Rheumatology*, 33, 1243–1246.
- Spiro, R. L., & Weitz, B. A. (1990). Adaptive selling conceptualization, measurement, and nomological validity. *Journal of Marketing Research*, 27, 61–69. <https://doi.org/10.1177/0022243790002700106>
- Straub, R. H. & Cutolo, M. (2007). Circadian rhythms in rheumatoid arthritis: Implications for pathophysiology and therapeutic management. *Arthritis and Rheumatism*, 56, 399–408. <https://doi.org/10.1002/art.22368>
- Uhlig, T., Haavardsholm, E. A., & Kvien, T. K. (2006). Comparison of the Health Assessment Questionnaire (HAQ) and the modified HAQ (MHAQ) in patients with rheumatoid arthritis. *Rheumatology*, 45, 454–458. <https://doi.org/10.1093/rheumatology/kei181>
- Uhm, D. C., Nam, E. S., Lee, H. Y., Lee, E. B., Yoon, Y. I., & Chai, G. J. (2012). Health-related quality of life in Korean patients with rheumatoid arthritis: Association with pain, disease activity, disability in activities of daily living and depression. *Journal of Korean Academy of Nursing*, 42(3), 434–442. <https://doi.org/10.4040/jkan.2012.42.3.434>
- van Tyul, L. H. D., Lems, W. F., & Boers, M. (2014). Measurement of stiffness in patients with rheumatoid arthritis in low disease activity or remission: A systematic review. *BMC Musculoskeletal Disorders*, 15(1), 28. <https://doi.org/10.1186/1471-2474-15-28>
- Westhoff, G., Buttgereit, F., Gromnica-Ihle, E., & Zink, A. (2008). Morning stiffness and its influence on early retirement in patients with recent onset rheumatoid arthritis. *Rheumatology*, 47, 980–984. <https://doi.org/10.1093/rheumatology/ken137>
- Yazici, Y., Pincus, T., Kautiainen, H., & Sokka, T. (2004). Morning stiffness in patients with early rheumatoid arthritis is associated more strongly with functional disability than with joint swelling and erythrocyte sedimentation rate. *The Journal of Rheumatology*, 31, 1723–1726.
- Young, A., Dixey, J., Kulinskaya, E., Cox, N., Davies, P., Devlin, J., Emery, P., Gough, A., James, D., Prouse, P., Williams, P., & Winfield, J. (2002). Which patients stop working because of rheumatoid arthritis? Results of five years' follow up in 732 patients from the Early RA Study (ERAS). *Annals of the Rheumatic Diseases*, 61, 335–340. <https://doi.org/10.1136/ard.61.4.335>

## ERRATUM

### Effect of Nonelastic Compression With an Adjustable Wrap After Total Knee Arthroplasty: Erratum.

The lead author of an article that appeared in the November/December 2020 issue of *Orthopaedic Nursing* reported an error in Table 3, page 381, and apologizes for this error. In the first table column, the outcome measurements are described (Total leg volume, Circumference infrapatellar, Circumference midpatellar, and Circumference suprapatellar) for the control group and the treatment group. In the last two outcome measurements (midpatellar and suprapatellar) the word “control” is used for both groups, whereas it should have read “treatment group” for the second row. The outcomes are correct. Please see a corrected version of this table at the following link: <http://links.lww.com/ONJ/A16>.

### REFERENCE

Hendrickx, A. A., Krijnen, W. P., Bimmel, R., van der Schans, C. P., & Damstra, R. J. (2020). Effect of nonelastic compression with an adjustable wrap after total knee arthroplasty. *Orthopaedic Nursing*, 39(6), 377–383.

DOI: 10.1097/NOR.0000000000000736