

University of Groningen

## Characterization of rare germline variants in familial multiple myeloma

Catalano, Calogerina; Paramasivam, Nagarajan; Blocka, Joanna; Giangioffe, Sara; Huhn, Stefanie; Schlesner, Matthias; Weinhold, Niels; Sijmons, Rolf; de Jong, Mirjam; Langer, Christian

*Published in:*  
Blood cancer journal

*DOI:*  
[10.1038/s41408-021-00422-6](https://doi.org/10.1038/s41408-021-00422-6)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Catalano, C., Paramasivam, N., Blocka, J., Giangioffe, S., Huhn, S., Schlesner, M., Weinhold, N., Sijmons, R., de Jong, M., Langer, C., Preuss, K-D., Nilsson, B., Durie, B., Goldschmidt, H., Bandapalli, O. R., Hemminki, K., & Foersti, A. (2021). Characterization of rare germline variants in familial multiple myeloma. *Blood cancer journal*, 11(2), [33]. <https://doi.org/10.1038/s41408-021-00422-6>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CORRESPONDENCE

Open Access

# Characterization of rare germline variants in familial multiple myeloma

Calogerina Catalano<sup>1,2</sup>, Nagarajan Paramasivam<sup>3</sup>, Joanna Blocka<sup>2</sup>, Sara Giangioffe<sup>1,2</sup>, Stefanie Huhn<sup>2,4</sup>, Matthias Schlesner<sup>5</sup>, Niels Weinhold<sup>2</sup>, Rolf Sijmons<sup>6</sup>, Mirjam de Jong<sup>6</sup>, Christian Langer<sup>7</sup>, Klaus-Dieter Preuss<sup>8</sup>, Björn Nilsson<sup>9</sup>, Brian Durie<sup>10</sup>, Hartmut Goldschmidt<sup>2,4</sup>, Obul Reddy Bandapalli<sup>1,11,12</sup>, Kari Hemminki<sup>1,13,14</sup> and Asta Försti<sup>1,11,12</sup>

Dear Editor,

Multiple myeloma (MM) is a malignancy of plasma cells, characterized by the presence of monoclonal immunoglobulin, known as M protein<sup>1</sup>. MM is preceded by monoclonal gammopathy of undetermined significance (MGUS) which is also a precursor of immunoglobulin light chain (AL) amyloidosis<sup>1</sup>. Previous studies have reported a 2- to 4-fold increased risk of MGUS or MM in first-degree relatives of MM or MGUS patients, suggesting the existence of inherited susceptibility<sup>2,3</sup>. For many years, high-risk germline predisposing genes have been lacking for MM. However, recent sequencing efforts have proposed a few novel candidates, most notably loss-of-function (LoF) variants in the tumor suppressor gene *DIS3* and in the histone demethylase gene *KDM1A*<sup>4–6</sup>, and others as recently reviewed in detail in Pertesi et al.<sup>7</sup>. In addition to the suspected rare, high-penetrance variants, genome-wide association studies have identified over 20 common, low-penetrance variants associated with the risk of MM; these were estimated to account for about 15% of the familial MM risk<sup>8</sup>.

As the genetic basis of most MM families remains unexplained, our study aimed at identifying germline predisposition genes in familial MM from Germany, Sweden, and the Netherlands, through whole genome and

exome sequencing<sup>9</sup>. Altogether, 21 families with 46 affected and 20 unaffected family members were recruited (Supplementary Information, Supplementary Fig. 1). Each family had at least two individuals diagnosed with MM or its precursors MGUS and smoldering MM (SMM). After sequencing, detailed bioinformatics analyses using our in-house developed Familial Cancer Variant Prioritization Pipeline version-2 (FCVPPv2) were conducted to prioritize the most likely candidates (Fig. 1 and Supplementary Methods). Gradual filtering of variants after sequencing is shown for each family in Supplementary Table 1. The functions of the gene products were collected from the UniProtKB database (<https://www.uniprot.org/>) and literature search. They are summarized in Fig. 2 and details are shown in Supplementary Information.

We identified 109 potential pathogenic missense variants; in most families, several candidates were found, and in four families none (Supplementary Table 2). All variants were private for each family, except for genes *KIF1B* and *DCSH1*, in which two different missense variants were found in two unrelated families (families 10 and 18 for *KIF1B* and 15 and 17 for *DCSH1*). *KIF1B* is involved in the transport of mitochondria and synaptic vesicles and *DCSH1* is a calcium-dependent cell adhesion protein. Among the other genes harboring missense variants, tumor suppressor function was indicated for *DAB2IP* and oncogene function for *ABL2*. The former had diverse signal transduction functions and it is implicated in immune processes, as are *TLN1*, *ZFAT*, *CLCF1*, *IL11RA*, *SEC14L1*, *SAMHD1*, *DCST1*, *TPP2*, and *MYO1G*.

Another group of genes with key regulatory functions constituted *FOXO1*, *B4GALT1*, and *NKX3-2*. Transcription factor *FOXO1* is a protein, which is the main target

Correspondence: Asta Försti ([a.foersti@kitz-heidelberg.de](mailto:a.foersti@kitz-heidelberg.de))

<sup>1</sup>Division of Molecular Genetic Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

<sup>2</sup>Department of Internal Medicine V, University of Heidelberg, Heidelberg, Germany


Full list of author information is available at the end of the article  
These authors contributed equally: Calogerina Catalano, Nagarajan Paramasivam.

These authors jointly supervised this work: Obul Reddy Bandapalli, Kari Hemminki and Asta Försti.

© The Author(s) 2021



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.




of insulin signaling, it increases osteoblast numbers and regulates B cell development. B4GALT1 is involved in the glycosylation of immunoglobulin G (IgG) and variants in this gene have been associated with IgG levels and hematological neoplasms, including MM. NKX3-2 (homeobox protein Nkx-3.2) is a member of the HOX gene transcription factors family, which are frequently dysregulated in hematologic malignancies.

Our candidate list included two genes, *KMT2A* and *USP28*, functionally related to the recently reported MM predisposing genes, *LSD1/KDM1A*, encoding a lysine-specific demethylase, and *USP45*, an apoptosis-related gene-regulating DNA repair<sup>5,6</sup>. *KMT2A* (alias MLL1) is a histone H3 lysine 4 (H3K4) methyltransferase, which plays an essential role in early development and hematopoiesis and which mediates chromatin modifications associated with epigenetic transcriptional activation. *USP28* is a deubiquitinase involved in DNA damage-induced apoptosis. It regulates MYC protein stability in response to DNA damage.

We checked our gene list also for the presence of the 82 somatically mutated driver genes in MM, described in Walker et al.<sup>10</sup> and Maura et al.<sup>11</sup>, but only *SAMHD1* passed all our in-house pipeline filters. *SAMHD1* is a somatic driver in MM and the protein plays a role in maintaining dNTP levels in regulating DNA replication and damage repair. It enhances immunoglobulin hypermutation in B-lymphocyte development.

We also identified 36 loss-of-function (LoF) variants in the MM families (Supplementary Table 3). If we would apply a MutPred-LOF (<http://mutpredlof.cs.indiana.edu/index.html>) score higher than 0.50 at a 5% false-positive rate, only two frameshift variants, in the genes *SLC30A5* and *LONP2*, and six stop codon variants would pass the threshold. None of these had an apparent relationship to MM. Of the eight splice site variants, five were predicted by Human Splicing Finder (<http://www.umd.be/HSF/HSF.shtml>) to alter the splicing motifs (indicated by “yes” in Supplementary Table 3), however with no link to MM. Many of the genes with LoF mutations encode



proteins with housekeeping functions, including LONP2, CSGALNACT2, HMGCLL1, and FUK.

We identified seven copy-number variants (CNVs) that segregated with MM in the families (Supplementary Table 4). These CNVs affected the coding regions of 11 genes. Duplication of chr4:15936942-16178663 in Family 5 covered the genes encoding fibroblast growth factor binding proteins *FGFBP1* and *FGFBP2*, prominin 1 (*PROM1*), and transmembrane anterior posterior transformation protein 1 homolog (*TAPT1*). One of the primary genetic events in MM is t(4:14) translocation, creating a fusion between the immunoglobulin heavy chain (*IGH*) enhancer and *FGFR3* and leading to overexpression of *FGFR3*<sup>12</sup>. *FGFBP1* and *FGFBP2* encode proteins that are involved in FGF ligand bioactivation by releasing them from the extracellular matrix. Thus, duplication of these two genes may lead to activation of the FGF signaling, enhanced MM cell proliferation and survival, and affect bone homeostasis. *PROM1* is involved in the suppression of cell differentiation and maintenance of stem cell properties.

All the identified variants were rare (allele frequency < 0.001) in the gnomAD database (<https://gnomad.broadinstitute.org>) and none of them was found as a germline variant in any cancer patient. However, COSMIC (<https://cancer.sanger.ac.uk/cosmic>) reported some of the variants as rare somatic mutations, mainly in cancer entities with a high mutation load, such as malignant melanoma and adenocarcinoma of the large intestine, but not in any hematological malignancies.

In a review of cancer-predisposing genes, it was observed that over 40% of germline variants were in genes that functioned also as somatic drivers<sup>13</sup>. In the above, we referred to some somatic drivers, and some of the observed genes are known to interact with key signaling pathways in MM, including PI3K/Akt/mTOR, Ras/Raf/MEK/MAPK, JAK/STAT, NF- $\kappa$ B, Wnt/ $\beta$ -catenin, and RANK/RANKL/OPG<sup>14</sup>. Among the relevant genes in our list, *DAB2IP*, encoding a Ras-GTPase activating protein, modulates key oncogenic pathways such as PI3K/Akt, NF- $\kappa$ B, and Wnt/ $\beta$ -catenin; *FOXO1* encodes for a downstream effector of Akt signaling; the *LRP1B* gene product negatively regulates the Wnt/ $\beta$ -catenin/TCF signaling, through its interaction with DVL2.

A somewhat surprising finding was that none of the 158 candidate genes matched with the genes linked to the 23 common, low-risk MM variants<sup>8</sup>. It is true that an overwhelming number of the published MM-related low-risk variants were located in the non-protein-coding region and some were distant from the coding regions<sup>8</sup>. The associated relative risks were mostly below 1.5, which would not be compatible with strong familial clustering. However, as many of the present families included only two affected members, chance clustering cannot be excluded. A further factor is that by the inclusion of persons with MGUS, which is more than one order of magnitude more prevalent than MM, certain genetic heterogeneity was introduced, in spite of the known shared genetic background<sup>3</sup>. Of note, the candidates that

passed the pipeline included two genes, *KMT2A* and *USP28*, functionally related to the recently proposed high/moderate penetrance MM predisposing genes, *LSD1/KDM1A*, *USP45*, *ARID1A*, and *DIS3*<sup>4–6</sup>.

In few families, no candidate variants were found using the present criteria. Some possible reasons were explained in the above paragraph, and there are more possible reasons. In families of two affected individuals, polygenic risks would be more likely than in multiplex families of many affected individuals. In a previous study, evidence of enrichment of the common MM risk alleles among familial cases compared to sporadic cases or population-based controls was reported<sup>15</sup>. Our search did not consider polygenic risk. Even though the controls were tested for the presence of the M protein, a negative result suggests that the person would remain disease free only the next decade or two<sup>9</sup>. Finally, the present bioinformatics analysis was limited to coding variants.

In conclusion, we report here curated sequencing results from 21 MM/MGUS families. While most of the 154 presented candidate genes are unlikely to have a causal relationship to MM, the identified genes could be a valuable contribution to forthcoming, pooled sequencing efforts. Familial clustering of MM is rare and the set of 21 families was only possible through multicenter efforts. Based on the functional characterization and the literature review the strong candidates included *DAB2IP*, *ABL2*, *SAMHD1*, *KMT2A*, *USP28*, *FOXO1*, *B4GALT1*, *NKX3-2*, several immune-related genes, and *FGFBP1*, *FGFBP2*, and *PROM1* within the CNV in chromosome 4. Interestingly, many of these are somatic driver genes in cancer.

#### Acknowledgements

We thank the Genomics and Proteomics Core Facility (GPCF) and the Omics IT and Data Management Core Facility (ODCF) of the German Cancer Research Center (DKFZ) for excellent technical support. This work was supported by the Black Swan Research Initiative/International Myeloma Foundation, Dietmar Hopp Foundation, and Transcan ERA-NET funding from the German Federal Ministry of Education and Research (BMBF). K.H. was supported by the European Union's Horizon 2020 research and innovation program, Grant No 856620. A.F. was supported by the German Jose Carreras Leukemia Foundation.

#### Author details

<sup>1</sup>Division of Molecular Genetic Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. <sup>2</sup>Department of Internal Medicine V, University of Heidelberg, Heidelberg, Germany. <sup>3</sup>Computational Oncology, Molecular Diagnostics Program, National Center for Tumor Diseases (NCT), Heidelberg, Germany. <sup>4</sup>National Center for Tumor Diseases Heidelberg (NCT), Heidelberg, Germany. <sup>5</sup>Bioinformatics and Omics Data Analytics, German Cancer Research Center (DKFZ), Heidelberg, Germany. <sup>6</sup>University Medical Center Groningen, University of Groningen, Groningen, The Netherlands. <sup>7</sup>Kempton Clinic, Kempton, Germany. <sup>8</sup>José Carreras Center for Immuno and Gene Therapy,

Department of Internal Medicine I, Saarland University Medical School, Homburg (Saar), Germany. <sup>9</sup>Hematology and Transfusion Medicine, Department of Laboratory Medicine, Lund University, Lund, Sweden. <sup>10</sup>Cedars Sinai Cancer Center, Los Angeles, CA, USA. <sup>11</sup>Hopp Children's Cancer Center (KiTZ), Heidelberg, Germany. <sup>12</sup>Division of Pediatric Neurooncology, German Cancer Research Center (DKFZ), German Cancer Consortium (DKTK), Heidelberg, Germany. <sup>13</sup>Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. <sup>14</sup>Faculty of Medicine and Biomedical Center in Pilsen, Charles University in Prague, Pilsen, Czech Republic

#### Data availability

Whole-genome sequencing data sets have been deposited to the European Genomephenome Archive (EGA) with accession number EGAS00001004734.

#### Conflict of interest

The authors declare no competing interests.

#### Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s41408-021-00422-6>.

Received: 2 October 2020 Revised: 4 January 2021 Accepted: 18 January 2021

Published online: 13 February 2021

#### References

- Kumar, S. K. et al. Multiple myeloma. *Nat. Rev. Dis. Primers* **3**, 17046 (2017).
- Frank, C. et al. Search for familial clustering of multiple myeloma with any cancer. *Leukemia* **30**, 627–632 (2016).
- Vachon, C. M. et al. Increased risk of monoclonal gammopathy in first-degree relatives of patients with multiple myeloma or monoclonal gammopathy of undetermined significance. *Blood* **114**, 785–790 (2009).
- Pertesi, M. et al. Exome sequencing identifies germline variants in *DIS3* in familial multiple myeloma. *Leukemia* **33**, 2324–2330 (2019).
- Waller, R. G. et al. Novel pedigree analysis implicates DNA repair and chromatin remodeling in multiple myeloma risk. *PLoS Genet.* **14**, e1007111 (2018).
- Wei, X. et al. Germline lysine-specific demethylase 1 (*LSD1/KDM1A*) mutations confer susceptibility to multiple myeloma. *Cancer Res.* **78**, 2747–2759 (2018).
- Pertesi, M. et al. Genetic predisposition for multiple myeloma. *Leukemia* **34**, 697–708 (2020).
- Went, M. et al. Identification of multiple risk loci and regulatory mechanisms influencing susceptibility to multiple myeloma. *Nat. Commun.* **9**, 3707 (2018).
- Blocka, J. et al. Familial cancer: how to successfully recruit families for germline mutations studies? Multiple myeloma as an example. *Clin. Lymphoma Myeloma Leuk.* **19**, 635–644 e632 (2019).
- Walker, B. A. et al. Identification of novel mutational drivers reveals oncogene dependencies in multiple myeloma. *Blood* **132**, 587–597 (2018).
- Maura, F. et al. Genomic landscape and chronological reconstruction of driver events in multiple myeloma. *Nat. Commun.* **10**, 3835 (2019).
- Foltz, S. M. et al. Evolution and structure of clinically relevant gene fusions in multiple myeloma. *Nat. Commun.* **11**, 2666 (2020).
- Rahman, N. Realizing the promise of cancer predisposition genes. *Nature* **505**, 302–308 (2014).
- Hu, J. et al. Targeting signaling pathways in multiple myeloma: Pathogenesis and implication for treatments. *Cancer Lett.* **414**, 214–221 (2018).
- Halvarsson, B. M. et al. Direct evidence for a polygenic etiology in familial multiple myeloma. *Blood Adv.* **1**, 619–623 (2017).