

University of Groningen

Immune evasion in primary testicular and central nervous system lymphomas

Minderman, Marthe; Amir, Avital; Kraan, Willem; Schilder-Tol, Esther J. M.; Oud, Monique E. C. M.; Scheepstra, Cornelis G.; Noorduyn, Arnold L.; Kluin, Philip M.; Kersten, Marie Jose; Spaargaren, Marcel

Published in:
Blood

DOI:
[10.1182/blood.2021011366](https://doi.org/10.1182/blood.2021011366)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Minderman, M., Amir, A., Kraan, W., Schilder-Tol, E. J. M., Oud, M. E. C. M., Scheepstra, C. G., Noorduyn, A. L., Kluin, P. M., Kersten, M. J., Spaargaren, M., & Pals, S. T. (2021). Immune evasion in primary testicular and central nervous system lymphomas: HLA loss rather than 9p24.1/PD-L1/PD-L2 alterations. *Blood*, 138(13), 1194-1197. <https://doi.org/10.1182/blood.2021011366>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

TO THE EDITOR:

Immune evasion in primary testicular and central nervous system lymphomas: HLA loss rather than 9p24.1/*PD-L1*/*PD-L2* alterations

Marthe Minderman,^{1,2} Avital Amir,^{1,2} Willem Kraan,^{1,2} Esther J. M. Schilder-Tol,^{1,2} Monique E. C. M. Oud,^{1,2} Cornelis G. Scheepstra,³ Arnold L. Noorduin,⁴ Philip M. Kluin,⁵ Marie José Kersten,^{2,6} Marcel Spaargaren,^{1,2,*} and Steven T. Pals^{1,2,*}

¹Department of Pathology Cancer Center Amsterdam (CCA), Academic Medical Center (AMC), Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands; ²Lymphoma and Myeloma Center Amsterdam–LYMMCARE, The Netherlands; ³Department of Pathology and Hematology, Onze Lieve Vrouwe Gasthuis, Amsterdam, The Netherlands; ⁴Pathology Laboratory, Dordrecht, The Netherlands; ⁵Department of Pathology, University Medical Center, Groningen, The Netherlands; and ⁶Department of Hematology, CCA, AMC, Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands

Cancer immune evasion is a major hurdle for effective anticancer therapy. Key strategies of tumor cells to avoid T-cell recognition include downregulation of major histocompatibility complex (MHC) molecules and activation of immune checkpoints.^{1,2} Particularly, the immune checkpoint protein programmed cell death 1 (PD-1) and its ligands, PD-L1 and -L2, play a key role in repressing T-cell activity in the tumor microenvironment, not only in solid cancers but also in hematological malignancies. Specifically in advanced classic Hodgkin lymphoma (cHL), blockade of the PD-1/PD-L1/L2 axis yields outstanding clinical responses.³ Most cHLs are infiltrated by PD-1⁺ T cells, and the tumor cells show strong expression of PD-L1/PD-L2 related to copy-number alterations (CNAs) at chromosome 9p24.1, containing the loci for *PD-L1/PD-L2* and *JAK2*,⁴ or alternatively, driven by Epstein-Barr virus (EBV) infection.⁵

Primary testicular lymphomas (PTLs) and primary central nervous system lymphomas (PCNSLs) are uncommon and aggressive large-B cell lymphomas with a poor response to therapies and prognosis and shared molecular characteristics. PTLs and PCNSLs that arise at sites considered to be immune privileged⁶ display a high prevalence of activating somatic *MYD88* mutations, often with a concurrent activating mutation in the immunoreceptor tyrosine-based activation motif of *CD79B*.^{7,8} Regarding immune evasion, PTLs and PCNSLs have been shown to exhibit frequent loss of HLA class I and II expression and/or loss of HLA loci.^{9,10} More recently, Chapuy et al reported frequent 9p24.1/*PD-L1/2* CNAs and translocations with concomitant protein overexpression in PTLs and PCNSLs.¹¹ A subsequent study by the same authors in a small series (n = 5) of patients with a PTL or PCNSL suggested clinical activity of PD-1 blockade with nivolumab.¹² Based on these reports, several clinical trials exploring the efficacy of PD-1/PD-L1 blockade have been initiated (registered on www.clinicaltrials.gov as NCT02779101, NCT03255018, NCT04401774, and NCT02857426). In the current study, we revisited the immune evasion mechanisms operating in PTLs and PCNSLs. We confirmed a high frequency of HLA class I and II expression loss. However, with the exception of EBV⁺ PCNSLs, which were PD-L1⁺, PTLs and PCNSLs seldom expressed PD-L1 and, accordingly, 9p24.1/*PD-L1/2* CNAs were rarely found.

We assessed HLA class I and II expression, PD-L1 expression, and 9p24.1/*PD-L1/2* CNAs in a panel of lymphomas diagnosed as PTLs (n = 25) or PCNSLs (n = 22) according to the World Health Organization classification,¹³ using immunohistochemistry and fluorescence in situ hybridization (FISH), respectively (Table 1). For comparison, a set of diffuse large B-cell lymphomas (DLBCLs; n = 22), enriched for cases with a non-germinal center B-cell (GCB) phenotype to better match the PTLs and PCNSLs, were studied. All lymphomas were classified as either GCB- or non-GCB-like, using the immunohistochemical algorithm of Hans et al.¹⁴ In addition, the presence of somatic mutations in *MYD88* and *CD79B*; of translocations of *cMYC*, *BCL2*, and *BCL6*; and of EBV status were assessed, as described previously.^{7,8} Consistent with previous reports,^{7,8,11} *MYD88* mutations were detected at high frequency in PTLs (76%) and PCNSLs (64%), and coexistent mutations in *CD79B* were often found. The presence of these *MYD88* and *CD79* mutations was almost mutually exclusive with translocations of *cMYC*, *BCL2*, and *BCL6* or with EBV expression (Figure 1A). Apropos of immune evasion, we observed a high prevalence of loss of HLA class I expression in PTLs (68%) and PCNSLs (64%). Similarly, HLA class II loss was common in PTLs (84%) and PCNSLs (59%). These findings are in line with previous studies by Riemersma et al¹⁰ and Booman et al,¹⁵ who also reported frequent HLA class I and II loss in PTLs and PCNSLs. In marked contrast to the study by Chapuy et al,¹¹ however, we barely detected expression of PD-L1 or 9p24.1/*PD-L1/2* CNAs in our cohort of PTLs and PCNSLs. A notable exception were EBV⁺ tumors, which showed strong PD-L1 expression (Figure 1A). In the entire group of EBV⁻ PCNSLs and PTLs (n = 42), only 3 cases with PD-L1 expression and 1 case with a 9p24.1/*PD-L1/2* copy number gain were found. The frequency of PD-L1 expression and 9p24.1/*PD-L1/2* CNA did not significantly differ from that in the DLBCLs. In approximately half of the cases, we observed variable, mostly weak, PD-L1 expression by macrophages in the tumor microenvironment. Representative PD-L1/PAX5 immunohistochemistry images are shown in supplemental Figure 1. All tissue samples were obtained during standard diagnostic procedures at the Academic Medical Center Amsterdam and affiliated hospitals and the University Medical Center

Groningen in accordance with the local institutional board requirements.

In view of the vast discrepancy between our finding and those of Chapuy et al,¹¹ we sought to further reinforce our data. First, to validate the techniques used to detect PD-L1 expression and 9p24.1/PD-L1/2 CNAs, a series of primary mediastinal B-cell lymphomas (PMBCLs; n = 25) were studied. Like cHLs, these lymphomas frequently show genetic aberrations in chromosome 9p24.1 with consequent PD-L1 overexpression.⁴ Indeed, we detected strong PD-L1 expression in a large proportion (68%) of the PMBCLs (supplemental Table 1), whereas 9p24.1/PD-L1/2 CNAs were observed in more than half of the cases (Figure 1B). Notably, in contrast to PTLs and PCNSLs, loss of HLA class I and II expression was uncommon in these PMBCLs (supplemental Table 1). Second, we studied *PD-L1* messenger RNA (mRNA) expression data extracted from publicly available databases. These mRNA data reveal enhanced expression of *PD-L1* in PMBCLs, but not in PTLs and PCNSLs, a finding consistent with the lack of PD-L1 expression in our current data set (Figure 1C). Notably, in line with our results, in a study of PCNSLs recently reported by Sethi et al,¹⁶ PD-L1 expression and/or 9p24.1/PD-L1/2 CNAs was barely found, except in EBV⁺ tumors, which were PD-L1⁺. A possible explanation for the discordantly high prevalence of 9p24.1/PD-

L1/2 CN gains reported by Chapuy et al¹¹ is an incorrectly low threshold setting in the quantitative polymerase chain reaction assays used to detect CNAs in the large "extension cohort" of their study. This error in threshold setting could also explain the strikingly lower percentage of CNAs in the initial "discovery cohort," which were identified by using high-density single nucleotide polymorphism arrays.

In summary, our study indicates that different large B-cell lymphoma subtypes use distinct immune evasion strategies: PTLs and PCNSLs seem to use the PD1/PD-L1/2 checkpoint rarely, but instead use the loss of MHC expression as a major mechanism of immune evasion; in PMBCLs PD-L1 overexpression, often caused by 9p24.1/PD-L1/2 CNAs, appears to play a major role. This finding suggests that patients with PMBCL, but not those with PTL or PCNSL, are likely to benefit from treatment with PD-1/PD-L1 immune checkpoint inhibitors. Indeed, although a preliminary clinical study suggested beneficial activity of nivolumab treatment,¹² the results of the phase 2 trial of nivolumab in patients with recurrent or refractory PCNSL or PTL (registered on www.clinicaltrials.gov as NCT02857426), which was initiated based on these initial findings, are highly disappointing: the objective response rate (ORR) was only 6.4% in 47 analyzed patients. Similarly, nivolumab monotherapy resulted in an ORR of only

Table 1. Clinical and molecular characteristics of PTLs, PCNSLs and DLBCLs

		PTL (n = 25)	PCNSL (n = 22)	DLBCL (n = 22)
Age, y	Median (range)	74 (58-89)	69 (49-83)	63 (4-89)
Gender	Male	100 (25/25)	67 (14/21)	64 (14/22)
	Female	0 (0/25)	33 (7/21)	36 (8/22)
EBV status	Positive	0 (0/25)	15 (3/20)	9 (2/22)
MYC	Rearranged	8 (2/25)	0 (0/15)	5 (1/22)
BCL2	Rearranged	0 (0/25)	6 (1/17)	0 (0/22)
BCL6	Rearranged	25 (6/24)	19 (3/16)	32 (7/22)
MYD88	Mutated	76 (19/25)	64 (14/22)	32 (7/22)**
CD79B	Mutated	20 (5/25)	23 (5/22)	14 (3/22)
COO	Non-GCB	76 (19/25)	95 (21/22)	86 (19/22)
	GCB	24 (6/25)	5 (1/22)	14 (3/22)
PD-L1 tumor	Negative	92 (23/25)	82 (18/22)	73 (16/22)
	Weak	0 (0/25)	4 (1/22)	9 (2/22)
	Positive	8 (2/25)	14 (3/22)	18 (4/22)
PD-L1/2 CNA	Gain	4 (1/23)	0 (0/13)	14 (3/22)
	Polysomy	0 (0/23)	0 (0/13)	0 (0/22)
HLA class I	Loss	68 (17/25)	64 (14/22)	32 (7/22)*
HLA class II	Loss	84 (21/25)	59 (13/22)	14 (3/22)**

Unless stated otherwise, data are expressed as percentage of cases that were positive (positive cases/total cases examined). The correlation between the clinical and molecular characteristics among the different tumor types was examined using the χ^2 test. P-values were 2-sided and $P < .05$ was considered statistically significant. * $P < .05$; ** $P < .01$. COO, cell of origin.

3% to 10% in patients with relapsed/refractory DLBCL.¹⁷ In contrast, interim analysis of the phase 2 KEYNOTE-170 study (www.clinicaltrials.gov, NCT02576990) showed an ORR of 45% in patients with relapsed/refractory PBMCL.¹⁸ Considering the ongoing clinical trials involving immune checkpoint inhibitors in patients with PCNSL or PTL (www.clinicaltrials.gov,

NCT04421560, NCT03770416, and NCT04609046), we believe that a critical reappraisal of the prevalence of PD-L1 expression and 9p24.1 alterations in these lymphomas is of crucial importance.




Figure 1. Oncoprint plot. Shown is the molecular analysis of the PTLs, PCNSLs, and DLBCLs (A) and PMBCLs (B). All tissue samples were obtained during standard diagnostic procedures. Immunohistochemistry was performed on formalin-fixed, paraffin-embedded tissue using anti-HLA class I (clone HC10, Nordic-MUbio), anti-HLA-DP,DQ,DR (clone CR3/43, DAKO), anti-PD-L1 (clone 22C3, DAKO), anti-PAX5 (clone SP43, Cell Marque), anti-CD10 (clone 56C6, ThermoFisher), anti-MUM1 (clone MUM1p, DAKO), anti-BCL2 (clone 124, Dako), and anti-BCL6 (clone PG-B6p, Dako) on a Labvision Autostainer 480S (ThermoFisher). Samples were scored positive for PD-L1 when membranous staining was observed in at least 5% of the malignant cells. Expression of EBV was determined by EBV-encoded RNA in-situ hybridization (EBER) probes (Biogenex). FISH for detection of *BCL2*, *BCL6*, and *cMYC* breaks was performed using probes and a FISH accessory kit (Dako). FISH for detection of *PD-L1/2* CNAs was performed with the ZytoLight CD274/PDCD1LG2/CEN 9 Dual Color Probe (ZytoVision). FISH slides were evaluated in the context of serial sections stained for PD-L1 and B-cell markers (CD20 and PAX5) to localize tumor infiltrates. Samples were scored as having 9p24.1 disomy, polysomy, copy gain, or amplification. The presence of 3 or 4 green signals was classified as gain and the presence of 5 or more green signals was classified as amplification. Testing for somatic *MYD88* and *CD79B* mutations was performed with allele-specific polymerase chain reaction, as described previously.⁷ Sanger sequencing was used to verify the presence of a mutation. (C) *PD-L1* mRNA expression analysis of publicly available microarray data sets derived from the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (accession numbers GSE10524, GSE61578, GSE34771 and GSE87371). All microarray data sets were generated with Affymetrix Human Genome U133 Plus 2.0 Array, and data analysis was performed with the R2: Genomics Analysis and Visualization Platform (<http://r2.amc.nl>). The horizontal line represents the median expression within each group. Differences among subtypes were tested by Kruskal-Wallis test with the post hoc Dunn's test. **P* < .05.

Acknowledgments

The authors thank M.F.M. de Rooij for helpful contributions to the data presentation.

This study was supported by a grant from Lymph&Co (M.J.K., S.T.P.).

Authorship

Contribution: M.M. performed the experiments, analyzed the data, and wrote the manuscript; A.A. designed the research, supervised the study, and analyzed the data; W.K. performed the experiments and analyzed the data; E.J.M.S.-T. and M.E.C.M.O. performed the experiments; C.G.S., A.L.N., and P.M.K. provided samples from patients with lymphoma; M.J.K., M.S., and S.T.P. designed the research, supervised the study, analyzed the data, and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profile: W.K., 0000-0003-1109-931X.

Correspondence: Steven T. Pals, Department of Pathology, Amsterdam University Medical Centers, loc. AMC, University of Amsterdam, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands; e-mail: s.t.pals@amc.uva.nl.

Footnotes

Submitted 18 February 2021; accepted 7 June 2021; prepublished online on *Blood* First Edition 14 June 2021.

*M.S. and S.T.P. are joint senior authors.

Original data are available in response to e-mail requests to Steven T. Pals (s.t.pals@amc.uva.nl).

The online version of this article contains a data supplement.

REFERENCES

1. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer*. 2012;12(4):252-264.
2. Mahoney KM, Rennert PD, Freeman GJ. Combination cancer immunotherapy and new immunomodulatory targets. *Nat Rev Drug Discov*. 2015;14(8):561-584.
3. Ansell SM, Lesokhin AM, Borrello I, et al. PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. *N Engl J Med*. 2015; 372(4):311-319.
4. Green MR, Monti S, Rodig SJ, et al. Integrative analysis reveals selective 9p24.1 amplification, increased PD-1 ligand expression, and further induction via JAK2 in nodular sclerosing Hodgkin lymphoma and primary mediastinal large B-cell lymphoma. *Blood*. 2010;116(17):3268-3277.

5. Chen BJ, Chapuy B, Ouyang J, et al. PD-L1 expression is characteristic of a subset of aggressive B-cell lymphomas and virus-associated malignancies. *Clin Cancer Res*. 2013;19(13):3462-3473.
6. Shechter R, London A, Schwartz M. Orchestrated leukocyte recruitment to immune-privileged sites: absolute barriers versus educational gates. *Nat Rev Immunol*. 2013;13(3):206-218.
7. Kraan W, Horlings HM, van Keimpema M, et al. High prevalence of oncogenic MYD88 and CD79B mutations in diffuse large B-cell lymphomas presenting at immune-privileged sites. *Blood Cancer J*. 2013;3(9): e139.
8. Kraan W, van Keimpema M, Horlings HM, et al. High prevalence of oncogenic MYD88 and CD79B mutations in primary testicular diffuse large B-cell lymphoma. *Leukemia*. 2014;28(3):719-720.
9. Booman M, Szuhai K, Rosenwald A, et al. Genomic alterations and gene expression in primary diffuse large B-cell lymphomas of immune-privileged sites: the importance of apoptosis and immunomodulatory pathways. *J Pathol*. 2008;216(2):209-217.
10. Riemersma SA, Jordanova ES, Schop RF, et al. Extensive genetic alterations of the HLA region, including homozygous deletions of HLA class II genes in B-cell lymphomas arising in immune-privileged sites. *Blood*. 2000;96(10):3569-3577.
11. Chapuy B, Roemer MG, Stewart C, et al. Targetable genetic features of primary testicular and primary central nervous system lymphomas. *Blood*. 2016;127(7):869-881.
12. Nayak L, Iwamoto FM, LaCasce A, et al. PD-1 blockade with nivolumab in relapsed/refractory primary central nervous system and testicular lymphoma. *Blood*. 2017;129(23):3071-3073.
13. Swerdlow SH, Campo E, Harris NL, et al, eds. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. 4th rev. ed. Geneva, Switzerland: World Health Organisation; 2016.
14. Hans CP, Weisenburger DD, Greiner TC, et al. Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood*. 2004;103(1):275-282.
15. Booman M, Douwes J, Glas AM, et al. Mechanisms and effects of loss of human leukocyte antigen class II expression in immune-privileged site-associated B-cell lymphoma. *Clin Cancer Res*. 2006;12(9):2698-2705.
16. Sethi TK, Kovach AE, Grover NS, et al. Clinicopathologic correlates of MYD88 L265P mutation and programmed cell death (PD-1) pathway in primary central nervous system lymphoma. *Leuk Lymphoma*. 2019; 60(12):2880-2889.
17. Ansell SM, Minnema MC, Johnson P, et al. Nivolumab for relapsed/refractory diffuse large B-cell lymphoma in patients ineligible for or having failed autologous transplantation: a single-arm, phase II study. *J Clin Oncol*. 2019;37(6):481-489.
18. Armand P, Rodig S, Melnichenko V, et al. Pembrolizumab in relapsed or refractory primary mediastinal large B-cell lymphoma. *J Clin Oncol*. 2019; 37(34):3291-3299.

DOI 10.1182/blood.2021011366

© 2021 by The American Society of Hematology