

University of Groningen

Hidden hunger in the hospital?

van Vliet, Iris

DOI:
[10.33612/diss.216397465](https://doi.org/10.33612/diss.216397465)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
van Vliet, I. (2022). *Hidden hunger in the hospital? Recognition of malnutrition and malnutrition risk in complex hospital care*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen. <https://doi.org/10.33612/diss.216397465>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CHAPTER 5

MALNUTRITION IN STABLE RENAL TRANSPLANT
RECIPIENTS ACCORDING TO GLIM CRITERIA:
REDUCED MUSCLE MASS AS PREDOMINANT
PHENOTYPE



KARIN BOSLOOPER-MEULENBELT* • IRIS M.Y. VAN VLIET* • ANTONIO W. GOMES-
NETO* • MARGRIET F.C. DE JONG • STEPHAN J.L. BAKKER • HARRIËT JAGER-
WITTENAAR • GERJAN J. NAVIS

published in Clin Nutr, 2021; 40(5): 3522-3530

** shared first authorship*

ABSTRACT

Background & aims Malnutrition has a negative impact on quality of life and survival in renal transplant recipients (RTR). Therefore, malnutrition detection is important in RTR, but this may be hampered by concomitant presence of weight gain and overweight. Recently, the Global Leadership Initiative on Malnutrition (GLIM) developed a set of diagnostic criteria for malnutrition. We aimed to assess the prevalence of malnutrition according to the GLIM criteria and the distribution of phenotypic criteria in RTR. Additionally, we examined the potential value of 24-h urinary creatinine excretion rate (CER) as alternative measure for the criterion reduced muscle mass.

Methods We used data from stable outpatient RTR included in the TransplantLines Cohort and Biobank Study (NCT02811835). Presence of weight loss and reduced intake or assimilation were derived from Patient-Generated Subjective Global Assessment (PG-SGA) item scores. Reduced muscle mass was assessed by multi-frequency bio-electrical impedance analysis (MF-BIA) and defined as an appendicular skeletal muscle mass index (ASMI) $< 7 \text{ kg/m}^2$ for men and $< 5.5 \text{ kg/m}^2$ for women, and in additional analysis defined as creatinine-height index (CHI, based on 24 h urine CER) $< 80\%$. Inflammation was present if C-reactive protein (CRP) was $> 5 \text{ mg/L}$. Malnutrition was defined as presence of at least one phenotypic (weight loss and/or low BMI and/or reduced muscle mass) and one etiologic criterion (reduced intake/assimilation and/or disease burden/inflammation).

Results We included 599 RTR (55 ± 13 years old, 62% male, BMI $27.2 \pm 4.7 \text{ kg/m}^2$) at a median of 3.1 years after transplantation. According to GLIM criteria, 14% was malnourished, of which 91% met the phenotypic criterion for reduced muscle mass. Similar results were found by using CHI as measure for muscle mass (13% malnutrition of which 79% with reduced muscle mass).

Conclusions Malnutrition is present in one in 7 stable RTR, with reduced muscle mass as the predominant phenotypic criterion. Assessment of nutritional status, most importantly muscle status, is warranted in routine care, to prevent malnutrition in RTR from remaining undetected and untreated. The diagnostic value of 24-h urinary CER in this regard requires further investigation.

Abbreviations

GLIM	Global Leadership Initiative on Malnutrition
QoL	quality of life
RTR	renal transplant recipients
MUST	Malnutrition Universal Screening Tool
DEXA	dual energy X-ray absorptiometry
BIA	bio-electrical impedance analysis
CT	computed tomography
MRI	magnetic resonance imaging
CER	creatinine excretion rate
PG-SGA	Patient-Generated Subjective Global Assessment
eGFR	estimated glomerular filtration rate
TSAT	transferrin saturation
CRP	C-reactive protein
ASM	appendicular skeletal muscle mass
ASMI	appendicular skeletal muscle mass index
CHI	creatinine-height index
SGA	Subjective Global Assessment
MIS	Malnutrition-Inflammation score

INTRODUCTION

Renal transplantation is considered the treatment of choice for patients with end-stage renal disease, resulting in a better quality of life (QoL) and survival compared with dialysis treatment¹⁻³. However, QoL is still lower and long-term outcomes are worse in renal transplant recipients (RTR) compared with the general population^{3,4}. Post-transplant weight gain and obesity are well-known and common health problems, related to transplant-specific factors such as cessation of dietary restrictions^{5,6}, side effects of prednisolone⁵ and low physical activity levels⁷, and are associated with adverse long-term outcomes in RTR⁸⁻¹². However, less attention is paid to the, possibly concomitant, high risk of malnutrition. Several transplant-related factors increase the malnutrition risk in RTR, including the use of immunosuppressive medications, infections, rejection of the graft, insulin resistance and the immune response to the graft¹³⁻¹⁵. Although RTR is an underrepresented population in the scientific literature on malnutrition in chronic kidney disease (CKD)¹⁶, previous research has shown that malnutrition also has a considerable negative impact on QoL and survival in RTR^{17,18}. Therefore, malnutrition should not be overlooked in this patient population.

Recently, the Global Leadership Initiative on Malnutrition (GLIM) published updated diagnostic consensus criteria for malnutrition¹⁹. According to the GLIM consensus, the diagnosis of malnutrition requires the presence of at least one phenotypic and one etiologic criterion. The phenotypic criteria include either weight loss, low Body Mass Index (BMI), or reduced muscle mass, and the etiologic criteria include either reduced food intake or assimilation, or disease burden/inflammation (**Table 1**). In RTR, the high prevalence of post-transplantation weight gain and obesity^{8,11} is accompanied by an altered body composition with an increased fat mass²⁰ and lower muscle mass²¹⁻²³. Therefore, using low BMI and weight loss alone may lack sensitivity to diagnose malnutrition in RTR, as excess fat mass may obscure underlying depletion of muscle mass. To our knowledge, however, the prevalence of malnutrition according to the GLIM criteria, and the relative contribution of each phenotypic criterion to the diagnosis of malnutrition, has not yet been investigated in this population.

Insight in the distribution of the different malnutrition phenotypes in RTR is important, as this may have implications for the assessment of malnutrition in clinical practice. Most validated malnutrition screening instruments, which are often used as a first step in the detection of malnutrition, include screening for low BMI and weight loss, but do not include screening for low muscle mass^{24,25}. Therefore, malnourished RTR with reduced muscle mass as phenotypic criterion may currently remain undetected in this consecutive two-step process of screening and assessment. Furthermore, while standard anthropometric measurements, such as weight and height, are routinely performed at outpatient visits, the assessment of muscle mass, for example with dual energy x-ray absorptiometry (DEXA) or bio-electrical impedance analysis (BIA) as recommended by GLIM, is usually not part of routine care due to practical and time constraints²⁶. In several renal transplant care centers, however, 24-h urine samples are routinely collected during outpatient visits, enabling measurement of 24-h urinary creatinine excretion rate (CER), which has shown

to be a reliable marker of muscle mass in RTR^{21,22}. Therefore, it is worthwhile to explore its potential value for the assessment of reduced muscle mass in diagnosing malnutrition in this population.

The aim of this study was to investigate the prevalence of malnutrition in stable outpatient RTR using the GLIM criteria and to examine the relative contribution of each phenotypic criterion, i.e., weight loss, low BMI and reduced muscle mass, to the diagnosis of malnutrition. Additionally, the prevalence of malnutrition and relative contribution of the phenotypic criterion low muscle mass with use of 24-h CER was examined.

METHODS

Study design and population

For this cross-sectional analysis of data from a cohort of RTR, data was extracted from the TransplantLines Biobank of the University Medical Center Groningen (UMCG) (ClinicalTrials.gov Identifier: NCT03272841). Rationale and design of this cohort study has been described previously²⁷. In brief, since June 2015, all eligible transplant candidates and transplant recipients were invited to participate in the TransplantLines study. Written informed consent was obtained before inclusion. The Medical Ethical Committee of the UMCG approved the TransplantLines study protocol (METc 2014/077) and all study procedures were performed in line with the principles of the Declaration of Helsinki. For the current study, we included adult RTR (≥ 18 years) enrolled in the TransplantLines study, with a functioning graft ≥ 1 year after transplantation and a scheduled study visit between June 2015 and December 2019. Participants with missing data in the variables that are used in the operationalization of the GLIM criteria for malnutrition ($N = 339$) were excluded for analyses.

Data collection

During the study visits, anthropometric measures and assessment of nutritional status were performed by trained student researchers. Height, weight, waist and hip circumference were assessed using a wall-secured stadiometer, a digital scale, and a retractable measurement tape, respectively. Body composition was assessed using a multi-frequency BIA device (Quadscan 4000, Bodystat, Douglas, British Isles). The Dutch version of the Patient-Generated Subjective Global Assessment (PG-SGA version 3.7 NL, as available on <http://www.pt-global.org/>) was used to assess nutritional status^{28,29}. The PG-SGA is a validated nutrition assessment tool and available in multiple language versions, based on translation and cultural adaptation processes using the International Society for Pharmacoeconomics and Outcomes Research's (ISPOR's) "Principles of Good Practice for the Translation and Cultural Adaptation Process for PRO Measures"^{30,31}. The first part of the PG-SGA (PG-SGA Short Form, SF) was completed by the patient and included items on weight history (Box 1), food intake (Box 2), nutrition impact symptoms (Box 3), activities and function (Box 4).

Prior to each TransplantLines study visit, fasted blood samples and 24-h urine samples were collected and analyzed by using standard laboratory procedures. Incomplete 24-h urine samples ($N = 53$), due to missed urine portions, were excluded for the data analysis. The serum creatinine-based Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) algorithm was used to calculate the estimated glomerular filtration rate (eGFR)³². Creatinine clearance in mL/min was calculated by dividing 24-h urinary creatinine excretion (mmol/24 h) by serum creatinine (mmol/L), multiplied by 694 to account for the translation from mL/24-h to mL/minutes. Second, creatinine clearance in mL/min/1.73 m² was calculated by multiplying creatinine clearance by 1.73 divided by the body surface area (BSA) according to the formula of du Bois³³. Proteinuria was defined as a urinary protein excretion 0.5 g/24 h. Other laboratory measures included for analysis in this study were hemoglobin, ferritin, transferrin saturation (TSAT), C-reactive protein (CRP), and albumin from blood samples, and urea and creatinine excretion from 24-h urine samples. Protein intake in grams per

day (g/d) was calculated from 24-h urinary urea excretion using the Maroni-equation³⁴. For protein intake in grams per kilogram per day (g/kg/d), in subjects with underweight (BMI < 20 kg/m²) and obesity (BMI >30 kg/m²) the body weight corresponding to a BMI of 20 and 27.5 kg/m², respectively was used^{35,36}.

Demographic variables and data on disease history, including the primary renal disease, dialysis treatment, medication and transplant characteristics, were extracted from the UMCG Renal Transplant Database. Other lifestyle parameters, including smoking behavior and alcohol consumption, were based on self-reported validated questionnaires administered during or prior to a TransplantLines study visit²⁷.

Diagnosis of malnutrition using the GLIM criteria

Criteria for the diagnosis of malnutrition according to GLIM criteria and operationalization of variables in the current study based on these criteria are shown in **Table 1**. Malnutrition diagnosis was based on presence of at least one phenotypic criterion (weight loss, low BMI, and/or reduced muscle mass) in combination with at least one etiologic criterion (reduced food intake or assimilation and/or disease burden/inflammation) as indicated by GLIM consensus¹⁹.

Weight loss. The phenotypic criterion for weight loss was defined as weight loss of > 5% within past 6 months and based on the self-reported weights in Box 1 of the PG-SGA.

Low BMI. BMI was calculated by dividing measured weight in kilograms by measured height in squared meter. For patients with Caucasian, African or unknown ethnicity, the phenotypic criterion for low BMI was defined as a BMI < 20 kg/m² in patients < 70 year old and BMI < 22 kg/m² in patients ≥ 70 year old. For patients with Asian ethnicity, the phenotypic criterion for low BMI was defined as a BMI < 18.5 kg/m² in patients < 70 year old and BMI < 20 kg/m² in patients ≥ 70 year old.

Reduced muscle mass. BIA measurements were used to assess muscle mass. Using resistance and reactance measured by BIA, appendicular muscle mass (ASM) was calculated using the formula by Sergi et al.³⁷. Appendicular muscle mass index (ASMI, kg/m²) was calculated by dividing ASM by height in squared meter. The phenotypic criterion for reduced muscle mass was defined as an ASMI < 7 kg/m² for men and < 5.5 kg/m² for women according to GLIM recommendations^{19,38}.

Reduced food intake or assimilation. Scores on PG-SGA Box 2 and Box 3 were used to assess food intake and assimilation. The etiologic criterion for reduced food intake was defined as either 1) PG-SGA Box 2 score ≥ 1 , indicating food intake less than usual over the past month, or 2) PG-SGA Box 3 score ≥ 1 , indicating presence of 1 or multiple nutrition impact symptoms (including gastro-intestinal symptoms, such as dysphagia, nausea, vomiting and diarrhea).

Disease burden/inflammation. As recommended by the GLIM Working Group, CRP was used as a supportive measure for the inflammation criterion^{19,39} for which we used a cut-

off of > 5 mg/L⁴⁰. However, as all RTR are considered to have chronic renal disease with low grade inflammation⁴¹ and therefore inherently meet the disease burden/ inflammation criterion as stated in the GLIM consensus¹⁹, we also assessed the prevalence of malnutrition in all RTR.

Table 1. Operationalization of the phenotypic and etiologic criteria for the diagnosis of malnutrition in the current study

	Phenotypic criteria			Etiologic criteria	
	Weight loss	Low BMI	Reduced muscle mass	Reduced food intake or assimilation	Disease burden/ inflammation
Current study	> 5% within the past 6 months based on self-reported weights in PG-SGA Box 1	<20 kg/m ² if age <70 years, or <22 kg/m ² if age ≥70 years Asia: <18.5 kg/m ² if age <70 years, or <20 kg/m ² if age ≥70 years	ASMI <7 kg/m ² (men) and <5.5 kg/m ² (women) Additional analysis: CHI <80%	PG-SGA Box 2 score ≥1 (reduced food intake in the last month) PG-SGA Box 3 score ≥1 (≥1 nutrition impact symptom)	1. CRP >5 mg/L 2. All RTR based on chronic disease burden

Abbreviations: ASMI: appendicular skeletal muscle mass index; BMI: Body Mass Index; CHI: creatinine-height index; CRP: C-reactive protein; RTR: renal transplant recipients; PG-SGA: Patient Generated Subjective Global Assessment

Use of 24-h CER in assessment of muscle mass

For the additional analysis using 24-h CER for the operationalization of the reduced muscle mass criterion, the creatinine-height index (CHI) as described by Blackburn et al.⁴² was used: CHI (%) measured total urinary creatinine excretion (mg)/(ideal total urinary creatinine excretion for patient of the same sex and height)* 100. Reduced muscle mass was defined as a CHI < 80%^{42, 43}.

Statistical analyses

Statistical analyses were performed using IBM SPSS Statistics for Windows, Version 23.0. Descriptive statistics are presented as means ± standard deviations (SD) for normally distributed variables, median [interquartile range] for non-normally distributed variables, and number (percentage) for categorical variables. The study population was then divided in two groups according to the GLIM malnutrition diagnosis to show differences in baseline characteristics between well-nourished and malnourished RTR. Differences in patient characteristics between the two groups (well- nourished versus malnourished)

were analyzed with the independent T-test for normally distributed continuous data, the Mann-Whitney U-test for non-normally distributed data, and the chi-square test for nominal data. A P-value < 0.05 (two-tailed) was considered as statistically significant. The BioVenn web application⁴⁴ was used to produce a Venn diagram, to visualize (overlap between) phenotypic criteria for the diagnosis of malnutrition.

RESULTS

Patient characteristics

A total of 599 RTR were included, with a mean age of 55 ± 13 years, of which 372 (62%) were male. The most common primary renal disease was cystic kidney disease or other congenital diseases in 151 RTR (25%), followed by glomerulonephritis in 146 RTR (24%), of which 2 patients underwent a re-transplantation because of recurrent glomerulonephritis. Median time after transplantation was 3.1 [1.0-10.0] years. In total, 237 (40%) of the patients received pre-emptive transplantation and 360 (60%) received dialysis treatment prior transplantation. Of all RTR 334 (56%) received a transplant from a living donor irrespective of prior dialysis treatment. Other patient characteristics and parameters are shown in **Table 2**.

Prevalence of malnutrition and distribution of phenotypic and etiologic GLIM criteria

Based on the presence of at least one phenotypic criterion and at least one etiologic criterion, 85 (14%) RTR were malnourished (**Table 3**). Of these malnourished patients, 77 (91%) met at least the phenotypic criterion for reduced muscle mass, and all but one (99%) met either the phenotypic criterion for reduced muscle mass or weight loss or both. Although mean BMI was lower in RTR with malnutrition ($p < 0.001$), 22% of malnourished RTR were classified as overweight or obese, compared with 72% of well-nourished RTR (**Table 2**).

Regarding the phenotypic criteria, 166 (28%) met at least one phenotypic criterion for the diagnosis of malnutrition (**Table 3**). Of these patients, 92% (153/166) met the criterion for reduced muscle mass, 9% (15/166) the phenotypic criterion for weight loss and 11% (19/166) the phenotypic criterion for low BMI. Most of the patients (17/19) that met the criterion for low BMI also met the criterion for reduced muscle mass (**Table 3** and **Figure 1**).

When considering all RTR as meeting the etiologic criterion for disease burden/inflammation based on presence of chronic disease, 166 (28%) were malnourished, of which 153 (92%) met the phenotypic criterion for reduced muscle mass, and all but two (99%) met either the phenotypic criterion for reduced muscle mass or weight loss or both.

Regarding the etiologic criteria, 295 (49%) met at least one etiologic criterion for the diagnosis of malnutrition, of which 226 (38%) patients met the etiologic criterion for reduced food intake or assimilation, and 119 (40%) met the etiologic criterion for disease burden/inflammation based on CRP.

Table 2. Characteristics of total population and across well-nourished and malnourished RTR following the GLIM-criteria

	Total	Well-nourished	Malnourished	P-value
N (%)	599 (100)	514 (86)	85 (14)	-
Sociodemographic characteristics				
Age (years)	55 ± 13	55 ± 13	55 ± 15	0.96
Male gender	372 (62)	313 (61)	59 (69)	0.13
Ethnicity				0.30
Caucasian	418 (70)	357 (69)	61 (72)	
African	6 (1)	4 (1)	2 (2)	
Asian	8 (1)	8 (2)	0 (0)	
Other	8 (1)	6 (1)	2 (2)	
Unknown	159 (27)	139 (27)	20 (24)	
Education level				0.03
Low	207 (34)	181 (35)	26 (30)	
Medium	189 (32)	168 (33)	21 (25)	
High	154 (26)	121 (23)	33 (39)	
Unknown	49 (8)	44 (9)	5 (6)	
Renal and transplant characteristics				
Primary renal disease				0.88
Glomerulonephritis	146 (24)	127 (25)	19 (22)	
Interstitial nephritis	51 (9)	42 (8)	9 (11)	
Cystic kidney disease and other congenital/hereditary	151 (25)	128 (25)	23 (27)	
Renal vascular disease, diabetes mellitus and other multisystem disease	129 (22)	113 (22)	16 (19)	
Other /unknown	122 (20)	104 (20)	18 (21)	
Time after transplantation (years)	3.1 [1.0 – 10.0]	3.1 [1.0 – 10.0]	2.5 [1.0 – 10.5]	0.95
Type of transplantation				0.11
Pre-emptive transplantation	237 (40)	197 (38)	40 (48)	
Dialysis prior transplantation	360 (60)	316 (62)	44 (52)	
Missing	2 (0)	1 (0)	1 (0)	
Type of donor				0.24
Living donor	334 (56)	282 (55)	52 (62)	
Deceased-donor donor	263 (44)	231 (45)	32 (38)	
Missing	2 (0)	1 (0)	1 (0)	
Immunosuppressive drugs				
Tacrolimus	402 (67)	342 (67)	60 (71)	0.48
Ciclosporin	93 (16)	82 (16)	11 (13)	0.47
Mycophenolic acid	459 (77)	397 (77)	62 (73)	0.37
Azathioprine	49 (8)	39 (8)	10 (12)	0.20
Prednisolone	584 (98)	502 (98)	82 (97)	0.43
eGFR (ml/min/1.73m ²)	51 ± 18	51 ± 17	56 ± 19	0.008
Creatinine clearance [§]				
in mL/min	70 ± 25	71 ± 26	66 ± 20	0.12
in ml/min/1.73m ²	62 ± 22	62 ± 22	62 ± 20	0.95

Proteinuria [§]	77 (13)	73 (14)	4 (4)	0.01
Hematological and inflammation parameters				
Hemoglobin (mmol/L)	8.4 ± 1.1	8.4 ± 1.1	8.4 ± 1.1	0.89
Ferritin (mmol/L)	88 [41 – 175]	89 [41 – 176]	84 [50 – 166]	0.88
TSAT (%)	24 ± 10	24 ± 10	23 ± 10	0.35
CRP (mg/L)	1.9 [0.7 – 4.6]	1.7 [0.7 – 4.3]	2.7 [0.8 – 7.0]	0.03
Albumin (g/L)	43.7 ± 2.9	43.6 ± 2.9	43.8 ± 3.2	0.75
Anthropometry and body composition				
Height (cm)	174 ± 9	174 ± 9	175 ± 10	0.40
BMI (kg/m ²)	27.2 ± 4.7	27.9 ± 4.6	23.4 ± 2.8	<0.001
BMI category				<0.001
Underweight	15 (3)	7 (1)	8 (9)	
Normal weight	195 (32)	136 (27)	59 (69)	
Overweight	230 (38)	215 (42)	15 (18)	
Obesity	159 (27)	156 (30)	3 (4)	
Waist circumference (cm)				
Male	102 ± 13	103 ± 13	94 ± 11	<0.001
Female	94 ± 14	96 ± 13	81 ± 9	<0.001
ASMI (kg/m ²)				
Male	7.46 ± 0.97	7.46 ± 0.93	6.50 ± 0.50	<0.001
Female	6.35 ± 0.86	6.47 ± 0.82	5.43 ± 0.60	<0.001
24 hour urinary creatinine excretion (mmol/24h) [§]				
Male	14.0 ± 3.8	14.3 ± 3.8	12.2 ± 3.4	<0.001
Female	10.4 ± 2.9	10.6 ± 2.9	9.1 ± 2.2	0.03
Nutritional intake[§]				
Protein intake*				
in g/kg/d**	1.08 ± 0.26	1.08 ± 0.26	1.07 ± 0.26	0.89
in g/d	84 ± 22	85 ± 22	76 ± 18	0.001
24-hour urinary sodium excretion rate (mmol/24h)	144 ± 59	147 ± 59	126 ± 55	0.005
24-hour urinary potassium excretion rate (mmol/24h)	72 ± 24	73 ± 24	67 ± 23	0.05
Other lifestyle parameters				
Smoking status				0.17
Current smoker	47 (8)	36 (7)	11 (13)	
Non-(current) smoker	450 (75)	390 (76)	60 (70)	
Unknown	102 (17)	88 (17)	14 (17)	
Alcohol consumption				0.27
Yes	385 (64)	336 (65)	49 (58)	
No	60 (10)	48 (9)	12 (14)	
Unknown	154 (26)	130 (26)	24 (28)	

Note: Data are presented as mean ± SD, number (%) or median [IQR].

*Protein intake was calculated from 24-h urinary urea excretion using the Maroni-equation 34. **Protein intake in g/kg/d with adjustment for underweight and obesity: a BMI <20 kg/m² is adjusted to a BMI of 20 kg/m² and a BMI >30 kg/m² is adjusted to a BMI of 27.5 kg/m². § Due to incomplete 24-hour urine samples or missing data the number (N) available for analysis for creatinine clearance

Association between malnutrition according to GLIM and clinical and nutritional parameters RTR that were classified as malnourished according to GLIM criteria had a higher education level ($p = 0.03$), and had proteinuria less often ($p = 0.01$). Whereas eGFR was significantly higher in malnourished patients ($p = 0.008$), creatinine clearance was not different for malnourished and well-nourished RTR. Regarding nutritional intake, a lower sodium intake (126 ± 55 mmol/24 h versus 147 ± 59 mmol/24 h, $p = 0.005$), potassium intake (73 ± 24 mmol/24 h versus 67 ± 23 mmol/24 h, $p = 0.05$) and total protein intake (76 ± 18 g/d versus 85 ± 22 g/d, $p = 0.001$), but not protein intake in g/kg/d, was observed in RTR with malnutrition compared with those without malnutrition (Table 2).

CHI as measure for reduced muscle mass

Data on CHI were available in 526 patients, with a mean age of 55 ± 13 years, of which 322 (61%) were male. No significant differences were found between characteristics of patients with and without CHI available with regard to age, sex, ethnicity, education level, primary renal disease, type of transplantation, renal function, BMI, smoking behavior, or alcohol consumption.

Mean CHI was 106 ± 29 , and 99 patients (19%) had a CHI < 80% and therefore met the criterion for reduced muscle mass based on the operationalization with CHI. Using CHI as a criterion for reduced muscle mass in the GLIM framework, 67 (13%) patients were classified as malnourished, of which 53 (79%) met the phenotypic criterion for reduced muscle mass, and 61 (91%) met either the phenotypic criterion for reduced muscle mass or weight loss or both. When considering all RTR as meeting the etiologic criterion for disease burden/inflammation based on presence of chronic disease, 121 (23%) patients were malnourished, of which 99 (82%) met the phenotypic criterion for reduced muscle mass and 110 (91%) met either the phenotypic criterion for reduced muscle mass or weight loss or both.

Table 3. GLIM criteria for malnutrition in RTR

	RTR N = 599		
	N	% within group	% of total
Phenotypic criterion present	166	-	28
1. Weight loss	15	9	3
2. Low BMI	19	11	3
3. Reduced muscle mass	153	92	26
<i>Combinations of phenotypic criteria present</i>			
Weight loss + low BMI	0	0	0
Weight loss + reduced muscle mass	3	2	1
Low BMI + reduced muscle mass	16	10	3
Weight loss + low BMI + reduced muscle mass	1	1	0
Etiologic criterion present	295	-	49
a. Reduced food intake/assimilation	226	77	38
Reduced food intake	180	61	30
Reduced assimilation	79	27	13
b. Inflammation, CRP>5 mg/L	119	40	20
<i>Combination of etiologic criteria present</i>			
Reduced food intake/assimilation + inflammation	50	17	8
Malnutrition according to GLIM criteria (at least 1 phenotypic + 1 etiologic criterion present)	85	-	14
1. Weight loss + reduced food intake/assimilation or inflammation	10	12	2
2. Low BMI + reduced food intake/assimilation or inflammation	11	13	2
3. Reduced muscle mass + reduced food intake/assimilation or inflammation	77	91	13

Abbreviations: BMI: Body Mass Index; CRP: C-reactive protein; GLIM: Global Leadership Initiative on Malnutrition; RTR: renal transplant recipients. * Number of patients, % within group of patients that either meets at least one phenotypic criterion, meets at least one etiologic criterion or is classified as malnourished, and % of total is shown.




Figure 1. Co-existence of phenotypic criteria for malnutrition in all RTR ⁴⁴

Abbreviations: BMI: Body Mass Index; RTR: renal transplant recipients. * Number of patients that meet either the phenotypic criterion for weight loss, low BMI, reduced muscle mass or combinations of those criteria, is shown.

DISCUSSION

In the present study, we assessed the prevalence of malnutrition and distribution of phenotypic criteria in RTR according to the GLIM criteria. Using these criteria, malnutrition is present in one in 7 stable outpatient RTR. Importantly, in the vast majority of malnourished RTR the phenotypic criterion for reduced muscle mass was present, while most malnourished RTR had a BMI within the normal range, and one fifth had a BMI within the overweight or obese range.

The malnutrition prevalence of 14% in this cohort is lower compared to previous reported prevalence rates of 28-52%, as measured by the Subjective Global Assessment (SGA) and the Malnutrition-Inflammation Score^{18, 45, 46}. Although absolute numbers differ between studies, due to the use of different instruments and differences in patient characteristics between the samples, these prevalence rates indicate that malnutrition is a common issue in RTR that warrants attention, including at routine outpatient visits. The current study is, to the best of our knowledge, the first to apply the GLIM criteria in RTR, contributing to the body of knowledge on both malnutrition in RTR and malnutrition according to the GLIM criteria in different patient populations⁴⁷⁻⁵¹.

Our findings indicate that the inclusion of the phenotypic criterion for reduced muscle mass is most important to detect malnutrition in stable outpatient RTR. Contribution of the criterion for weight loss and especially low BMI were very low in our study sample. This may be explained by the high prevalence of overweight and obesity (i.e., 65% in the current study) and post-transplant weight gain, which are known to specifically increase fat mass^{8, 11, 20, 23}. Underlying muscle mass depletion and malnutrition may therefore easily be overlooked, possibly negatively impacting health outcomes. Previous studies showed that reduced muscle mass, assessed by either serum creatinine or 24-h urinary CER, is associated with higher mortality rates and graft failure in RTR^{21, 52, 53}. Lower muscle mass as determined by BIA was also found to be associated with a worse renal function⁵⁴.

Although this study revealed reduced muscle mass as the dominant phenotype of malnutrition in RTR, diagnostic assessment of muscle mass has several limitations. Currently, no consensus exists about the optimal measurement tool for muscle mass in clinical practice¹⁹. Computed tomography (CT) and magnetic resonance imaging (MRI) are considered the gold standards for assessment of muscle mass, but are not often used for nutritional assessment in clinical practice because of the high costs and practical limitations^{38, 55}. The GLIM therefore recommends using other indirect body composition measures, such as DEXA or BIA¹⁹, and a previous study in RTR, with mostly a healthy BMI, showed that estimations of muscle mass by CT, DEXA and BIA were well-correlated⁵⁶. However, the use of BIA to estimate muscle mass may include errors caused by changes in hydration status and, importantly, may overestimate muscle mass in obesity^{57, 58}. Given the high rate of obesity in our cohort (27%), the prevalence of low muscle mass by use of BIA is probably underestimated in this study population. In addition to the recommended body composition measures, when applied properly, 24-h urinary CER is a reliable non-invasive method to assess muscle mass in the general population and RTR^{21, 22, 59}. However,

validated age- and gender specific cut-off values for low muscle mass based on 24-h CER are not yet available, which hampers its clinical use for the diagnosis of malnutrition. In the current study we therefore used the CHI, which was previously used to detect malnutrition in lung transplant candidates⁶⁰ and patients on prolonged mechanical ventilation⁴³, and we found very similar malnutrition prevalence numbers when using CHI as measure for reduced muscle mass.

Several clinical and nutritional parameters were associated with malnutrition in the present study. Regarding the clinical parameters, the higher eGFR in malnourished RTR seems counterintuitive, as worse renal function can impair nutritional status¹⁸. However, eGFR is overestimated in subjects with low muscle mass, as eGFR is based on serum creatinine level, which reflects not only removal (renal excretion) but also supply of creatinine which is predominantly derived from muscle turnover⁶¹. Creatinine clearance is not subject to this particular bias, and in fact, has been recommended as better measure of renal function in patients with low muscle mass or malnutrition⁶¹. In our cohort, the similar creatinine clearance in well-nourished and malnourished RTR supports the assumption that the higher eGFR in malnourished RTR is due to overestimation by low muscle mass. Currently, eGFR is a main parameter for renal monitoring in RTR. The overestimation of renal function by eGFR in malnourished patients may lead to underestimation of renal risk in these patients. Whereas this assumption requires further investigation, it nevertheless underscores the importance of routine nutritional assessment in RTR, to prevent malnutrition from remaining undetected, and to prevent bias in renal monitoring. Second, the lower proportion of proteinuric patients among malnourished RTR also seems counterintuitive, as proteinuria can compromise the nutritional status. However, in patients with overt proteinuria, reduction of dietary protein intake is known to reduce proteinuria⁶², and the same holds true for reduction of dietary sodium intake⁶³. Hence, the lower proportion of proteinuria in malnourished RTR may well be due to their lower intake of protein and sodium. Finally, the significant association between high education level and malnutrition was due to higher rates of the phenotypic criterion 'low muscle mass' in RTR with high education level (data not shown). With simultaneous lower rates of obesity in this group, this may well be explained by less overestimation of muscle mass by BIA in RTR with high education level⁵⁸. In additional analysis with use of CHI instead of BIA for muscle mass assessment, this significant association was indeed lost (data not shown).

With respect to the nutritional parameters, lower daily protein intake in g/d, but not in g/kg/d, was also independently associated with malnutrition. This discrepancy can be explained by differences in body weight between well-nourished and malnourished RTR, but this comparison may be obscured by not considering differences in body composition. A previous study found that the association between low protein intake and mortality in RTR was mediated by lower muscle mass⁶⁴. Although there are currently no recommendations for optimal protein intake for RTR, it can be hypothesized that, RTR with low muscle mass and/or malnutrition may benefit from increments in protein intake, similarly to the protein recommendations for elderly to maintain optimal muscle function⁶⁵. However, more research is required for determining optimal protein recommendations for RTR, as a causal relationship between protein intake and malnutrition cannot be determined from

the present study, and reversed causation cannot be excluded. As sodium and potassium intake were also significantly lower in malnourished RTR, the lower protein intake may well be a consequence of an overall lower food intake in the patients with malnutrition.

As reduced muscle mass is independently associated with increased mortality and graft failure in several populations, including RTR ^{21, 52, 53}, it may be an important factor to target with interventions to improve long-term outcomes in RTR. In addition to nutritional interventions, other types of interventions may be effective, including physical activity programs, which show some positive results in improving muscle performance and strength in RTR ⁶⁶⁻⁶⁸. However, most of these interventions were targeted at attenuating weight gain and reducing cardiovascular risk factors after transplantation rather than maintaining or increasing muscle mass. Future combined lifestyle intervention studies in RTR may benefit from targeting both sides of the coin, aiming for both reduction of fat mass and cardiovascular risk factors, as well as an increase of muscle mass and improvement or maintenance of an adequate nutritional status ⁶⁹.

This study has several limitations. Firstly, the information on weight loss was derived from patient-reported data, which is prone to bias and may result in either under- or overestimation of weight loss. Secondly, although the GLIM criteria has provided a useful framework to create more consistency in measurement of malnutrition, there is lack of consensus about the best way to measure muscle mass which hampers its assessment in both research and clinical setting. Especially for patient populations with high rates of obesity, including our population, it is important to find a reliable way to assess muscle mass, as BIA tends to overestimate muscle mass in obese individuals ^{57, 58} and malnutrition based on reduced muscle mass may consequently be overlooked. The use of CHI in the current study can be debated, since reference values for ideal CER are outdated, and not validated against currently used diagnostic techniques. Updated reference values and/or age- and gender specific cut-off values for 24-h CER validated against a golden standard for muscle mass assessment would be preferred, but are currently not available. Thirdly, only a limited amount of nutritional parameters were available for this study, and e.g. caloric intake was documented. Further studies are needed to better examine the role of nutritional intake in malnutrition in RTR. Lastly, with the lack of longitudinal data, we were not able to assess the prevalence of malnutrition over time and study its association with important health outcomes, e.g., graft and patient survival. However, the long-term follow-up of participants in TransplantLines cohort and biobank study will enable this in the coming years.

Several implications for clinical practice can be derived from our findings. Given the estimated prevalence of malnutrition of one out of 7 stable outpatient RTR, or a doubled number depending on the operationalization, structural nutritional assessment during outpatient visits is of paramount importance. Clinicians should be aware that BMI and critical weight loss alone lack sensitivity to detect malnutrition in RTR. General screening and assessment tools that only include these phenotypic criteria ^{24, 25}, when not followed by additional assessment of muscle mass, should be interpreted with caution in this population for the same reason. Instead, as low muscle mass is the most important driver of the malnutrition diagnosis in RTR, the assessment of muscle mass should be

incorporated in outpatient visits. However, more research is needed to determine the most reliable assessment tool for this specific population, in which overweight and obesity are highly prevalent. In our opinion, 24-h urinary CER is a promising tool for muscle mass assessment, which can easily be incorporated in a clinical and outpatient setting. We do acknowledge that more research is needed, e.g., to determine optimal cut-off values for CER, and to compare the predictive value of different existing screening and assessment tools and measurement techniques in RTR.

In conclusion, malnutrition is present in one in 7 stable outpatient RTR and is predominantly characterized by reduced muscle mass. Therefore, routine assessment and monitoring of muscle status is warranted in RTR, to prevent malnourished RTR from remaining undetected and untreated.

Author contributions

K.B. and I.M.Y.V. conceptualization, methodology, formal analysis, investigation and writing - original draft; A.G.N. resources and writing - review and editing; M.F.C.J.: writing - review and editing; S.J.L.B.: funding acquisition, resources and writing - review and editing; H.J.W.: writing - review and editing and supervision; G.J.N. conceptualization, writing - review and editing and supervision.

The final draft of the manuscript have been read and approved by all authors.

Funding

This study was part of the TransplantLines cohort and biobank study, which was supported by a grant from Astellas BV. Funders of this study had no role in the study design, collection of data, analysing the data, interpretation of results, writing the manuscript, or the decision to submit the manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank the TransplantLines team for their effort to collect the data and the participants for their participation in the study.

REFERENCES

- 1 Wolfe RA, Ashby VB, Milford EL, Ojo AO, Ettenger RE, Agodoa LYC, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med*. 1999; **341**: 1725-30.
- 2 Jofre R, Lopez-Gomez JM, Moreno F, Sanz-Guajardo D, Valderrabano F. Changes in quality of life after renal transplantation. *Am J Kidney Dis*. 1998; **32**: 93-100.
- 3 van Sandwijk MS, Al Arashi D, van de Hare FM, van der Torren JMR, Kersten MJ, Bijlsma JA, et al. Fatigue, anxiety, depression and quality of life in kidney transplant recipients, haemodialysis patients, patients with a haematological malignancy and healthy controls. *Nephrol Dial Transplant*. 2019; **34**: 833-8.
- 4 Oterdoom LH, de Vries AP, van Ree RM, Gansevoort RT, van Son WJ, van der Heide JJ, et al. N-terminal pro-B-type natriuretic peptide and mortality in renal transplant recipients versus the general population. *Transplantation*. 2009; **87**: 1562-70.
- 5 Stanfill A, Bloodworth R, Cashion A. Lessons learned: experiences of gaining weight by kidney transplant recipients. *Prog Transplant*. 2012; **22**: 71-8.
- 6 Boslooper-Meulenbelt K, Patijn O, Battjes-Fries MCE, Haisma H, Pot GK, Navis GJ. Barriers and Facilitators of Fruit and Vegetable Consumption in Renal Transplant Recipients, Family Members and Healthcare Professionals-A Focus Group Study. *Nutrients*. 2019; **11**: 2427.
- 7 Zelle DM, Kok T, Dontje ML, Danchell EI, Navis G, van Son WJ, et al. The role of diet and physical activity in post-transplant weight gain after renal transplantation. *Clin Transplant*. 2013; **27**: E484-90.
- 8 Johnson CP, Gallagher-Lepak S, Zhu Y, Porth C, Kelber S, Roza AM, et al. Factors influencing weight gain after renal transplantation. *Transplantation*. 1993; **56**: 822-7.
- 9 Cashion AK, Hathaway DK, Stanfill A, Thomas F, Ziebarth JD, Cui Y, et al. Pre-transplant predictors of one yr weight gain after kidney transplantation. *Clin Transplant*. 2014; **28**: 1271-8.
- 10 Hoogeveen EK, Aalten J, Rothman KJ, Roodnat JI, Mallat MJ, Borm G, et al. Effect of obesity on the outcome of kidney transplantation: a 20-year follow-up. *Transplantation*. 2011; **91**: 869-74.
- 11 Friedman AN, Miskulin DC, Rosenberg IH, Levey AS. Demographics and trends in overweight and obesity in patients at time of kidney transplantation. *Am J Kidney Dis*. 2003; **41**: 480-7.
- 12 Costa B, Moratelli L, Silva LB, Paiva AC, Silva AN, Carminatti M, et al. Body mass index in the first year after kidney transplantation. *Transplant Proc*. 2014; **46**: 1750-2.
- 13 Molnar MZ, Keszei A, Czira ME, Rudas A, Ujszaszi A, Haromszeki B, et al. Evaluation of the malnutrition-inflammation score in kidney transplant recipients. *Am J Kidney Dis*. 2010; **56**: 102-11.
- 14 Kalantar-Zadeh K, Cano NJ, Budde K, Chazot C, Kovesdy CP, Mak RH, et al. Diets and enteral supplements for improving outcomes in chronic kidney disease. *Nat Rev Nephrol*. 2011; **7**: 369-84.
- 15 Siew ED, Ikizler TA. Determinants of insulin resistance and its effects on protein metabolism in patients with advanced chronic kidney disease. *Contrib Nephrol*. 2008; **161**: 138-44.
- 16 Carrero JJ, Thomas F, Nagy K, Arogundade F, Avesani CM, Chan M, et al. Global Prevalence of Protein-Energy Wasting in Kidney Disease: A Meta-analysis of Contemporary Observational Studies From the International Society of Renal Nutrition and Metabolism. *J Ren Nutr*. 2018; **28**: 380-92.
- 17 Ujszaszi A, Czira ME, Fornadi K, Novak M, Mucsi I, Molnar MZ. Quality of life and protein-energy wasting in kidney transplant recipients. *Int Urol Nephrol*. 2012; **44**: 1257-68.
- 18 Molnar MZ, Czira ME, Rudas A, Ujszaszi A, Lindner A, Fornadi K, et al. Association of the malnutrition-inflammation score with clinical outcomes in kidney transplant recipients. *Am J Kidney Dis*. 2011; **58**: 101-8.
- 19 Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition - A consensus report from the global clinical nutrition community. *Clin Nutr*. 2019; **38**: 1-9.
- 20 van den Ham EC, Kooman JP, Christiaans MH, Leunissen KM, van Hooff JP. Posttransplantation weight gain is predominantly due to an increase in body fat mass. *Transplantation*. 2000; **70**: 241-2.
- 21 Oterdoom LH, van Ree RM, de Vries AP, Gansevoort RT, Schouten JP, van Son WJ, et al. Urinary creatinine excretion reflecting muscle mass is a predictor of mortality and graft loss in renal transplant recipients. *Transplantation*. 2008; **86**: 391-8.
- 22 Stam SP, Eisenga MF, Gomes-Neto AW, van Londen M, de Meijer VE, van Beek AP, et al. Muscle mass determined from urinary creatinine excretion rate, and muscle performance in renal transplant recipients. *J Cachexia Sarcopenia Muscle*. 2019; **10**: 621-9.
- 23 Woloszyk P, Malgorzewicz S, Chamiernia A, Debska-Slizien A. Obesity After Successful Kidney Transplantation. *Transplant Proc*. 2020.
- 24 Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for Undernutrition in Geriatric Practice: Developing the Short-Form Mini-Nutritional Assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci*. 2001; **56A**: M366-M72.
- 25 Stratton RJ, Hackston A, Longmore D, Dixon R, Price S, Stroud M, et al. Malnutrition in hospital outpatients and inpatients: prevalence, concurrent validity and ease of use of the 'malnutrition universal screening tool' ('MUST') for adults. *Br J Nutr*. 2004; **92**: 799-808.
- 26 Mulasi U, Kuchnia AJ, Cole AJ, Earthman CP. Bioimpedance at the bedside: current applications, limitations, and opportunities. *Nutr Clin Pract*. 2015; **30**: 180-93.
- 27 Eisenga MF, Gomes-Neto AW, van Londen M, Ziengs AL, Douwes RM, Stam SP, et al. Rationale and design of TransplantLines: a prospective cohort study and biobank of solid organ transplant recipients. *BMJ Open*. 2018; **8**: e024502.
- 28 Ottery FD. Definition of standardized nutritional assessment and interventional pathways in oncology. *Nutrition*. 1996; **12**: S15-9.
- 29 Sealy MJ, Hass U, Ottery FD, van der Schans CP, Roodenburg JLN, Jager-Wittenaar H. Translation and Cultural Adaptation of the Scored Patient-Generated Subjective Global Assessment: An Interdisciplinary Nutritional Instrument Appropriate for Dutch Cancer Patients. *Cancer Nurs*. 2018; **41**: 450-62.
- 30 Erickson N, Storck LJ, Kolm A, Norman K, Fey T, Schiffler V, et al. Tri-country translation, cultural adaptation, and validity confirmation of the Scored Patient-Generated Subjective Global Assessment. *Support Care Cancer*. 2019; **27**: 3499-507.
- 31 Henriksen C, Thoresen L, Fjoseide B, Lorentzen SS, Balstad TR, Ottery FD, et al. Linguistic and content validation of the translated and culturally adapted PG-SGA, as perceived by Norwegian cancer patients and healthcare professionals. *Clin Nutr ESPEN*. 2020; **38**: 178-84.
- 32 Levey AS, Stevens LA, Schmid CH, Zhang Y, Castro AF, GFeldman HI, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med*. 2009; **150**: 604-12.
- 33 Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if height and weight be known. *Arch Intern Med*. 1916; **17**: 863-71.
- 34 Maroni BJ, Steinman TI, Mitch WE. A method for estimating nitrogen intake of patients with chronic renal failure. *Kidney Int*. 1985; **27**: 58-65.
- 35 Weijs PJ, Sauerwein HP, Kondrup J. Protein recommendations in the ICU: g protein/kg body weight - which body weight for underweight and obese patients? *Clin Nutr*. 2012; **31**: 774-5.
- 36 Kruizenga H, Beijer S, Huisman-de Waal G, Jonkers-Schuitema C, Klos M, Remijnse-Meester W, et al. Richtlijn Ondervoeding: herkenning, diagnosestelling en behandeling van ondervoeding bij volwassenen. Stuurgroep Ondervoeding; 2019.
- 37 Sergi G, De Rui M, Veronese N, Bolzetta F, Berton L, Carraro S, et al. Assessing appendicular skeletal muscle mass with bioelectrical impedance analysis in free-living Caucasian older adults. *Clin Nutr*. 2015; **34**: 667-73.
- 38 Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyere O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019; **48**: 16-31.
- 39 de van der Schueren MAE, Keller H, Consortium G, Cederholm T, Barazzoni R, Compher C, et al. Global Leadership Initiative on Malnutrition (GLIM): Guidance on validation of the operational criteria for the diagnosis of protein-energy malnutrition in adults. *Clin Nutr*. 2020; **39**: 2872-80.
- 40 Evans WJ, Morley JE, Argiles J, Bales C, Baracos V, Guttridge D, et al. Cachexia: a new definition. *Clin Nutr*. 2008; **27**: 793-9.
- 41 Abedini S, Holme I, Marz W, Weihrauch G, Fellstrom B, Jardine A, et al. Inflammation in renal transplantation. *Clin J Am Soc Nephrol*. 2009; **4**: 1246-54.
- 42 Blackburn GL, Bistrian BR, Maini BS, Schlamm HT, Smith MF. Nutritional and metabolic assessment of the hospitalized patient. *The Hospitalized Patient*. 1977; **1**.
- 43 Datta D, Foley R, Wu R, Grady J, Scalise P. Can Creatinine Height Index Predict Weaning and

- Survival Outcomes in Patients on Prolonged Mechanical Ventilation After Critical Illness? *J Intensive Care Med.* 2018; **33**: 104-10.
- 44 Hulsen T, de Vlieg J, Alkema W. BioVenn - a web application for the comparison and visualization of biological lists using area-proportional Venn diagrams. *BMC Genomics.* 2008; **9**: 488.
- 45 Malgorzewicz S, Debska-Slizien A, Czajka B, Rutkowski B. Adipokines and nutritional status in kidney transplant recipients. *Transplant Proc.* 2014; **46**: 2622-6.
- 46 Sezer S, Ozdemir FN, Afsar B, Colak T, Kizay U, Haberal M. Subjective global assessment is a useful method to detect malnutrition in renal transplant patients. *Transplant Proc.* 2006; **38**: 517-20.
- 47 Allard JP, Keller H, Gramlich L, Jeejeebhoy KN, Laporte M, Duerksen DR. GLIM criteria has fair sensitivity and specificity for diagnosing malnutrition when using SGA as comparator. *Clin Nutr.* 2019.
- 48 Contreras-Bolivar V, Sanchez-Torralvo FJ, Ruiz-Vico M, Gonzalez-Almendros I, Barrios M, Padin S, et al. GLIM Criteria Using Hand Grip Strength Adequately Predict Six-Month Mortality in Cancer Inpatients. *Nutrients.* 2019; **11**.
- 49 Skeie E, Tangvik RJ, Nymo LS, Harthug S, Lassen K, Viste A. Weight loss and BMI criteria in GLIM's definition of malnutrition is associated with postoperative complications following abdominal resections - Results from a National Quality Registry. *Clin Nutr.* 2019.
- 50 Clark AB, Reijnierse EM, Lim WK, Maier AB. Prevalence of malnutrition comparing the GLIM criteria, ESPEN definition and MST malnutrition risk in geriatric rehabilitation patients: RESORT. *Clin Nutr.* 2020.
- 51 Matsumoto Y, Iwai K, Namikawa N, Matsuda S, Wakano C, Heya H, et al. The relationship between existing nutritional indicators and Global Leadership Initiative on Malnutrition (GLIM) criteria: A one-institution cross-sectional analysis. *Clin Nutr.* 2020.
- 52 Streja E, Molnar MZ, Kovcsdy CP, Bunnapradist S, Jing J, Nissenson AR, et al. Associations of pretransplant weight and muscle mass with mortality in renal transplant recipients. *Clin J Am Soc Nephrol.* 2011; **6**: 1463-73.
- 53 Di Micco L, Quinn RR, Ronksley PE, Bellizzi V, Lewin AM, Cianciaruso B, et al. Urine creatinine excretion and clinical outcomes in CKD. *Clin J Am Soc Nephrol.* 2013; **8**: 1877-83.
- 54 Tatal E, Sezer S, Uyar ME, Bal Z, Demirci BG, Acar FN. Evaluation of nutritional status in renal transplant recipients in accordance with changes in graft function. *Transplant Proc.* 2013; **45**: 1418-22.
- 55 Beaudart C, McCloskey E, Bruyere O, Cesari M, Rolland Y, Rizzoli R, et al. Sarcopenia in daily practice: assessment and management. *BMC Geriatr.* 2016; **16**: 170.
- 56 Yanishi M, Kinoshita H, Tsukaguchi H, Kimura Y, Koito Y, Jino E, et al. Dual Energy X-ray Absorptiometry and Bioimpedance Analysis are Clinically Useful for Measuring Muscle Mass in Kidney Transplant Recipients With Sarcopenia. *Transplant Proc.* 2018; **50**: 150-4.
- 57 Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Manuel Gomez J, et al. Bioelectrical impedance analysis-part II: utilization in clinical practice. *Clin Nutr.* 2004; **23**: 1430-53.
- 58 Jensen B, Braun W, Geisler C, Both M, Kluckmann K, Muller MJ, et al. Limitations of Fat-Free Mass for the Assessment of Muscle Mass in Obesity. *Obes Facts.* 2019; **12**: 307-15.
- 59 Heymsfield SB, Arteaga C, McManus C, Smith J, Moffitt S. Measurement of muscle mass in humans: validity of the 24-hour urinary creatinine method. *Am J Clin Nutr.* 1983; **37**: 478-94.
- 60 Schwebel O, Pin I, Barnoud D, Devouassoux G, Brichon PY, Chaffanjon P, et al. Prevalence and consequences of nutritional depletion in lung transplant candidates. *Eur Respir J.* 2000; **16**: 1050-5.
- 61 Zaman T, Filipowicz R, Beddhu S. Implications and importance of skeletal muscle mass in estimating glomerular filtration rate at dialysis initiation. *J Ren Nutr.* 2013; **23**: 233-6.
- 62 de Jong PE, Anderson S, de Zeeuw D. Glomerular Preload and Afterload Reduction as a Tool To Lower Urinary Protein Leakage: Will Such Treatments Also Help To Improve Renal Function Outcome? *J Am Soc Nephrol.* 1993; **3**: 1333-41.
- 63 Krikken JA, Laverman GD, Navis G. Benefits of dietary sodium restriction in the management of chronic kidney disease. *Curr Opin Nephrol Hypertens.* 2009; **18**: 531-8.
- 64 Said MY, Deetman PE, de Vries AP, Zelle DM, Gans RO, Navis G, et al. Causal path analyses of the association of protein intake with risk of mortality and graft failure in renal transplant recipients. *Clin Transplant.* 2015; **29**: 447-57.
- 65 Deutz NE, Bauer JM, Barazzoni R, Biolo G, Boirie Y, Bosy-Westphal A, et al. Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. *Clin Nutr.* 2014; **33**: 929-36.
- 66 Greenwood SA, Koufaki P, Mercer TH, Rush R, O'Connor E, Tuffnell R, et al. Aerobic or Resistance Training and Pulse Wave Velocity in Kidney Transplant Recipients: A 12-Week Pilot Randomized Controlled Trial (the Exercise in Renal Transplant [ExeRT] Trial). *Am J Kidney Dis.* 2015; **66**: 689-98.
- 67 Macdonald JH, Kirkman D, Jibani M. Kidney transplantation: a systematic review of interventional and observational studies of physical activity on intermediate outcomes. *Adv Chronic Kidney Dis.* 2009; **16**: 482-500.
- 68 Calella P, Hernandez-Sanchez S, Garofalo C, Ruiz JR, Carrero JJ, Bellizzi V. Exercise training in kidney transplant recipients: a systematic review. *J Nephrol.* 2019; **32**: 567-79.
- 69 Klaassen G, Zelle DM, Navis GJ, Dijkema D, Bemelman FJ, Bakker SJL, et al. Lifestyle intervention to improve quality of life and prevent weight gain after renal transplantation: Design of the Active Care after Transplantation (ACT) randomized controlled trial. *BMC Nephrol.* 2017; **18**: 296.