

University of Groningen

## Performance variability in perioperative sentinel events

Reijmerink, Iris M; Bos, Kelly; Leistikow, Ian P; Groeneweg, Jop; Cnossen, Fokie;  
Dongelmans, Dave A; van der Laan, Maarten J

*Published in:*  
British Journal of Surgery

*DOI:*  
[10.1093/bjs/znac067](https://doi.org/10.1093/bjs/znac067)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2022

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Reijmerink, I. M., Bos, K., Leistikow, I. P., Groeneweg, J., Cnossen, F., Dongelmans, D. A., & van der Laan, M. J. (2022). Performance variability in perioperative sentinel events: report on a nationwide data set. *British Journal of Surgery*, 109(7), 573-575. Article znac067. <https://doi.org/10.1093/bjs/znac067>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).



The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Performance variability in perioperative sentinel events: report on a nationwide data set

Iris M. Reijmerink<sup>1,2,\*</sup> , Kelly Bos<sup>1,3</sup>, Ian P. Leistikow<sup>4,5</sup>, Jop Groeneweg<sup>1,6,7</sup>, Fokie Cnossen<sup>1,8</sup>, Dave A. Dongelmans<sup>1,9</sup> and Maarten J. van der Laan<sup>1,2</sup> 

<sup>1</sup>Impulse Institute, Amsterdam, the Netherlands

<sup>2</sup>Department of Surgery, University Medical Centre Groningen, Groningen, the Netherlands

<sup>3</sup>Department of Surgery, Amsterdam University Medical Centres—location Academic Medical Centre, Amsterdam, the Netherlands

<sup>4</sup>Erasmus School of Health Policy and Management, Erasmus University, Rotterdam, the Netherlands

<sup>5</sup>Dutch Health and Youth Care Inspectorate, Utrecht, the Netherlands

<sup>6</sup>Centre for Safety in Healthcare, Delft University of Technology, Delft, the Netherlands

<sup>7</sup>Unit of Cognitive Psychology, Leiden University, Leiden, the Netherlands

<sup>8</sup>Department of Artificial Intelligence, Bernoulli Institute of Mathematics, Computer Science and Artificial Intelligence, University of Groningen, Groningen, the Netherlands

<sup>9</sup>Department of Intensive Care Medicine, Amsterdam University Medical Centres—location Academic Medical Centre, Amsterdam, the Netherlands

\*Correspondence to: Iris M. Reijmerink, University Medical Centre Groningen, Hanzeplein 1, 9713GZ Groningen, the Netherlands (e-mail: i.m.reijmerink@umcg.nl)

## Introduction

Sentinel events are unintended events causing death or serious harm to patients, and these remain widespread in healthcare across the globe<sup>1,2</sup>. Healthcare organizations in many countries are mandated to report sentinel events to national reporting systems, analyse them to determine root causes, and develop improvement measures to prevent recurrence<sup>3,4</sup>. Although the quality of sentinel event analysis has increased, similar events keep recurring<sup>4,5</sup>. Research suggests this might be due to the quality of the improvement measures<sup>4,6</sup>.

The capacity to develop effective improvement measures is impaired when sentinel event analyses overlook relevant contributing factors<sup>7</sup>. One such factor is performance variability<sup>4</sup>. Performance variability is the positive or negative variation in behaviour and performance caused by environmental, organizational, and work-related factors, as well as human and individual characteristics<sup>8,9</sup>. Reduction in performance variability is an important goal of quality management in industrial production and laboratory measurement<sup>10,11</sup>. Performance variability also has a profound impact on quality of care, but the integration of this as a factor in healthcare quality management remains scant<sup>12,13</sup>. Previously, Dutch hospitals have indicated that the teams performing sentinel event analyses do not usually have specific training or knowledge on performance variability, although the literature shows that this could contribute to addressing such variability adequately within a sentinel event analysis<sup>4</sup>.

The aim of this study was to establish how performance variability contributes to perioperative sentinel events, by examining how often performance variability is included in sentinel event analysis reports and forms the basis for suggested improvement measures.

## Methods

Ethical approval was not required for this study. All Dutch perioperative sentinel event analysis reports from July 2017 to July 2018 in the national database of the Dutch Health and Youth Care Inspectorate were analysed.

Whether performance variability was identified as a contributing factor to the sentinel event by the analysis team was first established, by examining whether the report explicitly mentioned performance variability or its synonyms (human factor(s), human error(s)). If not mentioned explicitly, performance variability as a factor contributing to the event was examined using the Human Factors Investigation Tool<sup>14</sup>. This tool identifies three levels in incidents caused by human factors: action errors occurring immediately before the incident (level 3), the thought processes leading to the action error (level 2), and the underlying causes (level 1). Performance variability was identified as a contributing factor if all three levels were found in the analysis report. If all three levels were not found, it was then established whether solely technical errors were stated as the cause of the sentinel event. Whether performance variability tools or literature were used in the analysis of the sentinel event was also analysed.

Finally, the suggested improvement measures were analysed to establish whether these addressed any performance shaping factors (PSFs) that may underlie performance variability. For this, the Standardized Plant Analysis Risk—Human Reliability Analysis method was used, which looks at eight PSFs<sup>15</sup>. Improvement measures were scored as good, adequate or insufficient, based on whether they addressed PSFs explicitly (good) or indirectly (adequate), or not at all (insufficient) (*Appendix S1*).

Received: November 15, 2021. Revised: January 10, 2022. Accepted: February 14, 2022

© The Author(s) 2022. Published by Oxford University Press on behalf of BJS Society Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

**Table 1 Quality scores and examples of improvement measures from Dutch perioperative sentinel event analysis reports between July 2017 and July 2018**

Improvement measure score	No. of measures (n = 442)	Example
Good	12 (2.7)	'Introduction of a whiteboard in the operating room for writing down prostheses sizes to prevent implanting the wrong prosthesis' Scored as good because directly aimed at reducing the complexity of the task as the mental effort required to memorize the prosthesis sizes is eliminated
Adequate	225 (50.9)	'Changing the introduction program for new employees' Although this measure was probably aimed at improving the experience and training of the operator(s) involved in the task, it did not mention what exactly needed to be changed nor the measured effect
Insufficient	205 (46.4)	'to take appropriate action'

Values in parentheses are percentages.

**Table 2 Frequency and examples of performance shaping factors addressed in improvement measures from Dutch perioperative sentinel event analysis reports between July 2017 and July 2018**

PSF	No. of measures covered by PSF (n = 442)	Example
Available time to diagnose and act upon abnormal situation	1 (0.2)	'The on-call surgeon will preferably not be running the outpatient clinic simultaneously in order to provide sufficient supervision to the Emergency department'
Stress and stressors	3 (0.7)	'Minimize phone use, and thus possible disturbance, during surgery'
Complexity of task at hand	2 (0.5)	'From now on, patients with an anatomical variation will be operated by two surgeons instead of one'
Experience and training of operator(s) involved in task	19 (4.3)	'Provide additional training in the recognition of complications following interventional cardiology procedures'
Formal procedures present for task	94 (21.3)	'Developing a checklist for pacemaker implementation'
Ergonomics and human-machine interaction	62 (14.0)	'Make a clearer distinction between the oxygen and medical air flow meter, by using a different type of connection for the access points to prevent mismatch'
Physical and mental fitness for duty	0 (0)	–
Work processes	36 (8.1)	'Incorporating an additional moment of verification during the time-out, to confirm that a particular procedure is known to all those present'
Unclear	20 (4.5)	–

Values in parentheses are percentages. PSF, performance shaping factor.

## Results

In total, 115 perioperative sentinel event analysis reports containing 442 improvement measures were evaluated. Characteristics of sentinel events are shown in [Table S1](#). In two reports (1.7 per cent), the emergence of the sentinel event was considered being a result of technical errors only. None of the reports explicitly mentioned performance variability or its synonyms. In 113 reports (98.3 per cent), however, performance variability was identified as a contributing factor in the emergence of the sentinel event. Only one analysis report referred to performance variability literature, and none of the reports used performance variability tools in their analyses.

Of the 442 improvement measures, 12 (2.7 per cent) addressed one or more PSFs explicitly and were thus scored as good. Some 225 (50.9 per cent) addressed one or more PSFs indirectly and were therefore scored as adequate. Finally, 205 improvements (46.4 per cent) were scored as insufficient as they did not mention any PSFs ([Table 1](#)).

Improvement measures addressing PSFs were mostly aimed at the formal procedures present (94 measures), ergonomics and human-machine interaction (62), and work processes (36) ([Table 2](#) and [Fig. S1](#)). For 20 improvement measures (4.5 per cent), it was unclear what PSF a measure was aimed to shape. As an example, 'the integration of a short time-out moment when an

operation takes longer than expected', could be aimed at the work processes, the complexity of the task or enhancing communication within the operating team.

## Discussion

Although performance variability was identified as an important contributing factor in almost all sentinel events, none of the analysis reports explicitly mentioned performance variability nor a synonym, and performance variability was under-represented in the improvement measures. These findings are alarming, as reducing performance variability is an important goal of quality management, and omitting it from improvement measures can therefore lead to suboptimal patient safety<sup>10,11</sup>.

Most of the improvement measures that addressed performance variability were aimed at procedures and work processes rather than at individual PSFs, such as the experience and training or physical and mental fitness of the operators. This resonates with the literature on root cause analysis, a frequently used approach to determine the cause of a sentinel event<sup>16</sup>. The root cause analysis literature suggests that analyses should focus primarily on systems and processes and not on individual performance, as system-level improvements are more effective<sup>6</sup>. Sentinel event analysis teams might also feel uncomfortable addressing individual performance

issues, as this can be interpreted as blaming and shaming. Previous research, however, showed that individual and organizational factors contribute equally to the development of sentinel events. Thus, both must be considered in order to truly understand the nature of sentinel events<sup>13</sup>. The Royal College of Surgeons of England mortality and morbidity meeting has already started to consider this, with questions being asked whether system, patient or staff factors contributed to an adverse event<sup>17</sup>. Recognition of staff-related PSFs, such as stress, does not equate to blaming individuals, but creates opportunities to address the underlying causes of such PSFs.

Although the present results have shown that performance variability plays an important role in nearly all perioperative sentinel events, knowledge of performance variability was underutilized in the reports. This confirms earlier studies in the healthcare sector reporting that specific knowledge or training in performance variability is scarce in sentinel event analysis teams and quality and safety departments of hospitals<sup>4,18</sup>. Knowledge of performance variability is of value in understanding why a sentinel event occurred, as well as in redesigning healthcare processes and systems to reduce the chances of recurrence. Understanding how the work environment influences human performance could also lead sentinel event analysis to address the causative factors, and may thus lead to more effective improvement measures.

The main goal of integrating performance variability knowledge into sentinel event analyses should, however, not solely be to identify causes of the event. It can also help understand things that go right: the safety II paradigm<sup>19</sup>. Integrating performance variability into sentinel event analyses will help to appreciate the conditions that strengthen the ability to function well despite difficult circumstances<sup>20</sup>. Explicitly addressing performance variability could improve sentinel event analyses, lead to more effective improvement measures that optimize human performance in the healthcare system, therefore having a larger potential to reduce preventable harm.

## Acknowledgements

The authors thank the Dutch Health and Youth Care Inspectorate (Utrecht, the Netherlands), G.J. Kamps, and A. van Wincoop, consultant and engineer at Intergo International Centre for Safety, Ergonomics & Human Factors (Amersfoort, the Netherlands), and A.F. The, medical student at the University of Groningen, for their contribution to this study. No preregistration exists for the studies reported in this article. Authors Iris M. Reijmerink and Kelly Bos contributed equally to this paper.

**Disclosure.** The authors declare no conflict of interest.

## Supplementary material

[Supplementary material](#) is available at BJS online.

## References

1. Langelan M, Broekens MA, de Bruijne MC, de Groot JF, Moesker MJ, Porte PJ et al. *Monitor Zorggerelateerde schade 2015/2016. Dossieronderzoek bij overleden patienten in Nederlandse ziekenhuizen. NIVEL 2017*. [https://www.nivel.nl/sites/default/files/bestanden/Rapport\\_Monitor\\_Zorggerelateerde\\_Schade\\_2017.pdf](https://www.nivel.nl/sites/default/files/bestanden/Rapport_Monitor_Zorggerelateerde_Schade_2017.pdf) (accessed 26 May 2020)
2. OECD/European Observatory on Health Systems and Policies. *State of Health in the EU. The Netherlands: Country Health Profile 2019*. [https://ec.europa.eu/health/sites/default/files/state/docs/2019\\_chp\\_nl\\_english.pdf](https://ec.europa.eu/health/sites/default/files/state/docs/2019_chp_nl_english.pdf) (accessed 5 June 2020)
3. Leistikow IP, Mulder SM, Vesseur J, Robben PBM. Learning from incidents in healthcare—the journey, not the arrival, matters. *BMJ Qual Saf* 2017;**26**:252–256
4. Bos K, Dongelmans DA, Greuters S, Kamps GJ, van der Laan MJ. The next step in learning from sentinel events in healthcare. *BMJ Open Qual* 2020;**9**:e000739
5. Nederlandse Federatie van Universitair medische centra (NFU). *Samenvatting voor raad van bestuur en leden analyseteam: beoordelen van verbetermaatregelen voortvloeiend uit calamiteitenanalyses*. [https://www.sturenopkwaliteit.nl/uploads/pdf/NFU\\_SoK\\_Calamiteiten\\_Samenvatting\\_Raad\\_van\\_Bestuur\\_en\\_Leden\\_analyseteam\\_maart\\_2019.pdf](https://www.sturenopkwaliteit.nl/uploads/pdf/NFU_SoK_Calamiteiten_Samenvatting_Raad_van_Bestuur_en_Leden_analyseteam_maart_2019.pdf) (accessed 5 June 2020)
6. Kellogg KM, Hettinger Z, Shah M, Wears RL, Sellers CR, Squires M et al. Our current approach to root cause analysis: is it contributing to our failure to improve patient safety? *BMJ Qual Saf* 2017;**26**:381–387
7. de Kam D, Kok J, Grit K, Leistikow I, Vlemminx M, Bal R. How incident reporting systems can stimulate social and participative learning: a mixed-methods study. *Health Policy* 2020;**124**:834–841
8. Reason J. Understanding adverse events: human factors. *Qual Health Care* 1995;**4**:80–89
9. Hirose T, Nomoto H, Hollnagel E, Hill R, Sawaragi T, Slater D. *Treating Variability Formally In FRAM*. 2020. <https://doi.org/10.13140/RG.2.2.26507.72485> (accessed 14 August 2020)
10. Fung V, Schmittiel JA, Fireman B, Meer A, Thomas S, Smider N et al. Meaningful variation in performance: what does variation in quality tell us about improving quality? *Med Care* 2010;**48**(2):113–139.
11. Shewhart WA. *The Economic Control of Quality of Manufactured Product* (7th edn). New York: D. van Nostrand Company, 1932
12. Selby JV, Schmittiel JA, Lee J, Fung V, Thomas S, Smider N et al. Meaningful variation in performance: what does variation in quality tell us about improving quality? *Med Care* 2010;**48**:133–139
13. Carayon P, Wood KE. Patient safety: the role of human factors and system engineering. *Stud Health Technol Inform* 2010;**153**:23–46
14. Gordon R, Flin R, Mearns K. Designing and evaluating a human factors investigation tool (HFIT) for accident analysis. *Saf Sci* 2005;**43**:147–171
15. Gertman D, Blackman H, Marble J, Byers J, Smith C. *The SPAR-H Human Reliability Analysis Method*. <https://www.nrc.gov/reading-rm/doc-collections/nuregs/contract/cr6883/cr6883.pdf> (accessed 11 January 2021)
16. Charles R, Hood B, Derosier JM, Gosbee JW, Li Y, Caird MS et al. How to perform a root cause analysis for workup and future prevention of medical errors: a review. *Patient Saf Surg* 2016;**10**:20
17. Royal College of Surgeons of England. *Morbidity and Mortality Meetings. A Guide to Good Practice*. London: Royal College of Surgeons of England, 2015
18. Canham A, Jun GT, Waterson P, Khalid S. Integrating systemic accident analysis into patient safety incident investigation practices. *Appl Ergon* 2018;**72**:1–9
19. Smith KM, Valenta AL. Safety I to safety II: a paradigm shift or more work as imagined? Comment on ‘false dawns and new horizons in patient safety research and practice’. *Int J Health Policy Manag* 2018;**7**:671–673
20. Hollnagel E, Wears RL, Braithwaite J. *From Safety-I to Safety-II: A White Paper*. <https://www.england.nhs.uk/signuptosafety/wp-content/uploads/sites/16/2015/10/safety-1-safety-2-white-papr.pdf> (accessed 2 May 2021)