

University of Groningen

## Neural correlates of acute post-traumatic dissociation

Mertens, Yoki Linn; Manthey, Antje; Sierk, Anika; Walter, Henrik; Daniels, Judith K.

*Published in:*  
BJPsych Open

*DOI:*  
[10.1192/bjo.2022.65](https://doi.org/10.1192/bjo.2022.65)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2022

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Mertens, Y. L., Manthey, A., Sierk, A., Walter, H., & Daniels, J. K. (2022). Neural correlates of acute post-traumatic dissociation: a functional neuroimaging script-driven imagery study. *BJPsych Open*, 8(4), Article e109. <https://doi.org/10.1192/bjo.2022.65>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Paper

# Neural correlates of acute post-traumatic dissociation: a functional neuroimaging script-driven imagery study

Yoki L. Mertens, Antje Manthey, Anika Sierk, Henrik Walter and Judith K. Daniels

## Background

Current neurobiological models of post-traumatic stress disorder (PTSD) assume excessive medial frontal activation and hypoactivation of cortico-limbic regions as neural markers of post-traumatic dissociation. Script-driven imagery is an established experimental paradigm that is used to study acute dissociative reactions during trauma exposure. However, there is a scarcity of experimental research investigating neural markers of dissociation; findings from existing script-driven neuroimaging studies are inconsistent and based on small sample sizes.

## Aims

The current aim was to identify the neural correlates of acute post-traumatic dissociation by employing the script-driven imagery paradigm in combination with functional magnetic resonance imaging.

## Method

Functional neuroimaging data was acquired in 51 female patients with PTSD with a history of interpersonal childhood trauma. Blood-oxygen-level-dependent response during the traumatic (versus neutral) autobiographical memory recall was analysed, and the derived activation clusters were correlated with dissociation measures.

## Results

During trauma recall, enhanced activation in the cerebellum, occipital gyri, supramarginal gyrus and amygdala was identified.

None of the derived clusters correlated significantly with dissociative symptoms, although patients reported increased levels of acute dissociation following the paradigm.

## Conclusions

The present study is one of the largest functional magnetic resonance imaging investigations of dissociative neural biomarkers in patients with PTSD undergoing experimentally induced trauma confrontation to elicit symptom-specific brain reactivity. In light of the current reproducibility crisis prominent in neuroimaging research owing to costly and time-consuming data acquisition, the current (null) findings highlight the difficulty of extracting reliable neurobiological biomarkers for complex subjective experiences such as dissociation.

## Keywords

Post-traumatic stress disorder; imaging; trauma confrontation; dissociation; biomarkers.

## Copyright and usage

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Post-traumatic dissociation – experiencing depersonalisation or derealisation sensations, often in response to stressors or trauma reminders – occurs frequently in people suffering from post-traumatic stress disorder (PTSD).<sup>1</sup> Despite the increased attention paid to post-traumatic dissociation following the inclusion of a dissociative PTSD subtype in the DSM-5,<sup>2</sup> neuroimaging research on the topic remains scarce. Recently published systematic reviews indicate that only a few dissociation-specific neural alterations could be identified in patients with PTSD, with most of them awaiting independent replication.<sup>3,4</sup> The majority of the existing literature reported brain activation during the resting state by comparing subgroups of patients based on their long-term patterns of responding without inducing a dissociative state during the actual data acquisition. Across studies, tentative support exists that patients with dissociative PTSD, compared with the classic PTSD group, depict increased connectivity between (pre-)frontal regions and subcortical regions, such as the amygdala, the periaqueductal gray<sup>5</sup> and the cerebellum.<sup>6</sup> The temporoparietal junction, a critical hub for multisensory integration, has shown increased connectivity to the superior colliculus<sup>7</sup> as well as the periaqueductal gray.<sup>8</sup> Altered brain circuitry was further reported between the right supramarginal gyrus and the thalamus,<sup>9</sup> of which the latter region depicted lower fractional anisotropy in a network with the left amygdala and hippocampus, correlating negatively with self-reported depersonalisation experiences.<sup>10</sup> Such brain activation

patterns during the resting state might detect differential connectivity networks, but cannot establish direct causal relationships between the observed alterations and the occurrence of dissociative symptoms. To achieve better causal attribution, symptom induction during data acquisition is needed.

## Script-elicited neural correlates of dissociation

The most well-established symptom provocation paradigm in PTSD neuroimaging research is script-driven imagery (SDI).<sup>11,12</sup> During SDI, participants are asked to repeatedly listen to and actively recall autobiographical memory audio scripts. According to a recent meta-analysis on the neural underpinnings of trauma-related autobiographical memory recall,<sup>13</sup> different mental processes may underly the retrieval of memory versus active engagement during imagery. These processes might correspond to different parts of the SDI paradigm: during the first 30 s, retrieval is prompted by passively listening to the script, while the patient is asked to actively imagine the scene described in the script for the following 30 s, and are assumed to correspond with differential brain activation patterns in PTSD.

To date, only a few studies have investigated neural activation patterns associated with the experience of post-traumatic dissociation by using the SDI paradigm. Early studies employing the SDI

were severely underpowered ( $n = 2^{14}$  and  $n = 7^{15}$  participants with PTSD), but introduced the idea of an absent amygdala activation on the backdrop of excessive prefrontal regulation as a correlate of dissociation. Lanius et al later suggested that such an overmodulation of the limbic system might be the causal mechanism differentiating the dissociative subtype of PTSD.<sup>16</sup> Hopper et al<sup>17</sup> employed SDI in 27 (mostly female) patients with chronic PTSD following (predominantly) motor vehicle accidents. The participants reported the severity of acute dissociation in response to the trauma confrontation directly following the functional neuroimaging paradigm. Dissociation correlated positively with activity in the left medial prefrontal and right superior temporal cortices, and negatively with the left superior temporal cortex and the right anterior insula. The anterior insula subserves interoception of bodily states and its underactivation in patients with dissociation was interpreted as 'biological substrate of extreme emotional underengagement'.<sup>17</sup> However, these correlates could not be replicated by the following two studies from the same laboratory also employing the SDI paradigm. One found positive correlations between acute dissociation and activation of the left middle frontal gyrus and right superior frontal gyrus ( $N = 21^{18}$ ) in a sample of participants with acute trauma. No significant neural correlates of acute dissociation were identified in a study probing pain processing with the SDI in PTSD ( $N = 17$ ).<sup>19</sup> Instead, a negative correlation between trait dissociation and activity in the amygdala, putamen, anterior cingulate cortex and left superior frontal gyrus was established.<sup>20,21</sup> In patients suffering from borderline personality disorder, dissociative processing during the SDI was associated with greater activation of the left superior frontal gyrus and lower activation of the right middle and inferior temporal gyri,<sup>22</sup> whereas Krause-Utz et al reported diminished bilateral amygdala activation as a correlate of acute dissociation after participants underwent a SDI task.<sup>23</sup>

In sum, there is some, albeit inconsistent, fMRI evidence suggesting that post-traumatic dissociation may be related to a fronto-limbic imbalance concurrent with increased feelings of depersonalisation, derealisation and emotional numbing reported by patients with PTSD. However, the heterogeneity of the results so far precludes the identification of a biomarker of post-traumatic dissociation. This heterogeneity may be because of various factors, including differences in sample composition (e.g. trauma event type, time since trauma exposure), usage of different measures capturing alternative aspects of dissociative experiencing (e.g. identity confusion) and small sample sizes. The field of neuroimaging has seen a lot of criticism regarding the low statistical power of most studies, because of their small sample sizes,<sup>24</sup> and simulation studies have shown that samples with less than 30 participants – such as the ones cited above – show particularly large variation on subsequent replications.<sup>25</sup> Because of the high costs of neuroimaging, concerted efforts are needed to (a) provide single studies with adequate sample sizes analysed, using power-enhancing analysis strategies; and (b) pool data across studies as recently done within large consortiums (e.g. Wang et al<sup>26</sup>). However, to date, neither option has been applied to the identification of biomarkers of post-traumatic dissociation.

## Study aims

The current study aims to identify neural activation patterns associated with acute post-traumatic dissociation during symptom provocation. To this end, a large sample ( $N > 50$ ) of patients with PTSD with coherent characteristics (female gender, history of interpersonal childhood trauma) will be analysed. The secondary aim of this study is to evaluate the paradigm most often used in the field, i.e. the SDI paradigm regarding the distinction of underlying memory

processes and its usefulness for the identification of biomarkers of dissociation.

## Method

### Participants

A total of  $N = 51$  female participants (mean age 40.29 years, s.d. 10.17) diagnosed with PTSD were included in the current study. Participants were recruited by means of public advertisements and mental health treatment centres. Interested individuals were first screened via telephone for magnetic resonance imaging (MRI) compatibility and medication status. They were fully informed about the study procedure and potential risks before signing the informed consent. Written informed consent was obtained from all participants. They received a booklet containing the self-report instruments and were asked to fill these in at home. At least one full day before the scanning session, interested patients were invited to the laboratory to undergo clinical diagnostics. They were included in the study if they met all inclusion criteria: 20–60 years, proficient in German, MRI compatible, no neurological disorder, no history of head injury, no substance dependency in the past 6 months, no intake of benzodiazepine or anticonvulsants, PTSD as primary disorder, PTSD symptoms for at least 3 months and experience of interpersonal childhood trauma before the age of 22 years. To account for the high comorbidity encountered in PTSD, participants with the following comorbid disorders as a secondary diagnosis were eligible: comorbid depressive, anxiety, eating, substance misuse and borderline personality disorders. Patients with other comorbidities (e.g. dissociative disorders) were excluded.

The current data stems from a larger trial investigating medication effects. Participants were scanned twice on two separate days. For the current article, only data from the placebo condition were used, which was counterbalanced across participants. During data collection, a single participant dropped out (owing to an accident) in between sessions, but their data from the first day of scanning (placebo condition) could be included in the current analysis. All other participants completed their participation; however, data for one participant could not be included for current analysis because of missing log files, resulting in a total sample of  $N = 51$ .

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. The study was approved by the medical ethics board of the University of Magdeburg [57/14] and the ethical committee of the Berlin Psychological University, and was preregistered (aspredicted.org: 78135).

### Procedure

After study inclusion, the scripts employed during the SDI paradigm were developed. To this end, participants were asked to provide short but detailed narratives (told from a first-person perspective in the present tense) of one neutral and one traumatic autobiographical memory containing specifically associations (e.g. time of day, bodily sensations, sounds, smell, colours, mood, cognition) to facilitate recollection and re-experiencing during script exposure. The trauma scripts were based on the traumatic memory that elicited the strongest intrusion symptoms at the point of testing (without being too overwhelming). In the current sample, the majority of trauma scripts related to experiences of childhood trauma (80.4%), of which 76.5% related to childhood sexual abuse, physical abuse or a combination of both (see Table 1). The

**Table 1** Demographic variables

|  | % (n)         | Mean (s.d.)   |
|--|---------------|---------------|
| Age, years                                   |               | 40.29 (10.17) |
| School education (highest degree)            |               |               |
| High school diploma                          | 82.3 (42)     |               |
| Junior high school diploma                   | 17.6 (9)      |               |
| Work experience (highest degree)             |               |               |
| University degree                            | 45.1 (23)     |               |
| Vocational training/apprenticeship           | 31.4 (16)     |               |
| Trainee                                      | 11.8 (6)      |               |
| None   | 11.8 (6)      |               |
| Relationship status                          |               |               |
| Single                                       | 51.0 (26)     |               |
| In partnership                               | 21.5 (11)     |               |
| Married                                      | 19.6 (10)     |               |
| Divorced                                     | 21.6 (11)     |               |
| Widowed                                      | 2.0 (1)       |               |
| Lifetime traumatic experiences (ETI-TL)      |               |               |
| Childhood sexual abuse <sup>a</sup>          | 84.3 (43)     |               |
| Violent attack <sup>a</sup>                  | 78.4 (40)     |               |
| Neglect                                      | 68.6 (35)     |               |
| Severe illness                               | 60.8 (31)     |               |
| Sexual abuse in adulthood <sup>a</sup>       | 52.9 (27)     |               |
| Death of close person                        | 49.0 (25)     |               |
| Accidental trauma                            | 43.1 (22)     |               |
| Torture                                      | 17.6 (9)      |               |
| Natural disaster                             | 17.6 (9)      |               |
| Imprisonment                                 | 11.8 (6)      |               |
| War combat                                   | 3.9 (2)       |               |
| Age at onset of worst traumatic experience   | 17.61 (12.37) |               |
| Interpersonal childhood trauma (CTQ)         |               |               |
| Childhood sexual abuse ( $\geq 8$ )          | 72.5 (37)     | 14.10 (7.25)  |
| Childhood physical abuse ( $\geq 8$ )        | 60.8 (31)     | 11.12 (5.73)  |
| Childhood emotional abuse ( $\geq 10$ )      | 82.4 (42)     | 19.06 (5.61)  |
| Childhood emotional neglect ( $\geq 15$ )    | 78.4 (40)     | 18.56 (4.79)  |
| Childhood physical neglect ( $\geq 8$ )      | 76.5 (39)     | 11.58 (4.70)  |
| Above cut-off on any abuse subscale          | 91.7 (44)     | 2.29 (0.97)   |
| Above cut-off on any neglect subscale        | 91.7 (44)     | 1.65 (0.64)   |
| Above any cut-off (combined)                 | 95.8 (46)     | 3.94 (1.39)   |
| Traumatic memory for script <sup>b</sup>     |               |               |
| Childhood trauma                             | 80.4 (41)     |               |
| Childhood sexual abuse                       | 54.9 (28)     |               |
| Childhood physical abuse                     | 11.8 (6)      |               |
| Childhood combined sexual and physical abuse | 9.8 (5)       |               |
| Other  | 3.9 (2)       |               |
| Adult trauma                                 | 19.6 (10)     |               |
| Adult sexual abuse                           | 5.9 (3)       |               |
| Adult physical abuse                         | 3.9 (2)       |               |
| Other  | 5.9 (3)       |               |
| Comorbid diagnosis (SCID-I and -II)          |               |               |
| Anxiety disorder                             | 80.4 (41)     |               |
| Affective disorders                          | 41.2 (21)     |               |
| Borderline personality disorder              | 15.7 (8)      |               |
| Eating disorders                             | 9.8 (5)       |               |
| Obsessive-compulsive disorder                | 6.0 (3)       |               |
| Substance misuse                             | 2.0 (1)       |               |

ETI-TL, Essen Trauma-Inventory – Trauma Checklist (multiple answers possible); CTQ, Childhood Trauma Questionnaire; SCID-I and -II, Structured Clinical Interview for DSM-IV Axis I and Axis II Disorders.  
a. Perpetrated by familiar person (e.g. family member), stranger or both.  
b. Defined as traumatic memory eliciting strongest intrusion symptoms at assessment.

neutral scripts were based on recent adult memories regarding everyday situations without a specific positive or negative valence (e.g. doing laundry). The narratives were converted into audio scripts with a neutral voice. In the days following script collection, participants were invited to the laboratory to undergo neuroimaging testing.

The functional imaging paradigm consisted of two condition blocks (neutral and trauma SDI), each comprising a 30-s baseline

period at the beginning as well as the end of the block, and three runs (repetitions of the script presentation). Each run lasted 3 min in total: 1 minute of imagery period (27 s of listening to the audio script and imaging event, and 33.75 s active recall) followed by 2 min of a recovery period, indicated by a bell sound. Congruent with previous research, the neutral condition always preceded the trauma condition to avoid carry-over effects (see Mickleborough et al<sup>19</sup>). At the end of each block, participants completed a self-report scale<sup>27</sup> to assess self-reported intrusive, avoidant, and dissociative responding elicited during the three recall periods across each block. Additionally, one re-experiencing ('During run X, did you re-experience part of the trauma involuntarily (intrusions)?') and one dissociation ('During run X, did you experience detachment sensations (dissociation)?') question was assessed per run (Fig. 1).

## Materials

### Clinician-administered diagnostic interviews


Potential participants were diagnosed with the gold standards of interview-based measures, i.e. the German version of the Structured Clinical Interview for DSM-IV<sup>28</sup> for Axis I disorders, and the Structured Clinical Interview for Dissociative Disorders.<sup>29</sup> PTSD diagnosis and symptom levels were interview-assessed by a trained clinician with the Clinician-Administered PTSD Scale (CAPS-IV<sup>30</sup>).

### Self-report measures

The participants completed various self-report questionnaires to assess traumatic experiences in childhood (Childhood Trauma Questionnaire (CTQ);<sup>31</sup> Cronbach's  $\alpha = 0.95$ ) and lifetime trauma (Essen Trauma Inventory (ETI)<sup>32</sup>), as well as the PTSD Checklist for DSM-IV<sup>33</sup> (Cronbach's  $\alpha = 0.72$ ), trait dissociation (German Version of the Dissociative Experiences Scale (FDS);<sup>34</sup> Cronbach's  $\alpha = 0.90$ ), depersonalisation (Cambridge Depersonalization Scale (CDS-30);<sup>35</sup> Cronbach's  $\alpha = 0.95$ ), somatoform dissociation (Somatoform Dissociation Questionnaire;<sup>36</sup> Cronbach's  $\alpha = 0.53$ ), peritraumatic dissociation (Peritraumatic Dissociative Experiences Questionnaire;<sup>37</sup> authorised German translation by A. Maercker<sup>38</sup>); Cronbach's  $\alpha = 0.45$ ), trait anxiety (State Trait Anxiety Inventory – Trait Version;<sup>39</sup> Cronbach's  $\alpha = 0.62$ ) and depression (Beck Depression Inventory-II;<sup>40</sup> Cronbach's  $\alpha = 0.93$ ). On the day of the scan, participants indicated their level of state dissociation (Clinician-Administered Dissociative States Scale (CADSS);<sup>41,42</sup> Cronbach's  $\alpha = 0.94$ ) twice, once before entering the MRI scanner and once immediately after exiting the scanner.

### Responses to Script-Driven Imagery Scale

The Responses to Script-Driven Imagery Scale (RSDI)<sup>27</sup> is an 11-item self-report scale to assess post-traumatic stress reactions, namely re-experiencing, avoidance and dissociation, in response to SDI during psychobiological symptom provocation research. Participants rated the severity level of each item (e.g. 'Did you feel disconnected from your body') on a seven-point Likert from 0 ('Not at all') to 6 ('A great deal'). For the current research, three subscale scores were computed by averaging the sum of self-reported re-experiencing (items 1–4), avoidance (items 5–7) and dissociation symptoms (items 8–11). The latter was employed to test how task-related brain activity corresponds to the acute dissociation experienced during fMRI. In the current sample, the internal consistency of the RSDI scale in the neutral condition was good, with a Cronbach's  $\alpha$  of 0.82 for the total scale, 0.45 for the re-experiencing subscale, 0.79 for the avoidance subscale and 0.85 for the dissociation subscale, of which the latter subscale depicted numerically



**Fig. 1.** Script-driven Imagery Paradigm. a, Schematic presentation of script-driven imagery paradigm. b, Script-elicited dissociation and intrusion for each run per script (Error bars: 95% Confidence Interval) CADSS, Clinician-Administered Dissociative States Scale; DIS, Dissociation; INT, Intrusion; RSDI, Responses to Script-driven Imagery Scale.

higher internal consistency compared with previous studies by Hopper et al.<sup>17,27</sup>

## Data analysis

### Script-evoked symptoms

To test whether the experimental manipulation successfully elicited trauma-related symptoms, the patients with PTSD responding to SDI was analysed as follows. First, three paired-sample *t*-tests were performed to compare the responses to SDI, immediately assessed after each condition block inside the scanner, and between the neutral and trauma conditions for each symptom subscale, respectively. Second, a paired-sample *t*-test was employed to compare the mean levels of state dissociation (measured with the CADSS) assessed before and after the neuroimaging paradigm. Third, two separate repeated-measures ANOVAs were employed to test for the impact of repeated presentation (i.e. potential habituation and sensitisation effects) of the autobiographical scripts on dissociation and intrusion, respectively, with script (trauma versus neutral) and repetition (first/second/third run) as within-participant factors.

### fMRI data acquisition

Participants were scanned on a 3T Siemens Magnetom TrioTim (syngo MR, Siemens, Erlangen, Germany) at the Berlin Institute of Health, Charité Universitätsmedizin Berlin, equipped with a 12-channel hybrid birdcage radiofrequency coil for magnetic resonance signal transmission and reception.

### Anatomical images

High-resolution T1-weighted anatomic images were acquired using magnetisation-prepared rapid acquisition with a gradient echo sequence scanned in sagittal orientation with the following parameters: field of view (FoV) 256, 192 slices, 1 mm isotropic voxel size, repetition time (TR) 1.9 ms, echo time (TE) 2.52 ms, flip angle 9°, time (TI) 900 ms, 50% distancing factor.

### Functional images

A set of 40 contiguous, transversely orientated, 3-mm thick functional slices were prescribed parallel to the anterior commissure-posterior plane. The T2\*-weighted Blood-oxygen-level-dependent functional brain volumes were obtained using gradient echo

planar imaging with an interleaved slice acquisition (64 × 64 matrix size, TR = 2.25 s, TE = 25 ms, flip angle 80°, FoV = 192 mm, distancing factor 20%).

## fMRI data analysis

### fMRI preprocessing

Image processing and statistical analyses were performed with Statistical Parametric Mapping (SPM for Mac; version 12; Wellcome Department of Neurology, London, UK; <https://www.fil.ion.ucl.ac.uk/spm/software/spm12/>) within MATLAB 9.9.0 (R2020b; MathWorks for Mac (The MathWorks, Inc., Natick, Massachusetts, United States; see [www.mathworks.com/products/matlab.html](http://www.mathworks.com/products/matlab.html)). DICOM images were converted to NIFTI format with dcm2niix. mac software (version 2 for Mac, 3 November 2020; Chris Rorden, University of South Carolina, South Carolina, United States; see <https://github.com/rordenlab/dcm2niix/releases>). Field map-derived voxel displacement maps were calculated per session to correct for motion distortions. The functional images corresponding to each condition (neutral versus traumatic imagery) were realigned and unwrapped to the mean image. Subsequent preprocessing steps included co-registration of anatomical image to the unwrapped mean image, segmentation of the coregistered anatomical image, normalisation (3 × 3 × 3 mm) of the realigned functional images to a Montréal Neurological Institute (MNI) anatomical template and spatial smoothing to a Gaussian kernel of 6 mm full-width half-maximum.

### First-level analyses

Voxel-wise general linear models were used to investigate the activation patterns during each condition. The following conditions were modelled as regressors: 'Retrieval' (27 s) and 'Re-experiencing' (33.75 s), present for the sessions 'Neutral' and 'Trauma'. Conditions within the tasks were contrasted voxel-wise to each other in first-level analyses, to identify areas that are less or more active in the trauma (versus neutral) condition at the participant level. The present study used the null periods before (30 s), in-between runs (i.e. 2 min of rest period in between script presentation), and after (30 s) the SDI paradigm as the implicit baseline measures. During these periods, participants had their eyes open staring at a fixation cross instructed to relax and 'let go' of the memory material.

### Second-level analyses

**Whole-brain analysis.** Whole-brain analyses were employed to identify task-elicited activation clusters of contrast trauma versus neutral in a three-step approach following Thome et al.<sup>13</sup> The main analysis consisted of the whole recall across the script and imagery period, as analysing all volumes acquired during the 1-min period in which participants actively recalled the memory granted maximum statistical power. Then, separate analyses for the retrieval period (during script presentation) and the re-experiencing phase (during imagery) were conducted to investigate potential brain activation differences, as suggested by Thome et al.<sup>13</sup>

**Region of interest approach.** To test the central tenets of conceptualisations of dissociation as the result of limbic overmodulation,<sup>16</sup> we supplemented our whole-brain analyses with a targeted analysis of amygdala activation using a region-of-interest (ROI) approach. We defined and extracted the bilateral amygdala complex as described in the Juelich Atlas,<sup>43</sup> employing the JuBrain SPM Anatomy Toolbox for Mac, version 3.0 (Institute of Neuroscience and Medicine (INM-1, INM-7) of the Forschungszentrum Jülich,

Germany; see [www.fil.ion.ucl.ac.uk/spm/ext/#Anatomy](http://www.fil.ion.ucl.ac.uk/spm/ext/#Anatomy)).<sup>44</sup> Small-volume and Bonferroni corrections were applied to reduce the risk of an inflated type 1 error (Bonferroni-adjusted  $P < 0.025$ , two-tailed).

**Association with dissociative processing.** To test whether the observed activation differences were (in part) being driven by the increased dissociation during the trauma condition, SPM voxel-wise multiple regression analyses were repeated with covaried dissociation measures. As such, participants' ratings on acute (RSDI dissociation subscale) and trait dissociation (FDS) and depersonalisation scores (CDS) were separately entered as covariates in the second-level analyses, to identify neural correlates of dissociative processing (whole brain, i.e. inside and outside the previously identified clusters). For the ROI analyses, mean activation parameters of the identified ROIs were extracted per participant, and correlated with their respective dissociation scores (i.e. bivariate correlation, two-tailed).

**Statistical thresholding.** Probabilistic threshold-free cluster enhancement (pTFCE;<sup>45</sup>) with family-wise error (FWE) correction was employed for statistical thresholding and multiple comparisons correction ( $P < 0.05$ ) for the identification of activation differences by condition. However, given the nature of the phenomenon studied in this investigation, we did not expect very strong brain activation correlates for post-traumatic dissociation. In line with the existing literature of dissociation, in a second step we thresholded the second-level analyses with dissociation scores as covariates at  $P < 0.001$  (uncorrected, cluster extent  $k \geq 10$ ). This was deemed useful to explore potential effects of small or medium effect size that did not survive the initial conservative thresholding.

**Exploratory analyses.** In case the observed activation patterns were not sufficiently explained by dissociative processing, alternative explanations such as the intensity of avoidance or re-experiencing during data acquisition were explored (see Hopper et al.<sup>17</sup>).

## Results

### Descriptive and correlational analyses

Demographic variables, including childhood and lifetime traumatic events, are summarised in Table 1. As expected, because of the targeted inclusion procedure, the sample reported a high frequency of interpersonal childhood trauma (CTQ), which was also predominantly labelled as the participant's worst traumatic experience. On average, participants reported six to seven traumatic events across the lifespan (ETI: mean 6.49, s.d. 2.45, range: 2–14) and exhibited high levels of PTSD severity (CAPS-IV: mean 67.45, s.d. 14.04, range: 40–95), as well as dissociation, depression and anxiety (see Table 2). Most dissociation measures were highly intercorrelated: script-elicited acute post-traumatic dissociative reactions were strongly related with state dissociation assessed after the experiment, depersonalisation and peritraumatic dissociation, but not trait dissociation. Severity of childhood trauma experiences correlated positively with trait dissociation, somatoform dissociation and acute re-experiencing, but not acute dissociation. Additional inspection of the CTQ subscales did not depict significant differential associations between childhood maltreatment type and dissociation levels evoked by the behavioural paradigm. Association strength of large effects was found between self-reported PTSD severity and trait, somatoform and peritraumatic dissociation.

**Table 2** Descriptive and correlational statistics

| Variable  | n  | Mean  | s.d.  | 1              | 2      | 3      | 4              | 5              | 6              | 7              | 8              | 9      | 10             | 11             | 12             | 13     | 14             | 15             | 16 |
|---|----|-------|-------|----------------|--------|--------|----------------|----------------|----------------|----------------|----------------|--------|----------------|----------------|----------------|--------|----------------|----------------|----|
| 1. Script-evoked dissociation (RSDI) <sup>a</sup>           | 51 | 2.40  | 1.47  | –              |        |        |                |                |                |                |                |        |                |                |                |        |                |                |    |
| 2. Script-evoked re-experiencing (RSDI)                     | 51 | 4.79  | 0.92  | –0.097         | –      |        |                |                |                |                |                |        |                |                |                |        |                |                |    |
| 3. Script-evoked avoidance (RSDI) <sup>a</sup>              | 51 | 2.45  | 1.39  | 0.327*         | –0.171 | –      |                |                |                |                |                |        |                |                |                |        |                |                |    |
| 4. State dissociation (CADSS)                               | 51 | 54.75 | 52.95 | <b>0.459**</b> | –0.132 | 0.161  | –              |                |                |                |                |        |                |                |                |        |                |                |    |
| 5. Trait dissociation (FDS) <sup>a</sup>                    | 51 | 26.74 | 15.98 | 0.273          | –0.057 | 0.117  | <b>0.579**</b> | –              |                |                |                |        |                |                |                |        |                |                |    |
| 6. Depersonalisation (CDS) <sup>a</sup>                     | 47 | 70.77 | 48.62 | <b>0.541**</b> | –0.144 | 0.138  | <b>0.805**</b> | <b>0.735**</b> | –              |                |                |        |                |                |                |        |                |                |    |
| 7. Somatoform dissociation (SDQ)                            | 47 | 34.34 | 11.71 | 0.226          | –0.150 | 0.035  | <b>0.668**</b> | <b>0.662**</b> | <b>0.774**</b> | –              |                |        |                |                |                |        |                |                |    |
| 8. Peritraumatic dissociation (PDEQ) <sup>a</sup>           | 47 | 22.60 | 9.59  | <b>0.518**</b> | –0.103 | 0.251  | <b>0.379**</b> | <b>0.420**</b> | <b>0.557**</b> | <b>0.523**</b> | –              |        |                |                |                |        |                |                |    |
| 9. Childhood Trauma (CTQ) <sup>a</sup>                      | 48 | 74.44 | 21.96 | 0.166          | –0.238 | 0.325* | 0.195          | 0.316*         | 0.235          | <b>0.412**</b> | 0.223          | –      |                |                |                |        |                |                |    |
| 10. Self-reported PTSD severity (PCL) <sup>a</sup>          | 51 | 37.92 | 6.58  | 0.270          | 0.023  | 0.206  | 0.282*         | <b>0.586**</b> | <b>0.520**</b> | <b>0.481**</b> | <b>0.531**</b> | 0.338* | –              |                |                |        |                |                |    |
| 11. Interview-assessed PTSD severity (CAPS-IV) <sup>a</sup> | 51 | 67.45 | 14.04 | 0.335*         | –0.017 | –0.069 | 0.150          | 0.228          | <b>0.418**</b> | 0.284          | 0.368*         | –0.071 | 0.231          | –              |                |        |                |                |    |
| 12. Re-experiencing (CAPS-IV Criterion B) <sup>a</sup>      | 51 | 19.73 | 5.47  | <b>0.387**</b> | –0.020 | –0.004 | 0.274          | 0.228          | 0.339*         | 0.192          | 0.187          | –0.074 | 0.098          | <b>0.712**</b> | –              |        |                |                |    |
| 13. Avoidance (CAPS-IV Criterion C) <sup>a</sup>            | 51 | 25.82 | 8.19  | 0.292*         | –0.151 | 0.088  | 0.302*         | 0.329*         | <b>0.497**</b> | 0.347*         | <b>0.460**</b> | 0.019  | 0.251          | <b>0.691**</b> | <b>0.419**</b> | –      |                |                |    |
| 14. Hyperarousal (CAPS-IV Criterion D) <sup>a</sup>         | 51 | 22.82 | 4.91  | 0.118          | 0.267  | –0.173 | –0.064         | 0.087          | 0.130          | 0.086          | –0.039         | –0.035 | 0.164          | <b>0.581**</b> | <b>0.373**</b> | 0.197  | –              |                |    |
| 15. Depression (BDI-II) <sup>a</sup>                        | 48 | 21.63 | 13.17 | –0.087         | 0.042  | –0.058 | 0.238          | <b>0.508**</b> | 0.315*         | 0.361*         | –0.128         | 0.126  | 0.200          | 0.152          | 0.139          | 0.055  | 0.266          | –              |    |
| 16. Trait anxiety (STAI-T) <sup>a</sup>                     | 47 | 55.32 | 10.50 | 0.068          | –0.013 | –0.082 | 0.324*         | <b>0.604**</b> | <b>0.523**</b> | <b>0.443**</b> | 0.178          | 0.158  | <b>0.427**</b> | <b>0.398**</b> | 0.244          | 0.352* | <b>0.429**</b> | <b>0.623**</b> | –  |

RSDI, Responses to Script-Driven Imagery Scale; FDS, German Version of the Dissociative Experiences Scale; CDS, Cambridge Depersonalization Scale; SDQ, Somatoform Dissociation Questionnaire; PDEQ, Peritraumatic Dissociative Experiences Questionnaire; CTQ, Childhood Trauma Questionnaire; PCL, PTSD Checklist for DSM-IV; CAPS-IV, Clinician-Administered PTSD Scale – Version IV; BDI-II, Beck Depression Inventory-II; STAI-T, State Trait Anxiety Inventory – Trait Version. \* $P < 0.05$  (two-tailed), \*\* $P < 0.01$  (two-tailed) in bold. a. Scores are normally distributed.

**Table 3** Script-elicited brain activity in 51 female patients with post-traumatic stress disorder

| x  | y   | z   | t    | P(FWE)      | k    | AAL                                     | Hemisphere |
|--|-----|-----|------|-------------|------|---|------------|
| Brain activation clusters during traumatic > neutral recall          |     |     |      |             |      |   |            |
| -6   | -70 | -25 | 7.24 | $P < 0.001$ | 4185 | Lobule VI of cerebellar hemisphere      | Left       |
| -45  | -76 | -7  | 7.18 | $P < 0.001$ |      | Inferior occipital gyrus                | Left       |
| 0  | -61 | -28 | 7.08 | $P < 0.001$ |      | Lobule VIII of vermis                   |            |
| -60  | -43 | 32  | 5.97 | $P = 0.006$ | 74   | Supramarginal gyrus                     | Left       |
| -57  | -34 | 23  | 5.69 | $P = 0.015$ |      | Supramarginal gyrus                     | Left       |
| -57  | -31 | 32  | 5.48 | $P = 0.028$ |      | Supramarginal gyrus                     | Left       |
| Brain activation clusters during traumatic > neutral retrieval       |     |     |      |             |      |   |            |
| -45  | -82 | 8   | 8.21 | $P < 0.001$ | 8863 | Middle Occipital gyrus                  | Left       |
| -54  | -64 | 11  | 8.19 | $P < 0.001$ |      | Middle Temporal gyrus                   | Left       |
| -60  | -58 | 5   | 8.09 | $P < 0.001$ |      | Middle Temporal gyrus                   | Left       |
| 60   | -40 | 26  | 6.75 | $P < 0.001$ | 1031 | Supramarginal gyrus                     | Right      |
| -3   | -28 | -7  | 6.74 | $P < 0.001$ |      | Dorsal raphe nucleus                    |            |
| 63   | -28 | 35  | 6.68 | $P = 0.001$ |      | Supramarginal gyrus                     | Right      |
| 9  | 5   | 50  | 5.87 | $P = 0.009$ | 61   | Supplementary motor area                | Right      |
| -6   | 8   | 41  | 5.70 | $P = 0.014$ |      | Middle cingulate and paracingulate gyri | Left       |
| 9  | 11  | 38  | 5.54 | $P = 0.023$ |      | Middle cingulate and paracingulate gyri | Right      |
| 3  | -1  | 62  | 5.86 | $P = 0.009$ | 52   | Supplementary motor area                | Right      |
| -6   | -7  | 68  | 5.66 | $P = 0.016$ |      | Supplementary motor area                | Left       |
| 9  | -7  | 65  | 5.50 | $P = 0.026$ |      | Supplementary motor area                | Right      |
| 21   | -1  | -1  | 5.78 | $P = 0.011$ | 140  | Pallidum                                | Right      |
| 42   | 5   | 11  | 5.71 | $P = 0.014$ |      | Rolandic operculum                      | Right      |
| 45   | 14  | 11  | 5.67 | $P = 0.016$ |      | Inferior frontal gyrus, opercular part  | Right      |
| 36   | 11  | 26  | 5.76 | $P = 0.012$ | 39   | Inferior frontal gyrus, opercular part  | Right      |
| -33  | 11  | 11  | 5.39 | $P = 0.036$ | 13   | Insula                                  | Left       |
| -39  | 2   | 8   | 5.35 | $P = 0.041$ |      | Insula                                  | Left       |
| Brain activation clusters during traumatic > neutral re-experiencing |     |     |      |             |      |   |            |
| -  | -   | -   | -    | -           | -    | -                                       | -          |

Coordinates (in Montréal Neurological Institute space) and anatomical labels (AAL) for overactivations in patients with post-traumatic stress disorder (PTSD) during the traumatic (versus neutral) memory (a) recall, and (b) retrieval during script presentation, and (c) re-experiencing during imagery (no statistically relevant findings) for a family-wise error (FWE)-corrected threshold of  $P_{\text{Probabilistic threshold-free cluster enhancement}} < 0.05$ , with cluster extent threshold of  $k \geq 10$ .

### Manipulation check: responses to SDI

Significantly stronger script-evoked dissociation was reported for the trauma (RSDI dissociation subscale mean 13.39, s.d. 5.91) compared with the neutral condition (mean 7.16, s.d. 4.89) ( $t(48) = -7.58$ ,  $P < 0.001$ ), illustrating successful symptom induction during the trauma condition. The successfulness of the paradigm was further supported by significantly increased state dissociation ( $t(49) = -5.08$ ,  $P < 0.001$ ), assessed before (CADSS mean 29.54, s.d. 35.08) and after (mean 54.75, s.d. 52.95) the neuroimaging experiment. The current sample further reported significantly increased re-experiencing ( $t(50) = -21.42$ ,  $P < 0.001$ ) and avoidance ( $t(89) = -7.58$ ,  $P < 0.001$ ) symptoms during the traumatic compared with the neutral condition.

### Repetition effect: dissociation

A significant interaction effect for script by repetition was found ( $F(1.42, 58.42) = 5.31$ ,  $P = 0.013$ ; because of sphericity (Mauchly's test  $\chi^2(2) = 19.50$ ,  $P < 0.001$ ), Greenhouse-Geisser correction was applied). *Post hoc* analyses showed that in the neutral condition, dissociation was not affected by repetition ( $F(1.49, 68.57) = 0.02$ ,  $P = 0.947$ ), whereas repeated presentation of the trauma script significantly increased dissociation levels ( $F(1.67, 76.76) = 5.86$ ,  $P = 0.007$ ). Thus, participants reported consistently low dissociation across repetitions in the neutral condition, but showed a linear increase in dissociation per repetition trial.

### Repetition effect: intrusion

Results indicated no evidence for a similar interaction effect for intrusions ( $F(1.64, 75.64) = 0.63$ ,  $P = 0.504$ ), but significant main effects for script ( $F(1, 46) = 196.59$ ,  $P < 0.001$ ) and repetition

( $F(2, 92) = 3.49$ ,  $P = 0.034$ ). *Post hoc* analyses confirmed that self-reported intrusion was consistently higher in the trauma compared with the neutral condition. No significant *post hoc* tests emerged for repetition in either condition.

### Functional neuroimaging results


#### Dissociation during trauma versus neutral recall

The contrast trauma versus neutral recall (i.e. retrieval and re-experiencing period combined) yielded a large activation cluster centred around the left cerebellum ( $-6$ ,  $-70$ ,  $-25$ ;  $t = 7.24$ , FWE  $P_{\text{PTFCE}} < 0.001$ ,  $k = 4185$ ) including the left inferior occipital cortex ( $-45$ ,  $-76$ ,  $-7$ ), and at the left supramarginal gyrus ( $-60$ ,  $-43$ ,  $32$ ;  $t = 5.97$ , FWE  $P_{\text{PTFCE}} < 0.001$ ,  $k = 74$ ) (Table 3; see Supplementary Fig. 1 available at <https://doi.org/10.1192/bjo.2022.65>). Acute post-traumatic dissociation (RSDI dissociation subscale) entered as covariate in the second-level analysis did not depict any significant correlation with brain activation, neither within nor outside the previously identified brain regions. Similarly, covaried analyses with trait dissociation (FDS) or depersonalisation (CDS) did not yield any significant correlations with brain activity.

#### Dissociation during trauma versus neutral retrieval

During script presentation (i.e. retrieval of autobiographical memory), participants exhibited significantly increased activation in clusters centred around the left middle occipital and temporal cortex, supramarginal cortex, bilateral supplementary motor area, pallidum and insula (Table 3). Covaried multiple regression models with dissociation measures (RSDI, FDS, CDS) with brain activity (inside and outside of these regions) did not result in any significant findings.





**Fig. 2.** Script-elicited signal activation clusters and corresponding brain-behaviour correlates ( $N = 51$ ).

#### Dissociation during trauma versus neutral re-experiencing

No activity clusters could be obtained under the FWE-corrected threshold for the re-experiencing period following script presentation. Likewise, multiple regression analyses with dissociation measures (RSDI, FDS, CDS) as covariates did not result in any significant association with brain activation.

#### ROI analyses

ROI analyses with small-volume correction for the left and right amygdala complex, as defined by the Juelich Atlas,<sup>43</sup> yielded increased activity in the left amygdala ( $-33, -7, 19, t = 4.47, k = 27, P_{FWE} = 0.002$ ) across the whole recall period (see Supplementary Fig. 1). During retrieval, elevated activity was observed in the left amygdala ( $-33, -7, 19, t = 4.98, k = 21, P_{FWE} < 0.001$ ) and right amygdala ( $24, -4, -16, t = 3.87, k = 15, P_{FWE} = 0.010$ ). No meaningful activation difference of the bilateral amygdala could be detected during the re-experiencing period alone. Correlational analyses between the derived activation clusters and acute post-traumatic dissociation (see Fig. 2), trait dissociation or depersonalisation symptoms did not yield significant findings. *Post hoc* exploration of a potential association between acute re-experiencing (RSDI) experienced during trauma script presentation and bilateral amygdala activation did not yield significant results.

#### Explorative correlational analyses and stability testing

To test whether the derived activation clusters could be explained by script-evoked acute re-experience or avoidance, the mean scores of these RSDI subscales were separately entered into the second-level

models as covariates for the contrast recall, retrieval and re-experiencing period. No activation correlations survived the statistical threshold corrected for multiple testing or emerged at an uncorrected alpha level of  $P_{TFCE} < 0.001$  level with a cluster extent of  $k \geq 10$ . To rule out that our contrasts were affected by carry-over effects (i.e. sustained brain activation during the subsequent rest period, see Lamke et al<sup>46</sup>), we employed an additional analysis technique (i.e. ‘scrubbing’<sup>47</sup>) to test the stability of our null results. Results stemming from the adjusted model did not depict meaningful divergences from the previous findings (see Supplementary material).

## Discussion

The current aim was to identify neural correlates of acute post-traumatic dissociation. To this end, brain activation was assessed during SDI in a large sample of female patients with PTSD with a history of interpersonal childhood trauma, and the derived activation was correlated with dissociation severity. The paradigm successfully induced dissociative and intrusive processing during the trauma condition, with increasing intensity per repetition as reported by the participants themselves (see Fig. 1). A subsequent state dissociation assessment confirmed a significant increase of dissociation severity following the paradigm. On the brain level, activation in the cerebellum, inferior occipital gyrus, left supramarginal gyrus and left amygdala significantly increased during traumatic memory recall compared with neutral memory recall. These regions are included in a wider network of brain regions subserving autobiographic memory recall.<sup>48</sup> Taken together, these results

suggest that the SDI paradigm was implemented successfully. However, inspection of neural correlates via covaried multiple regression analyses of the task-elicited brain activity with symptom measures of acute and trait dissociation/depersonalisation did not yield any meaningful results. We further inspected potential activation differences underlying retrieval (i.e. memory script presentation) and re-experiencing (i.e. imagery following script presentation). During trauma retrieval, the contrast yielded widespread elevated brain activation including the middle occipital gyrus, supramarginal gyrus, supplementary motor area, insula and amygdala, whereas no meaningful activation clusters survived statistical thresholding in the trauma re-experiencing period. Again, no distinct associations with dissociation severity emerged.

Post-traumatic dissociation is commonly conceptualised as a reactive defensive mechanism characterised by strong top-down inhibitory control to dampen a hypervigilant innate alarm system and numb physical and emotional distress often elicited by traumatic reminders, threat cues or stressors.<sup>49,50</sup> On a neurobiological level, excessive medial prefrontal activity is assumed to inhibit limbic regions (e.g. amygdala and insula) and alter processing in circuits involved in automatic physical responding and psychophysiological adaptations located in the brain stem. Present findings based on the brain activity of patients with PTSD reacting to personalised trauma memory scripts were unable to support the notion that post-traumatic dissociation corresponds with the proposed neural underpinnings of (pathological) emotional underengagement, namely excessive medial frontal hyperactivation and concurrent cortico-limbic hypoactivation (e.g. Lanius et al<sup>16</sup> and Hopper et al<sup>17</sup>). Although a ROI analysis suggested increased activity during trauma confrontation of the amygdala, a region associated with the processing of threat cues and trauma-specific stimuli,<sup>51,52</sup> the bilateral activation clusters were not large in volume or activation strength and did not correlate with any of the dissociation measures. Notably, none of the current analyses could replicate previous SDI studies indicating increased neural activity in the medial and frontal gyri<sup>18</sup> and medial prefrontal cortex,<sup>17</sup> all correlating significantly with acute dissociation.

The lack of identified neural correlates highlights the difficulty to extract reliable biomarkers for subtle subjective experiences such as dissociation via fMRI, especially when encountered in a heterogeneous psychiatric condition as PTSD. Considering that the present study tested almost the same number of patients with PTSD as the three aforementioned SDI studies together,<sup>17-19</sup> one could argue that previous significant findings are mostly spurious because of small sample sizes analysed with low statistical thresholds, which increase the risk for type 1 errors.<sup>25</sup> Alternatively, differences in data analyses approaches (e.g. ROI versus whole-brain analyses) or sample characteristics may partially account for the differential findings. For instance, Hopper et al's<sup>17</sup> sample suffered predominantly from a motor vehicle accident (as opposed to the current sample experiencing interpersonal childhood trauma) and reported numerically lower levels of trait dissociation (Dissociative Experiences Scale mean 9.3), but comparable levels of script-elicited acute dissociation (RSDI dissociation subscale mean 2.12). Childhood trauma, especially sexual and physical abuse, depicted a robust association with dissociation in both general population and psychiatric samples,<sup>53,54</sup> and has been related to alterations in neurobiological development.<sup>55</sup> Taking into account that the majority of the current sample reported moderate-to-severe levels of interpersonal traumatic childhood experiences, it is conceivable that the prolonged early stress may have indirectly affected (neural) responses to SDI. However, it should be noted that the severity of childhood trauma correlated significantly with trait and somatoform dissociation, but maltreatment sum (and subscale) scores were unrelated to the acute dissociation

experienced during the paradigm in the current sample (see Table 2).

With regard to the SDI paradigm, the current analyses revealed altered neural activity patterns that have already been observed by previous research. The existing PTSD literature has proposed various neural markers of intrusive re-experiencing, hyperarousal and hypervigilance, most prominently overactivation in brain regions related to emotional (e.g. amygdala), visual (e.g. occipital and parietal cortex) and memory processing (e.g. hippocampus) concurrent with a deficient hypoactivation in medial-frontal emotion regulation areas.<sup>16,56,57</sup> Again, the present findings did not find evidence for altered brain activity in frontal emotion regulation centres. Instead, the current sample depicted widespread elevated activity in the occipital cortex, posterior cerebellum and supramarginal gyrus when contrasting traumatic and neutral memory recall in patients with PTSD. Activation of the occipital cortex has traditionally been associated with both vivid imagery as well as autobiographical memory recall (see meta-analytic reviews by Winlove et al<sup>58</sup> and Svoboda et al<sup>48</sup>). New studies also implemented such activations in motor imagery.<sup>59</sup> It is conceivable that elevated brain activity in these regions following traumatic (versus neutral) memory recall corresponds to stronger sensory imagination, including recall of motor actions. Meanwhile, previous meta-analyses of symptom provocation studies and traumatic autobiographical memory recall in PTSD did not report such strong activation differences in the primary visual cortex.<sup>13,60</sup> Because of the inconsistent findings, it is currently unclear whether enhanced visual processing in the occipital and parietal regions, possibly associated with increased enhanced vividness and sensory imagination, are indeed neural markers of PTSD aetiology as postulated by Patel et al.<sup>57</sup>

The current study further sought to explore the activation differences underlying the recall of autobiographical memories: recall (retrieval and re-experiencing), retrieval only and re-experiencing only. According to a meta-analytical review by Thome et al,<sup>13</sup> within-group trauma-related recall in patients with PTSD elicited robust activation clusters in proximity to the precentral gyrus, caudate nucleus and right ventromedial prefrontal gyrus across the whole recall period. Likewise, enhanced activation in the ventromedial prefrontal cortices, supramarginal gyrus, precentral gyrus and caudate nucleus was found during retrieval; the latter two are suggestive an 'increase in procedural-based memory storage of patient's trauma memory'.<sup>13</sup> In contrast, the main activation clusters during re-experiencing were centred around the right insula (including amygdala and parahippocampal gyrus) and caudate nucleus, which correlated significantly with PTSD severity and may be indicative of enhanced (implicit) memory processing for (non-)associative fear responses during trauma imagery. However, these previous findings suggesting differential brain activation dependant on whether participants listen to the script or asked to imagine the traumatic event are based on a conjunction of studies employing different designs and contrasts. To our knowledge, the present investigation is one of the first to report findings across different processing stages within a single sample, and could only find limited support for the notion that distinct phases depicted differential brain activation.

Unexpectedly, the trauma re-experiencing period, which was the main contrast of interest in previous neuroimaging SDI studies,<sup>13</sup> elicited fewer and weaker brain activation clusters compared with the trauma retrieval phase. During script presentation (i.e. retrieval), the current sample depicted the strongest and most widespread activity in various brain regions (including the occipital gyri, supramarginal gyrus, supplementary motor area and amygdala), whereas no activation clusters survived statistical thresholding during re-experiencing, and were therefore not reported. It is

conceivable that the present sample did not comply sufficiently with the instructions and stopped reliving their traumatic memories once the script stopped playing. This notion remains purely speculative as self-report measures during the paradigm after each block only assessed mean symptom levels across the whole recall period, and not each retrieval and re-experiencing period separately. Still, the overall self-reported data indicated that the manipulation via SDI was successful, as participants reported higher PTSD symptom levels in the trauma compared with the neutral condition, and dissociation increased significantly per repetition of the traumatic script (i.e. sensitisation effect). In conjunction, current and previous findings do indicate that SDI is a valid paradigm to elicit trauma-related symptoms on a subjective level, but it still remains uncertain whether these symptom patterns correspond with robust and symptom-specific brain activation patterns, especially with regard to post-traumatic dissociation.

## Limitations

The current study presented data on the largest sample of patients with PTSD studied with the standard symptom provocation paradigm to date. However, the paradigm itself comes with certain limitations. Most importantly, it is correlational in nature, meaning that although it ensures that the symptom of interest is induced during data acquisition, making a causal attribution more plausible, it still relies on a correlational association between the subjective severity of this symptom and brain activation. An experimental manipulation allowing causal attribution might be able to move beyond this limitation, but is difficult to implement ethically. Another limitation of the present study is the lack of a trauma-exposed control group, which might have allowed us to determine whether the observed activation patterns are specific to PTSD development or are generally associated with exposure to a traumatic event. In addition, the current investigation did not assess whether participants had undergone trauma-focused treatment targeting the intrusive memory used in the script. One could argue that individual differences in habituation to trauma reminders (e.g. during exposure) may explain the intra-participant variability encountered across participants in their reactivity toward trauma script. However, in this case, the level of trauma-elicited reactivity should have correlated with self-reported PTSD symptoms, which was not the case in the present study. Notably, the homogeneity of the current patient sample can be regarded as both strength and weakness, as the focus on female participants with childhood trauma omits the need to control for gender and trauma type in the analyses, but also hampers the generalisation of our findings to other PTSD samples.

To conclude, the present study was unable to identify neural correlates of acute dissociation with the use of SDI and functional neuroimaging. Future research should explore alternative methods with a higher temporal resolution, such as electroencephalography, to analyse intra-participant variability over time (e.g. by using a sliding window approach to study shifts in of association networks over the course of the SDI paradigm) and test alterations of the study design (e.g. counterbalancing of order and comparison of short time intervals before, during and after trauma exposure, to explore potential anticipatory and carry-over effects). It would require an imaging method of both high spatial and temporal resolution to capture how brain patterns in people with PTSD differ within the milliseconds following trauma exposure; for instance, if people experiencing more dissociation depict increased bottom-up activation of prefrontal areas immediately followed by a top-down shutdown response of the initially elevated amygdala-steered threat response.<sup>50</sup> Additionally, we encourage inspection of brain activation patterns in stratified samples differing in trauma load (i.e. childhood maltreatment types

and severity,<sup>61</sup> multiple lifetime traumatic events, time since trauma onset) and current symptom severity. Arguably, PTSD profile differentiation based on patient history and individual item scores (instead of categorical diagnostic assessments) may lead to more precise brain-behaviour prediction modelling and enhanced ecological validity of theoretical frameworks on post-traumatic dissociation.

**Yoki L. Mertens** , Department of Clinical Psychology and Experimental Psychopathology, University of Groningen, The Netherlands; **Antje Manthey**, Charité University Clinic Berlin, Freie Universität Berlin, Humboldt-Universität zu Berlin and Berlin Institute of Health, Germany; **Anika Sierk**, Charité University Clinic Berlin, Freie Universität Berlin, Humboldt-Universität zu Berlin and Berlin Institute of Health, Germany; **Henrik Walter**, Charité University Clinic Berlin, Freie Universität Berlin, Humboldt-Universität zu Berlin and Berlin Institute of Health, Germany; **Judith K. Daniels**, Department of Clinical Psychology and Experimental Psychopathology, University of Groningen, The Netherlands

**Correspondence:** Judith K. Daniels. Email: [J.K.Daniels@rug.nl](mailto:J.K.Daniels@rug.nl)

First received 2 Nov 2021, final revision 12 Apr 2022, accepted 21 Apr 2022

## Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjo.2022.65>.

## Data availability

The data that support the findings of this study are available on request from the corresponding author, J.K.D. The data are not publicly available due to restrictions their containing information that could compromise the privacy of research participants.

## Author contributions

Y.L.M. conducted the formal analysis, wrote the original draft and reviewed and edited the manuscript. A.M. and A.S. were responsible for data acquisition and reviewed and edited the manuscript. H.W. conceptualised the study and reviewed and edited the manuscript. J.K. D. conceptualised the study, was responsible for funding acquisition and project administration, and reviewed and edited the manuscript.

## Funding

This work was funded by the German Research Foundation (Deutsche Forschungsgemeinschaft) (formerly grant DA 1222/4-1, now grant WA 1539/8-2, to J.K.D.), and the EU Rosalind-Franklin Fellowship Program awarded to J.K.D.

## Declaration of interest

None.

## References

- Lyssenko L, Vonderlin R, Kleindienst N, Alpers GW, Bohus M, Schmahl C. Dissociation in victims of childhood abuse or neglect: a meta-analytic review. *Psychol Med* 2018; **48**(15): 2467–76.
- Hansen M, Ross J, Armour C. Evidence of the dissociative PTSD subtype: a systematic literature review of latent class and profile analytic studies of PTSD. *J Affect Disord* 2017; **213**: 59–69.
- Lotfinia S, Soorgi Z, Mertens Y, Daniels J. Structural and functional brain alterations in psychiatric patients with dissociative experiences: a systematic review of magnetic resonance imaging studies. *J Psychiatr Res* 2020; **128**: 5–15.
- Roydeva Ml, Reinders AATS. Biomarkers of pathological dissociation: a systematic review. *Neurosci Biobehav Rev* 2021; **123**: 120–202.
- Nicholson AA, Friston KJ, Zeidman P, Harricharan S, McKinnon MC, Densmore M, et al. Dynamic causal modeling in PTSD and its dissociative subtype: bottom-up versus top-down processing within fear and emotion regulation circuitry. *Hum Brain Mapp* 2017; **38**(11): 5551–61.
- Rebellino D, Densmore M, Théberge J, McKinnon MC, Lanius RA. The cerebellum after trauma: resting-state functional connectivity of the cerebellum in posttraumatic stress disorder and its dissociative subtype. *Hum Brain Mapp* 2018; **39**(8): 3354–74.
- Olivé I, Densmore M, Harricharan S, Théberge J, McKinnon MC, Lanius R. Superior colliculus resting state networks in post-traumatic stress disorder and its dissociative subtype. *Hum Brain Mapp* 2018; **39**(1): 563–74.

- 8 Harricharan S, Rabellino D, Frewen PA, Densmore M, Théberge J, McKinnon MC, et al. fMRI functional connectivity of the periaqueductal gray in PTSD and its dissociative subtype. *Brain Behav* 2016; **6**(12): e00579.
- 9 Terpou BA, Densmore M, Théberge J, Frewen P, McKinnon MC, Lanius RA. Resting-state pulvinar-posterior parietal decoupling in PTSD and its dissociative subtype. *Hum Brain Mapp* 2018; **39**(11): 4228–40.
- 10 Sierk A, Manthey A, Brakemeier EL, Walter H, Daniels JK. The dissociative subtype of posttraumatic stress disorder is associated with subcortical white matter network alterations. *Brain Imaging Behav* 2021; **15**(2): 643–55.
- 11 Pitman RK, Orr SP, Forgue DF, Jong JBD, Claiborn JM. Psychophysiological assessment of posttraumatic stress disorder imagery in Vietnam combat veterans. *Arch Gen Psychiatry* 1987; **44**(11): 970–5.
- 12 Negreira AM, Abdallah CG. A review of fMRI affective processing paradigms used in the neurobiological study of posttraumatic stress disorder. *Chronic Stress* 2019; **3**: 247054701982903.
- 13 Thome J, Terpou BA, McKinnon MC, Lanius RA. The neural correlates of trauma-related autobiographical memory in posttraumatic stress disorder: a meta-analysis. *Depress Anxiety* 2020; **37**(4): 321–45.
- 14 Lanius RA, Hopper JW, Menon RS. Individual differences in a husband and wife who developed PTSD after a motor vehicle accident: a functional MRI case study. *Am J Psychiatry* 2003; **160**(4): 667–9.
- 15 Lanius RA, Williamson PC, Boksman K, Densmore M, Gupta M, Neufeld RWJ, et al. Brain activation during script-driven imagery induced dissociative responses in PTSD: a functional magnetic resonance imaging investigation. *Biol Psychiatry* 2002; **52**(4): 305–11.
- 16 Lanius RA, Vermetten E, Loewenstein RJ, Brand B, Schmahl C, Bremner JD, et al. Emotion modulation in PTSD: clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry* 2010; **167**(6): 640–7.
- 17 Hopper JW, Frewen PA, Van Der Kolk BA, Lanius RA. Neural correlates of re-experiencing, avoidance, and dissociation in PTSD: symptom dimensions and emotion dysregulation in responses to script-driven trauma imagery. *J Trauma Stress* 2007; **20**(5): 713–25.
- 18 Daniels JK, Coupland NJ, Hegadoren KM, Rowe BH, Densmore M, Neufeld RWJ, et al. Neural and behavioral correlates of peritraumatic dissociation in an acutely traumatized sample. *J Clin Psychiatry* 2012; **73**(4): 420–6.
- 19 Mickleborough MJ, Daniels JK, Coupland NJ, Kao R, Williamson PC, Lanius UF, et al. Effects of trauma-related cues on pain processing in posttraumatic stress disorder: an fMRI investigation. *J Psychiatry Neurosci* 2011; **36**(1): 6–14.
- 20 Lanius RA, Brand B, Vermetten E, Frewen PA, Spiegel D. The dissociative subtype of posttraumatic stress disorder: rationale, clinical and neurobiological evidence, and implications. *Depress Anxiety* 2012; **29**(8): 701–8.
- 21 Frewen PA, Lanius RA. Toward a psychobiology of posttraumatic self-dysregulation: reexperiencing, hyperarousal, dissociation, and emotional numbing. *Ann N Y Acad Sci* 2006; **1071**: 110–24.
- 22 Ludäscher P, Valerius G, Stiglmayr C, Mauchnik J, Lanius RA, Bohus M, et al. Pain sensitivity and neural processing during dissociative states in patients with borderline personality disorder with and without comorbid posttraumatic stress disorder: a pilot study. *J Psychiatry Neurosci* 2010; **35**(3): 177–84.
- 23 Krause-Utz A, Winter D, Schriener F, De CC, Lis S, Spinhoven P, et al. Reduced amygdala reactivity and impaired working memory during dissociation in borderline personality disorder. *Eur Arch Psychiatry Clin Neurosci* 2018; **268**(4): 401–15.
- 24 Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ, et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci* 2013; **14**(5): 365–76.
- 25 Cremers HR, Wager TD, Yarkoni T. The relation between statistical power and inference in fMRI. *PLoS One* 2017; **12**(11): e0184923.
- 26 Wang X, Xie H, Chen T, Cotton AS, Salminen LE, Logue MW, et al. Cortical volume abnormalities in posttraumatic stress disorder: an ENIGMA-psychiatric genomics consortium PTSD workgroup mega-analysis. *Mol Psychiatry* 2021; **26**(8): 4331–43.
- 27 Hopper JW, Frewen PA, Sack M, Lanius RA, Van Der Kolk BA. The responses to script-driven imagery scale (RSDI): assessment of state posttraumatic symptoms for psychobiological and treatment research. *J Psychopathol Behav Assess* 2007; **29**(4): 249–68.
- 28 Wittchen H-U, Wunderlich U, Gruschwitz S, Zaudig M. *SKID I. Strukturiertes Klinisches Interview für DSM-IV. Achse I: Psychische Störungen. Interviewheft und Beurteilungsheft. Eine deutschsprachige, erweiterte Bearb. d. amerikanischen Originalversion des SKID I [Structured Clinical Interview for DSM-IV: Axis I: Psychiatric Disorders]*. Hogrefe, 1997.
- 29 Gast U, Zündorf F, Hofmann A. Strukturiertes Klinisches Interview für DSM-IV-Dissoziative Störungen (SKID-D): Manual. [Structured Clinical Interview for DSM-IV Dissociative Disorders (SKID-D): Manual] Hogrefe, 2000.
- 30 Schnyder U, Moergeli H. German version of clinician-administered PTSD scale. *J Trauma Stress* 2002; **15**(6): 487–92.
- 31 Wingenfeld K, Spitzer C, Mensebach C, Grabe HJ, Hill A, Gast U, et al. die Deutsche version des Childhood Trauma Questionnaire (CTQ): erste Befunde zu den psychometrischen Kennwerten [The German version of the Childhood Trauma Questionnaire (CTQ): preliminary psychometric properties]. *Psychother Psychosom Medizinische Psychol* 2010; **60**(8): 442–50.
- 32 Tagay S, Erim Y, Möllering A, Stoelk B, Mewes R, Senf W. Das Essener Trauma-Inventar (ETI) – ein screeninginstrument zur identifikation traumatischer ereignisse und posttraumatischer störungen [The Essen Trauma Inventory (ETI) - a screening instrument for the identification of traumatic events and post-traumatic disorders]. *Psychother Psychosom Medizinische Psychol* 2007; **1**: 75–89.
- 33 Teegen F. *Deutsche Übersetzung der Posttraumatic Stress Disorder Checklist (PCL-C) des National Center for PTSD [German translation of the Posttraumatic Stress Disorder Checklist (PCL-C) from the National Center for PTSD]*. Universität Hamburg, Psychologisches Institut III, 1997.
- 34 Spitzer C, Mestel R, Klingelhöfer J, Gänsicke M, Freyberger HJ. Screening und veränderungsmessung dissoziativer psychopathologie: psychometrische charakteristika der kurzform des Fragebogens zu Dissoziativen Symptomen (FDS-20) [Screening and measurement of change of dissociative psychopathology: psychometric properties of the short version of the Fragebogen zu Dissoziativen Symptomen (FDS-20)]. *Psychother Psychosom Medizinische Psychol* 2004; **54**(3–4): 165–72.
- 35 Michal M, Sann U, Niebecker M, Lazanowsky C, Kernhof K, Aurich S, et al. Die erfassung des depersonalisations-derealisations-syndroms mit der Deutschen version der Cambridge Depersonalisation Scale (CDS). [The measurement of the depersonalisation-derealisation-syndrome with the German version of the Cambridge Depersonalisation Scale (CDS)]. *Bibliogr Psychother Psych Med* 2004; **54**: 367–74.
- 36 Mueller-Pfeiffer C, Schumacher S, Martin-Soelch C, Pazhenkottil AP, Wirtz G, Fuhrhans C, et al. The validity and reliability of the German version of the Somatoform Dissociation Questionnaire (SDQ-20). *J Trauma Dissociation* 2010; **11**(3): 337–57.
- 37 Marmar CR, Weiss DS, Schlenger WE, Fairbank JA, Jordan BK, Kulka RA, et al. Peritraumatic dissociation and posttraumatic stress in male Vietnam theater veterans. *Am J Psychiatry* 1994; **151**(suppl 6): 902–7.
- 38 Maercker A. *Unveröffentlichtes Manuskript [Unpublished Manuscript]*. Tu Dresden, 1994 ([https://www.psychologie.uzh.ch/dam/jcr:9a580e93-85b4-4b12-aa97-fb2778d9d855/PDEQ\\_Fragebogen.pdf](https://www.psychologie.uzh.ch/dam/jcr:9a580e93-85b4-4b12-aa97-fb2778d9d855/PDEQ_Fragebogen.pdf)).
- 39 Laux L, Spielberger CD. *Das State-Trait-Angstinventar: STAI [The State-Trait Anxiety Inventory: STAI]*. Beltz Test, 2001.
- 40 Hautzinger M, Keller F, Kühner C. *Beck Depressions-Inventar (BDI-II)* [Beck Depression Inventory (BDI-II)]. Harcourt Test Services, 2006.
- 41 Bremner JD, Krystal JH, Putnam FW, Southwick SM, Marmar C, Charney DS, et al. Measurement of dissociative states with the Clinician-Administered Dissociative States Scale (CADSS). *J Trauma Stress* 1998; **11**(1): 125–36.
- 42 Mertens YL, Daniels JK. The Clinician-Administered Dissociative States Scale (CADSS): validation of the German Version. *J Trauma Dissociation* [Epub ahead of print] 21 Oct 2021. Available from: <https://doi.org/10.1080/15299732.2021.1989111>.
- 43 Amunts K, Kedo O, Kindler M, Pieperhoff P, Mohlberg H, Shah NJ, et al. Cytoarchitectonic mapping of the human amygdala, hippocampal region and entorhinal cortex: intersubject variability and probability maps. *Anat Embryol (Berl)* 2005; **210**(5–6): 343–52.
- 44 Eickhoff SB, Stephan KE, Mohlberg H, Grefkes C, Fink GR, Amunts K, et al. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage* 2005; **25**(4): 1325–35.
- 45 Spisák T, Spisák Z, Zunhammer M, Bingel U, Smith S, Nichols T, et al. Probabilistic TFCE: a generalized combination of cluster size and voxel intensity to increase statistical power. *Neuroimage* 2019; **185**: 12–26.
- 46 Lamke JP, Daniels JK, Dörfel D, Gaebler M, Rahman RA, Hummel F, et al. The impact of stimulus valence and emotion regulation on sustained brain activation: task-rest switching in emotion. *PLoS One* 2014; **9**(3): e93098.
- 47 Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 2012; **59**(3): 2142.
- 48 Svoboda E, McKinnon MC, Levine B. The functional neuroanatomy of autobiographical memory: a meta-analysis. *Neuropsychologia* 2006; **44**(12): 2189–208.
- 49 Lanius RA, Rabellino D, Boyd JE, Harricharan S, Frewen PA, McKinnon MC. The innate alarm system in PTSD: conscious and subconscious processing of threat. *Curr Opin Psychol* 2017; **14**: 109–15.
- 50 Schauer M, Elbert T. Dissociation following traumatic stress. *Z Psychol* 2010; **218**(2): 109–27.
- 51 Felmingham K, Kemp AH, Williams L, Falconer E, Olivieri G, Peduto A, et al. Dissociative responses to conscious and non-conscious fear impact underlying brain function in post-traumatic stress disorder. *Psychol Med* 2008; **38**(12): 1771–80.
- 52 Rabellino D, Densmore M, Frewen PA, Théberge J, Lanius RA. The innate alarm circuit in post-traumatic stress disorder: conscious and subconscious processing of fear- and trauma-related cues. *Psychiatry Res Neuroimaging* 2016; **248**: 142–50.

- 53 Vonderlin R, Kleindienst N, Alpers GW, Bohus M, Lyssenko L, Schmahl C. Dissociation in victims of childhood abuse or neglect: a meta-analytic review. *Psychol Med* 2018; **48**(15): 2467–76.
- 54 Rafiq S, Campodonico C, Varese F. The relationship between childhood adversities and dissociation in severe mental illness: a meta-analytic review. *Acta Psychiatr Scand* 2018; **138**: 509–25.
- 55 Cross D, Fani N, Powers A, Bradley B. Neurobiological development in the context of childhood trauma. *Clin Psychol (New York)* 2017; **24**(2): 111–24.
- 56 Hayes JP, VanElzakker MB, Shin LM. Emotion and cognition interactions in PTSD: a review of neurocognitive and neuroimaging studies. *Front Integr Neurosci* 2012; **6**: 89.
- 57 Patel R, Spreng RN, Shin LM, Girard TA. Neurocircuitry models of posttraumatic stress disorder and beyond: a meta-analysis of functional neuroimaging studies. *Neurosci Biobehav Rev* 2012; **36**: 2130–42.
- 58 Winlove CIP, Milton F, Ranson J, Fulford J, MacKisack M, Macpherson F, et al. The neural correlates of visual imagery: a co-ordinate-based meta-analysis. *Cortex* 2018; **105**: 4–25.
- 59 Monaco S, Malfatti G, Culham JC, Cattaneo L, Turella L. Decoding motor imagery and action planning in the early visual cortex: overlapping but distinct neural mechanisms. *NeuroImage* 2020; **218**: 116981.
- 60 Sartory G, Cwik J, Knappertz H, Schü Rholt B, Lebens M. In search of the trauma memory: a meta-analysis of functional neuroimaging studies of symptom provocation in posttraumatic stress disorder (PTSD). *PLoS One* 2013; **8**(3): 58150.
- 61 Puetz VB, Puetz VB, Viding E, Gerin MI, Pingault JB, Sethi A, et al. Investigating patterns of neural response associated with childhood abuse v. childhood neglect. *Psychol Med* 2020; **50**(8): 1398–407.

